Title: Characterization of metabolically healthy and unhealthy obesity

through circulating proteins and metabolites

Giulia Pontali [1,2,7,a]

Christian X. Weichenberger [1,7] https://orcid.org/0000-0002-2176-0274

Johannes Rainer [1] https://orcid.org/0000-0002-6977-7147

Essi Hantikainen [1] https://orcid.org/0000-0002-4155-2894

Marilyn De Graeve [1] https://orcid.org/0000-0001-6916-401X

Fulvio Mattivi [3] https://orcid.org/0000-0003-4935-5876

Michael Kob [4,5]

Markus Ralser [6] https://orcid.org/0000-0001-9535-7413

Peter P. Pramstaller [1] https://orcid.org/0000-0002-9831-8302

Francisco S. Domingues [1,b] https://orcid.org/0000-0003-4567-4562

[1] Institute for Biomedicine, Eurac Research, Bolzano, Italy

[2] Department of Cellular Computational and Integrative Biology (CIBIO),

University of Trento, Povo, Italy

[3] Research and Innovation Center, Fondazione Edmund Mach, San Michele

all'Adige, Italy

[4] College of Health Care-Professions "Claudiana", 39100 Bolzano, Italy

[5] Division of Clinical Nutrition, Hospital of Bolzano (SABES-ASDAA), Bolzano, Italy

[6]Department of Biochemistry, Charité – Universitätsmedizin Berlin, Berlin,

Germany

[7]These authors contributed equally: Giulia Pontali, Christian X. Weichenberger

[a]Currently at: Fondazione Human Technopole, Milano, Italy

[b] email: francisco.domingues@eurac.edu

ABSTRACT

Background: Individuals affected by obesity present different health

trajectories and do not suffer from cardiometabolic complications all in the same

way. There is a need to better understand obesity subtypes and to develop

approaches for stratification. In this study we investigated both metabolomic and

proteomic signatures in serum and blood plasma samples discriminating

metabolically healthy from unhealthy obesity.

Methods: We investigated cross-sectional metabolomic and proteomic data from

participants of the Cooperative Health Research in South Tyrol (CHRIS) study.

Participants were grouped into metabolically healthy obesity (MHO) and

metabolically unhealthy obesity (MUO) based on available health data in the

study. A total of 461 individuals were included in the analysis, with n=130 MHO

and n=331 MUO. Random forest (RF) classifiers were used to discriminate

metabolically healthy from unhealthy obesity and to identify molecular features

characteristic of MHO/MUO. Linear regression models were used to assess

associations between each relevant metabolite/protein and MHO/MUO

phenotypes independently of age, sex and body composition.

Results: The MHO/MUO RF classifier achieved a performance of AUC = 0.709,

95% CI = (0.698,0.721). Three plasma proteins and 12 circulating metabolites

were identified as relevant predictors of MHO/MUO phenotypes. Linear

regression models confirmed the Apolipoprotein C-III (APOC3) association to be

independent of age, visceral fat composition, medication or serum triglyceride

levels.

Conclusion: APOC3 was identified as a novel predictor for obesity stratification,

highlighting the importance of circulating triglyceride levels in relation to

metabolic health.

INTRODUCTION

Obesity is a condition characterized by abnormal accumulation of body fat, that has been linked to type 2 diabetes, cardiovascular disease, and some types of cancer [1, 2]. It has been estimated that overweight and obesity accounts for a considerable proportion of chronic disease burden and premature death [3–5]. The global prevalence of obesity has been increasing. Recent estimates indicate that between 1990 and 2022 obesity prevalence in adults increased from 8.8% to 18.5% in women, and from 4.8% to 14.0% in men, overtaking the global prevalence of underweight in both genders in this period [6].

Obesity results from an interplay between genetics, lifestyle (specifically diet) and environmental factors [7], and is widely regarded as a multifactorial disease [8]. There is a substantial genetic component underlying individual variation in body weight, and different studies have estimated the heritability of obesity to be between 40%-70% [9]. Genetic studies also indicate that monogenic and polygenic obesity share a similar biology, where central nervous system pathways related to food intake play a key role [9]. Furthermore, the gut microbiome impacts obesity by modulating mood disorders, eating behaviors and detoxification processes [10].

Not all individuals affected by obesity suffer from metabolic complications, which cannot be explained only by the extent of adiposity. A considerable subgroup of individuals with obesity seems to exhibit a relative protection or a significantly lower risk of cardiometabolic disease. This phenotype has been described as metabolically healthy obesity (MHO), to differentiate from the more prevalent metabolically unhealthy obesity (MUO) [11]. Multiple definitions for MHO/MUO have been proposed, but in general they include the body mass index (BMI)

criteria for obesity, and the absence/presence of hyperglycemia, dyslipidemia,

hypertension and cardiovascular disease [11, 12]. MUO has been linked to

higher visceral body fat, a proinflammatory state and insulin resistance, while

MHO is characterized by a higher subcutaneous body fat, greater insulin

sensitivity and better cardiorespiratory fitness [11, 13]. MUO is potentially the

outcome of a long-term positive energy balance coupled with inability of

subcutaneous adipose tissue to expand [14–16]. Although MHO is associated

with a lower risk of cardiometabolic disease in comparison to MUO, it has long-

term negative impact on cardiometabolic health and should therefore not be

considered a benign condition [17, 18]. To some extent MHO is a transient

phenotype to MUO, although the MHO phenotype can be stable over years in a

subset of individuals [19–22]. However, the transition to MUO is not only

unidirectional, as conversion from MUO to MHO is also being observed [23].

High-throughput omics approaches capture detailed molecular profiles in

biological samples and allow us to investigate any relevant changes between

health conditions, providing insights into the relevant processes underlying

health and disease [24]. With such detailed molecular profiling is then possible

to better characterize disease subtypes, enabling enhanced stratification of

individuals for improved diagnosis, prognosis and more effective therapeutic

strategies [25]. These approaches have been successfully applied in the

investigation of metabolic health in obesity [26]. Extensive metabolic profiling of

the MHO phenotype [27–34] has highlighted differences between MHO and MUO

in the levels of several metabolites and identified altered metabolic pathways.

The results indicate differences in the levels of branched-chain amino acids and

aromatic amino acids, where the blood amino acid profile of MHO falls between

those of metabolically healthy individuals with normal weight and MUO [26].

Circulating proteomics profiles for MHO/MUO have also been investigated, with

results pointing towards increased inflammation and insulin resistance in MUO

[35–38]. These proteomics studies are based on relatively modest sample sizes,

limiting the power to determine relevant features for MHO and for stratification

of obesity.

In this study, we perform a large-scale analysis of serum metabolomic and

plasma proteomic signatures in MHO and MUO, utilizing the largest sample size

for combined omics analysis reported to date. This study aims to identify key

molecular features that differentiate these conditions, contributing to more

precise stratification of obesity.

METHODS

Study Cohort

The Cooperative Health Research in South Tyrol (CHRIS) study is a population-based cohort of 13,393 adults aged 18 and over, recruited from 13 municipalities in the alpine Val Venosta/Vinschgau district in the Bolzano-South Tyrol province of northern Italy [39]. Baseline visits were conducted from 2011 to 2018, collecting socio-demographic, health, lifestyle, medication and exposure data with questionnaires, interviews, and instrumental examinations, including electrocardiography, blood pressure and anthropometric measurements. Body composition, including body fat and visceral fat percentages were assessed by bioelectrical impedance (OMRON BF508). Medication information was collected by scanning the barcode of the boxes of the medication taken over the last seven days by each study participant and used to assign the respective Anatomical Therapeutic Chemical (ATC) codes. Blood and urine samples were also collected for biobanking, DNA extraction, and molecular characterization (genome and exome sequencing, metabolomics, and proteomics). The CHRIS resource was assigned the "Bioresource Research Impact Factor " (BRIF) code BRIF6107. Standard blood parameters were measured in blood samples at the Hospital of Merano using standardized clinical assays as described previously [40]. More specifically, the serum triglycerides were measured either with Cobas Triglyceride GPO-PAP assay on a Roche Modular PPE instrument or with a TRIGLYCERIDE assay on an Abbot Diagnostic ARCHITECT system.

Assessment of Obesity Phenotypes

Obesity was defined as having a BMI greater than or equal to 30 kg/m^2 [41].

Metabolically healthy obesity (MHO) and metabolically unhealthy obesity (MUO)

were defined according to previously established criteria [42] as summarized in

Table 1. Briefly, MUO is defined as individuals with obesity that are affected by

either hypertension, diabetes, dyslipidemia or cardiovascular disease, while MHO

is defined as all other individuals with obesity that are not MUO and where

metabolic health could be assessed (no missing data). The two phenotypes were

encoded in a single binary variable (MHO/MUO phenotypes).

Table 1. Criteria used in definition of MHO and MUO.

Phenotype	Criteria

MUO	• Pregnancy excluded **AND**
	
	• BMI ≥ 30 kg/m^2 (definition of obesity) **AND**
	
	• At least one of the following:
	
	    ○ Hypertension: Systolic blood pressure ≥ 140mmHg, Diastolic blood pressure ≥ 90mmHg, use of blood pressure lowering medication[a]
	
	    ○ Type 2 diabetes: reported type 2 diabetes, fasting blood glucose ≥126 mg/dl, glycated hemoglobin (HbA1c) ≥48 mmol/mol, use of diabetes medication[b]
	
	    ○ Dyslipidemia: blood HDL- cholesterol<40 mg/dl (in men), blood HDL- cholesterol < 50 mg/dl (in women), serum triglycerides level ≥150 mg/dl, using lipid medication[c]
	
	    ○ Cardiovascular disease: Any ischemic heart disease, stroke, transient ischemic attack, claudication, peripheral arterial disease, pulmonary embolism, deep venous thrombosis, atrial fibrillation, absence of P wave in 10s ECG

MHO	• Pregnancy excluded **AND** • BMI ≥ 30 kg/m^2 (definition of obesity) **AND** • Not MUO **AND** • Data used in MUO definition are not missing

155 [a]Medication with any of the following second level ATC codes: C02, C03, C04,

156 C07, C08, C09

157 [b]Medication with any of the following third level ATC codes: A10A, A10B

158 [c]Medication with second level ATC codes C10

Metabolite and protein quantification

Metabolomics and proteomics data were collected as previously described [43].

To summarize, the AbsoluteIDQ® p180 kit from Biocrates (Biocrates Life

Sciences AG, Innsbruck, Austria) was used for metabolite quantification in serum

samples collected in fasting participants. Samples were analyzed with a UHPLC-

MS/MS system consisting of an Agilent 1290 Infinity liquid chromatography

system (Agilent Technologies, CA, USA) and a Sciex QTRAP 6500 mass

spectrometer (AB Sciex LLC, MA, USA), and data were processed as described

previously [44]. In total, 174 metabolites were quantified, they are provided in

Supplementary Table 1. The plasma proteome was determined using Scanning

SWATH mass spectrometry [45] as described previously [46]. In total, 148

highly abundant proteins were quantified and included in this analysis, they are

listed in Supplementary Table 2.

Medication Adjustment

Metabolite and protein abundances were $\log_2$-transformed and adjusted for

frequent medication as previously described [43]. More specifically, linear

models were fitted separately for each metabolite or protein with their

abundance as response, and medication (encoded as individual binary variables)

as explanatory variables. Only medication taken at least twice per week was

considered for the adjustment. Medication used for MHO/MUO definitions were

excluded (blood pressure, diabetes or blood lipid medication). The residuals from

these models were used to construct medication-independent metabolite and

protein abundances for all subsequent analyses. Supplementary Table 3 lists the

types of medication for which abundances were adjusted, and the linear models

are described in Supplementary file section 4.1.

Random forest analysis

A random forest (RF) classification model was used to predict MHO vs. MUO,

encoded as a binary variable (MHO/MUO phenotypes), and following a similar

strategy to a previous analysis [43] using as predictors age, sex, body fat,

visceral fat, 174 metabolites and 148 proteins. Overall, 100 RF models were

generated, each containing 500 trees per model, with 80% training set sizes.

Predictions were validated 100 times by repeated random subsampling with 20%

test set sizes, while keeping the same ratio of MHO to MUO cases in the original

and subsampled sets. The statistical programming language R v4.2.0 was

utilized for computations and figure creation. More specifically, the randomForest

R package v4.7-1.1 [47] was used to perform the RF analysis. Model

performance was measured based on the receiver operating characteristic (ROC)

area under the curve (AUC), and the Matthew's correlation coefficient (MCC).

The MCC was calculated at the optimal threshold on the ROC curve,

corresponding to the true positive rate/false positive rate pair most distant from

the diagonal. The AUC and MCC values reported correspond to the means of the

100 test runs, provided with the respective 95% confidence interval (CI). The

Gini Index was used to measure feature importance. The significance of the Gini

Indices for the random forest models was estimated by permuting the response

variable using the rfPermute R package v2.5.1 [48]. This generates a

background distribution by calculating 100 random forests, one for each

permuted response variable, each with 500 trees. A background distribution for

each validation run was created, yielding 100 p-values for each predictor

variable. If in $\geq$ 50 out of the 100 models the predictor was significant at the

level $\alpha = 0.05$ it was identified as a relevant predictor.

Confirmation of relevant predictors

Linear regression models were applied to further investigate associations

between MHO/MUO and the relevant metabolites or proteins obtained from the

RF analysis. Separate multiple linear regression models were fitted with

abundances of each metabolite and protein as the response variable, and with

MHO/MUO status, age, sex and visceral fat as the explanatory variables

(Supplementary file section 4.2). The resulting p-values were adjusted for

multiple hypothesis testing using the Bonferroni method. After correction, results

were considered statistically significant if the adjusted p-value remained below

the conventional significance level of $\alpha = 0.05$. Additional linear models were

derived to investigate associations between relevant proteins and metabolites

and medication used in phenotype definition (Supplementary file section 4.3).

Logistic regression models were used to investigate if the association between

MHO/MUO phenotypes and plasma APOC3 is dependent on serum triglyceride

levels (Supplementary file section 4.4).

Ethical approval

The Ethics Committee of the Healthcare System of the Autonomous Province of

Bolzano-South Tyrol approved the CHRIS baseline protocol on 19 April 2011 (21-

2011). The CHRIS study conforms to the Declaration of Helsinki, and with

national and institutional legal and ethical requirements.

RESULTS

In this study, serum metabolomic and plasma proteomic signatures discriminating MHO from MUO are investigated on 461 subjects of the CHRIS study. First, the main characteristics of the study samples are provided. Second, the results from the random forest (RF) analysis are provided, including the prediction performance and the identification of relevant predictors for the classification of MHO/MUO. Third, the results of the multiple linear regression analyses are presented to support the identification of molecular predictors discriminating MHO from MUO.

Characteristics of the study sample

Data were available in the CHRIS study to assign obesity status for a total of 13,216 participants, with 2,238 of them being affected by obesity (16.93%). The data required to assign MHO/MUO phenotypes were available for a subset of this obesity group (n=2,170), and among these a total of 529 participants were classified as MHO (24.38%) and 1,641 as MUO (75.62%). All the subsequent analyses are based on subsets with available metabolomics and proteomics data, for a total size of n=130 for MHO and n=331 MUO groups. The characteristics of this analytic sample are provided in Table 2. A lower mean age and a larger proportion of females are observed in the MHO group in comparison to MUO, and hypertension is the most frequent condition in the MUO group.

Table 2. Characteristics of the analytic sample grouped by metabolically healthy (MHO) and unhealthy obesity (MUO) *

Characteristic	MHO (n=130)	MUO (n=331)
Age (years)	43.98 ± 13.94	57.10 ± 14.73
Male	47 (36.15%)	165 (49.85%)
Female	83 (63.85%)	166 (50.15%)
BMI (kg/m^2)	32.98 ± 3.14	33.43 ± 3.10
Visceral fat (%)	12.09 ± 4.07	15.21 ± 4.08
Body fat (%)	41.19 ± 8.62	39.65 ± 8.34
Hypertension	0 (0.00%)	245 (74.02%)
Type 2 diabetes	0 (0.00%)	58 (17.52%)
Dyslipidemia	0 (0.00%)	202 (61.03%)
Cardiovascular	0 (0.00%)	57 (17.22%)

*: Categorical variables are shown as number (percentage) and continuous variables are presented as mean ± standard deviation.

Random Forest Analysis

Random Forest models were trained and tested to discriminate MHO from MUO, based on the predictors age, sex, body fat, visceral fat, and medication-adjusted abundances of 174 metabolites and 148 proteins. Adjustment of concentrations for common medications ensured any results being independent of potential confounding by unrelated treatments, such as hormonal contraceptives [43, 46]. Test results indicate a moderate classifier performance of AUC = 0.709, 95% CI = (0.698,0.721) and MCC = 0.394, 95% CI = (0.377,0.411). The receiver operating characteristic curve and precision-recall curve are provided in Fig. 1.

Fig. 1 RF classification performance. ROC curve on left and precision recall
curve on the right. Testing results for 100 models, each shown as gray line with
average curve in black.

Individual predictors were selected based on their importance in the RF models

as determined by mean decrease in Gini Index. According to selection criteria

(described in the Methods section) 17 predictors were identified, including age,

visceral fat, three proteins and 12 metabolites (Fig. 2a). The most relevant

predictors are age, visceral fat and Apolipoprotein C-III (APOC3), followed by

Insulin-like growth factor-binding protein complex acid labile subunit (IGFALS)

and a phosphatidylcholine (PC aa C34:3). The distributions of the relevant

predictors in both MHO and MUO phenotypes are provided in Fig. 2b. In the MHO

phenotype we observe lower visceral fat and lower levels of APOC3, and higher

levels of IGFALS and PC aa C34:3.

Confirmation of relevant predictors

The relation between MHO/MUO phenotypes and the molecular predictors
identified by the RF models were further investigated by separate multiple
regression analyses to directly evaluate the differences in metabolite or protein
concentrations between the MHO vs MUO groups independently of age, sex or
body fat composition. Separate linear regression models were fitted having as
response variable the medication adjusted abundances of each of the three
proteins and 12 metabolites identified, and with the MHO/MUO phenotypes, sex,
age and visceral fat as the explanatory variables. Detailed results are presented
in Supplementary Table 4. The results confirm the association between
MHO/MUO phenotypes and APOC3, plasma serine protease inhibitor (SERPINA5)
and a phosphatidylcholine (PC aa C34:3), each with an adjusted p-value < 0.05.
The direction of these three statistically significant associations also agrees with
the observed median differences of the respective abundances in the two
phenotypes (Fig. 2b). Nevertheless, in the case of SERPINA5 the association with
sex is stronger than with the MHO/MUO phenotypes. The levels of insulin-like
growth factor-binding protein complex acid labile subunit (IGFALS) are instead
strongly associated with age and visceral fat but not with the MHO/MUO
phenotypes. Further, no significant associations were observed between the
remaining metabolites and MHO/MUO.

Given the strong association observed between APOC3 and MHO/MUO, we
conducted a first sensitivity analysis to further investigate whether this relation
could result from the intake of medication included in definition of MHO/MUO. To
do so a linear regression model was fitted with APOC3 levels as response
variable and MHO/MUO phenotypes and lipid modifier medication as explanatory
variables. A significant association is again observed with MHO/MUO phenotypes

but not with the lipid medication, indicating a robust relation between APOC3

and MHO/MUO phenotypes. A similar analysis for the PC aa C34:3 levels indicate

a stronger association to statin intake than to MHO/MUO phenotypes, suggesting

that the PC aa C34:3 relation to MHO/MUO phenotypes just reflects medication

intake. The results are provided in Supplementary Table 5.

Given that APOC3 is involved in triglyceride transport [49], the question arises

whether the relation between APOC3 levels and the MHO/MUO phenotypes just

reflects serum triglyceride concentration, which is also used in the phenotype

definition. To address this question, we conducted a second sensitivity analysis

by fitting a logistic regression model with MHO/MUO phenotypes as response

variable, and sex, age, APOC3 levels, visceral fat, and triglyceride levels as

explanatory variables (Supplementary Table 6). A strong association with the

phenotypes is observed for both age and triglyceride levels. A weaker but

significant association is observed for APOC3 levels as well, with a change in the

direction of the association after adjusting for triglyceride levels. These results

indicate that APOC3 is associated with MHO/MUO phenotypes independently of

triglyceride levels measured by an established clinical blood test.

Fig. 2 Relevant predictors for classification of MHO vs MUO phenotypes.
A Mean decrease in Gini Index for significant predictors. Predictor significance
was estimated by permutation testing, resulting in 100 p-values for each
predictor variable. Box plots are color-coded by the number of times a p-value
was significant for such a variable (p < 0.05): red if all 100 runs returned a
significant p-value; orange when between 80 and 99 runs returned a significant
p-value; blue when between 50 and 79 runs returned a significant p-value.
Panels are restricted to predictors that are significant in at least 50% of all runs,
and ordered by importance from left to right. **B** Violin plots presenting scaled
abundance distributions for MHO and MUO groups for the significant predictors
identified through RF, ordered by importance from left to right. Density plots of
MHO group are scaled to have the same width, a median of zero (horizontal bar)
and standard deviation of one. For comparability, the MUO group data have been
scaled to the standard deviation of the MHO group. The metabolite and protein
names are provided in Supplementary Tables 1 and 2. Labels for protein are in
bold.

DISCUSSION

A better understanding of obesity subtypes is necessary for the development of

more effective therapeutic strategies. In this study we investigated blood

metabolomic and proteomic signatures to discriminate metabolically healthy

from unhealthy obesity phenotypes.

Our results indicate that Apolipoprotein C-III (APOC3) is a robust molecular

predictor discriminating MHO from MUO, independently of age and visceral fat,

which are other strong predictors of these phenotypes. Higher abundances of

this apolipoprotein are observed in the MUO phenotype. APOC3 is found in

triglyceride-rich lipoproteins like chylomicrons and very-low-density lipoproteins

(VLDL) and inhibits hepatic uptake of VLDL [49, 50]. This inhibition contributes

to elevated levels of circulating triglycerides, influencing lipid metabolism.

Although the result is not unexpected, given that serum triglyceride levels are

used in the definition of MUO, further analysis with logistic regression indicates

that APOC3 levels are associated with MHO/MUO phenotypes independent of

serum triglyceride levels. The findings reported here are also in agreement with

previous work describing higher levels of VLDL in MUO individuals relative to

MHO [51], and are consistent with previous reports indicating triglyceride levels

to be a significant predictor for the transition from MHO to MUO [52]. Notably,

increased APOC3 levels have been observed in women with obesity [53], and

decreasing APOC3 levels have been linked to weight loss [54]. APOC3 deficiency

has triglyceride lowering effects and is associated with reduced cardiovascular

disease (CVD) risk [55, 56], and APOC3 polymorphisms have been associated

with metabolic syndrome [57] and have an impact on the coronary artery

disease risk [58].

The protein is a therapeutic target of volanesorsen, a drug approved for reduction of triglyceride levels, and is a candidate target for reducing CVD risk [49, 59].

To date, this work represents the largest study analyzing circulating proteome and metabolome signatures for the MHO/MUO phenotypes. Furthermore, the analysis includes proper adjustment of proteomics and metabolomics levels regarding medication intake.

There is no established consensus on the best criteria for MHO/MUO. Even though an established criterion was used in this work, the lack of a standard can be seen as a limitation. An alternative definition has been proposed for stratifying individuals according to mortality risk based on a large prospective study [60]. Although this new definition cannot be applied in the available CHRIS baseline study dataset, given the lack of a waist and hip circumference measurements, it would be of interest to replicate the analysis with this definition once the dataset from the ongoing CHRIS follow-up study is made available, where waist-to-hip ratios are being measured. Another limitation is the relatively limited set of blood metabolites and proteins profiled.

We foresee several opportunities for a follow-up to this work. We suggest further investigating APOC3 and VLDL levels, as well as the impact of triglyceride metabolism and transport in relation to health in obesity. The observed change in the association between APOC3 and MHO/MUO after adjustment for triglyceride levels needs further investigation. The replication of these findings in other studies and the exploration of causal relationships would be very valuable. Furthermore, it would be of great interest to investigate the relationship between

changes in APOC3 and lipoprotein levels, as well as the stability of MHO

phenotype and transition to MUO.

CONCLUSION

A model based on multi-omics was developed for discriminating MHO from MUO

achieving moderate performance. An analysis of the relevant features indicates

APOC3 as molecular predictor for the stratification of obesity. Higher levels of

plasma APOC3 are observed in MUO, independently of age, sex or medication.

The results highlight the role of lipoproteins and triglyceride metabolism and/or

transport in shaping metabolic health within obesity.

DATA AVAILABILITY

Data and samples can be requested for clearly defined research via the CHRIS

Portal (https://chrisportal.eurac.edu/).

REFERENCES

[1] Must A, Spadano J, Coakley EH, et al. The disease burden associated with overweight and obesity. *JAMA* 1999; 282: 1523–1529.

[2] Lauby-Secretan B, Scoccianti C, Loomis D, et al. Body Fatness and Cancer--Viewpoint of the IARC Working Group. *N Engl J Med* 2016; 375: 794–798.

[3] GBD 2015 Obesity Collaborators, Afshin A, Forouzanfar MH, et al. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med* 2017; 377: 13–27.

[4] Dai H, Alsalhe TA, Chalghaf N, et al. The global burden of disease attributable to high body mass index in 195 countries and territories, 1990-2017: An analysis of the Global Burden of Disease Study. *PLoS Med* 2020; 17: e1003198.

[5] GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Lond Engl* 2020; 396: 1223–1249.

[6] NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222 million children, adolescents, and adults. *Lancet Lond Engl* 2024; 403: 1027–1050.

[7] Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019; 15: 288–298.

[8] Diels S, Vanden Berghe W, Van Hul W. Insights into the multifactorial causation of obesity by integrated genetic and epigenetic analysis. *Obes Rev Off J Int Assoc Study Obes* 2020; 21: e13019.

[9] Loos RJF, Yeo GSH. The genetics of obesity: from discovery to biology. *Nat Rev Genet* 2022; 23: 120–133.

[10] Choi BS-Y, Daoust L, Pilon G, et al. Potential therapeutic applications of the gut microbiome in obesity: from brain function to body detoxification. *Int J Obes 2005* 2020; 44: 1818–1831.

[11] Blüher M. Metabolically Healthy Obesity. *Endocr Rev* 2020; 41: bnaa004.

[12] Smith GI, Mittendorfer B, Klein S. Metabolically healthy obesity: facts and fantasies. *J Clin Invest* 2019; 129: 3978–3989.

[13] Stefan N, Kantartzis K, Machann J, et al. Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med* 2008; 168: 1609–1616.

[14] Goossens GH. The Metabolic Phenotype in Obesity: Fat Mass, Body Fat Distribution, and Adipose Tissue Function. *Obes Facts* 2017; 10: 207–215.

[15] Iacobini C, Pugliese G, Blasetti Fantauzzi C, et al. Metabolically healthy versus metabolically unhealthy obesity. *Metabolism* 2019; 92: 51–60.

[16] Ghaben AL, Scherer PE. Adipogenesis and metabolic health. *Nat Rev Mol Cell Biol* 2019;
20: 242–258.

[17] Magkos F. Metabolically healthy obesity: what's in a name? *Am J Clin Nutr* 2019; 110: 533–
539.

[18] Tanriover C, Copur S, Gaipov A, et al. Metabolically healthy obesity: Misleading phrase or
healthy phenotype? *Eur J Intern Med* 2023; 111: 5–20.

[19] Kouvari M, Panagiotakos DB, Yannakoulia M, et al. Transition from metabolically benign to
metabolically unhealthy obesity and 10-year cardiovascular disease incidence: The
ATTICA cohort study. *Metabolism* 2019; 93: 18–24.

[20] Camhi SM, Must A, Gona PN, et al. Duration and stability of metabolically healthy obesity
over 30 years. *Int J Obes 2005* 2019; 43: 1803–1810.

[21] Elías-López D, Vargas-Vázquez A, Mehta R, et al. Natural course of metabolically healthy
phenotype and risk of developing Cardiometabolic diseases: a three years follow-up study.
BMC Endocr Disord 2021; 21: 85.

[22] Zhou Z, Macpherson J, Gray SR, et al. Are people with metabolically healthy obesity really
healthy? A prospective cohort study of 381,363 UK Biobank participants. *Diabetologia*
2021; 64: 1963–1972.

[23] Appleton SL, Seaborn CJ, Visvanathan R, et al. Diabetes and cardiovascular disease
outcomes in the metabolically healthy obese phenotype: a cohort study. *Diabetes Care*
2013; 36: 2388–2394.

[24] Babu M, Snyder M. Multi-Omics Profiling for Health. *Mol Cell Proteomics MCP* 2023; 22:
100561.

[25] Johansson Å, Andreassen OA, Brunak S, et al. Precision medicine in complex diseases-
Molecular subgrouping for improved prediction and treatment stratification. *J Intern Med*
2023; 294: 378–396.

[26] Paczkowska-Abdulsalam M, Kretowski A. Obesity, metabolic health and omics: Current
status and future directions. *World J Diabetes* 2021; 12: 420–436.

[27] Batch BC, Shah SH, Newgard CB, et al. Branched chain amino acids are novel biomarkers
for discrimination of metabolic wellness. *Metabolism* 2013; 62: 961–969.

[28] Gao X, Zhang W, Wang Y, et al. Serum metabolic biomarkers distinguish metabolically
healthy peripherally obese from unhealthy centrally obese individuals. *Nutr Metab* 2016;
13: 33.

[29] Bagheri M, Farzadfar F, Qi L, et al. Obesity-Related Metabolomic Profiles and
Discrimination of Metabolically Unhealthy Obesity. *J Proteome Res* 2018; 17: 1452–1462.

[30] Cirulli ET, Guo L, Leon Swisher C, et al. Profound Perturbation of the Metabolome in
Obesity Is Associated with Health Risk. *Cell Metab* 2019; 29: 488-500.e2.

[31] Ojwang AA, Smuts CM, Zec M, et al. Comparison of dietary and plasma phospholipid fatty
acids between normal weight and overweight black South Africans according to metabolic
health: The PURE study. *Prostaglandins Leukot Essent Fatty Acids* 2020; 158: 102039.

[32] Lind L, Salihovic S, Sundström J, et al. Metabolic Profiling of Obesity With and Without the
Metabolic Syndrome: A Multisample Evaluation. *J Clin Endocrinol Metab* 2022; 107: 1337–
1345.

[33] Orozco-Ruiz X, Anesi A, Mattivi F, et al. Branched-Chain and Aromatic Amino Acids Related
to Visceral Adipose Tissue Impact Metabolic Health Risk Markers. *J Clin Endocrinol Metab*
2022; 107: e2896–e2905.

[34] Wei D, González-Marrachelli V, Melgarejo JD, et al. Cardiovascular risk of metabolically
healthy obesity in two european populations: Prevention potential from a metabolomic
study. *Cardiovasc Diabetol* 2023; 22: 82.

[35] Doumatey AP, Zhou J, Zhou M, et al. Proinflammatory and lipid biomarkers mediate
metabolically healthy obesity: A proteomics study. *Obes Silver Spring Md* 2016; 24: 1257–
1265.

[36] Doulamis IP, Konstantopoulos P, Tzani A, et al. Visceral white adipose tissue and serum
proteomic alternations in metabolically healthy obese patients undergoing bariatric
surgery. *Cytokine* 2019; 115: 76–83.

[37] Cheng Q, Yuan X, Lin S, et al. Serum proteome profiling reveals differentially expressed
proteins between subjects with metabolically healthy obesity and nonalcoholic fatty liver
disease. *J Proteomics* 2022; 260: 104556.

[38] Liao J, Goodrich JA, Chen W, et al. Cardiometabolic profiles and proteomics associated
with obesity phenotypes in a longitudinal cohort of young adults. *Sci Rep* 2024; 14: 7384.

[39] Pattaro C, Gögele M, Mascalzoni D, et al. The Cooperative Health Research in South Tyrol
(CHRIS) study: rationale, objectives, and preliminary results. *J Transl Med* 2015; 13: 348.

[40] Noce D, Gögele M, Schwienbacher C, et al. Sequential recruitment of study participants
may inflate genetic heritability estimates. *Hum Genet* 2017; 136: 743–757.

[41] Obesity: preventing and managing the global epidemic. Report of a WHO consultation.
World Health Organ Tech Rep Ser 2000; 894: i–xii, 1–253.

[42] van Vliet-Ostaptchouk JV, Nuotio M-L, Slagter SN, et al. The prevalence of metabolic
syndrome and metabolically healthy obesity in Europe: a collaborative analysis of ten large
cohort studies. *BMC Endocr Disord* 2014; 14: 9.

[43] Hantikainen E, Weichenberger CX, Dordevic N, et al. Metabolite and protein associations
with general health in the population-based CHRIS study. *Sci Rep* 2024; 14: 26635.

[44] Verri Hernandes V, Dordevic N, Hantikainen EM, et al. Age, Sex, Body Mass Index, Diet and
Menopause Related Metabolites in a Large Homogeneous Alpine Cohort. *Metabolites*
2022; 12: 205.

[45] Messner CB, Demichev V, Bloomfield N, et al. Ultra-fast proteomics with Scanning SWATH.
Nat Biotechnol 2021; 39: 846–854.

[46] Dordevic N, Dierks C, Hantikainen E, et al. Pervasive Influence of Hormonal
Contraceptives on the Human Plasma Proteome in a Broad Population Study. 2023;
2023.10.11.23296871.

[47] Liaw A, Wiener M. Classification and Regression by randomForest. *R News* 2002; 2: 18–22.

[48] Archer E. rfPermute: Estimate Permutation p-Values for Random Forest Importance
Metrics, https://cran.r-project.org/package=rfPermute (2023).

[49] Borén J, Packard CJ, Taskinen M-R. The Roles of ApoC-III on the Metabolism of Triglyceride-
Rich Lipoproteins in Humans. *Front Endocrinol* 2020; 11: 474.

[50] Mendivil CO, Zheng C, Furtado J, et al. Metabolism of very-low-density lipoprotein and low-
density lipoprotein containing apolipoprotein C-III and not other small apolipoproteins.
Arterioscler Thromb Vasc Biol 2010; 30: 239–245.

[51] Telle-Hansen VH, Christensen JJ, Formo GA, et al. A comprehensive metabolic profiling of
the metabolically healthy obesity phenotype. *Lipids Health Dis* 2020; 19: 90.

[52] Navarro-González D, Sánchez-Íñigo L, Fernández-Montero A, et al. Are all metabolically
healthy individuals with obesity at the same risk of diabetes onset? *Obes Silver Spring Md*
2016; 24: 2615–2623.

[53] Bertran L, Rusu EC, Guirro M, et al. Circulating proteomic profiles in women with morbid
obesity compared to normal-weight women. *J Proteomics* 2025; 310: 105317.

[54] Geyer PE, Wewer Albrechtsen NJ, Tyanova S, et al. Proteomics reveals the effects of
sustained weight loss on the human plasma proteome. *Mol Syst Biol* 2016; 12: 901.

[55] Wulff AB, Nordestgaard BG, Tybjærg-Hansen A. APOC3 Loss-of-Function Mutations,
Remnant Cholesterol, Low-Density Lipoprotein Cholesterol, and Cardiovascular Risk:
Mediation- and Meta-Analyses of 137 895 Individuals. *Arterioscler Thromb Vasc Biol* 2018;
38: 660–668.

[56] Tall AR, Thomas DG, Gonzalez-Cabodevilla AG, et al. Addressing dyslipidemic risk beyond
LDL-cholesterol. *J Clin Invest* 2022; 132: e148559.

[57] Povel CM, Boer JMA, Reiling E, et al. Genetic variants and the metabolic syndrome: a
systematic review. *Obes Rev Off J Int Assoc Study Obes* 2011; 12: 952–967.

[58] Olivieri O, Bassi A, Stranieri C, et al. Apolipoprotein C-III, metabolic syndrome, and risk of
coronary artery disease. *J Lipid Res* 2003; 44: 2374–2381.

[59] Post N, Yu R, Greenlee S, et al. Metabolism and Disposition of Volanesorsen, a 2'-O-(2
methoxyethyl) Antisense Oligonucleotide, Across Species. *Drug Metab Dispos Biol Fate*
Chem 2019; 47: 1164–1173.

[60] Zembic A, Eckel N, Stefan N, et al. An Empirically Derived Definition of Metabolically
Healthy Obesity Based on Risk of Cardiovascular and Total Mortality. *JAMA Netw Open*
2021; 4: e218505.

ACKNOWLEDGEMENTS

CHRIS study investigators thank all study participants, the Healthcare System of the Autonomous Province of Bolzano-South Tyrol, and all Eurac Research staff involved in the study (https://www.eurac.edu/chrisack).

AUTHOR CONTRIBUTIONS

Conceptualization: CXW, GP, FM and FSD. Methodology: GP, CXW, JR and FSD. Software: GP and CXW. Formal analysis and investigation: GP, CXW and FSD. Resources: JR, MR, MK, FSD and PPP. Data curation: CXW, JR and MR. Writing: GP, CXW, EH, JR, MDG and FSD. Visualization: CXW. Supervision: FM and FSD. Funding acquisition: FM, MK, FSD and PPP. All authors approved the final version of the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

The CHRIS study was funded by the Autonomous Province of Bolzano - Department of Innovation, Research, University and Museums and supported by the European Regional Development Fund (FESR1157). The research was funded by the European Region Tyrol South Tyrol Trentino, project EUREGIO EFH (decision 2059 from 1 December 2017, Autonomous Province of Trento).

594 **ADDITIONAL INFORMATION**

595 **Supplementary Information**

596 Supplementary File, PDF