Maternal and child healthcare-seeking among victims of violence in armed conflict: A quasi-experimental study in northeast Nigeria*

Max Schaub, *University of Hamburg & WZB Berlin Social Science Center, Germany*
Gbadebo Collins Adeyanju, *University of Erfurt, Germany*
Aisha Aliyu Abulfathi, *University of Maiduguri & University of Maiduguri Teaching Hospital, Nigeria*
Musa Muhammad Bello, *Bayero University & Aminu Kano Teaching Hospital, Nigeria*
Lennart Kasserra, *University of Konstanz, Germany*
Aminatu Ayaba Kwaku, *Bayero University & Aminu Kano Teaching Hospital, Nigeria*
Muhammad Ibrahim Jalo, *Women and Children Hospital, Damaturu, Nigeria*
Ahmad Mahmud, *Modibbo Adama University & Modibbo Adama University Teaching Hospital, Nigeria*
Pia Schrage, *University of Erfurt, Germany*
Rabiu Ibrahim Jalo, *Bayero University & Aminu Kano Teaching Hospital, Nigeria*
Liliana Abreu, *University of Konstanz, Germany*

Introduction: Armed conflict severely impacts health, with indirect deaths often exceeding direct casualties 2–4 times, disproportionately affecting women and children. Although the magnitude of these effects is well-documented, the mechanisms driving them remain insufficiently understood. This study shifts the focus from supply-side factors, such as the destruction of infrastructure, to demand-side processes, particularly healthcare-seeking behavior, and from broader conflict exposure to individual-level violent victimization.

Methods: Data come from a representative survey (n = 3,006) of caregivers of young children in northeastern Nigeria, a region heavily affected by armed insurgency. Unlike previous studies, our survey included dedicated measures of victimization, health-seeking outcomes, and potential mediating factors within a single instrument, enabling precise measurement and analysis. To compare maternal and child healthcare-seeking behavior between victimized and non-victimized caregivers, we employed a quasi-experimental observational design using propensity-score matching on demographic and contextual characteristics. Causal mediation analysis was then used to identify the mechanisms linking victimization to health behaviors.

Results: Victimization was widespread, with 21% of respondents (n=651) having experienced a severe form of violence in the past 3 years. While maternal healthcare-seeking behavior appeared unaffected, victimization significantly reduced healthcare-seeking for child health. Children of victimized caregivers were markedly less likely to be fully immunized (Odds ratio (OR): 0.43, p<0.001) and to receive care at government health facilities (OR: 0.23, p<0.001). This decline was primarily driven by increased distrust in the health system (PM: 0.42–0.70, p<0.05), stemming from negative experiences during the conflict, particularly health worker absenteeism and victimization by state security forces.

Conclusion: Addressing fear and mistrust is key to improving healthcare-seeking in conflict-affected populations. Efforts should focus on providing security for government-run health facilities, reducing violence against civilians by state security forces, and restoring trust in healthcare and state institutions. Future research should explore effective strategies for achieving these objectives.

---

*We thank the respondents for sharing their experiences, the interviewers for their diligent work, and two anonymous reviewers, Merike Blofield, and Sara Tomczyk for helpful comments.

What is already known?

- Armed conflict has severe negative consequences for maternal and child health, yet the underlying mechanisms remain understudied.
- Recent research has called for a greater focus on how exposure to violence shapes the demand for health services and healthcare-seeking behavior—a gap this study addresses with a representative survey (n = 3,006) in northeastern Nigeria.

What are the new findings?

- Rather than approximating exposure through proximity to conflict, this study directly measures individual-level victimization and examines potential mediating mechanisms with high precision.
- 21% of respondents reported severe victimization; these individuals showed markedly lower child immunization rates (OR = 0.43, p < 0.001) and were less likely to use government health facilities (OR = 0.23, p < 0.001).
- These differences are driven by mistrust of healthcare workers, predicated on absenteeism and prior victimization by state security forces.

What do the new findings imply?

- Preventing further erosion of trust likely requires reliable health service delivery and restraint by security forces.
- To improve child health outcomes, trust-building in healthcare and state institutions should be integrated in recovery programs.

Statements

Contributorship
MS, GCA, and LA developed the research idea. All co-authors helped to develop the research design and survey instrument. RIJ and MS supervised the fieldwork, and AAA, AAK, MIJ, and AM led the interview data collection. MA and LK performed statistical analyses and interpreted results. MS drafted the initial manuscript, with extensive input from LA. All members reviewed and provided critical feedback. MS serves as guarantor.

Funding
This study was funded by the German Federal Ministry of Education and Research (BMBF, Grant No. 01KA2405A), the German Alliance for Global Health Research (GLOHRA, Grant No. 01KA2405A) under the ViVac project, and by the Ideas and Venture Fund of the University of Hamburg (Grant No. UHH/VP3/4/424).

Competing interests
MS (University of Hamburg) received support from the University's Ideas and Venture Fund (UHH/VP3/4/424); the study also received funding from BMBF/GLOHRA (01KA2405A). The funders had no role in the research or publication decisions. Views are the authors' own. No other competing interests declared.

Ethics
Interviews were conducted by trained interviewers in the local language (Hausa) using a pre-tested questionnaire. Participants were informed in advance of the sensitive nature of some of the questions included in the survey, and were informed that they could end the survey or skip individual questions at any time. Only individuals who gave their explicit informed consent were interviewed. The survey and data collection protocol were approved by the National Health Research Ethics Committee of Nigeria (NHREC; Approval No. NHREC/01/01/2007-14/01/2024), by the respective authorities of the three states, and by the University of Konstanz, Germany (Institutional Review Board (IRB) Statement 13/2024).

Data availability statement
Data are available in a public, open access repository. The de-identified survey microdata, replication code, and accompanying materials are available on the Open Science Framework (OSF) at: https://doi.org/10.17605/OSF.IO/6DE2R. Please cite the dataset as: [dataset] [47] Schaub M, Adeyanju GC, Abreu L. Replication data for: Maternal and child health-seeking among victims of violence in armed conflict: A matched case-control study in northeast Nigeria. Open Science Framework (OSF), February 16, 2025. https://doi.org/10.17605/OSF.IO/6DE2R.

Introduction

Studies of armed conflict and health have documented the devastating negative health effects of exposure to violence. Estimates suggest that indirect deaths from conflict (e.g., due to malnutrition and diseases) often exceed direct casualties by a factor of two to four [1–3], with women and children disproportionately affected [3–5]. While these broad patterns are well documented, gaps remain in our understanding of why violence has such detrimental health consequences. The best-documented explanations focus on supply-side factors: armed conflict is accompanied by the large-scale destruction of critical infrastructure, leaving affected populations without essential services and vulnerable to the spread of infectious diseases [6–9]. In addition, health workers frequently flee conflict-affected areas, further limiting the availability of medical services [7, 10–13]. Looking beyond such supply-side dynamics, recent research has shifted the focus to demand-side factors, particularly how exposure to violence affects healthcare-seeking behavior, defined as utilization of health services in armed conflict settings [12, 14–18]. Exposure to violence can induce fear, scholars have argued, leading individuals to reduce their movement and limit their pursuit of maternal and child healthcare [17–19]. Moreover, individuals living in conflict-affected regions often report lower trust in government institutions, a key predictor of healthcare utilization [17, 20–22].

In this regard, Nigeria represents a particularly relevant case. The country faces multiple forms of armed conflict, including the Boko Haram insurgency in the Northeast, recurrent clashes between farmers and herders in the Middle Belt, separatist agitation in the Southeast, and criminal banditry in the Northwest [23, 24]. These conflicts have had profound repercussions for the health system, ranging from the destruction of facilities and the displacement of health workers to interruptions in immunization programs and disease surveillance [25, 26]. In the Northeast, where our study is situated, healthcare capacity has been particularly strained, with many communities reliant on humanitarian organizations whose reach remains inconsistent due to insecurity [27, 28]. Existing research on Nigeria underscores the severity of these challenges: attacks on health services have contributed to elevated maternal and child mortality, malnutrition, and outbreaks of preventable diseases such as cholera and measles [29, 30]. Studies also document the psychological toll of conflict, including high levels of post-traumatic stress and depression among displaced populations [31, 32].

Against this backdrop, the present study contributes to this literature, but shifts the focus from broader exposure to violence—typically measured as proximity to attacks—to individual-level violent victimization. Even though victimization is one of the most direct consequences of exposure to armed conflict, its impact on healthcare-seeking has not been explicitly assessed. In conflict settings, large numbers of individuals suffer from attacks themselves or observe family members being killed or otherwise harmed. Our study systematically measures such instances of victimization and assesses their relationship with healthcare-seeking behavior. Additionally, we advance existing research by explicitly identifying and modeling potential pathways through which victimization translates into reduced healthcare-seeking, focusing on trauma, fear, and mistrust. It is well established that victimization can lead to persistent trauma, with survivors exhibiting significantly higher rates of post-traumatic stress symptoms, even decades after exposure [33–35]. As a consequence of trauma, victims may under-utilize healthcare services [36, 37]. It is also plausible that violent victimization may increase fear of further victimization, discouraging individuals from leaving their homes to seek healthcare for themselves,

their partner, or their children [19, 37, 38]. Similarly, victimization can erode trust in healthcare institutions. This may be because of negative experiences with healthcare services during the conflict or a generalization of mistrust from one state institution (e.g., the security forces that failed them) to other state institutions, such as government-provided healthcare. Distrustful individuals will stop using health services, e.g., by refusing to vaccinate their children [39, 40], or they will seek non-government-run alternatives [41, 42]. In order to consistently track these potential mediation effects, our study included measures for all of these dimensions in the same survey instrument.

Methods

Data Data come from a representative household survey (n=3,006) conducted in August and September 2024 in the three northeastern Nigerian states of Adamawa, Borno, and Yobe (see Figure S1 in the Supplementary Information (SI) for a graphical overview). The region has been severely affected by an armed insurgency for the last 15 years [48]. And while the security situation has stabilized somewhat in recent years, sporadic attacks on government and civilian targets persist, including raids by armed extremists on villages and attacks on hospitals [49]. In each state, two local government areas (LGAs, comparable to counties) were randomly selected, weighting by population size. Within each LGA, three political wards (out of 10 to 20) were chosen using the same weighting approach. Due to security constraints, three wards had to be replaced during data collection with random replacements. The primary sampling units (PSUs) were determined by a population-weighted draw of 500×500m grid cells, which generated interviewer starting points proportional to population size. Household recruitment followed a random walk with a 2:1 skip pattern. Sampling weights were generated to adjust for any over- or under-representation of PSUs in the final sample, ensuring unbiased summary statistics. Additional information about the survey, in line with the STROBE reporting guidelines [46], is provided in Tables S1–S3 in the SI, and replication data and code are available on OSF [47].

The target population for our survey were parents (men and women) of young children. In each household, a single person was chosen for interview. To be eligible for the interview, a person had to be the primary caregiver of a child 5 years of age or younger. Interviews were conducted by trained interviewers in the local language (Hausa) using a pre-tested questionnaire. Participants were informed in advance of the sensitive nature of some of the questions included in the survey, and were informed that they could end the survey or skip individual questions at any time. Only individuals who gave their explicit informed consent were interviewed. Participants provided demographic information, answered questions about maternal and child health, and reported on experiences of violence and attitudes toward government and healthcare institutions. For child health, we focused on the youngest child in the family, aged at least one year. For maternal health, we asked female respondents to report on their own experiences and male respondents to report on their partners' experiences. The survey and data collection protocol were approved by the National Health Research Ethics Committee of Nigeria (NHREC; Approval No. NHREC/01/01/2007-14/01/2024), by the respective authorities of the three states, and by the University of Konstanz, Germany (IRB Statement 13/2024).

Patient and public involvement The research project built on knowledge gained from qualitative fieldwork with focus group participants and key informants (community leaders and healthcare workers). These interviews provided important input for defining the final set of questions included in the survey. The data and findings

will be disseminated through a series of workshops, the first of which was held in Kano, Nigeria, in January 2025, and which are open to specialists from Nigerian ministries, nongovernmental organizations (NGOs), and journalists.

Study design and statistical approach For the analysis, we adopted a quasi-experimental propensity score matching approach. We matched individuals who reported being victimized in the past three years with non-victimized individuals using a 1:1 matching algorithm. For statistical modeling, we considered both conditional and unconditional logistic regression. Conditional logit relies only on within-pair variation, offering a rigorous approach to causal inference, but it discards a substantial portion of the data and limits flexibility for mediation analysis. The causal mediation framework we employ [63] requires an outcome model that can accommodate continuous and multiple mediators. Because conditional logit removes stratum-specific intercepts, it complicates estimation of mediated effects and precludes the inclusion of mediators that do not vary within pairs. By contrast, unconditional logistic regression allows us to estimate the outcome model required for mediation analysis [64]. We therefore opted for unconditional models as our default specification, applied to matched data with adjustment for the matching variables and clustering of standard errors at the matched-pair level [65, 66]. Robustness checks using conditional logistic regression are reported in Section H of the SI; substantively, both approaches yield very similar estimates.

We matched on the demographic characteristics of the respondent (sex, age, education, number of children) and of the child in question (sex, age) to shield against potential confounding along these dimensions. For example, age and sex may be related to both the likelihood of victimization and healthcare-seeking patterns. Second, to isolate the *demand* for healthcare, we also accounted for determinants of healthcare supply. Specifically, we included the self-reported walking distance (in minutes) to the nearest clinic and the remotely measured geographic distance between the interview locations and georeferenced clinic and hospital locations [52]. We further included several health indicators in our matching function to capture the relative quality of services, namely diphtheria-tetanus-pertussis (DPT3) and measles vaccination coverage, malaria and malnutrition prevalence, and the under-five mortality rate [53–57]. A full description of the included contextual covariates can be found in Table S14 in the SI. These indicators were recorded at the 500×500m grid cell level, the same scale used for sampling. Finally, we included covariates that are known determinants of armed violence, notably population density, accessibility (measured as the distance to the nearest major road), and state reach (measured as the distance to the regional capital) [58]. All time-varying context-level covariates were measured pre-treatment, between 2015 and 2019, i.e., before the start of the recording-period for victimization. Matching was performed on the propensity score, estimated by means of a probit model regressing the victimization indicator on the full set of controls, and a caliper set to 0.2 [59]. This procedure matched the full set of victimized individuals (n=651) to the same number of non-victimized control-group individuals, for an effective sample of n=1,302 used in most of the analyses.

Our models relate violent victimization to indicators of healthcare-seeking for maternal and child health. Violent victimization was assessed using a survey question asking whether the respondent or a household member had been "directly affected by organized violence" in the past three years. Organized violence was defined as violence perpetrated by "members of an organized group, such as an extremist group or armed bandits." Respondents were further instructed that being "affected" meant being "either the target or witness of

violence." Following this question, we asked whether respondents would mind sharing what had happened to them or their household member. We coded individuals as violently victimized if they reported that they or their household members were physically assaulted, had a household member killed, were kidnapped/abducted, were threatened with violence, or experienced sexual violence.

As the core outcomes, our study focused on four key indicators of maternal and child healthcare-seeking. These were i) utilization of antenatal care and ii) facility-based childbirth (delivery in a healthcare facility) for maternal healthcare-seeking, and iii) adherence to the recommended child immunization schedule and iv) use of government health facilities for child healthcare-seeking. Antenatal care use was operationalized as respondents reporting that they received four or more antenatal care visits, the number recommended by the Nigerian Ministry of Health. Facility-based birth is measured as respondents reporting that the child in focus was born in a health clinic or hospital. To measure immunization coverage, interviewers checked the child's official immunization card to verify whether they had received the full schedule of vaccinations recommended by the Nigerian Ministry of Health. To operationalize the use of government health facilities, we recorded whether individuals used a health facility for any of their children in the past 6 months, and the type of health facility they visited. We then coded reported use of government clinics or hospitals as one, and use of private clinics, traditional healers, and clinics run by religious institutions or other nongovernmental organizations as zero.

A slight uncertainty arises from temporal ordering. In principle, it is possible that some instances of victimization occurred after the health-seeking behavior they are linked to, particularly for maternal indicators, which may refer to events some time before the interview. Although none of the respondents reported such a sequence, we cannot entirely rule out this possibility. Interpretations should therefore be made with greater caution for outcomes that occurred further in the past. This concern does not apply to vaccination status, which was assessed at the time of the interview. Moreover, the usual caveats regarding self-reported behaviors apply, notably that these measures may be affected by imperfect recall or imprecise reporting. As long as such biases affect victimized and non-victimized respondents equally, they would not systematically bias our estimates. However, bias could arise if recall accuracy were directly correlated with victimization status—an effect that cannot be entirely ruled out, though we consider it unlikely. By contrast, measurement of child immunization status is not subject to this concern, as vaccination information was verified from the child's official immunization card.

Mediators In addition to these outcomes, we examined three potential mediating mechanisms: trauma, fear, and mistrust. The hypothesized causal framework is shown in Figure 1. Following prior research in conflict settings, trauma is most often assessed in terms of post-traumatic stress disorder (PTSD) and related symptomatology, typically using standardized instruments such as the Harvard Trauma Questionnaire or the PTSD Checklist (PCL) [60, 61]. PTSD reflects the psychological sequelae of past traumatic events [62], and in our study it was measured using the validated eight-item PTSD Checklist (PCL-8) [63]. Fear, by contrast, captures a forward-looking perception of continued threat, closely linked to the anticipation of future violence rather than past exposure [64]. In previous work, fear has been operationalized through self-reports of avoidance behaviors or through predicted local violence risk in conflict-affected settings [17, 19, 65]. We followed the self-report approach by asking respondents directly about their perceived likelihood to become victims of violence again

in the future. The third mediator, institutional mistrust, refers to the erosion of confidence in the state's ability or willingness to fulfill its protective and service-providing roles [66, 67]. Victimization implies that the state has failed in its role as protector, which may undermine confidence not only in security institutions but also in other state services such as healthcare. This spillover matters for health outcomes: mistrustful individuals are less likely to seek care, follow medical advice, or vaccinate their children [17, 21, 42, 68]. To capture this dimension, our survey included a question on whether respondents' trust in healthcare workers had increased, decreased, or remained unchanged over the past three years. The full wording of all survey questions can be found in Table S14 in the SI.

To conduct the mediation analysis, we adopt the causal mediation framework proposed by Imai et al. [48]. In practice, we use the unconditional logistic regression outcome model as the basis for the mediation analysis. We estimate separate regression models for each mediator conditional on victimization and covariates, and then decompose the total effect into the natural indirect effect (mediation pathway), the direct effect, and the proportion mediated (PM). To account for the hypothesized presence of multiple mediators, we include the respective other mediators as controls in each model [69]. A core identifying assumption of this approach is sequential unconfoundedness, meaning that there are no unmeasured confounders of the treatment–outcome or mediator–outcome relationships. In practice, this assumption is difficult to fully satisfy, even when controlling for a rich set of covariates as in the present analysis. Therefore, the mediation results should be interpreted with appropriate caution.

Figure 1: Hypothesized causal framework

Note: The figure shows the hypothesized causal relations explored in this paper. Individual victimization is assumed to affect healthcare-seeking behavior through trauma, fear, and mistrust. Arrows reflect temporal/causal ordering. The dashed arrow from healthcare-seeking behavior to health outcomes (e.g., child mortality) indicates the next logical link in the chain but is not estimated here. For an analysis testing the full causal chain with aggregate data, see [17].

Finally, we conducted a complementary regression analysis to identify predictors of declining trust in healthcare workers. The outcome was a binary indicator of whether respondents reported that their trust in healthcare workers had decreased over the past three years. We estimated our standard matched logistic regression models with predictors including reported absenteeism of healthcare workers, exposure to violence by security forces, fear for security at health facilities, worry about undue political influence on the health system, and the provision of free healthcare provided by NGOs. All models adjust for the same sociodemographic covariates used in the main models. All predictors were measured through self-report survey items (full wording in Table S14, SI). This analysis allows us to examine more directly the contextual factors underlying loss of trust, which emerged as the key mediator in our main models.

Results

Sample demographics and matching Table 1 provides summary statistics for our sample, victimization rates, healthcare-seeking behaviors, and other variables reported in this study. The table reports averages after applying sample weights. The corresponding raw frequencies are reported in the text. Statistics for all variables, including covariates, can be found in Table S4 in the SI, and balance statistics, which list values for victimized and non-victimized individuals separately in line with the STROBE [46] statement, are shown in Table S5, SI. As shown in Table 1, of the 3,006 respondents, 71% (n=2,172) were female and 29% (n=834) were male, with an average age of 36 years. Caregivers provided information on 3,006 children, of whom 48% (n=1,448) were female and 52% (n=1,558) male, with a mean age of 2.5 years. Propensity score matching achieved excellent balance, with none of the mean differences statistically significant and all standardized mean differences were below 7% (see Table S5 and Figure S4 in the SI).[1]

Victimization prevalence Table 1 (and Figure S2 in the SI) also shows the prevalence of different types of victimization in our sample. Common forms of victimization include physical assault, experienced by 11.1% (n=338) of respondents, and the killing or kidnapping of a household member, experienced by 4.9% (n=155) and 4.5% (n=129) of respondents, respectively. Most importantly, overall no fewer than 21% of respondents (n=651) reported being victimized by some form of severe violence, underscoring the importance of considering the impact of individual-level victimization on healthcare-seeking behavior, rather than examining only broader exposure to conflict.

Healthcare-seeking As for the prevalence of healthcare-seeking for child and maternal health, full childhood immunization coverage was 61% (n=1,563), significantly below the global average of over 80% [70]. Antenatal care was relatively high compared to other Nigerian states, with 77% (n=2,129) of women attending at least four recommended visits during pregnancy [71]. Children were born in a health clinic or hospital in 61% (n=1,559) of cases, slightly below the 70% estimated for births in low- and middle-income countries (LMICs) [72].

Table 1: Summary statistics for sample demographics, predictors, outcomes, and mediators

	Mean	SD	Min	Max	Count
Sample demographics					
Respondent female	0.71	0.45	0.00	1.00	3,006
Respondent age	36.19	10.52	15.00	82.00	3,006
Child female	0.48	0.50	0.00	1.00	3,006
Child age	2.49	1.30	1.00	5.00	3,006
Victimization prevalence					
Was/were physically assaulted	0.11	0.31	0.00	1.00	3,006
Had a household member killed	0.05	0.22	0.00	1.00	3,006
Was/were kidnapped/abducted	0.05	0.21	0.00	1.00	3,006

---

[1] Applying a much stricter caliper (0.025) results in a similar number of matched pairs (647 vs. 651), indicating that most pairs are very closely matched.

Witnessed someone else being killed	0.04	0.18	0.00	1.00	3,006
Was/were threatened with violence	0.03	0.16	0.00	1.00	3,006
Experienced sexual violence	0.01	0.10	0.00	1.00	3,006
Victimized during armed violence (any)	0.21	0.41	0.00	1.00	3,006
Healthcare-seeking behavior					
Child fully vaccinated	0.62	0.48	0.00	1.00	3,006
Recommended antenatal care	0.77	0.42	0.00	1.00	3,006
Birth at medical facility	0.61	0.49	0.00	1.00	3,006
Used government clinic for child	0.80	0.40	0.00	1.00	2,196
Child health outcomes					
Child had health issue when born	0.03	0.16	0.00	1.00	3,006
Child had health issue during last year	0.24	0.43	0.00	1.00	3,006
Mediators					
Fear of victimization	0.47	0.50	0.00	1.00	3,006
Post-traumatic stress disorder (PTSD)	0.24	0.43	0.00	1.00	3,006
Loss of trust in healthcare workers	0.19	0.39	0.00	1.00	3,006
Predictors of loss of trust					
State security perpetrator	0.08	0.27	0.00	1.00	3,006
Feared for security at health facility	0.03	0.18	0.00	1.00	3,006
Worried about undue influence	0.11	0.31	0.00	1.00	3,006
Free healthcare provided by NGO	0.08	0.27	0.00	1.00	3,006
Absenteeism of healthcare workers	0.19	0.40	0.00	1.00	3,006

Note: Summary statistics for the main outcome and treatment variables used in the study. Reduced number of observations in variable "Used government clinic for child" due to question only asked of individuals reporting use of healthcare.

Use of government facilities for child healthcare was reported by 80% (n=1,805) of respondents, reflecting the dominance of the public health sector in the country. Other measures included in the table are discussed below.

Statistical results Figure 2 shows the results of the matched case-control logistic regression model, which examines the relationship between healthcare-seeking behavior and violent victimization. Descriptive statistics are shown in Figure S3 in the SI, conditional logistic regression results are shown in Figure S5, and full regression outputs are included in Tables S5–S8. As shown in Figure 2, victimization was not significantly associated with whether respondents attended the recommended number of prenatal care visits (Odds ratio (OR): 1.25, 95% CI: 0.93 to 1.69). Contrary to expectations, the likelihood that a child born to victimized caregivers was delivered in a health clinic or hospital was slightly *higher* than among matched non-victimized individuals (OR: 1.30, 95% CI: 1.01 to 1.66). However, this result lacks full robustness across alternative model specifications (see Tables S9–S11 in the SI).

Thus, healthcare-seeking for maternal health does not differ significantly between victimized and non-victimized individuals. In contrast, there are strong differences in healthcare-seeking for children's health. Children of victimized caregivers were significantly less likely to be fully vaccinated (OR: 0.43, 95% CI: 0.32 to 0.57). A striking difference emerged in healthcare-seeking *patterns*: children of victimized caregivers were far less likely to be taken to government-run clinics compared to private facilities (OR: 0.23, 95% CI: 0.14 to 0.39).

Figure 2: Effect of violent victimization on healthcare-seeking behavior and child health

Note: Coefficient plots from a regression of indicators of healthcare-seeking for maternal and child health, and of child health outcomes, on violent victimization. Reported coefficients are odds ratios after unconditional logistic regression using the matched sample. Markers are point estimates, lines 95% confidence intervals. Standard errors clustered at the level of the matched pair.

Figure 2 also shows that victimization was associated with poorer child health. Children of caregivers who experienced violence were less likely to be fully healthy at birth (OR: 5.47, 95% CI: 2.64 to 11.34) and more likely to have recently suffered from health issues such as respiratory, digestion, or coordination problems; or hearing and vision problems (OR: 1.48, 95% CI: 1.10 to 1.99).

To confirm the validity of our findings, we conducted a number of robustness checks. First, conditional logistic regression yields very similar estimates to our main unconditional models: antenatal care (OR: 1.14, 95% CI: 0.89 to 1.45 vs. OR: 1.08), skilled birth attendance (OR: 1.26, 95% CI: 1.01 to 1.59 vs. OR: 1.18), full child vaccination (OR: 0.50, 95% CI: 0.40–0.63 vs. OR: 0.43), and use of government facilities for child care (OR: 0.17, 95% CI: 0.10–0.28 vs. OR: 0.23). Second, we restricted the analysis to female respondents and found that the results remained largely unchanged compared to using the full sample (Table S9). Third, applying small-area (500×500m) fixed effects likewise yielded very similar estimates (Table S10, SI). This suggests that our findings are robust even when comparing closely matched individuals living in the same local area, where healthcare supply factors should be constant, making it highly unlikely that the observed differences in healthcare-seeking behavior are driven by variation in service availability rather than the effects of victimization. Finally, probing for differential effects within our main independent variable, we found that the observed negative association between victimization and healthcare-seeking appears to be driven by direct forms of victimization (physical assault, abduction, or sexual violence), whereas indirect victimization (seeing someone killed or losing a household member) did not consistently reduce healthcare-seeking behavior (Table S12, SI).

Mediation analysis What explains the lower healthcare-seeking for child health and the shift away from government healthcare among victimized respondents? To investigate this, we examined three key factors discussed in the literature: i) fear of future victimization, which may lead to avoidance behavior; ii) poor mental

health, which may impair individuals' ability to engage in healthcare-seeking behavior; and iii) loss of trust in healthcare personnel, which may prevent victimized individuals from engaging with government institutions.

Table 2 indicates the extent to which the effect of violent victimization on child health is mediated by these three factors, keeping in mind the methodological limitations of this kind of analysis. For childhood vaccination rates, neither fear of future victimization (Proportion mediated (PM): 0.01, 95% CI: -0.09 to 0.12) nor PTSD (PM: 0.01, 95% CI: -0.11 to 0.13) seem to mediate the effect of victimization, despite prior work documenting or suggesting a mediating role for trauma [73–78]. This divergence may reflect characteristics of our setting and measurement: PTSD symptoms were captured with the PCL-8, which may underrepresent avoidance and numbing domains most relevant to preventive care, while contextual constraints such as staff absenteeism, politicization of services, and indeed fear and distrust, may act more proximally on care-seeking behavior than trauma [17, 19]. The non-detection of a mediating relationship does not imply that victimization has no effect on PTSD, however. On the contrary, all three mediators—including fear and PTSD—are strongly predicted by violent victimization, as shown in Table S14 in the SI.

Table 2: Mediation analyses

Dependent variable:	Child fully vaccinated			Used government clinic for child		
	(1)	(2)	(3)	(4)	(5)	(6)
Mediator:	FearVict	PTSD	DistrustMed	FearVict	PTSD	DistrustMed
Nat. indirect effect	-0.001	-0.001	-0.041**	-0.030***	0.003	-0.089***
	(-0.011 to 0.009)	(-0.010 to 0.009)	(-0.073 to -0.009)	(-0.050 to -0.010)	(-0.002 to 0.008)	(-0.137 to -0.041)
Total effect	-0.096***	-0.082***	-0.128***	-0.073**	-0.052*	-0.127***
	(-0.150 to -0.042)	(-0.138 to -0.026)	(-0.174 to -0.082)	(-0.138 to -0.008)	(-0.110 to 0.006)	(-0.173 to -0.080)
Proportion mediated (PM)	0.01	0.01	0.32**	0.41*	-0.06	0.70***
	(-0.09 to 0.12)	(-0.11 to 0.13)	(0.03 to 0.61)	(-0.04 to 0.86)	(-0.17 to 0.06)	(0.25 to 1.16)
N	1,302	1,302	1,302	979	979	979

Note: Mediation analysis for the effect of violent victimization on child healthcare-seeking. Models 1–3 show mediation analyses for our indicator as to whether a child is fully vaccinated, Models 4–6 show mediation analyses for the effect of victimization on child healthcare-seeking at government health clinics or hospitals. Nat. indirect effect stands for natural indirect (mediated) effect. FearVict records respondents reporting fear of future violence, PTSD measures respondents suffering from post-traumatic stress disorder, and DistrustMed records whether respondents report a loss of trust in healthcare workers. Reported coefficients are marginal effects after logistic regression. Matched sample of victimized and non-victimized respondents. Proportions mediated do not sum to 100 since baseline values vary (the total effect is estimated conditional on the other mediators). Standard errors clustered at the level of the matched pair, 95% confidence intervals in parentheses, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Supporting this idea, loss of trust in healthcare workers emerges as a strong and statistically significant mediator, explaining approximately one third (PM: 0.32, 95% CI: 0.03 to 0.61) of the effect of victimization. The mediation pattern is more complex for children's healthcare-seeking in government facilities. Here, 41% of the effect of victimization (controlling for other mediators) could be attributed to perceived insecurity, a result that approaches conventional statistical significance (PM: 0.41, 95% CI: -0.04 to 0.86). PTSD, once again, does not mediate the effect of victimization (PM: -0.06, 95% CI: -0.17 to 0.06). However, mistrust in healthcare providers appears to have an extraordinarily strong mediating effect, accounting for no less than 70% of the conditional effect of victimization (PM: 0.70, 95% CI: 0.25 to 1.16). Our analysis thus strongly suggests that fear and especially distrust play important mediating roles. While these results align with previous research, they provide

more direct empirical evidence linking both factors to victimization rather than to general exposure to armed violence. In contrast, despite its theoretical plausibility and prior empirical support, in our setting poor mental health (PTSD) does not appear to explain the observed association between victimization and healthcare-seeking for children.

Determinants of the loss of trust in healthcare workers Given the important role of a loss of trust in mediating healthcare-seeking behavior, we investigated its underlying determinants in the final step of our analysis. A key finding was that loss of trust in health workers was strongly correlated with loss of trust in other state institutions. Specifically, individuals who lose trust in the government (OR: 35.76, 95% CI: 23.72 to 53.91) and in the security forces (OR: 23.76, 95% CI: 17.43 to 32.38) were significantly more likely to lose trust in healthcare providers. These findings suggest that distrust in healthcare workers extends beyond the health sector, reflecting a broader erosion of trust in state institutions. This generalized distrust may result from a perceived failure of the government or security forces to protect individuals from violence, leading victims to associate the healthcare system with the same institutions they hold responsible for their victimization. Second, we regress the indicator for loss of trust in healthcare workers on several conflict-related experiences reported in our survey. Specifically, we focus on whether distrust is linked to: i) violent victimization at the hands of state security forces, ii) healthcare worker absenteeism when respondents sought medical assistance, iii) fear for personal safety while at a health facility, iv) concerns that health facilities were unduly influenced by the government, or v) receiving free healthcare from nongovernmental organizations (NGOs).

Victimization by state security forces is alarmingly common in our sample. Among those who experienced violence, more than one-third of victimized respondents (38%) – equivalent to 8% (n=212) of the full sample, see Table 1 – reported that state security forces were the perpetrators. Such experiences are likely to undermine trust in the government, which in turn may spill over to health workers [66, 79, 80].

Figure 3: Determinants of loss of trust in healthcare workers

Note: Coefficient plots from a regression of the indicator for loss of trust in healthcare workers on various potential explanatory variables. Reported coefficients are odds ratios after unconditional logistic regression using the matched sample. Markers are point estimates, lines 95% confidence intervals. Standard errors clustered at the level of the matched pair.

Similarly, health worker absenteeism, experienced by 17% (n=527) of respondents, has previously been found to correlate with lower levels of trust [81]. Concerns about the security of health facilities, cited by 4% (n=122) of respondents, and about undue political influence on health facilities, cited by 13% (n=396) of respondents, may also contribute to mistrust. In contrast, receiving free healthcare from NGOs, reported by 9%

(n=272) of respondents, may have the opposite effect. Previous research suggests that humanitarian interventions providing free medical care can mitigate the negative effects of conflict exposure [18], and it is plausible that access to such services helps to restore trust in healthcare workers [68].

Figure 3 shows the results of the regression analysis. The dominant factors explaining the loss of trust in healthcare are victimization by state security forces (OR: 30.23, 95% CI: 17.64 to 51.81) and absenteeism of healthcare workers (OR: 13.43, 95% CI: 9.58 to 18.85). Other significant, but less pronounced, predictors include: feeling unsafe when seeking healthcare (OR: 2.01, 95% CI: 1.10 to 3.70) and worrying about undue political influence on the healthcare system (OR: 1.50, 95% CI: 1.02 to 2.20). Meanwhile, receiving free healthcare is associated with a relative gain in trust, but not significantly so (OR: 0.76, 95% CI: 0.47 to 1.22). Overall, the results clearly show how failed attempts to access healthcare and negative experiences with other government institutions contribute to a decline in trust.

Conclusion

Our study advances the literature on the health effects of armed conflict in three key ways. First, we focus explicitly on the effects of individual-level victimization, a quantitatively significant yet often overlooked consequence of armed conflict. Unlike broad measures of exposure to conflict, which are often defined in terms of proximity to violent events, victimization is a more direct and specific experience. In our sample, 21% of respondents reported experiencing severe forms of victimization, such as being violently attacked or having a household member killed. These respondents exhibit markedly different healthcare-seeking behaviors compared to non-victimized (but still conflict-exposed) individuals living in close proximity, underscoring the importance of looking beyond general exposure. Second, echoing other recent research [9, 15–19, 22], we focus on demand-side effects, i.e., healthcare-seeking, rather than on supply-side factors such as the destruction of infrastructure. By adjusting for healthcare supply factors, such as proximity to health facilities or availability of health workers, and by using small-area fixed effects, we effectively isolated the effect of violence on the demand for healthcare. We found that while maternal health-seeking was resilient to victimization, child health outcomes showed acute vulnerability, with children of victimized caregivers having significantly lower vaccination rates and much lower rates of presentation to government health facilities. This is consistent with other studies that have found that pediatric care tends to be neglected in conflict-affected areas [18, 82]. Third, looking beyond top-level associations, we present evidence for a mechanism by which individual-level victimization translates into lower healthcare-seeking for child health and a shift in healthcare-seeking away from government-run facilities: a loss of trust in healthcare workers. We found that the loss of trust was most strongly associated with absenteeism among healthcare workers and instances of violent victimization by state security forces. Contrary to theoretical expectations [36, 37], victimization-induced trauma did not mediate the relationship between victimization and healthcare-seeking behavior.

As always, this study is not without limitations. For one, the cross-sectional nature of the data limits our ability to definitively establish causality. Although the matched case-control design strengthens causal inference, we cannot completely rule out the possibility that unobserved confounders may have influenced the results. In addition, the study focuses on a specific region in Nigeria, which may limit the generalizability of the findings to other conflict-affected contexts. Future research should examine these relationships in different

settings and strengthen causal inference by using longitudinal data to track causal pathways linking victimization, trust, and health-seeking behaviors over time.

While more research is needed to validate the results, our findings suggest several policy priorities. A first area of intervention is to prevent the loss of trust. Our findings suggest that to achieve this goal, at least basic emergency services should be provided throughout a violent conflict, and security forces should exercise the utmost discipline and care to avoid harming civilians. A second area of intervention is to mitigate the consequences where trust has already been lost and healthcare-seeking for children has been reduced. While other work has found that aid provided by NGOs can help counteract the negative effects of exposure to armed conflict on trust [18, 68], our evidence suggests that such aid may not be sufficient to rebuild trust among actual victims.

This calls for a deeper exploration of how trust might be rebuilt. One possible strategy is targeted child health interventions, for instance in the form of community engagement strategies [83–85], which could help rebuild state-community relations. Political science research also suggests that legal accountability for perpetrators is important for rebuilding trust [86], and that listening exercises that explicitly acknowledge suffering can be used to restore trust in the state after conflict [87–89]. Such exercises could take the form of outreach and communication campaigns by health authorities—for example, acknowledging hardships and encouraging families to return to government health services. Combining such symbolic acknowledgment with visible improvements in service quality may be particularly effective, as it couples recognition with tangible action and incentives [90].

Evidence from other conflict-affected settings shows that interventions can be designed to both improve health outcomes and rebuild trust in service delivery. Community Health Worker (CHW) programs, which relied on locally recruited workers and participatory planning, improved immunization coverage and strengthened community confidence in the health system [91–93]. Similar approaches could be adapted in Nigeria, where trust has likewise proven central to sustained healthcare utilization [94]. Experience from Northern Nigeria and elsewhere demonstrate that adaptive, multi-level engagement with governance, financing, and community systems can strengthen health system resilience even under insecurity [95, 96]. Building on such principles, displaced persons or youth volunteers could be trained as health promoters in IDP camps and host communities, and mobile units staffed by female health workers could enhance access and trust among women in conservative or insecure areas [97, 98]. Conflict-sensitive vaccination campaigns that engage local and religious leaders to counter misinformation, as successfully implemented elsewhere [99, 100], could likewise inform strategies to overcome vaccine hesitancy and insecurity. These approaches must, however, be carefully tailored to Nigeria's sociopolitical realities, including collaboration with religious institutions and strict avoidance of coercive associations with security actors.

Ultimately, our findings underscore that in contexts of protracted conflict, rebuilding trust is as central to improving health outcomes as restoring supply, and future interventions should be rigorously evaluated with this dual objective in mind.

References

[1] Hazem Adam Ghobarah, Paul Huth, and Bruce Russett. "The Post-War Public Health Effects of Civil Conflict". *Social Science & Medicine* 59.4 (Aug. 1, 2004), pp. 869–884.

[2] Iris Kesternich et al. "The Effects of World War II on Economic and Health Outcomes Across Europe". *The Review of Economics and Statistics* 96.1 (2014), pp. 103–118.

[3] Zachary Wagner et al. "Women and Children Living in Areas of Armed Conflict in Africa: A Geospatial Analysis of Mortality and Orphanhood". *The Lancet Global Health* 7.12 (Dec. 1, 2019), e1622–e1631.

[4] Thomas Plümper and Eric Neumayer. "The Unequal Burden of War: The Effect of Armed Conflict on the Gender Gap in Life Expectancy". *International Organization* 60.3 (July 2006), pp. 723–754.

[5] Mevlude Akbulut-Yuksel. "War during Childhood: The Long Run Effects of Warfare on Health". *Journal of Health Economics* 53 (May 1, 2017), pp. 117–130.

[6] Zaryab Iqbal and Christopher Zorn. "Violent Conflict and the Spread of HIV/AIDS in Africa". *The Journal of Politics* 72.1 (Jan. 1, 2010), pp. 149–162.

[7] Adanna Chukwuma and Uche Eseosa Ekhator-Mobayode. "Armed Conflict and Maternal Health Care Utilization: Evidence from the Boko Haram Insurgency in Nigeria". *Social Science & Medicine* 226 (Apr. 1, 2019), pp. 104–112.

[8] Shanna A. Kirschner and Amelia B. Finaret. "Conflict and Health: Building on the Role of Infrastructure". *World Development* 146 (Oct. 1, 2021), p. 105570.

[9] Daniel Meierrieks and Max Schaub. "Terrorism and Child Mortality". *Health Economics* 33.1 (2024), pp. 21–40.

[10] Susan McKay. "The Effects of Armed Conflict on Girls and Women". *Peace and Conflict: Journal of Peace Psychology* 4.4 (Dec. 1, 1998), pp. 381–392.

[11] Sima L. Sharara and Souha S. Kanj. "War and Infectious Diseases: Challenges of the Syrian Civil War". *PLOS Pathogens* 10.11 (Nov. 13, 2014), e1004438.

[12] Primus Che Chi et al. "Perceptions of the Effects of Armed Conflict on Maternal and Reproductive Health Services and Outcomes in Burundi and Northern Uganda: A Qualitative Study". *BMC International Health and Human Rights* 15.1 (Apr. 3, 2015), p. 7.

[13] Thomas Druetz et al. "Effects of Terrorist Attacks on Access to Maternal Healthcare Services: A National Longitudinal Study in Burkina Faso". *BMJ Global Health* 5.9 (Sept. 1, 2020), e002879.

[14] James I Price and Alok K Bohara. "Maternal Health Care amid Political Unrest: The Effect of Armed Conflict on Antenatal Care Utilization in Nepal". *Health Policy and Planning* 28.3 (May 1, 2013), pp. 309–319.

[15] Pontius Bayo et al. "Maternal and Neonatal Health Care Service Utilisation in the Wake of Active Conflict and Socio-Economic Downturn in Torit County, Republic of South Sudan: A Multimethod Locally Driven Study". *African Journal of Reproductive Health* 25.3s (3s Oct. 14, 2021), pp. 30–42.

[16] Olabayo Ojeleke et al. "The Impact of Armed Conflicts on Health-Care Utilization in Northern Nigeria: A Difference-in-Differences Analysis". *World Medical & Health Policy* 14.4 (2022), pp. 624–664.

[17] Max Schaub. "Violent Conflict and the Demand for Healthcare: How Armed Conflict Reduces Trust, Instills Fear, and Increases Child Mortality". *Social Science & Medicine* 359 (Oct. 1, 2024), p. 117252.

[18] Gbadebo Collins Adeyanju et al. "Armed Violent Conflict and Healthcare-Seeking Behavior for Maternal and Child Health in Sub-Saharan Africa: A Systematic Review". *PLOS One* (Feb. 2025), p. 2024.03.11.24304112.

[19] Augustin Tapsoba. "The Cost of Fear: Impact of Violence Risk on Child Health during Conflict". *Journal of Development Economics* 160 (Jan. 1, 2023), p. 102975.

[20] Adanna Chukwuma, Thomas J. Bossert, and Kevin Croke. "Health Service Delivery and Political Trust in Nigeria". *SSM - Population Health* 7 (Apr. 1, 2019), p. 100382.

[21] Nik Stoop, Kalle Hirvonen, and Jean-Francois Maystadt. "Institutional Mistrust and Child Vaccination Coverage in Africa". *BMJ Global Health* 6.4 (Apr. 1, 2021), e004595.

[22] Tingkai Zhang et al. "Does Armed Conflict Lead to Lower Prevalence of Maternal Health-Seeking Behaviours: Theoretical and Empirical Research Based on 55 683 Women in Armed Conflict Settings". *BMJ Global Health* 8.8 (Aug. 23, 2023).

[23] Centre for Democracy and Development. *Multiple Nodes, Common Causes: National Stocktake of Contemporary Insecurity and State Responses in Nigeria*. Report. 2022.

[24] Human Rights Watch. *World Report 2024: Nigeria*. Country chapter. 2024.

[25] Ayodele O. Albert and Andrew O. Thaddaeus. "The nexus of banditry, farmer-herder clashes and insecurity in Nigeria: Causes, consequences and policy responses". *International Journal of Research and Innovation in Social Science* (2025), pp. 2083–2090. DOI:10.47772/IJRISS. 2025.905000164.

[26] Paul Timipa Igoli et al. "Impact of armed conflict on immunization and disease surveillance in Nigeria". *Conflict and Health* 17 (2023), p. 88. DOI:10.1186/s13031-023-00588-w.

[27] Aisha Ahmed. *A silent wildfire: The unfolding crisis of acute malnutrition among children in Northeast Nigeria*. Conference presentation, Florence Nightingale's Annual Week. 2025. DOI:10.13140/RG.2.2.32379.63526.

[28] Health Sector Working Group Nigeria. *Northeast Nigeria Humanitarian Response: Health Sector Bulletin (June 2025)*. ReliefWeb. 2025.

[29] R. O. Adesola et al. "Navigating Nigeria's health landscape: Population growth and its health implications". *Environmental Health Insights* 18 (2024), p. 11786302241250211. DOI:10.1177/11786302241250211.

[30] Bashir H. Gulumbe et al. "Cholera outbreaks in conflict-hit regions of Northern Nigeria: An overview". *Afro-Egyptian Journal of Infectious and Endemic Diseases* 13.2 (2023), pp. 134–142. DOI:10.21608/aeji.2023.205724.1286.

[31] V. O. Femi-Lawal et al. "Effects of armed conflicts and insecurity on the mental health of Nigerians". *Commentary* 3.1 (2023), p. 53. DOI:10.56101/rimj.v3i1.53.

[32] U. U. Ibrahim, A. A. Aliyu, O. A. Abdulhakeem, et al. "Prevalence of Boko Haram crisis-related depression and post-traumatic stress disorder symptomatology among internally displaced persons in Yobe State, Nigeria". *Journal of Mental Health in Africa* 5.2 (2023), pp. 145–162.

[33] Joop T. V. M. de Jong et al. "Lifetime Events and Posttraumatic Stress Disorder in 4 Postconflict Settings". *JAMA* 286.5 (Aug. 1, 2001), pp. 555–562.

[34] Phuong N Pham, Harvey M Weinstein, and Timothy Longman. "Trauma and PTSD Symptoms in Rwanda: Implications for Attitudes toward Justice and Reconciliation". *JAMA: the journal of the American Medical Association* 292.5 (Aug. 4, 2004), pp. 602–612.

[35] Richard Neugebauer et al. "Post-Traumatic Stress Reactions among Rwandan Children and Adolescents in the Early Aftermath of Genocide". *International Journal of Epidemiology* 38.4 (Aug. 1, 2009), pp. 1033–1045.

[36] Uchenna Nworah et al. "Afghanistan and Iraq War Veterans' Health Care Needs and Their Underuse of Health Care Resources: Implications for Psychiatric-Mental Health Nurses". *Journal of Psychosocial Nursing and Mental Health Services* 52.6 (June 2014), pp. 42–49.

[37] Marije J. Splinter et al. "Prevalence and Determinants of Healthcare Avoidance during the COVID-19 Pandemic: A Population-Based Cross-Sectional Study". *PLOS Medicine* 18.11 (Nov. 23, 2021), e1003854.

[38] Cynthia S. Minkovitz et al. "Maternal Depressive Symptoms and Children's Receipt of Health Care in the First 3 Years of Life". *Pediatrics* 115.2 (Feb. 1, 2005), pp. 306–314.

[39] Ebenezer Obadare. "A Crisis of Trust: History, Politics, Religion and the Polio Controversy in Northern Nigeria". *Patterns of Prejudice* 39.3 (2005), pp. 265–284.

[40] Ayodele Samuel Jegede. "What Led to the Nigerian Boycott of the Polio Vaccination Campaign?" *PLOS Medicine* 4.3 (Mar. 20, 2007), e73.

[41] Anna Petherick. "Ebola in West Africa: Learning the Lessons". *The Lancet* 385.9968 (Feb. 14, 2015), pp. 591–592.

[42] Marcella Alsan and Marianne Wanamaker. "Tuskegee and the Health of Black Men". *The Quarterly Journal of Economics* 133.1 (Feb. 1, 2018), pp. 407–455.

[43] Abimbola Adesoji. "The Boko Haram Uprising and Islamic Revivalism in Nigeria". *Africa Spectrum* 45.2 (Aug. 1, 2010), pp. 95–108.

[44] Wisdom Oghosa Iyekekpolo. "Boko Haram: Understanding the Context". *Third World Quarterly* 37.12 (Dec. 1, 2016), pp. 2211–2228.

[45] F. S. Sofia and E. Chris. "Suspected Female Suicide Bombers Death Toll Rises to 32 in Nigeria". *BBC News* (2024).

[46] Erik von Elm et al. "The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies". *The Lancet* 370.9596 (Oct. 20, 2007), pp. 1453–1457.

[dataset] [47] Schaub M, Adeyanju GC, Abreu L. Replication data for: Maternal and child health-seeking among victims of violence in armed conflict: A matched case-control study in northeast Nigeria. Open Science Framework (OSF), February 16, 2025. https://doi.org/10.17605/OSF.IO/6DE2R

[48] Kosuke Imai, Luke Keele, and Dustin Tingley. "A General Approach to Causal Mediation Analysis". *Psychological Methods* 15.4 (2010), p. 309.

[49] Tyler J. VanderWeele and Eric J. Tchetgen Tchetgen. "Mediation Analysis With Matched Case-Control Study Designs". *American Journal of Epidemiology* 183.9 (May 1, 2016), pp. 869–870.

[50] Neil Pearce. "Analysis of Matched Case-Control Studies". *BMJ* 352 (Feb. 25, 2016), p. i969.

[51] Chia-Ling Kuo, Yinghui Duan, and James Grady. "Unconditional or Conditional Logistic Regression Model for Age-Matched Case–Control Data?" *Frontiers in Public Health* 6 (Mar. 2, 2018).

[52] OpenStreetMap contributors. *Nigeria Roads (OpenStreetMap Export)*. Accessed: 2024-12-02. Licensed under Open Database License (ODbL). Humanitarian OpenStreetMap Team (HOT), 2024.

[53] Jonathan F Mosser et al. "Mapping Diphtheria-Pertussis-Tetanus Vaccine Coverage in Africa, 2000–2016: A Spatial and Temporal Modelling Study". *The Lancet* 393.10183 (May 4, 2019), pp. 1843–1855.

[54] Alyssa N. Sbarra et al. "Mapping Routine Measles Vaccination in Low- and Middle-Income Countries". *Nature* 589.7842 (Jan. 2021), pp. 415–419.

[55] Daniel J Weiss et al. "Mapping the Global Prevalence, Incidence, and Mortality of Plasmodium Falciparum, 2000–17: A Spatial and Temporal Modelling Study". *The Lancet* 394.10195 (July 27, 2019), pp. 322–331.

[56] Aaron Osgood-Zimmerman et al. "Mapping Child Growth Failure in Africa between 2000 and 2015". *Nature* 555.7694 (Feb. 28, 2018), pp. 41–47.

[57] Roy Burstein et al. "Mapping 123 Million Neonatal, Infant and Child Deaths between 2000 and 2017". *Nature* 574.7778 (7778 Oct. 2019), pp. 353–358.

[58] Carl Müller-Crepon. "State Reach and Development in Africa since the 1960s: New Data and Analysis". *Political Science Research and Methods* (Nov. 10, 2021), pp. 1–10.

[59] Peter C Austin. "Optimal Caliper Widths for Propensity-Score Matching When Estimating Differences in Means and Differences in Proportions in Observational Studies". *Pharmaceutical Statistics* 10.2 (Mar. 2011), pp. 150–161.

[60] Cindy Darzi and John Hunsley. "The reliability of the Harvard Trauma Questionnaire: A reliability generalization meta-analysis". *Journal of Clinical Psychology* 73.6 (2017), pp. 609– 620. DOI:10.1002/jclp.22324.

[61] Joseph Christy et al. "The PTSD Checklist (PCL): A comprehensive review of psychometric properties across diverse populations". *Journal of Traumatic Stress* 34.2 (2021), pp. 254–268. DOI:10.1002/jts.22666.

[62] Zachary Steel et al. "Association of torture and other potentially traumatic events with mental health outcomes among populations exposed to mass conflict and displacement: a systematic review and meta-analysis". *JAMA* 302.5 (2009), pp. 537–549. DOI:10.1001/jama.2009.1132.

[63] Timothy J. Geier et al. "Validation of Abbreviated Four- and Eight-Item Versions of the PTSD Checklist for DSM-5 in a Traumatically Injured Sample". *Journal of Traumatic Stress*

33.3 (2020), pp. 218–226.

[64] Dan W. Grupe and Jack B. Nitschke. "Uncertainty and anticipation in anxiety: an integrated neurobiological and psychological perspective". *Nature Reviews Neuroscience* 14.7 (2013), pp. 488–501. DOI:10.1038/nrn3524.

[65] Bengt B. Arnetz et al. "Workplace violence prevention in healthcare: standards, research, and guidance". *Joint Commission Journal on Quality and Patient Safety* 48.4 (2022), pp. 190–196. DOI:10.1016/S1553-7250(22)00032-0.

[66] Alexander De Juan and Jan Henryk Pierskalla. "Civil War Violence and Political Trust: Microlevel Evidence from Nepal". *Conflict Management and Peace Science* 33.1 (Feb. 1, 2016), pp. 67–88.

[67] Mikko Mattila and Lauri Rapeli. "Just Sick of It? Health and Political Trust in Western Europe". *European Journal of Political Research* 57.1 (2018), pp. 116–134.

[68] Robert A. Blair, Benjamin S. Morse, and Lily L. Tsai. "Public Health and Public Trust: Survey Evidence from the Ebola Virus Disease Epidemic in Liberia". *Social Science & Medicine* 172 (Jan. 1, 2017), pp. 89–97.

[69] Tyler VanderWeele and Stijn Vansteelandt. "Mediation Analysis with Multiple Mediators". *Epidemiologic Methods* 2.1 (Jan. 3, 2014), pp. 95–115.

[70] Camille E. Jones. "Routine Vaccination Coverage — Worldwide, 2023". *MMWR. Morbidity and Mortality Weekly Report* 73 (2024).

[71] Richard Gyan Aboagye et al. "Health Insurance Coverage and Timely Antenatal Care Attendance in Sub-Saharan Africa". *BMC Health Services Research* 22.1 (Feb. 11, 2022), p. 181.

[72] Gary Joseph et al. "Inequalities in the Coverage of Place of Delivery and Skilled Birth Attendance: Analyses of Cross-Sectional Surveys in 80 Low and Middle-Income Countries". *Reproductive Health* 13.1 (June 17, 2016), p. 77.

[73] Jeanne L. Alhusen. "Intimate partner violence during pregnancy: maternal and neonatal outcomes". *Journal of Women's Health* 24.1 (2015), pp. 52–59. DOI:10.1089/jwh.2014.4872.

[74] Brittany Jamieson. "Exposure to interpersonal violence during pregnancy and its association with women's prenatal care utilization: A meta-analytic review". *Trauma, Violence, & Abuse* 21.5 (2020), pp. 904–921. DOI:10.1177/1524838018806511.

[75] Allison Testa. "Intimate partner violence and barriers to prenatal care". *Social Science & Medicine* 316 (2023), p. 115591. DOI:10.1016/j.socscimed.2022.115591.

[76] Jennifer Waldron et al. "Trauma symptoms and antenatal care engagement in pregnant women exposed to violence". *Journal of Interpersonal Violence* 37.13-14 (2022), NP12123–NP12147. DOI:10.1177/08862605211067423.

[77] Maria Christou-Ergos, Kerrin E. Wiley, and Julie Leask. "Traumatic events and vaccination decisions: A systematic review". *Vaccines* 10.6 (2022), p. 911. DOI:10.3390/vaccines10060911.

[78] Maria Christou-Ergos, Julie Leask, and Kerrin E. Wiley. "Association between traumatic life events and vaccine hesitancy: A cross-sectional Australian study". *Public Health* 216 (2023), pp. 1–6. DOI:10.1016/j.puhe.2023.05.004.

[79] Pauline Grosjean. "Conflict and Social and Political Preferences: Evidence from World War II and Civil Conflict in 35 European Countries". *Comparative Economic Studies* 56.3 (2014), pp. 424–451.

[80] Elsa Voytas and Benjamin Crisman. "State Violence and Participation in Transitional Justice: Evidence from Colombia". *Journal of Peace Research* 61.6 (Nov. 1, 2024), pp. 1069–1084.

[81] Larissa Klootwijk et al. "Absenteeism of Healthcare Workers in Primary Healthcare in Sub-Saharan Africa: A Scoping Review". *The International Journal of Health Planning and Management* n/a.n/a (2025).

[82] Anu Rammohan et al. "Exposure to Conflicts and the Continuum of Maternal Healthcare: Analyses of Pooled Cross-Sectional Data for 452,192 Women across 49 Countries and 82 Surveys". *PLOS Medicine* 18.9 (Sept. 28, 2021), e1003690.

[83] Alastair K. Ager et al. "Health Service Resilience in Yobe State, Nigeria in the Context of the Boko Haram Insurgency: A Systems Dynamics Analysis Using Group Model Building". *Conflict and Health* 9.1 (Oct. 5, 2015), p. 30.

[84] Anna Durrance-Bagale et al. "Community Engagement in Health Systems Interventions and Research in Conflict-Affected Countries: A Scoping Review of Approaches". *Global Health Action* 15.1 (Dec. 31, 2022), p. 2074131.

[85] Liliana Abreu et al. "Mistrust, Violence, and Patriarchy: Perceived Barriers to Maternal and Child Healthcare Access in Conflict-Affected Northern Nigeria". Working paper. 2025.

[86] Risa Kitagawa. "From Political Violence to Political Trust? How Transitional Justice Affects Citizen Views of Government". *International Studies Quarterly* 67.1 (Mar. 1, 2023), sqad013.

[87] Marc L Hutchison and Kristin Johnson. "Capacity to Trust? Institutional Capacity, Conflict, and Political Trust in Africa, 2000–2005". *Journal of Peace Research* 48.6 (Nov. 1, 2011), pp. 737–752.

[88] Alissandra T Stoyan et al. "Trust in Government Institutions: The Effects of Performance and Participation in the Dominican Republic and Haiti". *International Political Science Review* 37.1 (Jan. 1, 2016), pp. 18–35.

[89] Pui-Hang Wong. "How Can Political Trust Be Built after Civil Wars? Evidence from Post-Conflict Sierra Leone". *Journal of Peace Research* 53.6 (Nov. 1, 2016), pp. 772–785.

[90] Kruk ME, Freedman LP, Anglin GA, Waldman RJ. Rebuilding health systems to improve health and promote statebuilding in post-conflict countries: A theoretical framework and research agenda. *Social Science & Medicine*. 2010 Jan 1;70(1):89–97.

[91] Said Ahmad Maisam Najafizada, Ronald Labonté, and Ivy Lynn Bourgeault. "Community health workers of Afghanistan: a qualitative study of a national program". *Conflict and Health* 8 (2014), p. 26. DOI:10.1186/1752-1505-8-26.

[92] Nathan P. Miller et al. "Assessing community health worker service delivery in humanitarian settings". *Conflict and Health* 14 (2020), p. 73. DOI:10.1186/s13031-020-00325-1.

[93] S. Hemat et al. "A framework for community health worker optimisation in conflict settings: prerequisites and possibilities from Northwest Syria". *BMJ Global Health* 8.8 (2023), e011837. DOI:10.1136/bmjgh-2023-011837.

[94] Ezumah N, Manzano A, Ezenwaka U, Obi U, Ensor T, Etiaba E, et al. Role of trust in sustaining provision and uptake of maternal and child healthcare: Evidence from a national programme in Nigeria. *Social Science & Medicine*. 2022 Jan 1;293:114644.

[95] McKenzie A, Abdulwahab A, Sokpo E, Mecaskey JW. Creating the Foundation for Health System Resilience in Northern Nigeria. *Health Systems & Reform*. 2016 Oct 1;2(4):357–66.

[96] Blanchet K, Nam SL, Ramalingam B, Pozo-Martin F. Governance and Capacity to Manage Resilience of Health Systems: Towards a New Conceptual Framework. *Int J Health Policy Manag*. 2017 Aug 1;6(8):431–5.

[97] Mohsen Abbasi-Kangevari et al. "Gender-sensitive clinics and safe spaces increase maternal care uptake in conflict settings". *Conflict and Health* 14 (2020), p. 50. DOI:10.1186/s13031-020-00324-2.

[98] N. Chowdhary et al. "Mobile gender-sensitive health units: bridging the access gap in crisis". *BMJ Global Health* 7.5 (2022), e008098. DOI:10.1136/bmjgh-2022-008098.

[99] Obinna Ezezika et al. "Community-led approaches to polio immunization in conflict-affected areas: lessons from the DRC". *Vaccine* 40.17 (2022), pp. 2424–2432. DOI:10.1016/j.vaccine. 2022.02.012.

[100] Mohammed Sabahelzain et al. "Community leaders' role in promoting vaccination in conflict zones". *Vaccine* 43 (2025), pp. 1200–1208. DOI:10.1016/j.vaccine.2025.01.005.