A precision health approach to medication management in neurodevelopmental conditions: a model development and validation study using four international cohorts

Marlee M. Vandewouw[1,2,3], Kamran Niroomand[1], Harshit Bokadia[1], Sophia Lenz[1], Jesiqua Rapley[1], Alfredo Arias[4], Jennifer Crosbie[5,6], Elisabetta Trinari[7], Elizabeth Kelley[8], Robert Nicolson[9], Russell J. Schachar[5,6], Paul D. Arnold[10], Alana Iaboni[1], Jason P. Lerch[11,12,13], Melanie Penner[1], Danielle Baribeau[1, 5,6,14,15], Evdokia Anagnostou[1,14,16], Azadeh Kushki[1,17]

1. Autism Research Center, Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Toronto, Canada
2. Center for Precision Psychiatry, Massachusetts General Hospital, Boston, United States of America
3. Department of Psychiatry, Harvard Medical School, Boston, United States of America
4. Province of Ontario Neurodevelopmental (POND) Network, Toronto, Canada
5. Department of Psychiatry, The Hospital for Sick Children, Toronto, Canada
6. Department of Psychiatry, University of Toronto, Toronto, Canada
7. Faculty of Health Sciences, McMaster University, Hamilton, Canada
8. Departments of Psychology and Psychiatry, Queen's University, Kingston, Canada
9. Department of Psychiatry, University of Western Ontario, London, Canada
10. Department of Psychiatry, Cumming School of Medicine, University of Calgary, Calgary, Canada
11. Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom
12. Department of Medical Biophysics, University of Toronto, Toronto, Canada
13. Program in Neurosciences & Mental Health, The Hospital for Sick Children, Toronto, Canada
14. Institute of Medical Sciences, University of Toronto, Toronto, Canada
15. Institute of Health Policy, Management, and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, Canada
16. Department of Paediatrics, University of Toronto, Toronto, Canada
17. Institute of Biomedical Engineering, The University of Toronto, Toronto, Canada

Corresponding author:
Dr. Azadeh Kushki
Bloorview Research Institute
Holland Bloorview Kids Rehabilitation Hospital
150 Kilgour Road Toronto, Ontario M4G 1R8
Tel: 416-425-6220, extension 3563

Keywords: artificial intelligence, medication management, neurodevelopmental conditions, autism

Summary

Background: Psychotropic medications are commonly used for children with

neurodevelopmental conditions, but their effectiveness varies, making prescribing

challenging and potentially exposing individuals to multiple medication trials. We

developed artificial intelligence (AI) models to predict medication success for stimulants,

anti-depressants, and anti-psychotics. We first demonstrate feasibility using cross-

sectional data from three research cohorts, then use a cohort of patients from a

pharmacology clinic to predict medication choice by class, longitudinally, from electronic

medical records (EMRs).

Methods: Models were built to predict cross-sectional medication usage from the Child

Behaviour Checklist. Data from the Province of Ontario Neurodevelopmental (POND)

network (*N*=598) trained and tested the models, while data from the Healthy Brain

Network (HBN; *N*=1,764) and Adolescent Brain Cognitive Development (ABCD;

N=2,396) studies were used for external validation. For the EMR cohort, data from the

Psychopharmacology Program (PPP; *N*=312) at Holland Bloorview Kids Rehabilitation

Hospital were used to predict longitudinal success. Stacked ensemble models were built

separately for each medication class, and area under the receiving operating

characteristic curve (AU-ROC) evaluated performance.

Findings: The research cohorts demonstrated feasibility, with internal testing (POND)

achieving an AU-ROC (median [IQR]) of 0.75 [0.73,0.80] for stimulants, 0.83 [0.78,0.87]

for anti-depressants, and 0.79 [0.72,0.86] for anti-psychotics. Performance in external

testing sets (HBN and ABCD) confirmed generalizability. In the EMR cohort (PPP), AU-

ROCs were high: 0.87 [0.83,0.91] for anti-psychotics, 0.84 [0.81,0.88] for stimulants and

0.82 [0.77,0.87] for anti-depressants.

Interpretation: This study demonstrates the feasibility of using AI to enhance

medication management for children with neurodevelopmental conditions, with expert

clinician decisions learned with high accuracy. These findings support the potential for

AI decision aids in community settings, promoting faster access to personalized care

while highlighting the complexity of clinical and sociodemographic factors influencing

medication decisions.

Funding: Funding was provided by the Canadian Institutes of Health Research

(Operating Grant #527447) and Ontario Brain Institute.

Research in context

Evidence before this study: Current medication management practices for neurodevelopmental conditions, particularly for children who have not responded to first-line options, are based clinical best guess approaches that can have negative effects on children, their caregivers, and the health system. Precision health tools using artificial intelligence, which are suitable for community use, have the potential to improve the health system's capacity for providing timely and impactful care. We searched PubMed on October 17, 2024, for studies published in English evaluating artificial intelligence approaches for medication management in neurodevelopmental conditions using the terms ("autism" OR "neurodevelopmental condition" OR "neurodevelopmental disorder" OR "neurodivergence") AND ("artificial intelligence" OR "machine learning" OR "deep learning" OR "prediction") AND ("medication management" OR "medication response" OR "medication appropriateness" OR "medication decision support") also including terms where "drug" is substituted for "medication". We did not find any relevant studies.

Added value of this study: This study demonstrates the feasibility of using AI to assist in medication management for children with neurodevelopmental conditions, with a strong ability to learn expert clinical decisions. These findings show AI may be able to support access to faster, more personalized treatment decisions regarding psychotropic medications. We also identified the relevant clinical and demographic features to the model's medication recommendations, as well as several biases with respect to sociodemographic factors, highlighting the complexity of factors that contribute to clinical decision-making.

Implications of all the available evidence: This work highlights AI's potential to improve medication management for children with neurodevelopmental conditions, by offering personalized treatment recommendations. However, identified biases underscore the need for addressing existing inequities. Future research should focus on prospective validation, integration into clinical practice, and bias mitigation to ensure equitable access to effective treatments for all children.

Introduction

Neurodevelopmental conditions, such as autism spectrum disorder (autism),

intellectual disability (ID), and attention-deficit/hyperactivity disorder (ADHD), are highly

heterogeneous and overlapping in their biology and phenotypic presentation[1,2], which

includes hundreds of associated genes[3], and significant variability in neurobiological

correlates[2]. Despite decades of research, no markers exist to indicate which biological

pathway may be implicated for an individual child to guide individualized care[4,5], and

current approaches to care provision remain reliant on broad diagnostic labels. This lack

of precision is particularly challenging for pharmacological interventions in

neurodevelopmental conditions. These interventions are highly prevalent with greater

than 50% of children with neurodevelopmental conditions prescribed psychotropic

medications to address behavioural and/or mental health concerns (e.g., anxiety,

hyperactivity, aggressive or self-injurious behavior), and high rates of polypharmacy

during childhood (~30%)[6].

Despite the prevalent use of psychotropic medications in neurodevelopmental

conditions, there is tremendous variability in individual-level responses to psychotropic

treatment for co-occurring mental health and behavioural condition[7], reflecting the

heterogeneity of biological pathways impacted, diverse clinical presentations, as well as

sociodemographic complexities that influence the acceptability and uptake of

medications. This variability poses a significant challenge in determining the appropriate

medication for an individual child, especially in cases where co-occurring conditions

(e.g., autism, ADHD, anxiety) can interact and increase the complexity of presentation

and response. In these cases, prescribing practices in the children and adolescents

remain almost entirely off-label[6,8–12] and based on trial-and-error approaches that can

expose children to multiple medication trials with the potential for severe and life-long

adverse effects for some classes of medication[13]. This highlights the critical need for

precision health tools that can provide medication recommendations based on an

individual's unique presentation.

To fill this gap, we propose an artificial intelligence (AI) model that uses an

individual's characteristics to predict the medication choice among the commonly

prescribed classes in neurodevelopmental conditions: stimulants, anti-depressants, and

anti-psychotics. We first establish the feasibility of this approach by building models to

predict medication usage using a research cohort of children and youth with

neurodevelopmental conditions, using two international research cohorts for external

validation. Then, to examine how this approaches generalizes to real-world clinical

settings, we leveraged a patient cohort to predict continuation of a medication class

after 3 months from electronic medical record (EMR) data of children and youth with

neurodevelopmental conditions who visited a specialized psychopharmacology clinic.

Methods

Study design and cohorts

For the research cohorts, training and internal testing was conducted using data

from 598 children and youth with neurodevelopmental conditions extracted from the

Province of Ontario Neurodevelopmental (POND) network (Ontario, Canada). External

testing was performed using two international cohorts: 1,764 children and youth with

neurodevelopmental conditions from the Healthy Brain Network (HBN[14]; New York,

United States), and 2,396 children and youth with neurodevelopmental conditions from

the Adolescent Brain Cognitive Development study (ABCD[15]; United States).

For clinical validation, longitudinal EMRs from the Psychopharmacology Program

(PPP) at Holland Bloorview Kids Rehabilitation Hospital (Toronto, Ontario, Canada)

were used. PPP sees children with neurodevelopmental conditions who have tried at

least one psychoactive medication without adequate success. Data were extracted

from electronic medical records of 312 children who visited the PPP over a three-year

period (2019 – 2022) and returned for a follow-up approximately 3 months later.

Participant demographics are summarized in

Table 1. Further details on the cohorts, including eligibility and consent

procedures, and a comparison of the research cohorts can be found in the

Supplementary Material. This study was approved by the Research Ethics Board at

Holland Bloorview Kids Rehabilitation Hospital.

Table 1. Cohort demographics and medication outcomes.

		Research cohorts			EMR cohort
		POND	**HBN**	**ABCD**	**PPP**
N		598	1,764	2,396	312
Age range (years)		5 – 19	5 – 17	8 – 11	4 – 17
Mean age [SD] (years)		11.4 [3.2]	10.1 [2.7]	9.9 [0.6]	10.4 [2.9]
Sex [%]	Male	424 [71]	1,213 [69]	1,523 [64]	262 [84]
	Female	174 [29]	551 [31]	873 [36]	50 [16]
Autism [%]		338 [57]	325 [18]	0 [0]	292 [94]
ADHD [%]		193 [32]	1,381 [78]	1,751 [73]	181 [58]
OCD [%]		67 [11]	90 [5]	1,048 [44]	16 [5]
Intellectual disability [%]	Yes	84 [14]	86 [5]	149 [6]	105 [34]
	No	514 [86]	1,678 [95]	2,247 [94]	135 [43]
	Unknown	0 [0]	0 [0]	0 [0]	72 [23]
Race/ ethnicity [%]	White	327 [55]	871 [49]	1,257 [52]	NA
	Minoritized	156 [26]	866 [49]	1,137 [47]	NA
	Not collected	115 [19]	27 [2]	2 [1]	312 [100]
Annual household income [%]	Low	86 [14]	307 [17]	770 [32]	NA
	Medium	98 [16]	345 [20]	608 [25]	NA
	High	170 [28]	786 [45]	798 [33]	NA
	Not collected	244 [41]	326 [18]	220 [9]	312 [100]
Outcome [%]	Stimulant	171 [29]	177 [10]	551 [23]	114 [37]
	Anti-depressant	96 [16]	72 [4]	142 [6]	82 [26]
	Anti-psychotic	52 [9]	26 [1]	43 [2]	123 [39]

EMR: electronic medical record; PPP: Psychopharmacology Program; POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study; SD: standard deviation; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder. For POND, primary diagnosis is reported; for ABCD, HBN, and PPP, refer to supplemental.

Predictors and outcome

For the PPP cohort, the set of predictors were co-created with the clinicians at the PPP program to reflect features relevant to the medication classes considered, and the availability of variables in the electronic medical records at the first visit. These predictors were broadly grouped into five categories: demographics, clinical, family-related, child-related, and care-related. The predicted outcome was the medication decision at the follow-up visit (

Table 1).

For the research cohorts, demographics (age and sex), neurodevelopmental

diagnoses (autism, ADHD, and OCD), IQ as a continuous score, and data from the

Child Behaviour Checklist[16] were used as predictors. The predicted outcome was the

medication class the child was on at the time of study (

Table 1). A full description of the predictors and outcome are presented in the
Supplementary Material.

Model

AI models were built to predict the outcome from the set of predictors, separately
for each medication class (Figure 1). For the research cohorts, the models were trained
and internally tested using the POND dataset, while HBN and ABCD were withheld for
external testing. By validating with two external datasets, we assessed the
generalizability and robustness of the models, ensuring results are not specific to a
single dataset and can be replicated across diverse populations. For the EMR cohort
(PPP), only internal testing was performed.

Figure 1. Study design.

To address class imbalance, we applied random under-sampling of the majority
class to achieve approximately balanced 1:1 class ratios. This algorithm-agnostic
approach reduces majority-class dominance and strengthens the minority-class learning
signal without relying on synthetic data or class weighting. To limit dependence on any
single balanced draw, we repeated the under-sampling for 10 independent iterations,
noting that findings were stable across 5, 10, and 20 iterations.

For the AI models, we implemented a Super Learner approach[17], combining

CatBoost gradient boosting[18] and deep neural networks as base learners, with logistic

regression meta-learner trained on out-of-fold predictions to generate final predictions.

We used nested cross-validation with inner 3-fold loops for hyperparameter optimization

using Optuna[19] and outer 5-fold loops where base learners generated out-of-fold

predictions for the logistic regression meta-learner. Further details are provided in the

Supplementary material.

Performance evaluation

Model performance was assessed using the area under the receiving operating

characteristic curve (AU-ROC), reporting medians and interquartile ranges (IQR) across

folds and under-sampling iterations. Additional metrics (area under the precision-recall

curve (AU-PR), sensitivity, and specificity) are provided in the Supplementary Material.

To interpret which features were most important for predicting longitudinal medication

success, Shapley Additive exPlanations (SHAP) values[20] were computed for the EMR

(PPP) models. SHAP values were computed for each of the base models and combined

by taking their weighted average using the coefficients from the logistic regression

meta-learner.

The demographic parity and equalized odds ratios[21,22] were used to evaluate

whether the model predictions were independent of sensitive group status for sex,

intellectual disability (ID), race/ethnicity, and annual household income. The metric was

computed at the default probability threshold (0.50) across folds, under-sampling

iterations, and datasets for the research cohorts; the 80% rule was used to determine

any biases[23].

Role of the funding source

The funders of this study had no role in the collection, analysis, and interpretation

or data, in the writing of the report, nor in the decision to submit the report for

publication.

Results

Model performance

The research cohorts demonstrated the feasibility of AI predicting treatment

status cross-sectionally for each medication class (Table 2 and Figure 2), with internal

testing (POND) demonstrating good-to-excellent AU-ROC that was significantly higher

than chance for stimulants (median [IQR]: 0.75 [0.73, 0.80]), anti-depressants (0.83

[0.78, 0.87]), and anti-psychotics (0.79 [0.72, 0.86]). The performance of the external

testing sets (HBN and ABCD) indicated good generalizability, with the AU-ROCs slightly

reduced but still in the good-to-excellent range.

Table 2. Model performance.

| | Research cohorts | | | EMR cohort |
| | Internal testing | External testing | | Internal testing |
Medan AU-ROC [IQR]	**POND**	**HBN**	**ABCD**	**PPP**
Stimulant	.75 [.73, .80]	.70 [.67, .71]	.70 [.70, .71]	.84 [.81, .88]
Anti-depressant	.83 [.78, .87]	.75 [.74, .77]	.67 [.65, .69]	.82 [.77, .87]
Anti-psychotic	.79 [.72, .86]	.79 [.75, .81]	.81 [.79, .83]	.87 [.83, .91]

EMR: electronic medical record; PPP: Psychopharmacology Program; POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study; CI: confidence interval; AU-ROC: area under the receiving operating characteristic curve; IQR: interquartile range

Figure 2. ROC curves for the research cohorts, for the internal testing (A; POND) and external testing (B; HBN and ABCD) sets for each of the stimulant classes (green: stimulants, purple: anti-depressants, pink: anti-psychotics); AU-ROCs are also presented.

POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study; ROC: receiving operating characteristic curve; AU-ROC: area under the ROC

184   For the EMR cohort (PPP; Table 2 and Figure 3A), the AU-ROC across the

185   cross-validation folds was excellent for anti-psychotics (0.87 [0.83, 0.91]), stimulants

186   (0.84 [0.81, 0.88]) and anti-depressants (0.82 [0.77, 0.87]).

EMR cohort

Figure 3. (A) ROC curves for the EMR cohort (PPP) for each of the stimulant classes (green: stimulants, purple: anti-depressants, pink: anti-psychotics); AU-ROCs are also presented. (B) Feature importance for the EMR cohort (PPP) assessed using SHAP values. The SHAP values are plotted for each participant for the 10 most important features for each medication class, sorted by importance (mean absolute SHAP value) from top to bottom. Colours represent the feature value (red for high values, blue for low values), indicating whether higher or lower values contribute positively or negatively to prediction scores.
EMR: electronic medical record; PPP: Psychopharmacology Program; ROC: receiving operating characteristic curve; AU-ROC: area under the ROC; SHAP: Shapley Additive exPlanations; ADHD: attention-deficit/hyperactivity disorder

Feature importance

SHAP values were computed for the EMR cohort (PPP), to identify which features were most important for predicting medication choice (Figure 3B). Prior use of a medication class was the most important feature across all four classes. In particular, prior use of a medication within the same class, increased the likelihood of a medication choice in that class. Age was also a salient feature: younger, middle, and older participants were more likely to be recommended stimulants, anti-psychotics, and anti-depressants, respectively. Other important features included ADHD and communication abilities for stimulants, anxiety and aggression for anti-depressants, and intellectual disability and prior prescription of stimulants for anti-psychotics.

Bias

The demographic parity and equalized odds ratios were computed to evaluate

the bias of the models (Table 3), where values exceeding 0.80 indicate bias according

to the four-fifths rule[24]. For the research cohorts, parity was mixed. Individuals who were

female, and who did not have an intellectual disability (ID) were more likely to be

recommended a stimulant (male, female: 39%, 56%; ID, no-ID: 37%, 52%). Individuals

who were male, did not have an ID, who were white, and who were of high income class

were more likely to be recommended an anti-depressant (male, female: 40%, 27%; ID,

no-ID: 23%, 32%; white, minority: 38%, 22%; low, middle, high income: 31%, 25%,

35%). On the other hand, those with an ID, and those with a lower annual household

income were more likely to be recommended anti-psychotics (ID, no-ID: 54%, 31%; low,

middle, high income: 45%, 22%, 30%). These observations extended to the EMR

cohort, with the exception of sex, with males being more likely to be recommended

stimulants (male, female: 45%, 36%) and females being more likely to be recommended

anti-depressants (male, female: 44%, 55%). This was in line with trends in the data.

Table 3. Demographic parity ratio (DPR) and equalized odds ratio (EOR) for sensitive groups.

		Research cohorts		EMR cohort	
		DPR	EOR	DPR	EOR
Sex	Stimulant	0.70	0.70	0.79	0.49
	Anti-depressant	0.69	0.68	0.79	0.72
	Anti-psychotic	0.81	0.83	0.91	0.81
Intellectual disability	Stimulant	0.72	0.63	0.45	0.29
	Anti-depressant	0.71	0.65	0.62	0.41
	Anti-psychotic	0.57	0.74	0.62	0.59
Race/ethnicity	Stimulant	0.95	0.94	–	
	Anti-depressant	0.57	0.71	–	
	Anti-psychotic	0.83	0.90	–	
Annual household income	Stimulant	0.94	0.85	–	
	Anti-depressant	0.72	0.77	–	
	Anti-psychotic	0.50	0.56	–	

EMR: electronic medical record

212 **Discussion**

213 Medication management for children with neurodevelopmental conditions remains a

214 significant challenge, particularly in community settings where subspecialty expertise

215 may not be available. To address this gap, this study aimed to develop and evaluate an

216 AI algorithm to predict medication choices in neurodevelopmental conditions. Ultimately,

217 this type of tool could be used as a decision-aid to support medication management

218 where specialized expertise is not available.

219 We showed that AI can learn expert clinician medication decisions with high

220 accuracy. We also demonstrated that the results generalize across independent,

221 international datasets, which is notable given the drastically different characteristics of

222 these cohorts, in terms of demographics, diagnosis, phenotype, and region-specific

223 access to service and supports (see Supplemental Table 1 and Supplemental Table 2).

224 While the accuracy in these research cohorts is lower compared to the EMR model, this

225 is likely due to the cross-sectional nature of the prediction in these cohorts. For

226 example, use of stimulant medications to target ADHD behaviours confounds the

227 measures of ADHD traits when considered cross-sectionally, in turn impacting

228 predictive performance.

229 Overall, these results support the feasibility of using an AI-based decision aid tool.

230 This can be particularly advantageous for community practitioners, where subspecialty

231 expertise required for medication management is often not available. Ultimately, an AI-

232 based decision aid can bring precision to current prescribing practices in the community

233 which may expose children to multiple medication trials that may have severe and life-

234 long adverse effects[25]. Prospective trials are needed to further validate this type of

235 algorithms in clinical settings.

236 Explainability and interpretability are critical features of responsible AI systems,

237 contributing to enhancing trust and transparency with both clinicians and patients[26]. To

238 promote explainability, we characterized the relative contributions of clinical and

239 demographic features to the AI algorithm's medication recommendations. For the EMR

240 cohort, these contributions were closely aligned with indications for each medication

241 class (e.g., stimulants for ADHD, SSRIs for anxiety). Interestingly, the top contributors

for all classes were prior use of a medication in that class. This finding reflects the fact

that medication choices often remain within the same class of medication but entail

changes in the medication type and dosing within that class (e.g., amphetamines,

methylphenidate within the stimulant class). Future work will focus on adding this type of

resolution to the AI predictions. Other contributing features to medication decision

reflected sociodemographic and environmental factors that may contribute to clinical

decision making beyond primary indications. For example, school environment and

funding eligibility were significant contributors to the decision to recommend an

antipsychotic. These finding highlight the complexity of the clinical and

sociodemographic factors that contribute to value-laden decisions around medication

management for children with neurodevelopmental conditions.

Another aspect of responsible AI is potential algorithmic biases that may

disproportionately disadvantage subgroups of individuals. Our analysis revealed several

biases including increased recommendation of antipsychotics to children who had

intellectual disability and were from lower income households. On the other hand, anti-

depressants recommendations were increased for children who did not have an

intellectual disability, and who were white. It is important to note that these biases are

not related to any specific feature of the AI algorithm, but mirror previously reported

biases in prescription patterns which exist across the system[27]. For example, decreased

access to early diagnosis and behavioural interventions for children from lower income

household may contribute to a higher likelihood of crisis situations that necessitate

prescription of antipsychotic medications. Similarly, it is well established that depression

can be challenging to detect in individuals with an intellectual disability[28]. While

computational approaches to bias mitigation can be applied to equalize predicted

prescription rates across groups, these will not solve the underlying system-level

inequities that impact children with marginalized identities.

The results of this study should be interpreted considering several limitations. Firstly,

for the longitudinal prediction, we defined outcome as the medication choice at the

follow-up visit. Treatment success in neurodevelopmental conditions is a complex,

subjective, and value-laden construct. The use of medication choice was co-created

with clinicians as an outcome as it reflects the aggregate effects of the multitude of

factors contributing to a clinical decision to not stop a medication (e.g., ineffectiveness,

side-effects, acceptability or other factors). These children also visited the specialized

clinic after at least one psychoactive medication was tried without success, and thus this

may introduce biases when the models are used to make a first-line medication

decision. Although gender identity and race/ethnicity were not included as predictors

due to incomplete capture in the primary EMR cohort, their omission may contribute to

model misspecification bias. These variables remain important contextual factors for

understanding medication access and prescribing patterns. Future work should

incorporate reliably captured measures and evaluate model performance across gender

and racial/ethnic groups, alongside fairness assessments, updating the model as

needed to mitigate potential inequities. The models were trained using data exclusively

from North America; future work should investigate their translation to other global

populations to account for potential differences in healthcare systems, prescribing

practices, and sociocultural influences on medication usage. Finally, this study

exclusively focused on predicting medication classes; these classes are broad, and

future work should examine within-class predictability.

In summary, the present study focused on AI model development and evaluation,

demonstrating the feasibility of this approach for medication recommendation for

children with neurodevelopmental conditions. To enable effective clinical

implementation of this type of AI-tool, future studies are needed to evaluate the

algorithm in a prospective design at the point of care as well as acceptability, and

barriers and facilitators to uptake.

Declaration of interests

EA has received research funding from Roche and Anavex, consulting fees from Roche,

Quadrant Therapeutics, Ono, and Impel Pharmaceuticals, in-kind support from AMO

Pharma and CRA-Simons Foundation, royalties from APPI and Springer, an editorial

honorarium from Wiley, and has a patent for holly^{TM}. DB has received research funding

from MapLight Therapeutics. MP has received consulting fee from Addis &

Associates/Roche, and MedCounsel. AK has a patent for holly^{TM} with royalties paid

from Awake Labs and has received consulting fees from DNAStack and Shaftesbury.

The remaining authors have no conflicts to report.

Acknowledgements

Funding was provided by the Canadian Institutes of Health Research (Operating Grant

#527447). Acknowledgements for each of the research cohorts (POND, HBN, and

ABCD) used in this study are provided below.

POND: This research was conducted with the support of the Ontario Brain Institute

(POND, PIs: Dr. Evdokia Anagnostou and Dr. Jason P. Lerch), an independent non-

profit corporation, funded partially by the Ontario government. The opinions, results and

conclusions are those of the authors and no endorsement by the Ontario Brain Institute

is intended or should be inferred.

HBN: This manuscript was prepared using a limited access dataset obtained from the

Child Mind Institute Biobank, the Healthy Brain Network (HBN). This manuscript reflects

the views of the authors and does not necessarily reflect the opinions or views of the

Child Mind Institute.

ABCD: Data used in the preparation of this article were obtained from the Adolescent

Brain Cognitive DevelopmentSM (ABCD) Study (https://abcdstudy.org), held in the

NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit

more than 10,000 children age 9-10 and follow them over 10 years into early adulthood.

The ABCD Study® is supported by the National Institutes of Health and additional

federal partners under award numbers U01DA041048, U01DA050989, U01DA051016,

U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174,

U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988,

U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038,

Data sharing statement

Data from four cohorts were used in this study, each governed by their own data

sharing restrictions; full details can be found in the Supplementary Material. Code used

in this manuscript is available at GitHub (https://github.com/marlvan/med_prediction).

Supplementary material

PPP

Dataset description

The clinical cohort used for this study was longitudinal electronic medical records

from the Psychopharmacology Program (PPP) at Holland Bloorview Kids Rehabilitation

Hospital (Toronto, Canada). The PPP is a program with provincial catchment in Ontario

that provides medication management for children and youth with neurodevelopmental

conditions. Children were eligible for this study if (1) their first visit was between 2019-

2022, (2) they were at least four years of age at their first visit, and (3) they returned to

the clinic for a follow-up visit. As the data was extracted from electronic medical records,

no consent was obtained. Data from PPP is not publicly available due to ethics

restrictions. Clinical data from all longitudinal clinical visits were extracted into a

RedCap database from semi-structured medical record templates by trained chart

reviewers. Two chart reviewers co-reviewed each medical record in sets of five and

reviewed discrepancies until 85% agreement on all items was achieved. Reviewers

proceeded independently after that, and any uncertainties were discussed with the

study team.

Predictors

For PPP, the predictors were grouped into five categories: demographics,

clinical, family-related, child-related, and care-related. Categorical variables with

unknown or undocumented values were dummy-coded with 'Unknown' as the implicit

reference category. This retains all participants and prevents missing data from being

conflated with a substantive level. Missingness ranged between 0 and 28% across

predictors, and 65% of participants had at least one missing predictor.

Demographic variables included age, sex (male or female), and region (urban

(Toronto, General Toronto Area, or other Ontario city) or rural (rural or northern

community)). Due to missing data, gender and race/ethnicity could not included as

predictors. To capture socioeconomic status, whether a parent was receiving social

assistance or means-tested disability support as a source of income was included as a

predictor.

Clinical variables included neurodevelopmental diagnoses (attention-

deficit/hyperactivity disorder (ADHD), autism, obsessive-compulsive disorder (OCD),

intellectual disability, learning disorder, other psychiatric condition), mental health

symptoms (anxiety, depression, irritability), and chronic conditions (epilepsy, cerebral

palsy, asthma, migraines, gastroesophageal reflux disease, persisting constipation,

obesity, and other).

Family-related factors included a family history of psychiatric or developmental

diagnoses in a first degree relative, and family reports of conflict between family

members in the household, caregiver burnout, and financial barriers to care, as reported

by families and documented by the treating clinician in the child's chart.

Child-related factors included school type (public, mainstream classroom, private,

mainstream classroom, public, specialized classroom, private, specialized classroom,

homeschool, or other), school attendance (full days, part days or withdrawal, frequent

calls home for pick-up, frequent school refusal or frequent absences, not attending or

demitted from school, or other), communication abilities (speaks fluidly in sentences,

speaks a few words or short phrases, non-speaking and uses communication

technique, non-speaking and no other formal communication form, and other). History

of trauma (children's aid involvement, witnessing violence, experiencing violence,

physical abuse, sexual abuse, or neglect, other significant stressor as perceived by the

child, or other stressor), police involvement, hospitalization, emergency department

visits, intensive crisis service involvement due to behavioural or mental health concerns,

aggression, self-injury, and suicidal thoughts were also included.

Care-related factors included receiving government funding for care (specifically,

the Ontario Assistance for Children with Severe Disabilities Program), which was coded

as "yes", "has not applied", "not eligible", or "other". Data on current, past, and

recommended supports were collected using the following categories: (1) behavioural

therapy, Applied Behaviour Analysis (ABA), or parent coaching by a behaviour

therapist, (2) child therapy (e.g., Cognitive Behavioural Therapy (CBT) or group

therapy), (3) case management, (4) allied health, (5) respite, (6) other, or (7) none.

Prior use of five commonly prescribed medication classes (stimulants, non-stimulants,

hypnotics, anti-depressants, and anti-psychotics) was determined as whether a child

was taking a medication at the time of the first visit to the clinic. The behaviour that was

targeted for treatment at the first visit was collected using the following categories:

physical aggression, swearing/verbal threats, elopement, self-injury, verbal refusal,

repetitive behaviours, property destruction, tantrums/meltdowns, and/or other. Data on

side effects reported at the first visit were categorized as: headache, stomach-ache,

constipation, nausea/vomiting, weight gain, weight loss, irritability,

drowsy/fatigue/sedation, insomnia, appetite decrease, appetite increase, other, and/or

none. Whether adherence problems for medications were discussed at the first visit was

also included as a predictor.

Outcome

The predicted outcome was the medication class the child was on at the follow-

up visit.

Bias

Bias was assessed on sex (male versus female) and intellectual disability (yes

versus no).

POND

Dataset description

The POND network is a research network spanning several institutions across

Ontario, Canada that collects demographic, phenotypic, and biological data from

children and youth with and without neurodevelopmental conditions. Participants were

eligible for the current study if they received a diagnosis of autism, ADHD, or OCD, if

they were between 5 and 19 years of age, and if diagnostic, medication usage, IQ, and

the Child Behaviour Checklist (CBCL[16]) was available. Children and youth provided

written informed consent if they were able; otherwise, they provided verbal assent, and

their caregiver provided written consent. Data from POND is managed by the Ontario

Brain Institute and is available through a controlled data release. All requests for access

should be directed to the Brain-CODE platform

(https://www.braincode.ca/content/controlled-data-releases#dr006).

Predictors

Predictors for POND included demographics (age and sex), diagnosis (autism,

ADHD, or OCD), IQ, and the CBCL. Diagnoses were confirmed with expert clinical

judgement alongside the Autism Diagnostic Observation Schedule, 2 (ADOS-2[29]) and

Autism Diagnostic Interview, Revised (ADI-R[30]) for autism, the Parent Interview for Child

Symptoms (PI/orCS[31]) and/or the Kiddie Schedule for Affective Disorders and

Schizophrenia (K-SADS[32]) for ADHD, and the Children's Yale-Brown Obsessive-

Compulsive Scale (YBOCS[33]) for OCD. Full-scale IQ was collected using age-

appropriate Wechsler[34–38] or Standford-Binet[39] scales. All the individual questions,

syndromic scale *T*-scores (anxious/depressed, withdrawn/depressed, somatic

complaints, social problems, thought problems, attention problems, rule-breaking

behaviour, and aggressive behaviour), and summary *T*-scores (internalizing problems,

externalizing problems, and total problems) from the CBCL were included as predictors.

Outcome

Caregivers were asked to provide a list of medication their child was taking.

These medications were classified as stimulants, non-stimulants, anti-depressants, and

anti-psychotics with assistance from clinicians. The predicted outcome was the

medication class(es) the child was on.

Bias

Bias was assessed for sex, intellectual disability, race/ethnicity, and

socioeconomic status. For sex, biological sex (male versus female) was used.

Intellectual disability (yes versus no) was determined using the full-scale IQ cutoff of 70.

For race/ethnicity, participants/caregivers were asked to select all groups that applied:

Aboriginal, Arab, Black, Chinese, East Asian, Filipino, Japanese, Korean, Latin

American/Hispanic, South Asian, Southeast Asian, West Asian, White, other; this data

was used to categorize individuals as white or minoritized. For socioeconomic status,

we used annual household income categorized as low (<$50,000), middle ($50,000-

$99,999), and high (≥$100,000).

HBN

Dataset description

HBN is a mental health study of children and adolescents across the New York

city (United States) area that has collected demographic, phenotypic, and biological

data; families who had psychiatric concerns about their child were recruited for the

study. Participants were eligible for the current study if they received a diagnosis of

autism, ADHD, or OCD, if they were between 5 and 19 years of age, and if diagnostic,

medication usage, IQ, and the CBCL[16] was available. Participants at least 18 years of

age provided written informed consent; otherwise, they provided written assent, and

their caregiver provided written consent. Access to the data used in this study is

protected by a Data Usage Agreement (DUA; see

https://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network for more details),

which was awarded to AK for this study.

Predictors

Aligning with POND, predictors for HBN included demographics (age and sex),

diagnosis (autism, ADHD, and/or OCD), full-scale IQ, and the CBCL. For diagnosis, the

consensus DSM-5[40] diagnoses provided by the HBN clinical team were used. Full-scale

IQ was assessed using Wechsler scales[37,38] or the Kaufman Brief Intelligence Test[41].

Outcome

Caregivers were asked to provide a list of medication their child had taken the

day of the visit. These medications were classified as stimulants, non-stimulants, anti-

depressants, and anti-psychotics with assistance from clinicians. The predicted

outcome was the medication class(es) the child was on.

Bias

Bias was assessed on sex, intellectual disability, race/ethnicity, and

socioeconomic status. For sex, biological sex (male versus female) was used.

Intellectual disability (yes versus no) was determined using the full-scale IQ cutoff of 70.

For race/ethnicity, participants/caregivers indicated their ethnicity (not Hispanic or

Latino, Hispanic or Latino, decline to specify, or unknown) and race (White/Caucasian,

Black/African American, Hispanic, Asian, Indian, Native American Indian, American

Indian/Alaskan Native, Native Hawaiian/Other Pacific Islander, two or more races, or

other race), which were used to categorize individuals as white or minoritized. For

socioeconomic status, we used annual household income categorized as low

(<$50,000), middle ($50,000-$99,999), and high (≥$100,000).

ABCD

Dataset description

The ABCD study[42] is a longitudinal study of adolescents from the general

population across the United States that has collected demographic, phenotypic, and

biological data. For the purposes of this study, only the first time point (collected at

approximately 9-11 years of age) was used. Participants were eligible for the current

study if medication usage, IQ, the Kiddie Schedule for Affective Disorders and

Schizophrenia (K-SADS), and the CBCL was collected. Caregivers provided written

informed consent, and participants provided written assent. Access to the data used in

this study is protected by an NIMH Data Archive DUA for the Adolescent Brain

Cognitive Development (ABCD) permission group (see

https://nda.nih.gov/user/dashboard/data_permissions.html for more details), which was

awarded under DUA #15619 for this study.

Predictors

Aligning with POND, predictors for ABCD included demographics (age and sex),

diagnosis (autism, ADHD, and/or OCD), IQ, and the CBCL. Diagnoses were assigned

using a computerized version of the K-SADS[43] completed by parents and/or youth. IQ

was assessed using the NIH Toolbox[44] Cognition Battery's age-corrected total cognition

composite score, which shows a high degree of alignment with the Wechsler scales

used in POND[45].

Outcome

Caregivers were asked to provide a list of medication their child had taken in the

two weeks preceding the visit. These medications were classified as stimulants, non-

stimulants, anti-depressants, and anti-psychotics with assistance from clinicians. The

predicted outcome was the medication class(es) the child was on.

Bias

Bias will be assessed on sex, intellectual disability, race/ethnicity, and

socioeconomic status. For sex, biological sex (male versus female) was used.

Intellectual disability (yes versus no) was determined using the IQ cutoff of 70. For

race/ethnicity, participants/caregivers indicated their race/ethnicity (Hispanic/Latino,

non-Hispanic Asian, non-Hispanic Black, non-Hispanic White, and Multi-Racial/Other),

which were used to categorize individuals as white or minoritized. For socioeconomic

status, we used annual household income categorized as low (<$50,000), middle

($50,000-$99,999), and high (≥$100,000).

AI models

We implemented a Super Learner approach[17] in Python (version 3.12),

combining CatBoost gradient boosting[18] (version 1.2.7) and deep neural networks

(Keras[46] (version 3.7.0) with TensorFlow[47] (version 2.18.0)) as base learners, with a

logistic regression model trained on out-of-fold predictions to generate final predictions.

Hyperparameters for both base learners were optimized using Optuna's Tree-

structured Parzen Estimator[19] with 100 trials and nested 3-fold cross-validation within

each training fold, with both models optimizing AU-ROC. The deep neural network

architecture was tuned across 1-3 hidden layers, 48-96 units per layer, and dropout

rates of 0.20-0.40. Networks used ReLU activation and the Adam optimizer, with

learning rates tuned between 3×10^{-4} to 3×10^{-3}. Training used a fixed batch size of 32

with early stopping (patience=12) monitoring AU-ROC over a maximum of 200 epochs.

CatBoost models were configured with 50-400 iterations, learning rates of 0.01-0.30,

tree depths of 3-6, and early stopping after 200 rounds without improvement.

The Super Learner consisted of a logistic regression model (L2 penalty = 1.0,

liblinear solver) to combine predicted probabilities from both base learners. We

employed nested cross-validation, with inner 3-fold loops for hyperparameter

optimization and outer 5-fold loops in which base learners trained on 4 folds generated

out-of-fold predictions for the logistic regression model. This design followed standard

Super Learner methodology to prevent overfitting, avoid data leakage, and yield

unbiased performance estimates.

Comparison of research cohorts

The demographic variables (age, sex, diagnosis, intellectual disability,

race/ethnicity, and socioeconomic status) were compared amongst the research cohorts

(Supplemental Table 1). Chi-squared tests were used for categorical variables, holding

significance at *p*<.050 and reporting Cramer's *v* effect sizes. Shapiro-Wilks tests

indicated that all continuous variables were non-normally distributed; as such, Kruskal-

Wallis tests were used, holding significance at *p*<.050, and reporting eta-squared effect

sizes.

Supplemental Table 1. Demographic comparison of the research cohorts.

Measure	p-value	Effect size	Post-hocs
Age	<.0001	.05	ABCD > POND > HBN
Sex	.0001	.06	Females: ABCD > POND, HBN
Autism	<.0001	.53	POND > HBN > ABCD
ADHD	<.0001	.32	HBN > ABCD > POND
OCD	<.0001	.43	ABCD > POND > HBN
Intellectual disability	<.0001	.11	POND > HBN, ABCD
Race/ethnicity	<.0001	.10	White: POND > HBN, ABCD
Annual household income	<.0001	.13	HBN > POND > ABCD

POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder

554   The CBCL summary *T*-scores were also compared amongst the datasets

555   (Supplemental Table 33), using Kruskal-Wallis tests due to non-normality of all variables

556   (Shapiro-Wilks tests), holding significance at $p<.050$, and reporting eta-squared effect

557   sizes.

Supplemental Table 2. CBCL comparison of the research cohorts.

Measure	Median [IQR] ABCD	HBN	POND	p-value	Effect size	Post-hocs
Anxious/depressed	54 [50, 63]	55 [50, 64]	62 [53, 68]	<.0001	.03	POND > HBN, ABCD
Withdrawn/depressed	54 [60, 52]	56 [50, 64]	60 [54, 68]	<.0001	.04	POND > HBN > ABCD
Somatic complaints	57 [50, 64]	56 [50, 61]	57 [53, 64]	.0004	<.01	ABCD > POND, POND > HBN
Social problems	54 [51, 60]	58 [51, 64]	61 [54, 68]	<.0001	.06	POND > HBN, ABCD
Thought problems	58 [51, 64]	58 [51, 67]	66 [58, 71]	<.0001	.05	POND > HBN, ABCD
Attention problems	59 [53, 60]	64 [57, 69]	65 [59, 71]	<.0001	.05	POND > HBN > ABCD
Rule-Breaking behaviour	53 [50, 60]	53 [51, 60]	54 [51, 62]	.0008	<.01	ABCD > HBN, POND > ABCD
Aggressive behaviour	54 [50, 62]	57 [50, 66]	58 [51, 66]	<.0001	.02	POND > HBN > ABCD
Internalizing problems	56 [48, 64]	57 [50, 65]	63 [55, 69]	<.0001	.03	POND > HBN > ABCD
Externalizing problems	54 [46, 62]	56 [48, 65]	58 [50, 66]	<.0001	.01	POND > HBN > ABCD
Total problems	56 [50, 64]	60 [52, 68]	64 [56, 70]	<.0001	.04	POND > HBN > ABCD

CBCL: Child Behaviour Checklist; POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study

558   ***Performance evaluation***

559   Additional metrics are provided in the Supplemental Table 3, including the area

560   under the precision–recall curve (AU-PR; where recall = sensitivity and precision =

561   positive predictive value [PPV]), along with sensitivity and specificity.

Supplemental Table 33. Model performance.

	Research cohorts			Clinical cohort
	Internal testing	External testing		Internal testing
	POND	HBN	ABCD	PPP
Median AU-PR [IQR]				
Stimulant	.70 [.67, .72]	.68 [.68, .69]	.65 [.65, .66]	.86 [.86, .87]
Anti-depressant	.82 [.79, .84]	.75 [.74, .76]	.66 [.65, .67]	.84 [.82, .85]

Anti-psychotic	.81 [.78, .84]	.79 [.77, .81]	.81 [.80, .82]	.90 [.89, .92]
Median sensitivity [IQR]				
Stimulant	.72 [.70, .73]	.82 [.80, .83]	.77 [.76, .79]	.70 [.68, .72]
Anti-depressant	.73 [.70, .76]	.51 [.49, .52]	.44 [.41, .46]	.69 [.66, .72]
Anti-psychotic	.71 [.66, .76]	.60 [.57, .64]	.58 [.54, .61]	.83 [.68, .72]
Medan specificity [IQR]				
Stimulant	.63 [.61, .65]	.46 [.44, .47]	.53 [.51, .54]	.78 [.77, .80]
Anti-depressant	.76 [.73, .79]	.83 [.82, .85]	.76 [.73, .78]	.80 [.78, .83]
Anti-psychotic	.71 [.66, .76]	.80 [.77, .84]	.86 [.84, .89]	.90 [.89, .92]

PPP: Psychopharmacology Program; POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; ABCD: Adolescent Brain Cognitive Development study; CI: confidence interval; AU-ROC: area under the receiving operating characteristic curve

References

Astle DE, Holmes J, Kievit R, Gathercole SE. Annual Research Review: The transdiagnostic revolution in neurodevelopmental disorders. *Journal of Child Psychology and Psychiatry* 2021; **n/a**. DOI:10.1111/jcpp.13481.

Vandewouw MM, Brian J, Crosbie J, *et al.* Identifying Replicable Subgroups in Neurodevelopmental Conditions Using Resting-State Functional Magnetic Resonance Imaging Data. *JAMA Network Open* 2023; **6**: e232066.

Trost B, Thiruvahindrapuram B, Chan AJS, *et al.* Genomic architecture of Autism Spectrum Disorder from comprehensive whole-genome sequence annotation. Genetic and Genomic Medicine, 2022 DOI:10.1101/2022.05.05.22274031.

Persico AM, Ricciardello A, Lamberti M, *et al.* The pediatric psychopharmacology of autism spectrum disorder: A systematic review - Part I: The past and the present. *Prog Neuropsychopharmacol Biol Psychiatry* 2021; **110**: 110326.

Lord C, Charman T, Havdahl A, *et al.* The Lancet Commission on the future of care and clinical research in autism. *Lancet* 2022; **399**: 271–334.

Feroe AG, Uppal N, Gutiérrez-Sacristán A, *et al.* Medication Use in the Management of Comorbidities Among Individuals With Autism Spectrum Disorder From a Large Nationwide Insurance Database. *JAMA Pediatrics* 2021; **175**: 957–65.

Baribeau D, Vorstman J, Anagnostou E. Novel treatments in autism spectrum disorder. *Curr Opin Psychiatry* 2022; **35**: 101–10.

Murray ML, Hsia Y, Glaser K, *et al.* Pharmacological treatments prescribed to people with autism spectrum disorder (ASD) in primary health care. *Psychopharmacology (Berl)* 2014; **231**: 1011–21.

Lake JK, Weiss JA, Dergal J, Lunsky Y. Child, parent, and service predictors of psychotropic polypharmacy among adolescents and young adults with an autism spectrum disorder. *J Child Adolesc Psychopharmacol* 2014; **24**: 486–93.

Spencer D, Marshall J, Post B, *et al.* Psychotropic medication use and polypharmacy in children with autism spectrum disorders. *Pediatrics* 2013; **132**: 833–40.

Mellahn OJ, Knott R, Tiego J, *et al.* Understanding the Diversity of Pharmacotherapeutic Management of ADHD With Co-occurring Autism: An Australian Cross-Sectional Survey. *Front Psychiatry* 2022; **13**: 914668.

Ritter C, Hewitt K, McMorris CA. Psychotropic Polypharmacy Among Children and Youth with Autism: A Systematic Review. *J Child Adolesc Psychopharmacol* 2021; **31**: 244–58.

Howes OD, Rogdaki M, Findon JL, *et al.* Autism spectrum disorder: Consensus guidelines on assessment, treatment and research from the British Association for Psychopharmacology. *J Psychopharmacol* 2018; **32**: 3–29.

Alexander LM, Escalera J, Ai L, *et al.* An open resource for transdiagnostic research in pediatric mental health and learning disorders. *Sci Data* 2017; **4**: 170181.

Barch DM, Albaugh MD, Avenevoli S, *et al.* Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Dev Cogn Neurosci* 2018; **32**: 55–66.

Achenbach TM, Rescorla LA. Manual for the ASEBA school-age forms and profiles. Burlington, VT: University of Vermont Research Center for Children, Youth, and Families, 2001.

van der Laan MJ, Polley EC, Hubbard AE. Super learner. *Stat Appl Genet Mol Biol* 2007; **6**: Article25.

Prokhorenkova L, Gusev G, Vorobev A, Dorogush AV, Gulin A. CatBoost: unbiased boosting with categorical features. In: Proceedings of the 32nd International Conference on Neural Information Processing Systems. Red Hook, NY, USA: Curran Associates Inc., 2018: 6639–49.

Xu L, Liu S, Lin A, Su Z, Liang D. Interpretable Prediction and Analysis of PVA Hydrogel Mechanical Behavior Using Machine Learning. *Gels* 2025; **11**: 550.

Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. In: Proceedings of the 31st International Conference on Neural Information Processing Systems. Red Hook, NY, USA: Curran Associates Inc., 2017: 4768–77.

Bird S, Dudík M, Edgar R, *et al.* Fairlearn: A toolkit for assessing and improving fairness in AI. Microsoft, 2020 https://www.microsoft.com/en-us/research/publication/fairlearn-a-toolkit-for-assessing-and-improving-fairness-in-ai/.

Agarwal A, Beygelzimer A, Dudik M, Langford J, Wallach H. A Reductions Approach to Fair Classification. In: Dy J, Krause A, eds. Proceedings of the 35th International Conference on Machine Learning. PMLR, 2018: 60–9.

Feldman M, Friedler SA, Moeller J, Scheidegger C, Venkatasubramanian S. Certifying and Removing Disparate Impact. In: Proceedings of the 21th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. New York, NY, USA: Association for Computing Machinery, 2015: 259–68.

Holzer H, Neumark D. Assessing Affirmative Action. *Journal of Economic Literature* 2000; **38**: 483–568.

Howes OD, Rogdaki M, Findon JL, *et al.* Autism spectrum disorder: Consensus guidelines on assessment, treatment and research from the British Association for Psychopharmacology. *J Psychopharmacol* 2018; **32**: 3–29.

Cutillo CM, Sharma KR, Foschini L, *et al.* Machine intelligence in healthcare-perspectives on trustworthiness, explainability, usability, and transparency. *NPJ Digit Med* 2020; **3**: 47.

Antoniou T, McCormack D, Kitchen S, *et al.* Geographic variation and sociodemographic correlates of prescription psychotropic drug use among children and youth in Ontario, Canada: a population-based study. *BMC Public Health* 2023; **23**: 85.

Buckley N, Glasson EJ, Chen W, *et al.* Prevalence estimates of mental health problems in children and adolescents with intellectual disability: A systematic review and meta-analysis. *Aust N Z J Psychiatry* 2020; **54**: 970–84.

Lord C, Rutter M, DiLavore, P, Risi S, Gotham K, Bishop S. Autism diagnostic observation schedule: ADOS-2, 2nd edn. Los Angeles, CA: Western Psychological Services, 2012.

Rutter M, Le Couteur A, Lord C. Autism diagnostic interview revised. Los Angeles, CA: Western Psychological Services, 2003.

Ickowicz A, Schachar RJ, Sugarman R, Chen SX, Millette C, Cook L. The parent interview for child symptoms: a situation-specific clinical research interview for attention-deficit hyperactivity and related disorders. *Can J Psychiatry* 2006; **51**: 325–8.

Kaufman J, Birmaher B, Brent D, *et al.* Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): Initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry* 1997; **36**: 980–8.

Scahill L, Riddle MA, McSwiggin-Hardin M, *et al.* Children's Yale-Brown Obsessive Compulsive Scale: Reliability and validity. *J Am Acad Child Adolesc Psychiatry* 1997; **36**: 844–52.

Wechsler D. Wechsler abbreviated scale of intelligence, 1st edn. San Antonio, TX: The Psychological Corporation, 1999.

Wechsler D. Wechsler preschool and primary scales of intelligence, 3rd edn. San Antonio, TX: The Psychological Corporation, 2002.

Wechsler D. Wechsler intelligence scale for children, 4th edn. San Antonio, TX: The Psychological Corporation, 2003.

Wechsler D. Wechsler abbreviated scale of intelligence, 2nd edn. San Antonio, TX: NCS Pearson, 2011.

Wechsler D. Wechsler intelligence scale for children, 5th edn. Bloomington, MN: Pearson, 2014.

Roid GH. Stanford-Binet intelligence scales, 5th edn. Itasca, IL: Riverside Publishing, 2003.

Diagnostic and statistical manual of mental disorders, 5th edn. Arlington, TX: American Psychiatric Association, 2013.

Kaufman AS, Kaufman NL. Kaufman Brief Intelligence Test, 2nd edn. Pearson.

Volkow ND, Koob GF, Croyle RT, *et al.* The conception of the ABCD study: From substance use to a broad NIH collaboration. *Dev Cogn Neurosci* 2018; **32**: 4–7.

Townsend L, Kobak K, Kearney C, *et al.* Development of Three Web-Based Computerized Versions of the Kiddie Schedule for Affective Disorders and Schizophrenia Child Psychiatric Diagnostic Interview: Preliminary Validity Data. *J Am Acad Child Adolesc Psychiatry* 2020; **59**: 309–25.

Weintraub S, Dikmen SS, Heaton RK, *et al.* Cognition assessment using the NIH Toolbox. *Neurology* 2013; **80**: S54-64.

Heaton RK, Akshoomoff N, Tulsky D, *et al.* Reliability and validity of composite scores from the NIH Toolbox Cognition Battery in adults. *J Int Neuropsychol Soc* 2014; **20**: 588–98.

Chollet F, others. Keras. 2015. https://keras.io.

Martín Abadi, Ashish Agarwal, Paul Barham, *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems. 2015. https://www.tensorflow.org/.