Optimising COVID-19 Episode Identification Using Serology and PCR/Rapid Antigen Testing: Insights from the BRACE Trial

Ellie McDonald[a], Laure F Pittet[a,b,c], Marc Bonten[d,e], Anthony Byrne[f], John Campbell[g], Julio Croda[h,i,j], Margareth Pretti Dalcolmo[k], Andrew Davidson[l], Glauce dos Santos[k], Kaya Gardiner[a,m], Amanda Gwee[b,n,o], Bruno Araújo Jardim[p], Marcus Vinícius Guimaraes Lacerda[q,r,s], Michaela Lucas[t,u,v,w], David J Lynn[x,y], Laurens Manning[w,z,aa], Helen S Marshall[bb], Kirsten P Perrett[b,cc,dd], Cristina Prat-Aymerich[d,e], Marco AM Puga[i,j], Peter C Richmond[w,aa,ee], Jesús Rodríguez-Baño[ff,gg], Ushma Wadia[w,aa], Adilia Warris[hh,ii], Nicholas J Wood[jj,kk,ll], Nigel Curtis[a,b,n,‡], Nicole L Messina[a,b,‡], and the *BRACE Trial Consortium Group*[†]

[‡] Co-last authors, equal contribution.

[†] Members of the *BRACE trial Consortium* are listed in the Supplementary Appendix

[a]Infectious Diseases Group, Murdoch Children's Research Institute, Parkville, Victoria, Australia

[b]Department of Paediatrics, The University of Melbourne, Parkville, Victoria, Australia.

[c]Immunology, Vaccinology, and Infectious Diseases Unit, Department of Paediatrics, Gynaecology and Obsterics, Faculty of Medicine, University of Geneva and University Hospitals of Geneva, Geneva, Switzerland.

[d]European Clinical Research Alliance on Infectious Diseases, Utrecht, the Netherlands

[e]Julius Center for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht University, the Netherlands.

[f]St Vincent's Hospital, Heart Lung Stream, Darlinghurst, Sydney Australia and The University of New South Wales, Faculty of Medicine, St Vincent's Clinical School, Sydney Australia.

[g]Exeter Collaboration for Academic Primary Care, University of Exeter Medical School, Exeter, United Kingdom.

[h]Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, CT, USA.

[i]Fiocruz Mato Grosso do Sul, Fundação Oswaldo Cruz, Campo Grande, Brazil.

[j]Universidade Federal de Mato Grosso do Sul, Campo Grande, Brazil.

[k]Centro de Referência Professor Hélio Fraga, ENSP/FIOCRUZ (Fundação Oswaldo Cruz), Rio de Janeiro, Brazil.

[l]Melbourne Children's Trial Centre, Murdoch Children's Research Institute, Parkville, Victoria, Australia.

[m]Research Operations, The Royal Children's Hospital Melbourne, Parkville, Victoria, Australia.

[n]Infectious Diseases, Royal Children's Hospital Melbourne, Parkville, Victoria, Australia.

[o]Antimicrobials Group, Murdoch Children's Research Institute, Parkville, Victoria, Australia.

[p]Institute of Clinical Research Carlos Borborema, Doctor Heitor Vieira Dourado Tropical Medicine Foundation, Manaus, Brazil.

[q]Fundação de Medicina Tropical Doutor Heitor Vieira Dourado, Manaus, Brazil.

[r]Instituto Leônidas & Maria Deane, Oswaldo Cruz Foundation, Ministry of Health, Manaus, Brazil.

[s]University of Texas Medical Branch, Galveston, TX, USA.

[t]Department of Immunology, Pathwest, Queen Elizabeth II Medical Centre, Nedlands, Western Australia, Australia.

[u]Department of Immunology, Perth Children's Hospital, Nedlands, Western Australia, Australia.

[v]Department of Immunology, Sir Charles Gairdner Hospital, Nedlands, Western Australia, Australia.

[w]School of Medicine, University of Western Australia, Perth, Western Australia, Australia

[x]Flinders Health and Medical Research Institute, Flinders University, Bedford Park, South Australia, Australia.

[y]Precision Medicine Theme, South Australian Health and Medical Research Institute, Adelaide, South Australia, Australia.

[z]Department of Infectious Diseases, Fiona Stanley Hospital, Murdoch, Western Australia, Australia.

[aa]Wesfarmers Centre for Vaccines and Infectious Diseases, The Kids Research Institute, Nedlands, Western Australia, Australia

[bb]Adelaide Medical School and Robinson Research Institute, The University of Adelaide and the Women's and Children's Research Centre, Women's and Children's Health Network, Adelaide, SA, Australia.

[cc]Department of Allergy and Immunology, Royal Children's Hospital Melbourne, Parkville, Victoria, Australia.

[dd]Population Allergy Group, Murdoch Children's Research Institute, Parkville, Victoria, Australia.

[ee]Department of Immunology and General Paediatrics, Perth Children's Hospital, Nedlands, Western Australia, Australia.

[ff]Division of Infectious Diseases and Microbiology, Department of Medicine, Hospital Universitario Virgen Macarena, University of Seville, Biomedicines Institute of Seville-Consejo Superior de Investigaciones Científicas, Seville, Spain.

[gg]CIBER de Enfermedades Infecciosas, Instituto de Salud Carloss III, Madrid, Spain.

[hh]Department of Infectious Diseases, Great Ormond Street Hospital, London, UK.

[ii]Medical Research Council Centre for Medical Mycology, Department of Biosciences, Faculty of Health and Life Sciences, University of Exeter, Exeter, United Kingdom.

[jj]Faculty of Medicine and Health, University of Sydney, Sydney, New South Wales, Australia.

[kk]National Centre for Immunisation Research and Surveillance of Vaccine Preventable Disease, Westmead, New South Wales, Australia.

[ll]Sydney Children's Hospital Network, Westmead, New South Wales, Australia.

Correspondence to: Dr Nicole Messina, Department of Paediatrics, Royal Children's Hospital, 50 Flemington Rd., Parkville, VIC 3052, Australia. nicole.messina@unimelb.edu.au

Author name	Email	ORCiD
Ellie McDonald	ellie.mcdonald@mcri.edu.au	0000-0003-2295-4641
Laure F Pittet	laure.pittet@mcri.edu.au	0000-0002-2395-4574
Marc Bonten	mbonten@umcutrecht.nl	
Anthony Byrne	anthony.byrne@svha.org.au	0000-0003-0789-5161
John Campbell	jlc80@aol.com	0000-0002-6752-3493
Julio Croda	juliocroda@gmail.com	0000-0002-6665-6825
Margareth Pretti Dalcolmo	margarethdalcolmo@gmail.com	0000-0002-6820-1082
Andrew Davidson	andrew.davidson@mcri.edu.au	0000-0002-7050-7419
Glauce dos Santos	glaucedossantos@gmail.com	
Kaya Gardiner	kaya.gardiner@rch.org.au	0000-0001-9796-4567
Amanda Gwee	amanda.gwee@rch.org.au	0000-0003-4016-8986
Bruno Araújo Jardim	brunojardim89@hotmail.com	0000-0003-3465-7556
Marcus Vinícius Guimaraes Lacerda	marcuslacerda.br@gmail.com	0000-0003-3374-9985
Michaela Lucas	michaela.lucas@uwa.edu.au	0000-0001-8881-9990
David J Lynn	david.lynn@sahmri.com	0000-0003-4664-1404
Laurens Manning	laurens.manning@uwa.edu.au	0000-0003-4334-5351
Helen S Marshall	helen.marshall@adelaide.edu.au	0000-0003-2521-5166
Kirsten P Perrett	kirsten.perrett@rch.org.au	0000-0002-5683-996X
Cristina Prat-Aymerich	cristina.prat@ecraid.eu	0000-0001-6974-9165
Marco AM Puga	marco.m.puga@gmail.com	0000-0002-6514-537X
Peter C Richmond	peter.richmond@uwa.edu.au	0000-0001-7562-7228
Jesús Rodríguez-Baño	jesusrb@us.es	0000-0001-6732-9001
Ushma Wadia	ushma.wadia@telethonkids.org.au	0000-0002-8097-5072

Adilia Warris   a.warris@exeter.ac.uk   0000-0001-6586-3358

Nicholas J Wood   nicholas.wood@health.nsw.gov.au   0000-0002-3830-5307

Nigel Curtis   nigel.curtis@rch.org.au   0000-0003-3446-4594

Nicole L Messina   nicole.messina@unimelb.edu.au   0000-0001-8404-4462

Abstract

Background

Accurately identifying COVID-19 episodes was crucial during the pandemic for evaluating interventions. Results from diagnostic tools like PCR, rapid antigen test (RAT) and serology are affected by factors such as timing of tests and vaccination status. The BRACE trial developed an algorithm integrating these diagnostic tools for illness episode classification.

Methods

In the BRACE trial, 3988 participants reported 5512 febrile/respiratory illness episodes and provided longitudinal blood samples over one year. SARS-CoV-2 diagnosis relied on a three-component algorithm: (1) a serology algorithm assessing anti-SARS-CoV-2 nucleocapsid antibody seroconversion, (2) a PCR/RAT algorithm, and (3) an episode interpretation algorithm combining serology and PCR/RAT results to categorise episodes as COVID-19, Not COVID-19 or Uncertain. The algorithms accounted for vaccination status and timing of testing relative to symptom onset to refine episode classifications.

Results

Of 5512 illness episodes, 890 (16%) were classified as COVID-19, 3852 (70%) as Not COVID-19, and 770 (14%) as Uncertain. Compared to relying solely on PCR/RAT results, integrating serology in the algorithm reduced the proportion of Uncertain classifications by more than half. Among the COVID-19 episodes, 89% were identified by positive PCR/RAT results, and the remaining 11% (with missing or negative PCR/RAT tests) were identified by serology. Discordance between PCR/RAT and serology occurred in 13% of episodes.

Conclusion

An algorithm integrating PCR/RAT and serology results in the context of test timing and vaccine

status enabled the accurate identification of COVID-19 episodes and minimised the number of

episodes that would otherwise have been classified as Uncertain.

Background

Accurately identifying COVID-19 episodes was vital for evaluating interventions in the pandemic. The performance of diagnostic tests for SARS-CoV-2, such as respiratory swab testing and serology, is influenced by the timing of testing relative to symptom onset, as well as prior COVID-19 vaccinations.

In the BRACE trial, comprehensive symptom data were collected alongside SARS-CoV-2 testing.[1-3] Here, we describe the algorithm developed during the BRACE trial, intended to more precisely identify COVID-19 episodes using illness episode data and diagnostic test results.

To account for test timing in relation to symptom onset and any effect of COVID-19 vaccination, a three-component interpretation algorithm was developed. This incorporated serological result interpretation, PCR/rapid antigen test (RAT) respiratory swab test interpretation, and finally episode classification to COVID-19, Not COVID-19 or Uncertain categories.

In addition to evaluating the sensitivity and specificity of PCR/RAT testing compared to serology for SARS-CoV-2, and the frequency of discordant results, we assessed the impact of each component of the algorithm on episode classification.

Methods

Participants and study design

The BRACE trial (NCT04327206) was a phase 3 international randomised controlled trial to assess the impact of BCG vaccination on the prevalence of COVID-19 among healthcare workers.[1-3] Trial outcomes relied on accurate ascertainment of the onset of a participant's first episode of COVID-19.

The trial was designed early in the pandemic, with recruitment starting in March 2020. This study included the 3988 participants recruited to the BRACE trial from May 2020 to April 2021 from Australia, the Netherlands, Spain, the UK and Brazil. Selection of COVID-19 symptoms to be collected and diagnostic tests for SARS-CoV-2 were informed by best practice and government policies at the time.

Illness episode ascertainment

Recent illness symptoms were ascertained weekly via a custom-built smartphone app or direct communication (phone call or text). During illness episodes, symptoms were recorded daily, and participants were prompted to undergo SARS-CoV-2 respiratory swab testing using PCR (via government testing centres) or RAT (self-administered). Additionally, more comprehensive questionnaires were administered at the start of the trial and quarterly throughout follow-up. These collected information about COVID-19-specific vaccinations (date and vaccine type). Blood samples were collected at baseline and 3, 6, 9 and 12 months post-randomisation to measure anti-SARS-CoV-2 nucleocapsid (NCP) antibodies using the Roche Cobas Elecsys anti-SARS-CoV-2 assay [4, 5] at the Victorian Infectious Diseases Reference Laboratory. For seroconversion, a cut-off index (COI) of ≥1.0 was used as per manufacturer's instructions.[6, 7]

Serological interpretation algorithm

The serology algorithm produced one of four possible outcomes: Seroconversion, No seroconversion, Indeterminate or No result (where a blood sample was unavailable). For each illness episode, the most recent blood sample collected at least seven days prior to symptom onset (pre-episode sample), and the first and second blood samples collected after symptom onset were identified. The NCP serology result from these samples were used in the serological interpretation algorithm. The algorithm considered the timing of blood sample in relation to the illness episode, any prior administration of the *CoronaVac* (Sinovac Biotech) COVID-19 vaccine (a whole inactivated virus vaccine which can induce anti-NCP antibodies) and change in anti-NCP antibody titre (in cases of

positive-to-positive serology results). A detailed description of the serological algorithm is provided in Supplementary Figure 1.

PCR/RAT interpretation algorithm

The PCR/RAT interpretation algorithm yielded three possible outcomes: Positive, Negative, or No result. Both PCR and RAT positive results were considered confirmation of SARS-CoV-2 infection if the timing of the test fell within defined symptomatic illness parameters. Positive PCR results were considered valid if the sample was tested within three days prior to 10 days (RAT) or 21 days (PCR) after the start, or within seven days from the end, of an illness episode. A negative PCR was considered 'Negative', but a negative RAT was classified as 'No result' due to the lower sensitivity of RAT testing.[8] A detailed description of the PCR/RAT interpretation algorithm is provided in Supplementary Figure 2.

Episode interpretation algorithm

The episode interpretation algorithm produced three possible classifications for each illness episode: COVID-19, Not COVID-19 or Uncertain (which included episodes for which results were missing, or indeterminate). The results of the serological and PCR/RAT algorithms were combined to categorise illness episodes using the episode interpretation algorithm, detailed in Figure 1. Additional conditions were then applied which evaluated other COVID-19 tests and episode results within the same timeframe.

Results

Of the 3988 participants, 2559 reported at least one episode of illness in the year post-randomisation, with a total of 5512 episodes.

Of the 5512 episodes, 890 (16%) were classified as COVID-19, 3852 (70%) as Not COVID-19 and 770 (14%) as Uncertain (Figure 1). The participant flow from the combined PCR/RAT result and from the

pre-episode NCP serology to the final episode classification are shown in Figure 2 and Supplementary Figure 3 respectively.

Of the 890 episodes defined as COVID-19, 792 (89%) had a positive PCR or RAT test result. Serology results were concordant (seroconversion) in 375 (47.4%) episodes, discordant (no seroconversion) in 38 (4.8%), and Indeterminate or No result in 379 (47.9%) (Figure 2). The remaining 98 COVID-19 episodes comprised 56 (6.3%) with a missing PCR/RAT result and 42 (4.7%) with a negative PCR result (Supplementary Figure 4). The discordant PCR/RAT and serology results were not attributable to post-episode NCP titres being just above the threshold for seroconversion (Figure 3). The post-episode NCP titres in relation to the time interval between the episode start and post-episode blood collection for the seroconverted PCR negative and PCR/RAT positive episodes are shown in Supplementary Figure 5. Median post-episode NCP titres were higher in seroconverting episodes with negative PCR results compared to those with positive PCR/RAT results (median 62.2, IQR: 15.9–122.9 vs. 41.3, IQR: 14.3–101.4; p=0.3). Relevant to this, median time between episode onset and post-episode serological testing was higher in the negative PCR episode group compared to the positive PCR/RAT group (median 76.5 days, IQR: 33-102 days vs. 61 days, IQR: 37-82 days; p=0.1).

There were 305 episodes that had a negative RAT test. Of the RAT-negative episodes without a PCR result, 11/179 (6.2%) were categorised as COVID-19 based on seroconversion (Supplementary Figure 6). In comparison, of the episodes without a PCR result, 56/858 (7.0%) were categorised as COVID-19 based on seroconversion (Figure 3).

Among the 3852 episodes classified as Not COVID-19 by the final episode algorithm, 3050 (77%) had a negative PCR result (Figure 2 & Supplementary Figure 3). Of these, 1622 (53.2%) had concordant negative serology and PCR results (Figure 2). Additionally, 1351 (44.3%) episodes had a missing NCP serology result, but a negative PCR result. The remaining 77 (2.5%) episodes were discordant (seroconversion with a negative PCR result). However, the positive NCP serology result in these cases was attributed to participants' other PCR- or RAT-positive episodes within the same timeframe. For

the remaining 802 (20%) Not COVID-19 episodes with a missing PCR result, 744 had no

seroconversion between pre- and post-episode blood samples. These episodes would have been

classified as Uncertain if the trial had relied on PCR/RAT results only. In the remaining 58 (7.2%)

episodes, participants seroconverted, but the episode algorithm attributed this to another PCR- or

RAT-positive episode.

Of the 770 episodes classified as Uncertain in the final episode result, 738 (95.8%) had both missing

or indeterminant serology and PCR/RAT results. Of the remaining 32 episodes, participants

seroconverted but the episodes were categorised as Uncertain because the participants had multiple

episodes with missing PCR/RAT results within the same seroconversion timeframe. This precluded

the attribution of the COVID-19 diagnosis to one of the episodes.

At the participant level, Uncertain result episodes were similarly distributed between every

combination of COVID-19 and Not COVID-19 episodes (Figure 4). The sequential distribution of

COVID-19, Not COVID-19 and Uncertain result episodes for each participant demonstrates the

progression from algorithm processing to the final trial outcomes. It highlights the points at which

participants were censored, which occurred at one of two events: their first COVID-19 episode or

their first episode with an Uncertain result (Figure 5).

Discussion

The algorithms developed for the BRACE trial provided a novel and robust method for the

classification of illness episodes and identification of COVID-19 cases. Unlike other trials that relied

exclusively on PCR/RAT [9-11] or serological testing,[12] often classifying cases individually,[13, 14]

our three-step algorithm combined these test results with real-world considerations. These included

timing of testing relative to symptom onset, variations in the sensitivity and specificity of PCR/RAT

and serology, together with the effects of COVID-19 vaccination on serological results.[15] Critically,

we accounted for the lower reliability of a negative RAT result due to its lower sensitivity and the impact of *CoronaVac*-induced anti-NCP antibodies.[8]

Post-hoc analysis showed that less than 10% of participants with RAT-negative, PCR-missing episodes seroconverted and were classified as COVID-19, while nearly 15% of participants with RAT-negative, PCR-missing episodes were excluded from primary analysis due to missing serological results. Discordance between PCR/RAT and serology test results was observed in a small proportion (11%) of episodes. Possible explanations for the 38 (5%) PCR/RAT-positive episodes without seroconversion include mild infections that did not elicit a systemic response, early waning of immunity, limitations in the sensitivity of serology, or false positive PCR/RAT. Possible explanations for the 42 (6%) PCR-negative episodes with seroconversion include inadequate swabbing, technical issues, participant reporting errors, and undetected asymptomatic infections during the interval between blood samples.

Serological testing proved especially valuable when PCR/RAT results were unavailable. Negative serology reclassified 744 episodes with missing PCR/RAT results as Not COVID-19 by demonstrating an absence of seroconversion. Additionally, serology alone identified 11% (98/890) of COVID-19 episodes in which the PCR/RAT result was negative or missing. The algorithm was particularly beneficial in mitigating the impact of incomplete data by reducing the proportion of episodes for which classification would not have been possible (and therefore classified as Uncertain) by more than half, from 30% (1628/5512) to 14% (770/5512).

The addition of serology to PCR/RAT testing involves considerable cost and resources, including staff, consumables, shipment, laboratory assays and data management. However, the benefits included in-person follow-ups that encouraged trial retention, enhanced data accuracy, clarified discordances and the facilitation of additional research opportunities, such as additional immunological studies.[16-21]

Our analyses have some limitations. First, our definition of symptomatic COVID-19 relied on the original case criteria, excluding non-febrile episodes without respiratory symptoms. Second, the timing and type of PCR/RAT testing were not standardised and varied depending on individual settings. Third, serological testing was done quarterly rather than at a consistent interval following illness episodes, which may have affected the detection of seroconversion. Finally, a proportion of illness episodes had incomplete data. However, it is among the COVID-19-related trials with the most complete data available.

Our three-component algorithm developed in the BRACE trial has potential for adaption to diverse epidemiological scenarios beyond COVID-19. Infectious diseases such as influenza or respiratory syncytial virus also require accurate longitudinal monitoring to track transmission, assess vaccine efficacy, and evaluate intervention and treatment outcomes.

In summary, despite robust data collection efforts, COVID-19 trials faced practical challenges, particularly during the early pandemic. Testing availability and participant adherence contributed to the 14% of episodes still classified as missing. Adding serological testing demanded substantial time, financial resources, and participant effort but effectively halved the number of unclassified episodes. Our study underscores the important role of algorithms in achieving comprehensive classification, particularly in complex real-world study settings.

Authors' Contributions

EM, NC and NM were responsible for the original draft of this paper, with EM responsible for Visualization and analysis. All authors contributed to administration and/or investigation in the BRACE trial, and all were involved in the review and editing of this manuscript. NC, NM, LP and KG were responsible for Conceptualization, Funding acquisition and Supervision. NC, NM, LP and EM were responsible for Methodology. All authors had final responsibility for the decision to submit for

publication. NC and NM attest that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Declaration of Interest Statements

PCR has received investigator-initiated research grants to his institution from Sanofi outside this work and has received institutional funding from GlaxoSmithKline and Sanofi for local and international lectures and from AstraZeneca, Clover Biopharmaceuticals, GlaxoSmithKline, Novavax, Sanofi, and Pfizer for participation in vaccine scientific advisory boards unrelated to this work. He has been an investigator in sponsored, multicentre vaccine trials for AstraZeneca, Moderna, Novavax, Pfizer, and Sanofi with funding to his institution. He is Chief investigator on an investigator led COVID-19 vaccine trial and institution has received funding from the Medical Research Future Fund. HSM has received investigator-initiated research grants to her institution from Sanofi and Pfizer outside this work. She has been an investigator in sponsored, multicentre vaccine trials for Iliad, Seqirus and Pfizer, with funding to her institution. She is a Chief investigator on an investigator led COVID-19 vaccine trial and institution has received funding from the Medical Research Future Fund. All other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data sharing statement

Deidentified participant data and data dictionary are available to others on request and on completion of a signed data access agreement. Requests can be made in writing to braceresearch@mcri.edu.au.

Ethics Committee approval

The study was approved by the Royal Children's Hospital Melbourne Human Research Ethics Committee (No.62586); the protocol was approved by the ethics committee at each site and all participants provided informed consent. The trial was overseen by a steering committee and a data safety and monitoring board.

Acknowledgements

We thank the trial participants and site personnel who were involved in assisting the trial (detailed in the Supplementary File). AJ Vaccines (Copenhagen) for facilitating BCG vaccine supplies; Devon Freewheelers (UK) for UK transport; Catalent (UK) for drug management; Fiocruz Clinical Research Platform for coordination and monitoring of sites in Brazil; the Pharmacy, Pathology and Immunisation Service teams from The Royal Children's Hospital Melbourne, and the Orygen Group for their support.

Funding sources

The trial is supported by the Bill & Melinda Gates Foundation [INV017302], the Minderoo Foundation [COV-001], Sarah and Lachlan Murdoch, the Royal Children's Hospital Foundation [2020-1263 BRACE Trial], Health Services Union NSW, the Peter Sowerby Foundation, SA Health, the Insurance Advisernet Foundation, the NAB Foundation, the Calvert-Jones Foundation, the Modara Pines Charitable Foundation, the UHG Foundation Pty Ltd, Epworth Healthcare and individual donors. The Murdoch Children's Research Institute (MCRI) leads the BRACE trial across 36 sites in five countries. It is supported by the Victorian Government's Operational Infrastructure Support Programme. NC is supported by a National Health and Medical Research Council (NHMRC) Investigator Grant (GNT1197117). LFP is supported by the Swiss National Science Foundation (Early

Postdoc Mobility Grant, P2GEP3_178155). The funders of this study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

References

[1] Pittet LF, Messina NL, Gardiner K, Orsini F, Abruzzo V, Bannister S, et al. BCG vaccination to reduce the impact of COVID-19 in healthcare workers: Protocol for a randomised controlled trial (BRACE trial). BMJ Open. 2021;11:e052101.

[2] Pittet LF, Messina NL, Orsini F, Moore CL, Abruzzo V, Barry S, et al. Randomized Trial of BCG Vaccine to Protect against Covid-19 in Health Care Workers. N Engl J Med. 2023;388:1582-96.

[3] Pittet LF, Messina NL, McDonald E, Orsini F, Barry S, Bonten M, et al. Bacille Calmette-Guérin vaccination to prevent febrile and respiratory illness in adults (BRACE): secondary outcomes of a randomised controlled phase 3 trial. eClinicalMedicine. 2024;72.

[4] Bond KA, Williams E, Nicholson S, Lim S, Johnson D, Cox B, et al. Longitudinal evaluation of laboratory-based serological assays for SARS-CoV-2 antibody detection. Pathology. 2021;53:773-9.

[5] Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nat Microbiol. 2020;5:562-9.

[6] Elecsys® Anti-SARS-CoV-2. Immunoassay for the qualitative detection of antibodies against SARS-CoV-2. In: Ltd RDI, editor.2020.

[7] Noble C ME, Nicholson S, Biering-Sørensen S, Pittet LF, Byrne AL, Croda J, Dalcolmo M, Lacerda M, Lucas M, Lynn DJ, Prat Aymerich C, Richmond PC, Warris A, Curtis N, Messina NL; BRACE Trial Consortium Group. . Characterising the SARS-CoV-2 nucleocapsid (N) protein antibody response. Journal of Infection. 2025.

[8] Mak GC, Cheng PK, Lau SS, Wong KK, Lau CS, Lam ET, et al. Evaluation of rapid antigen test for detection of SARS-CoV-2 virus. J Clin Virol. 2020;129:104500.

[9] Czajka H, Zapolnik P, Krzych L, Kmiecik W, Stopyra L, Nowakowska A, et al. A Multi-Center, Randomised, Double-Blind, Placebo-Controlled Phase III Clinical Trial Evaluating the Impact of BCG Re-Vaccination on the Incidence and Severity of SARS-CoV-2 Infections among Symptomatic Healthcare Professionals during the COVID-19 Pandemic in Poland-First Results. Vaccines (Basel). 2022;10.

[10] Upton CM, van Wijk RC, Mockeliunas L, Simonsson USH, McHarry K, van den Hoogen G, et al. Safety and efficacy of BCG re-vaccination in relation to COVID-19 morbidity in healthcare workers: A double-blind, randomised, controlled, phase 3 trial. EClinicalMedicine. 2022;48:101414.

[11] Koekenbier EL, Fohse K, van de Maat JS, Oosterheert JJ, van Nieuwkoop C, Hoogerwerf JJ, et al. Bacillus Calmette-Guerin vaccine for prevention of COVID-19 and other respiratory tract infections in older adults with comorbidities: a randomized controlled trial. Clin Microbiol Infect. 2023;29:781-8.

[12] Dos Anjos LRB, da Costa AC, Cardoso A, Guimaraes RA, Rodrigues RL, Ribeiro KM, et al. Efficacy and Safety of BCG Revaccination With M. bovis BCG Moscow to Prevent COVID-19 Infection in Health Care Workers: A Randomized Phase II Clinical Trial. Front Immunol. 2022;13:841868.

[13] Faustman DL, Lee A, Hostetter ER, Aristarkhova A, Ng NC, Shpilsky GF, et al. Multiple BCG vaccinations for the prevention of COVID-19 and other infectious diseases in type 1 diabetes. Cell Rep Med. 2022;3:100728.

[14] Santos AP, Werneck GL, Dalvi APR, Dos Santos CC, Tierno P, Condelo HS, et al. The effect of BCG vaccination on infection and antibody levels against SARS-CoV-2-The results of ProBCG: a multicenter randomized clinical trial in Brazil. Int J Infect Dis. 2023;130:8-16.

[15] Noble C, McDonald E, Nicholson S, Biering-Sorensen S, Pittet LF, Byrne AL, et al. Characterising the SARS-CoV-2 nucleocapsid (N) protein antibody response. J Infect. 2025:106436.

[16] Dos Santos PCP, Messina NL, de Oliveira RD, da Silva PV, Puga MAM, Dalcolmo M, et al. Effect of BCG vaccination against Mycobacterium tuberculosis infection in adult Brazilian health-care workers: a nested clinical trial. Lancet Infect Dis. 2024.

[17] Messina NL, Germano S, McElroy R, Bonnici R, Grubor-Bauk B, Lynn DJ, et al. Specific and off-target immune responses following COVID-19 vaccination with ChAdOx1-S and BNT162b2 vaccines-an exploratory sub-study of the BRACE trial. EBioMedicine. 2024;103:105100.

[18] Messina NL, Sperotto MG, Puga MAM, da Silva PV, de Oliveira RD, Moore CL, et al. Impact of vaccine platform and BCG vaccination on antibody responses to COVID-19 vaccination. Front Immunol. 2023;14:1172851.

[19] Messina NL, Germano S, McElroy R, Rudraraju R, Bonnici R, Pittet LF, et al. Off-target effects of bacillus Calmette-Guerin vaccination on immune responses to SARS-CoV-2: implications for protection against severe COVID-19. Clin Transl Immunology. 2022;11:e1387.

[20] Stevens NE, Ryan FJ, Messina NL, Blake SJ, Norton TS, Germano S, et al. No evidence of durable trained immunity after two doses of adenovirus-vectored or mRNA COVID-19 vaccines. J Clin Invest. 2023;133.

[21] Ryan FJ, Norton TS, McCafferty C, Blake SJ, Stevens NE, James J, et al. A systems immunology study comparing innate and adaptive immune responses in adults to COVID-19 mRNA and adenovirus vectored vaccines. Cell Rep Med. 2023;4:100971.

Figure 1. Episode interpretation algorithm: the detail

Figure 2. Episode interpretation outcome

Figure 3. Pre and post NCP titre serology results by PCR/RAT and episode result

Figure 4. Overall episode interpretation

Figure 5. Sequential episode interpretation: COVID-19, Not COVID-19 and Uncertain outcome episodes per participant