The epidemiology of pathogens with pandemic potential: A review of key parameters and clustering analysis

Authors

Jack Ward[1], Oswaldo Gressani[2], Sol Kim[1,3], Niel Hens[2,4], W. John Edmunds[1]

Affiliations

[1]Centre for the Mathematical Modelling of Infectious Diseases, Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, UK.

[2]Data Science Institute, I-BioStat, Hasselt University, Hasselt, Belgium.

[3]Department of Infectious Disease Epidemiology and Dynamics, Institute of Tropical Medicine, Nagasaki University (or) Institute of Tropical Medicine, Nagasaki University

[4]Centre for Health Economic Research and Modelling Infectious Diseases, Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

Corresponding author
Jack Ward (jack.ward@lshtm.ac.uk)

Abstract

Introduction
In the light of the COVID-19 pandemic many countries are trying to widen their pandemic planning from its traditional focus on influenza. However, it is impossible to draw up detailed plans for every pathogen with epidemic potential. We set out to try to simplify this process by reviewing the epidemiology of a range of pathogens with pandemic potential and seeing whether they fall into groups with shared epidemiological traits.

Methods
We reviewed the epidemiological characteristics of 19 different pathogens with pandemic potential (those on the WHO priority list of pathogens, different strains of influenza and Mpox). We extracted data on key parameters (basic reproduction number (R_0), serial interval, proportion of presymptomatic transmission, case fatality risk (CFR) and transmission route) and applied an unsupervised learning framework. This combined Monte Carlo sampling with ensemble clustering to classify pathogens into distinct epidemiological archetypes based on their shared characteristics.

Results
From 154 articles we extracted 302 epidemiological parameter estimates. The clustering algorithms categorise these pathogens into six archetypes (1) Vector-borne pathogens with secondary transmission routes, (2) Airborne, high transmission pathogens with high presymptomatic transmission, (3) Airborne, high transmission pathogens with low

presymptomatic transmission, (4) Contact zoonoses with high CFR, (5) Respiratory zoonoses with high CFR and (6) Contact zoonoses with evidence of presymptomatic transmission.

Conclusion

Unsupervised learning on epidemiological data can be used to define distinct pathogen archetypes. This method offers a valuable framework to allocate emerging and novel pathogens into defined groups to evaluate common approaches for their control.

Keywords

Pandemic preparedness; Blueprint pathogens; infectious diseases; epidemiology

---

1. Introduction

Recent global epidemics of COVID-19 and Mpox have illustrated that we remain vulnerable to global biological incidents. Historically, pandemic preparedness strategies have been limited in scope. For instance, prior to COVID-19, the UK government's sole pandemic plan was the 2011 Influenza Pandemic Preparedness Strategy [1,2]. This narrow focus left critical gaps in threat readiness that have been exploited by non-influenza pathogens such as SARS-CoV-2 and Mpox virus (MPV).

Given the potential health and economic impacts of pandemics, the way in which we plan for such risks needs to be revised. There are 26 viral families known to infect humans [3], but only a fraction of these viruses will possess the ability for widespread transmission in the community [4]. Historically, this fraction has been listed, based on historical outbreaks and ranked to inform policy makers on which pathogens possess the highest pandemic potential. A list based approach, while useful, is inflexible and is rooted in responding to yesterday's pandemic rather than proactively planning.

Categorising pathogens based on shared epidemiological traits rather than using historical lists offers a more flexible and inclusive framework for pandemic planning [4]. A trait based approach would facilitate proactive planning for emerging threats by categorising pathogens by characteristics, allowing planners to assess a wider breadth of scenarios and control measures rather than specific historical examples [4].

To address these gaps, we propose classifying pathogens into archetypes based on epidemiological traits. Using data collected from previous systematic reviews where possible, and individual papers and parameter estimation where necessary, we implement

84 an unsupervised machine learning framework combining Monte Carlo sampling with
85 ensemble clustering to identify relevant pathogen archetypes. By categorising pathogens by
86 their reproduction number, serial interval, case fatality ratio, proportion of presymptomatic
87 transmission and transmission route, we show how pathogens can be grouped by shared
88 characteristics which may point to common approaches for their control.
89
90

91 ____

92
93

94 2. Methods

95 2.1 Review of epidemiological parameters

We selected 19 pathogens for review based on their epidemic or pandemic potential. This
list was primarily guided by the World Health Organisation's (WHO) R&D Blueprint (as of
June 2024) [5] and included: SARS-CoV-2 (Wild-Type, Alpha, Delta, Omicron), SARS-CoV-
1, MERS-CoV, Crimean-Congo hemorrhagic fever orthonairovirus (CCHFV), Ebola virus
(EBOV), Marburg virus (MARV), Lassa virus (LASV), Nipah virus (NiV), Rift Valley fever
virus (RVFV), Zika virus (ZIKV), Mpox virus (MPV), and several influenza A viruses (H1N1,
H2N2, H3N2, H1N1pdm09, and A/H5N1).
103
104 We sought to identify quantitative estimates for parameters related to each pathogen's
105 transmission route(s), infection timeline, and severity. Key parameters included the basic
106 reproduction number (R_0), overdispersion parameter for the reproduction number (k),
107 incubation period, latent period, infectious period, serial interval, case fatality risk (CFR), and
108 infection fatality risk (IFR).
109
Our search strategy involved systematic queries of PubMed for peer-reviewed articles and
preprints. Search terms were structured around three components: pathogen name,
parameter type, and "systematic review" (**Supplementary Table S1**). Studies were required
to report quantitative estimates derived from primary epidemiological data within systematic
reviews or meta-analyses. For pathogens lacking comprehensive systematic reviews, we
conducted targeted searches using pathogen-specific terminology, without a fixed strategy.
We also included articles that provided datasets allowing for parameter estimation.
117

118 2.2 Parameter estimation

We estimated key parameters that were not available from the literature review. Where
appropriate, we estimated values for the incubation period, serial interval, R_0 and proportion
of presymptomatic transmission for selected pathogens. For the incubation period, we used
the {EpiLPS} package [6–8] where publicly available data permitted. We estimated serial
intervals by fitting lognormal and gamma distributions to the number of onsets for a given
day, accounting for double censoring using the R package {primarycensored} [9,10]. For R_0,
we used the package {epichains} [10,11], to provide an estimate for CCHFV on the basis of

data collected by the European Centre for Disease Prevention and Control [11]. Full
methodological details are provided in the Supplementary Information.
128

129 2.3 Clustering of epidemiological parameters

130 We employed a Monte Carlo based ensemble clustering method to classify pathogens
131 according to their epidemiological and transmission characteristics. An ensemble method
132 was chosen to aggregate results from multiple clustering algorithms which may be potentially
133 epidemiologically conflicting [12].
134

For each pathogen, five key characteristics were considered, representing its transmissibility
(R_0), the speed of its spread (serial interval), its severity (CFR), its mode of transmission
(transmission route), and the timing of its infectiousness (proportion of presymptomatic
transmission). To account for parameter uncertainty, a Monte Carlo simulation of 5,000
iterations was performed. In each iteration, a new set of continuous parameters (R_0, serial
interval, CFR) was drawn for each pathogen from specified probability distributions. The
categorical transmission route data was converted into binary features using one-hot
encoding and held constant across simulations. The proportion of presymptomatic
transmission was estimated for each iteration as the probability that the serial interval is
shorter than the incubation period ($P(SI < IP)$), calculated by generating 5,000 correlated
quantile pairs from the respective distributions.
146

147 Following parameter generation, the full 5,000-iteration dataset was standardised to a mean
148 of zero and a standard deviation of one. A K-means clustering algorithm, with a fixed K of 6
149 and 25 random starts, was then applied to each of the 5,000 simulated parameter sets.
150

The results of these individual runs were synthesised into a stable consensus grouping using
an ensemble approach. First, a co-assignment matrix was constructed, tallying the
frequency with which each pair of pathogens was assigned to the same cluster across all
5,000 iterations. This was converted into a consensus-based dissimilarity matrix, to which an
agglomerative hierarchical clustering algorithm with "average" linkage was applied. The final
number of six consensus clusters was determined by silhouette analysis, with the results
visualised as a dendrogram.
158

159 2.3.1 Sensitivity analysis

We conducted a series of sensitivity analyses to evaluate both the robustness and the
generalisability of our clustering method. First, to test the stability of the original pathogen
archetypes, we assessed how the clusters changed with the inclusion of additional
epidemiological parameters, this included *k*, the incubation, latent and infectious period. This
allowed us to determine how robust our initial classifications were to changes in the
parameter space.

Second, to explore the flexibility and potential further application of our method, we
performed an analysis on a core set of parameters: R_0, incubation period, and CFR. This
enabled the incorporation of a more diverse range of pathogens, including those with

different transmission dynamics, and to investigate how they fit within this classification. We
extended this to include pathogens with had been excluded from the original analysis
(RVFV), food and water borne pathogens (*Vibrio cholerae* and Norovirus), viruses common
in paediatrics (Measles virus, Enterovirus A71 and Human metapneumovirus), bioterrorism
related pathogens (*Yersinia pestis*, Variola virus and *Bacillus anthracis*), other vector borne
pathogens (SFTS virus and Chikungunya), and a retrovirus (Human immunodeficiency virus)
(**Supplementary Table S2**).

176
177

3. Results

3.1 Parameter review

A total of 154 articles were retrieved (**Supplementary Figure S1**). This included 43 articles
obtained through the initial search of systematic reviews and 69 articles obtained from
supplementary sources. Among these, one was a grey literature report published by the
WHO. We extracted 302 parameter estimates from the articles identified (**Figure 1**).

28 articles were identified that provided sufficient data to estimate the incubation period with
the {EpiLPS} package. Additionally, 14 articles contributed pre-existing datasets that were
incorporated into the analysis.

Figure 1. Distribution of parameter estimates from the literature search
From 154 articles we extracted 302 parameter estimates. 63 for the reproduction number, 15 for the dispersion parameter, 51 for the incubation period, 33 for the serial interval, 27 for the latent period, 33 for the infectious period, 59 for the CFR, and 21 for the IFR.

3.2 Epidemiological characteristics

3.2.1 Transmission dynamics

We extracted 63 estimates for the reproduction number, which varied across pathogens and outbreaks due to differences in settings and the implementation of control measures (**Table 1**). One additional estimate was derived, with the median R☐ for CCHFV estimated at 0.03 (95% CrI: 0.004–0.09) based on cases reported in the European Union between 2013 and 2014 (**Supplementary figure S4**).

A total of 15 estimates for the dispersion parameter (k) were extracted, revealing varying degrees of heterogeneity in transmission. SARS-CoV-1, SARS-CoV-2, MERS-CoV and NiV exhibited particularly low k values, indicating a high potential for superspreading events, whereas influenza displayed greater uniformity in transmission. Pathogens such as EBOV were reported to have a wide range of estimates suggesting the degree of superspreading may be dependent on outbreak setting (**Table 1**).

3.2.2 Time to key events

We extracted 51 estimates for the incubation period and provided 13 additional estimates
based on publicly available data. The length of the incubation period varied across
pathogens, with pandemic influenza exhibiting the shortest incubation period. In contrast,
pathogens primarily transmitted through contact with infected body fluids, such as LASV,
EBOV, and CCHF, had the longest incubation periods (**Table 1**).

For the serial interval, we extracted 33 estimates. As with the incubation period, serial
intervals were shortest for pandemic influenza and SARS-CoV-2 and longest for contact-
transmitted pathogens (**Table 1**). Additionally, we estimated serial intervals for LASV and
CCHF, with mean values of 11.5 days (95% CrI: 0.9–34.6) (**Supplementary Figure S2**) and
12.0 days (95% CrI: 3.0–27.2) (**Supplementary Figure S3**), respectively.

For the latent and infectious periods, we extracted 27 and 33 estimates, respectively. These
durations showed notable variation between pathogens, highlighting differences in disease
progression **(Table 1)**.

219 3.2.3 Severity and Mortality Risk

We extracted 59 estimates for the case fatality rate (CFR) and 21 for the infection fatality
rate (IFR). CFR estimates exhibited a broad range across pathogens, reflecting varying
levels of disease severity (**Table 1**). Additionally, CFR varied within pathogens, with
substantial heterogeneity between studies, outbreaks and age groups. For instance, NiV
outbreaks with implemented control measures reported lower CFR estimates (**Table 1**).

Compared to CFR, fewer IFR estimates were reported. IFR values were consistently lower
than CFR estimates for the same pathogen and demonstrated age-dependent variation
(**Table 1**)

> **Table 1. for Epidemiological parameters of pandemic potential pathogens. Located after conclusion**

230 3.3 Ensemble clustering

A different number of final consensus clusters was modelled (K=2-10)**.** The six-cluster
solution was ultimately considered the most appropriate according to silhouette analysis
(**Supplementary figure S5**) as well as heuristically. In the final clustering, CCHFV and ZIKV
were grouped into cluster 1; SARS-CoV-2 Alpha, Delta and Omicron into cluster 2; all
pandemic influenza subtypes (except H5N1), SARS-CoV-2 WT and SARS-CoV-1 into
cluster 3; EBOV, MARV and NiV into cluster 4; A/H5N1, MERS-CoV into cluster 5; and
LASV and MPV into cluster 6 (**Figure 2**). The characteristics of each archetype are detailed
in **Table 2**.
239

Figure 2. Pathogen archetypes by consensus hierarchical clustering.
The final, stable classification of 18 pathogens based on their epidemiological and transmission characteristics. The horizontal branch lengths represent the dissimilarity between clusters, with shorter branches indicating a higher frequency of co-assignment in the underlying Monte Carlo simulations.

240
241
242
243

Table 2. Archetype characterisation.						
Archetype	**Pathogens**	**Archetype parameters (mean (95% CI))**				
		R_0	Serial interval (d)	CFR (%)	Proportion of presymptomatic transmission (%)	Transmission route
1	CCHFV, ZIKV	0.07 (0.00– 0.39)	12.03 (3.85– 23.66)	5.9 (0.0–13.4)	0.09 (0.0– 0.2)	Vector (n=2) Sexual contact (n=1) Direct contact (n=1) Animal to human (n=1)
2	COVID-19 Alpha, Delta,	6.33 (3.58– 11.13)	3.42 (2.78–4.19)	1.8 (0.4–3.4)	87.5 (61.5– 100.0)	Respiratory (n=3)

	Omicron					
3	COVID-19 WT, A/H1N1/09, A/H1N1, A/H2N2, A/H3N2, SARS-CoV-1	2.03 (1.38–3.19)	4.41 (2.75–13.06)	3.1 (0.0–10.6)	7.7 (0.4– 50.0)	Respiratory (n=6)
4	EBOV, MARV, NiV	0.93 (0.12–2.10)	12.53 (3.77–17.60)	61.3 (58.3–64.4)	0.0 (0.0–0.0)	Direct contact (n=3) Animal to human (n=3)
5	A/H5N1, MERS-CoV	0.50 (0.02–1.22)	9.69 (2.38–13.28)	46.3 (36.5–56.1)	0.02 (0.0–0.10)	Respiratory (n=2) Animal to human (n=2)
6	LASV, MPV	1.89 (0.07 - 7.03)	10.12 (0.99 - 33.23)	18.6 (3.0– 35.5)	76.2 (73.1–79.2)	Direct contact (n=2) Animal to human (n=2)

3.4 Archetype characterisation

3.4.1 Vector-borne pathogens with secondary transmission routes

Archetype 1 consists of two vector-borne pathogens, CCHFV and ZIKV. This group is characterised by a very low reproductive number, with a mean human-to-human R_0 of 0.07 (95% CI: 0.00–0.39), and a long serial interval of 12.03 days (95% CI: 3.85–23.66). CFR is moderately low at 5.9% (95% CI: 0.0–13.4), and presymptomatic transmission is very low (0.09%).

3.4.2 Airborne, high transmission pathogens with high presymptomatic transmission
Archetype 2 is composed of the SARS-CoV-2 variants Alpha, Delta, and Omicron. This group has the highest reproductive number of all archetypes, with a mean R_0 of 6.33 (95% CI: 3.58–11.13), and the shortest serial interval at 3.42 days (95% CI: 2.78–4.19). It has a low CFR of 1.8% (95% CI: 0.4–3.4) and a high proportion of presymptomatic transmission, estimated at 87.5% (95% CI: 61.5–100.0).

3.4.3 Airborne, high transmission pathogens

Archetype 3 brings together SARS-CoV-1, the wild-type strain of SARS-CoV-2, and four influenza strains (A/H1N1/09, A/H1N1, A/H2N2, A/H3N2). These pathogens are characterised by a moderately high reproductive number (mean R_0 of 2.03; 95% CI: 1.38–3.19) and a short serial interval of 4.41 days (95% CI: 2.75–13.06). The CFR is low at 3.1% (95% CI: 0.0–10.6). In contrast to Archetype 2, presymptomatic transmission is lower at 7.7% (95% CI: 0.4–50.0).

3.4.4 Contact zoonoses with high CFR

Archetype 4 includes the filoviruses (EBOV, MARV) and NiV. This group is defined by a high CFR, with a mean of 61.3% (95% CI: 58.3–64.4). These pathogens have limited human-to-human transmissibility, with an average R_0 of 0.93 (95% CI: 0.12–2.10), and a long serial interval of 12.53 days (95% CI: 3.77–17.60). Presymptomatic transmission is absent (0.0%).

3.4.5 Respiratory zoonoses with high CFR

Archetype 5 contains two respiratory zoonotic pathogens, A/H5N1 and MERS-CoV. This group is distinguished by a high CFR of 46.3% (95% CI: 36.5–56.1) combined with a respiratory transmission route. Human-to-human transmission is limited, as indicated by a mean R_0 of 0.50 (95% CI: 0.02–1.22). The serial interval is long, at 9.7 days (95% CI: 2.38–13.28), and the proportion of presymptomatic transmission is negligible (0.02%).

3.4.6 Contact zoonoses with evidence of presymptomatic transmission

Archetype 6 is composed of LASV and MPV. These contact-based zoonoses are characterised by a high CFR of 18.6% (95% CI: 3.0–35.5) and a mean R_0 of 1.89 (95% CI: 0.07–7.03). The serial interval is long, at 10.1 days. The most notable feature of this archetype is the significant proportion of presymptomatic transmission, estimated at 76.2% (95% CI: 73.1–79.2), which distinguishes it from the other high-CFR zoonotic groups.

3.4.7 Sensitivity Analysis

To evaluate the robustness of our clustering method we tested the stability of the original pathogen archetypes against changes in the included parameters. We performed two analyses: one that included additional epidemiological parameters (*k,* incubation, latent and infectious period) alongside the original set, and another that excluded the proportion of presymptomatic transmission (**Supplementary Figures S6** and **S7**, respectively). The results demonstrate that the clusters are robust to these modifications. EBOV and MARV, SARS-CoV-2 (Delta) and SARS-CoV-2 (Omicron), and the A/H1N1 influenza strains, remained consistently grouped together across both analyses. This would indicate that the generated archetypes reflect fundamental similarities in the pathogens' epidemiology.

In addition, we assessed the flexibility of the framework by applying it to a more diverse range of pathogens using a core set of widely applicable parameters: R0, incubation period, CFR, and transmission route. The framework incorporated and classified a wide variety of additional pathogens (**Supplementary Figure S9**).

---

4. Discussion

We reviewed key epidemiological parameters for 19 pathogens with pandemic potential and applied clustering algorithms to identify pathogen archetypes that share similar characteristics. Our findings suggest that grouping pathogens based on transmission traits such as R_0, serial interval, CFR, presymptomatic transmission, and transmission route could provide a pragmatic approach to pandemic preparedness.

The most frequently reported parameters were the incubation period, reproduction number,
and CFR. However, data availability was uneven, with SARS-CoV-2 and influenza
accounting for nearly half of all estimates. Parameter estimates varied both across
pathogens and within studies of the same pathogen, aligning with previous reviews [15].
Notably, R_0, serial interval, and CFR estimates were highly context dependent. For
example, MERS-CoV R estimates ranged from 5.4 (95% CI: 4.61-6.19) in an uncontrolled
hospital outbreak to 0.14 (95% CI: 0.04-0.26) with control measures in place [16]. Similarly,
influenza A/H1N1 R_0 estimates were higher in confined settings compared to overall
estimates [17]. Likewise, EBOV estimates varied by country during the 2013-2016 epidemic
[18]. Serial interval estimates also decreased when control measures were implemented,
with the serial interval of SARS-CoV-2 decreasing post epidemic peak in China correlating
with decreased time to isolation [19].

There were considerable differences in CFR estimates between outbreaks of the same
pathogen. Influenza A/H5N1 varies by clade [20]. MPV varies when hospital care is available
[21] or when comparing outcomes from outbreaks in Africa to outbreaks in the United States
[22]. Varying estimates for NiV highlights how CFR can vary depending on country, strain
and the control measures implemented, with the CFR being lower in Singapore (1999)
compared to Malaysia (1998-1999) [23]. These examples illustrate that parameter estimates
are generated across a wide range of contexts, and the importance of contextualising
parameter estimates when applying them to modelling efforts.

Our clustering analysis identified six pathogen archetypes, each with shared characteristics
that could inform the development of group-based, rather than pathogen-specific, control
strategies. Given that key interventions like contact tracing and case isolation are directly
influenced by parameters such as R_0, the serial interval, and the proportion of
presymptomatic transmission [14,24]. Pathogen-specific planning however, is and will
remain useful. Clustering pathogens with shared epidemiology would allow for specific plans
to be adapted for pathogens that share similar characteristics. A dual approach would ideally
allow for both in depth preparation for known risks and breadth in preparedness for a wider
range of pathogens.

A limitation of our approach is that its classifications are entirely dependent on the quality
and context of the input parameters. The resulting archetypes may group pathogens whose
real-world transmission dynamics differ in crucial ways; therefore, interpretation of clusters
should always be done with caution. This is particularly evident in the model's estimates for
presymptomatic transmission. For the high-transmission COVID-19 variants (Archetype 2),
our estimate of 87.5% (95% CI: 61.5–100.0) is on the higher end of published estimates of
40-80% [25,26]. The values for cluster 6 (MPV and LASV) is markedly higher than published
estimates. There is some evidence to suggest presymptomatic transmission is a factor in
MPV transmission dynamics, with estimates around 53% [27], its significance for LASV
however remains uncertain.

This discrepancy likely stems from the definition used in our analysis: the probability that the
serial interval is shorter than the incubation period (P(SI < IP)). This calculation is a proxy
and is highly sensitive to the underlying distributions [28]. The available distribution data for
some pathogens, potentially skewed by inconsistent symptom onset reporting in source

studies, may not perfectly reflect the true timeline of infectiousness. This can lead to a P(SI < IP) calculation that may overestimate the true biological extent of presymptomatic spread.

Furthermore, the context of the source data is highly important. The R□ estimate for LASV used in this analysis was derived from a hospital outbreak setting [29]. Nosocomial R□ values often fail to reflect community-level transmissibility. Therefore, the clustering of LASV with MPV a pathogen with more documented sustained community transmission may imply a higher risk for human-to-human LASV transmission than is typically observed in endemic settings, where transmission is primarily zoonotic with occasional, self-limiting outbreaks [30]. Historically however, the majority introductions of MPV into human populations failed to generate a secondary case, which would be more in line with LASV [31]. Therefore, this grouping is not as counterintuitive as it might seem in light of the 2022 Mpox outbreaks. These discrepancies highlight the sensitivity to input data and underscore the need for robust; parameter estimates to refine these classifications.

Our preference was to use synthesised estimates from systematic reviews wherever available. This approach provides a generalised baseline for each pathogen, preventing the analysis from being skewed by the unique characteristics of a single outbreak. However, this method may inadvertently incorporate biases inherent in the aggregated data, such as those from early-outbreak reporting, or mask context-specific variations.

The estimates for the serial interval and R0 for LASV and CCHFV respectively are subject to considerable uncertainty. Human-to-human transmission for these pathogens is best studied in hospital settings, where nosocomial outbreaks have been documented. Identifying transmission for these pathogens in community or household settings is particularly difficult as it is difficult to distinguish between vector-to-human from human-to-human transmission. As a result, estimation of transmission parameters will largely rely on hospital-based outbreaks, as used in this study. Whilst these estimates are subject to significant uncertainty, they provide a feasible basis for inference.

These findings highlight the potential for this framework to serve as an adaptable tool for classifying and assessing pathogens beyond those included in this study. We extended the number of pathogens in the sensitivity analysis (**Supplementary Figure S9**) to include pathogens with different characteristics to the original selection. Highlighting that the framework could be used to classify food/water-borne pathogens, bioterrorism-related pathogens and pathogens with extended delay periods such as HIV. Future work should address the impact of key interventions across each archetype. Moreover, the framework should be continuously updated as new epidemiological data emerge, refining the clustering methods and integrating additional pathogens, including both novel and emerging threats. To that end, the development of a global, standardised repository for epidemiological parameters would enable the rapid integration of data on novel and emerging threats, allowing for actionable guidance in regard to pandemic preparedness.

5. Conclusion

Documenting epidemiological parameters is crucial for effective outbreak risk analysis. We provide 302 parameter estimates for 19 pathogens, offering a valuable foundation for modelling their spread and containment. However, key transmission parameters such as the dispersion parameter and latent period remain underreported, highlighting the need for further research to strengthen outbreak preparedness. Our clustering approach demonstrates a practical framework for evaluating plausible parameter ranges across groups of similar pathogens. By maintaining a dynamic classification system, public health preparedness efforts can shift away from a reactive, pathogen-specific focus toward a more anticipatory, trait-based strategy for managing future infectious disease risks.

Table 1. Epidemiological parameters of pandemic potential pathogens				
Pathogen	**Parameter**	**Description**	**Value**	**Reference**
A/H1N1	Basic reproduction number	Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. All waves	Median: 1.8 (IQR: 1.47-2.27)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. Confined settings	Median: 3.82 (IQR: 2.68-4.84)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 1st wave	Median: 1.81 (IQR: 1.50-2.28)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 2nd wave	Median:1.73 (IQR: 1.39-2.33)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median	Median:1.70 (IQR: 1.55-1.76)	Biggerstaff et al. 2014 [17]

		estimates. 3rd wave		
	Dispersion parameter	Modelling, estimated using time series analysis, using a large household survey dataset conducted in Maryland late 1918	0.94 (95% CI: 0.59-1.72)	Fraser et al. 2011 [32]
	Serial interval (d)	Median of the mean generation time or serial interval used to estimate reproduction number.	Median: 3.3 (range: 1.5-6.0)	Biggerstaff et al. 2014 [17]
		Systematic review for serial interval estimates for influenza up to 2013.	6 studies with a range of 1.9 to 8.28 days	Vink et al. 2014 [33]
	Incubation period (d)	Modelling study, estimated through the re-analysis of daily incidence data of cases on ships departing Australia in 1919	Lognormal distribution with mean of 1.34	Nishiura 2007 [34]
	Latent period (d)	Modelling study, viral excretion profile over time used directly to estimate latent period. (H1N1/H3N2)	Weibull distribution with mean of 1.60 (95% CI: 1.50-1.70)	Cori et al. 2012 [35]
	Infectious period (d)	Modelling study, viral excretion profile over time used directly to estimate infectious period. (H1N1/H3N2)	Weibull distribution with mean of 1.0 (95% CI: 0.50-1.70)	Cori et al. 2012 [35]
	Case fatality risk (%)	Modelling, referenced range of CFR's for H1N1 pandemic.	1-4	Carrat et al. 2006 [36]
		Pandemic Influenza Risk Management

WHO Guidance. Estimated value (for 1918 pandemic) | 2-3 | WHO 2017 [37] |
| A/H1N1pdm09 | Basic reproduction number | Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. | Median: 1.46 (IQR: 1.30-1.70) | Biggerstaff et al. 2014 [17] |
| | | Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median | Median: 1.96 (IQR: 1.50-2.23) | Biggerstaff et al. 2014 [17] |

		estimates. Confined settings		
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 1st wave	Median:1.47 (IQR: 1.31-1.71)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 2nd wave	Median: 1.48 (IQR: 1.30-1.66)	Biggerstaff et al. 2014 [17]
	Dispersion parameter	Modelling seasonal influenza in Switzerland from 2003 to 2015	Mean: 7.38 (95% CI: 5.30–10.64)	Brugger and Althaus 2020 [38]
	Serial interval (d)	Median of the mean generation time or serial interval used to estimate reproduction number.	Median: 2.8. (range: 1.90-7.0)	Biggerstaff et al. 2014 [17]
		Systematic review for serial interval estimates for influenza up to 2013.	36 studies ranging from 1.9 to 5 days	Vink et al. 2014 [33]
		Estimates from household datasets with information on symptom onset dates.	Mean ranged from 1.7 to 3.7 days, with a pooled mean of 2.8	Vink et al. 2014 [33]
		Serial interval estimated from model fit of the serial interval to index case–to–case interval data	Normal distribution with mean of 2.1	Vink et al. 2014 [33]
	Incubation period (d)	Modeling study, estimated using laboratory-confirmed swine influenza case-information in the UK 2009	Weibull distribution with mean of 1.66 (95% CI: 1.42-1.90)	Tom et al. 2010 [39]
		Modeling study, estimated using laboratory-confirmed swine influenza case-information in the UK 2009	Gamma distribution with mean of 1.65 (95% CI: 1.41-1.89)	Tom et al. 2010 [39]
		EpiLPS - Dataset from Lessler et al. 2009 [40]	Semipar. Distribution with mean of 2.0 (95%	Estimated

			CrI:1.80-2.10)	
		Modelling study - Outbreak of 2009 Pandemic Influenza A (H1N1) at a New York City School	Median: 1.4 days (95% CI: 1.0–1.8)	Lessler et al. 2009 [41]
	Latent period (d)	Modelling study, estimated using data on laboratory-confirmed cases of pandemic H1N1 influenza reported in Ontario, Canada, between Apr. 13 and June 20, 2009	Mean: 2.62 (95% CI: 2.28-3.12)	Tuite et al. 2010 [42]
	Infectious period (d)	Modelling study, estimated using data on laboratory-confirmed cases of pandemic H1N1 influenza reported in Ontario, Canada, between Apr. 13 and June 20, 2009	Mean: 3.38 (95% CI: 2.06-4.69)	Tuite et al. 2010 [42]
	Case fatality risk (%)	Pandemic Influenza Risk Management WHO Guidance. Estimated value. 2009 pandemic	0.02	WHO 2017 [37]
		Sero-epidemiological study. Prevalence of cross‑reactive antibodies to H1N1pdm virus and rates of H1N1pdm infection	0.02	Van Kerkhove et al. 2013 [43]
	Infection fatality risk	Systematic review of published estimates of the case fatality risk of H1N1pdm09 up to 2013. Laboratory-confirmed cases	Point estimates 0–13,500 deaths per 100,000 cases	Wong et al. 2013 [44]
		Symptomatic cases	Point estimates 0–1,200 per 100,000 cases	Wong et al. 2013 [44]
		Infections	Point estimates 1–10 per 100,000 infections	Wong et al. 2013 [44]
		Symptomatic cases in children	One death per 100,000 symptomatic cases in children	Wong et al. 2013 [44]

		Symptomatic cases in the elderly	Approximately 1,000 deaths per 100,000 symptomatic cases in the elderly,	Wong et al. 2013 [44]
A/H2N2	Basic reproduction number	Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates.	Median: 1.65 (IQR: 1.53-1.70)	Biggerstaff et al. 2014 [17]
	Serial interval (d)	Median of the mean generation time or serial interval used to estimate reproduction number.	Median: 3.5 (Range: 2.60-4.10)	Biggerstaff et al. 2014 [17]
	Incubation period (d)	Systematic review of volunteer challenge studies	Median: 2 (IQR: 2.00-2.50)	Carrat et al. 2008 [45]
	Latent period (d)	Modelling study, assumed value	Mean: 1.9	Elveback et al. 1976 [46]
	Infectious period (d)	Modelling study, assumed value	Mean: 4.1	Elveback et al. 1976 [46]
	Case fatality risk (%)	Pandemic Influenza Risk Management WHO Guidance. Estimated value. (1957 pandemic)	0.2	WHO 2017 [37]
A/H3N2	Basic reproduction number	Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates.	Median: 1.80 (IQR: 1.56-1.85)	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. Confined settings	Median: 1.39	Biggerstaff et al. 2014 [17]
		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 1st wave	Median: 1.56	Biggerstaff et al. 2014 [17]

		Systematic review of the reproduction number for pandemic influenza. Combined estimates for the basic and effective reproductive numbers and presenting median estimates. 2nd wave	Median: 1.68	Biggerstaff et al. 2014 [17]
	Serial interval (d)	Median of the mean generation time or serial interval used to estimate reproduction number.	Median: 4.0 (Range: 2.95-4.10)	Biggerstaff et al. 2014 [17]
		Systematic review for serial interval estimates for influenza up to 2013.	Mean values of 3.1 and 3.4	Vink et al. 2014 [33]
		Estimates from household datasets with information on symptom onset dates.	Normal distribution with mean of 2.2	Vink et al. 2014 [33]
	Incubation period (d)	Modelling study, estimated through analysis of an outbreak of influenza aboard a commercial airliner	Weibull distribution with mean of 1.48	Ferguson et al. 2005 [47]
	Latent period (d)	Modelling. Assumed value. Assumed to be same length as incubation period	Weibull distribution with mean of 1.48	Ferguson et al. 2005 [47]
	Infectious period (d)	Modelling. Transmission model to estimate the main characteristics of influenza transmission in households	Gamma distribution with mean of 3.80 (95% CI: 3.10-4.60)	Cauchemez et al. 2004 [48]
		Modelling. Transmission model to estimate the main characteristics of influenza transmission in households (Children)	Gamma distribution with mean of 3.60 (95% CI: 2.30-5.20)	Cauchemez et al. 2004 [48]
		Modelling. Transmission model to estimate the main characteristics of influenza transmission in households (Adults)	Gamma distribution with mean of 3.9 (95% CI: 3.20-4.90)	Cauchemez et al. 2004 [48]
	Case fatality risk (%)	Pandemic Influenza Risk Management WHO Guidance. Estimated value. (1964 pandemic)	0.2	WHO 2017 [37]
A/H5N1	Basic reproduction number	Modelling study, estimated Re in Vietnam 2004–2006.	0.0 (95% CI: 0.0-0.42)	Bettencourt and Ribeiro 2008 [49]

		Modelling study, estimated Re in Indonesia 2004–2006.	0.0 (95% CI: 0.0-0.0)	Bettencourt and Ribeiro 2008 [49]
		Modelling study, estimated Re in Indonesia 2005–2009.	0.1-0.25 (95% CI: 0.0-0.40)	Aditama et al. 2012 [50]
		Modelling study, estimated lower limit on the local R0 in a household outbreak in Indonesia 2006.	1.14 (95% CI: 0.61-2.14)	Yang et al. 2007 [51]
		Modelling. Estimated from previous human to human cases.	0.06 (95% CI: 0.01-0.2)	Ferguson et al. 2004 [52]
	Serial interval (d)	Median of the mean value of generation time or serial interval used to estimate reproduction number.	Mean: 7.8 (range: 6.0-9.5)	Biggerstaff et al. 2014 [17]
		Modelling. Estimated by fitting case data from Aditama et al. 2012. Preprint	Median: 6.80 (95% CrI: 0.3–13.3)	Ward et al. 2024 [53]
	Incubation period (d)	Review, median value from outbreaks in Thailand and Vietnam 2004	Median: 4.0 (range: 2.0-8.0)	Beigel et al. 2005 [54]
		Estimated value. Retrospective descriptive study of 24 human cases in China 1997-2008	Median: 5.0 (range: 2.0-9.50)	Huai et al. 2008 [55]
		Estimated value, using survival analysis techniques. 43 human cases in China	Weibull distribution with mean of 3.30 (95% CI: 2.70-3.90)	Cowling et al. 2013 [56]
		Estimated value, 8 human cases in Eastern Turkey in 2006	Mean: 5 (Range: 4.0-7.0)	Oner et al 2006 [57]
	Infectious period (d)	Modelling. Assumed value. Family Cluster, Indonesia 2006	Uniform distribution with mean of 9 (range 5-13)	Yang et al. 2007 [51]
	Case fatality risk (%)	Systematic Review. Crude CFR 1997-2009	Median: 56.3% (IQR: 32.5–77.8)	Van Kerkhove et al. 2011 [58]
		Systematic review of individual case data. 1997-2015 (Overall)	53.5	Lai et al. 2016 [20]

		Systematic review of individual case data. 1997-2015 (Clade 0)	31.6	Lai et al. 2016 [20]
		Systematic review of individual case data. 1997-2015 (Clade 1)	58.6	Lai et al. 2016 [20]
		Systematic review of individual case data. 1997-2015 (Clade 2.1)	84.6	Lai et al. 2016 [20]
		Systematic review of individual case data. 1997-2015 (Clade 2.2)	33.2	Lai et al. 2016 [20]
		Systematic review of individual case data. 1997-2015 (Clade 2.3)	61.8	Lai et al. 2016 [20]
		Systematic review of individual case data. 1997-2015 (Clade 7)	100	Lai et al. 2016 [20]
	Infection fatality risk (%)	Adjusted CFR based on surveillance and seroprevalence studies	14-33	Li et al. 2008 [59]
CCHFV	Basic reproduction number	Estimated using epichains. Using data from cases of CCHFV infected in the European union/European Economic Area from 2013–2024.	Median: 0.03 (95% CrI: 0.004–0.09)	Estimated
	Serial interval (d)	Estimated from fitting outbreak data reporting the interval between the onset of illness in successive cases.	Gamma distribution with mean of 12.0 days (95% CrI: 3.0–27.2)	Estimated
	Incubation period (d)	EpiLPS – Dataset collected from review.	Gamma distribution with mean of 5.70 (95% CrI: 5.30-6.00)	Estimated
	Case fatality risk (%)	Review of published reports of CCHF in Europe, Asia, middle east and Africa. 1944-2010	Mean: 30.6	Bente et al. 2013 [60]
		Systematic review, CFR worldwide of confirmed cases 1948-2018	Mean: 19.9 (IQR: 8-32)	Belhadi et al. 2022 [61]

		Systematic review, Overall CFR (%) with ongoing CCHF infection up to 2020	11.7% (95% CI: 9.1-14.5)	Belobo et al. 2021 [62]	
		Systematic review, fatality rate in the Arab world 1978 to 2021	Mean: 29 (range: 24-61)	Perveen and Gulfaraz Khan 2022 [63]	
		Systematic review, CFR calculated from annual cases from 1944 to 2017	Mean: 32.2	Nasirian 2020 [64]	
EBOV	Basic reproduction number	Systematic review. R0 reported as a range of central estimates, from database inception up to July 2023	0.05-12.00	Nash et al. 2024 [15]	
		Systematic review. Pooled mean of Ebola R0 in African countries from 1976 to February 2023.	Mean: 1.95 (95% CI: 1.74-2.15)	Muzembo et al. 2024 [18]	
		Transmission in hospitals and funeral rites during the 2013–2016 Ebola epidemic in West Africa (overall basic reproductive number)	Mean: 1.8 (range: 1.5-2)	Muzembo et al. 2024 [18]	
		Systematic review, Zaire ebolavirus,2013-2016 epidemic in Nigeria (pooled mean).	Mean: 9.38 (95% CI: 4.16-14.59)	Muzembo et al. 2024 [18]	
		Systematic review, Zaire ebolavirus,2013-2016 epidemic in DRC (pooled mean)	Mean: 3.31 (95% CI: 2.3-4.32)	Muzembo et al. 2024 [18]	
		Systematic review, Sudan ebolavirus, 2000 outbreak in Uganda (pooled mean Ebola).	Mean: 2.0 (95% CI: 1.25-2.76)	Muzembo et al. 2024 [18]	
		Systematic review, Zaire ebolavirus, 2013-2016 epidemic in Liberia (pooled mean)	Mean: 1.83 (95% CI: 1.61-2.05)	Muzembo et al. 2024 [18]	
		Systematic review, Zaire ebolavirus,2013-2016 epidemic in Sierra Leonne (pooled mean)	Mean: 1.73 (95% CI: 1.47-2.0)	Muzembo et al. 2024 [18]	
		Systematic review, Zaire ebolavirus,2013-2016 epidemic in Guinea (pooled mean	Mean: 1.44 (95% CI: 1.29-1.6)	Muzembo et al. 2024 [18]	
		Systematic review of early modelling studies. 2013-2016 Epidemic in four West African	Median: 1.78 (IQR: 1.44-	Wong et al. 2017 [65]	

		countries. median of the R0 means reported.	1.78)	
	Dispersion parameter	Systematic review. K reported as a range of central estimates, up to July 2023	0.02-2.20	Nash et al. 2024 [15]
	Serial interval (d)	Systematic review, up to July 2023 (Pooled value).	Mean: 15.4 (95% CI: 13.20-17.50)	Nash et al. 2024 [15]
		Systematic review of early modelling studies. 2013-2016 Epidemics in four West African countries. median of the means reported.	Median: 14.35 (IQR: 12.28-16.35)	Wong et al. 2017 [65]
	Incubation period (d)	Systematic review up to July 2023. Pooled random effect	Mean: 8.5 (95% CI: 7.70-9.20)	Nash et al. 2024 [15]
	Latent period (d)	Systematic review- 1976 to 2000. Data from individuals with single-day exposures	Mean: 12.27	Velásquez et al. 2015 [66]
		Systematic review of early modelling studies. 2013-2016 Epidemic in four West African countries. median of the means reported.	Median: 9.7 (IQR: 8.8-10.38)	Wong et al. 2017 [65]
		Systematic review up to July 2023. Pooled random effect	0.10-31.20	Nash et al. 2024 [15]
	Infectious period (d)	Systematic review up to July 2023. Pooled random effect and range of central estimates	Mean: 5.0 (95% CI: 3.70-6.30)	Nash et al. 2024 [15]
		Systematic review of early modelling studies. 2013-2016 Epidemic in four West African countries. median of the means reported	Median: 7 (IQR: 4-10)	Wong et al. 2017 [65]
		Systematic review- 1976 to 2000. data from individuals with single-day exposures (Survivors)	Mean:9.4	Velásquez et al. 2015 [66]
		Systematic review- 1976 to 2000. data from individuals with single-day exposures (fatal infections)	Mean: 5.33	Velásquez et al. 2015 [66]
	Case fatality risk (%)	Systematic review up to July 2023. Mean CFR across all estimates	Mean: 57.8	Nash et al. 2024 [15]

			Systematic review of Asian and African countries between 1999 and June 2021, estimated pooled case fatality rates	61.1 (95% CI: 50.26-71.85)	Khan et al. 2022 [67]
LASV	Effective reproduction number		Modelling. Estimated from published outbreaks and the number of LF hospitalized patients to Kenema Government Hospital in Sierra Leone. Pure human to human transmission	0.73	Lo Iacono et al. 2015 [29]
	Serial interval (d)		Estimated using Lo Iacono et al. 2015 [29] data combined with outbreak data. fitted to a lognormal distribution	Lognormal distribution with mean of 11.5 days (95% CrI: 0.9-34.6)	Estimated
			Modelling study, referenced value (Lo Iacono et al. 2015 [29])	Gamma distribution with mean of 7.8	Zhao et al. 2020 [68]
	Incubation period (d)		Systematic review, range of central estimates reported	Range: 7.0-12.80	Doohan et al. 2024 [69]
			EpiLPS – Dataset from Akhmetzhanov et al. 2019 [70]	Lognormal distribution with mean of 12.6 (95% CrI: 11.8-13.8)	Estimated
	Latent period (d)		Modelling study, Referenced value	Lognormal distribution with mean of 10.0 (Range: 5-21)	Tuite et al. 2019 [71]
	Infectious period (d)		Modelling study, Referenced value	Lognormal distribution with mean of 10.0 (Range: 6-17)	Tuite et al. 2019 [71]
	Case fatality risk (%)		Systematic review, CFR for imported Lassa fever cases in non-endemic countries outside West Africa. 1969-2019	35.1	Wolf et al. 2020 [30]
			Systematic review, overall fatality rate in sub-Saharan Africa 1972-2020	29.7 (95% CI: 22.3-37.5)	Kenmoe et al. 2020 [72]
			Systematic review, pooled estimate up to 2023	33.1 (95% CI: 25.7-41.5)	Doohan et al. 2024 [69]
	Infection fatality		Crude estimate of the overall	1%	Dwalu et al.

	risk (%)	case-fatality rate		2024 [73]
MARV	Basic reproduction number	Modelling study, R0 estimated using previous Marburg outbreaks	Median: 0.81 (95% CI: 0.08-1.83)	Qian et al. 2023 [74]
		Modelling study, R0 estimated for the 2005 epidemic in Angola	1.59 (95% CI: 1.53–1.66)	Ajelli and Merle 2012 [75]
	Dispersion parameter	Modelling study, K estimated from 18 chains of transmission from the outbreak in DRC.	negative binomial distribution with a range of 0.52-0.67	Qian et al. 2023 [74]
	Serial interval (d)	Modelling. Estimated using Identified discernible infector-infectee pairs from line list data and obtained the difference between the dates of infection of each pair.	Gamma distribution with mean of 9.2	Qian et al. 2023 [74]
	Incubation period (d)	Modelling study, estimated range of cental values	5.0-10.0	Qian et al. 2023 [74]
		EpiLPS - Dataset from Pavlin 2014 [76]	Weibull distribution with mean of 6.90 (95% CrI: 6.20-7.60)	Estimated
		Modelling study, estimated from pooled data from all Marburg cases between 1967 and 2008	Median: 7 (range: 2.0-13.0)	Pavlin 2014 [76]
	Latent period (d)	Modelling study, latent period estimated using the average viral load in non-human primates	Mean: 3	Ajelli and Merle 2012 [75]
		Modelling study, estimated by fitting the epidemic curve of MARV cases during the epidemic in Angola (assumes incubation period is same length as latent period)	Mean: 6.50 (95% CI: 6.0-7.0)	Bettencourt 2009 [77]
	Infectious period (d)	Modelling study, estimated by fitting the epidemic curve of MARV cases during the epidemic in Angola	Mean: 3.0 (95% CI: 3.0-4.0)	Bettencourt 2009 [77]
	Case fatality risk (%)	Systematic review, from database inception to March 2023 (estimated pooled total random)	61.9 (95% CI: 38.8-80.6)	Cuomo-Dannenburg et al 2024 [78]

MERS-CoV	Basic reproduction number	Systematic review, Saudi Arabia or Middle East area data	Range: 0.45-0.98	Park et al. 2018 [16]
		Systematic review, South Korea data (Early stage)	Range: 2.5–8.09	Park et al. 2018 [16]
		Modelling. Referenced value derived from Cauchemez et al. 2014	Median: 0.95 (95% CI: 0.6-1.3)	Peak et al. 2017 [79]
	Dispersion parameter	Systematic review, point estimates of k	Range: 0.06 (95% CI: 0.03-0.09) to 2.94 (95% CI: 0.23-infinity)	Wang et al. 2021 [80]
	Serial interval (d)	Modelling. Outbreak in a hospital in Saudi Arabia. Estimated using symptom onset times in the patient of infected–infector pairs. Outbreak period covers interventions being introduced	Lognormal distribution with mean of 7.60 (95% CI: 2.50-23.10)	Assiri et al. 2013 [81]
		Modelling study, Outbreak in two hospitals in South Korea. Estimated using symptom onset times in the patient of infected–infector pairs. Outbreak period covers interventions being introduced	Gamma distribution with median of 14.60 (95% CI: 12.90-16.50)	Park et al. 2016 [82]
		Modelling study, estimated by analysing 119 cases in South Korea	Gamma distribution with mean of 12.60 (95% CI: 12.10-13.10)	Cowling et al. 2015 [83]
	Incubation period (d)	EpiLPS – Dataset from Cauchemez et al. 2014 [84]	Lognormal distribution with mean of 5.40 (95% CrI: 4.50-6.50)	Estimated
		Systematic review up to 2017	Range: 4.50-7.80	Park et al. 2018 [16]
	Latent period (d)	Modelling assumed value. Deemed unlikely that a case will cause any subsequent infections prior to 7 days after infection	7.0	Lessler et al. 2014 [85]

	Infectious period (d)	Modelling study. Assumed value. Maximum duration of infectiousness	Uniform distribution with median of 16.43 (95% CI: 9.59-24.5)	Peak et al. 2017 [79]
	Case fatality risk (%)	Systematic review, up to 2017. Mortality rate in south Korea	Range: 14.5–47.8	Park et al. 2018 [16]
		Systematic review, up to 2017. Mortality rate in Saudi Arabia	Range: 22–69.2	Park et al. 2018 [16]
		Systematic review, up to 2017. Mortality rate from multiple areas.	Range: 26.6 - 59.4	Park et al. 2018 [16]
Mpox	Basic reproduction number	Systematic review of the 2022 Mpox outbreak. Pooled mean	Mean: 1.8 (95% CI: 1.7-1.9)	Okoli et al. 2024 [86]
		Systematic review. Analysis of active surveillance data collected in the DRC between 1980 and 1984.	0.8	Beer et al. 2019 [87]
	Dispersion parameter	Modelling study, DRC 1980-1984. Estimated by analysing chain size data.	0.36 (95% CI: 0.14-1.47)	Blumberg and Lloyd-Smith 2013 [88]
		Modelling study 2022 Mpox outbreak. Estimated using genomic and epidemiological metadata.	0.3 (95% CI: 0.18-0.54)	Paredes et al. 2024 [89]
	Serial interval (d)	Systematic review of the 2022 Mpox outbreaks. Pooled mean.	8.5 (95% CI: 7.3-9.9)	Okoli et al. 2023 [90]
		Modelling study 2022 Mpox outbreak USA. May–August 2022.	Gamma distribution with mean of 8.5 (95% CI: 7.3-9.9)	Madewell et al. 2023 [91]
		Modelling study 2022 Mpox outbreak UK. PCR confirmed cases between 6 May and 1 August 2022.	Gamma distribution with mean of 8.0 (95% CI: 6.5-9.8)	Ward et al. 2022 [27]
	Incubation period (d)	Systematic review, analysis of 18,275 Mpox cases during the 2022 outbreak	Median: 7.0 (IQR: 3-21)	Chenchula et al. 2023 [92]
		Systematic review of the 2022 Mpox outbreaks. Pooled mean	Mean: 7.8 (95% CI: 6.6-	Okoli et al.

		(up to Dec 2022)	9.0)	2023 [90]
		Systematic review of the 2022 Mpox outbreaks. Pooled mean (up to May 2022)	Mean: 7.4 (95% CI: 6.4-8.4)	Okoli et al. 2023 [90]
		Systematic review of previous Mpox outbreaks. Pooled mean	Mean: 12.9 (95% CI: 10.4-15.5)	Okoli et al. 2023 [90]
		Systematic Review, analysis of outbreaks pre and post 2022	Mean: 7.9 (Range: 1-21)	Hatami et al. 2023 [93]
		Modelling study 2022 Mpox outbreak USA. May–August 2022	Lognormal distribution with mean of 5.6 (95% CI: 4.3-7.8)	Madewell et al. 2023 [91]
		EpiLPS – dataset from Miura et al. 2022 [94]	Weibull distribution with mean of 7.6 (95% CrI: 6.5-9.9)	Estimated
		Modelling study 2022 Mpox outbreak UK. PCR confirmed cases between 6 May and 1 August 2022.	Lognormal distribution with mean of 8.9 (95% CI: 7.9-9.9)	Ward et al. 2022 [27]
	Latent period (d)	Modelling study, assumed value from viral shedding data (Preprint)	3.0	Asakura et al. 2024 [95]
	Infectious period (d)	Modelling study, assumed value from documented duration of illness	21	Endo et al. 2022 [96]
		Modelling study, May-June 2022 MSM. Estimated infectious period while not refraining from sexual contacts (Netherlands)	6.0 (95% CI: 4.4-7.8)	Xiridou et al. 2023 [97]
		Modelling study, July 2022 MSM. Estimated infectious period while not refraining from sexual contacts (Netherlands)	2.6 (95% CI: 2.0-4.3)	Xiridou et al. 2023 [97]
		Modelling study, assumed value based on viral shedding data (Preprint)	10.0	Asakura et al. 2024 [95]
	Case fatality risk (%)	Systematic review - CFR of outbreaks. From discovery to 2019	Mean: 8.7 (95% CI: 7.0-10.8)	Bunge et al. 2022 [22]

		Systematic review - CFR of Central African clade outbreaks. From discovery to 2019	Mean:10.6 (95% CI: 8.4-13.3)	Bunge et al. 2022 [22]
		Systematic review - CFR of West African clade outbreaks. From discovery to 2019	Mean: 3.6 (95% CI: 1.7-6.8)	Bunge et al. 2022 [22]
		Systematic review - CFR of West African clade, African countries only. From discovery to 2019	Mean: 4.6 (95% CI: 2.1-8.6)	Bunge et al. 2022 [22]
		Systematic review 1950 to 2022. CFR when hospital care is available	0.03 (95% CI: 0.0-0.44)	DeWitt et al. 2022 [21]
		Systematic review 1980 to 2022, CFR in hospitalised patients	4 (95% CI:1-9%)	Benites-Zapata et al. 2022 [98]
NiV	Basic reproduction number	Review of hospital-based surveillance and outbreak investigations in Bangladesh from 2001 to 2014. Hospital-based surveillance implemented in 2007	0.33 (95% CI: 0.19-0.59)	Nikolay et al. 2019 [99]
		Review of hospital-based surveillance and outbreak investigations in Bangladesh. R0 for cases identified 2007-2014. Hospital-based surveillance implemented in 2007	0.23 (95% CI: 0.11-0.46)	Nikolay et al. 2019 [99]
		Review of hospital-based surveillance and outbreak investigations in Bagladesh. R0 for primary cases between 2001 and 2014. Hospital-based surveillance implemented in 2007	0.30 (95% CI: 0.15-0.61)	Nikolay et al. 2019 [99]
		Review of hospital-based surveillance and outbreak investigations in Bangladesh 2001-2014. R0 for cases hospitalized >7 days since symptom onset or not hospitalised. Hospital-based surveillance implemented in 2007	0.60 (95% CI: 0.07-4.97)	Nikolay et al. 2019 [99]
	Dispersion parameter	Modelling study, referenced value	0.06	Bradbury et al. 2023 [100]

	Serial interval (d)	Review of hospital-based surveillance and outbreak investigations in Bangladesh from 2001 to 2014. Serial interval estimated using epidemiologically linked transmission pairs. Bangladesh 2001-2014. Hospital-based surveillance implemented in 2007	Gamma distribution with mean of 12.7	Nikolay et al. 2019 [99]
	Incubation period (d)	EpiLPS – Dataset from Nikolay et al. 2019 [99]	Gamma distribution with mean of 9.4 (95% CrI: 8.7-10.1)	Estimated
		Systematic review up to 30 May 2019. Incubation period in the Philippines 2014	Median: 8.0 (Range: 4-20)	Hegde et al. 2023 [23]
		Systematic review up to 30 May 2019. Incubation period in Bangladesh	Median: 9.0 (Range: 6-14)	Hegde et al. 2023 [23]
		Systematic review up to 30 May 2019. Incubation period in India	Median: 10.0 (Range: 6-18)	Hegde et al. 2023 [23]
	Infectious period (d)	Review of hospital-based surveillance and outbreak investigations in Bangladesh from 2001 to 2014. Assumed maximum infectious period for contact tracing	15	Nikolay et al. 2019 [99]
	Case fatality risk (%)	Systematic review, overall random effect meta-analysis. 1999-2014 (Bangladesh, India, Malaysia, Singapore, Philippines)	61 (95% CI: 45.7-75.4)	Kenmoe et al. 2019 [101]
		Systematic review, Bangladesh random effect meta-analysis from Bangladesh 2001-2014	67.9 (95% CI: 47.7-85.4%)	Kenmoe et al. 2019 [101]
		Systematic review, random effect meta-analysis from India 2001	71.3 (95% CI: 63.2-78.8)	Kenmoe et al. 2019 [101]
		Systematic review, random effect meta-analysis from Malaysia 1998-1999	32.6% (95% CI: 25.8-39.8)	Kenmoe et al. 2019 [101]
		Systematic review, random effect meta-analysis from the Philippines 2014	81.8 (95% CI: 52.7-99.5)	Kenmoe et al. 2019 [101]
		Systematic review, random	2.9 (0.0-11.9)	Kenmoe et al.

		effect meta-analysis from Singapore 1999		2019 [101]
		Systematic review, CFR in Singapore 1999. Pig imports from Malaysia banned; abattoirs closed; preventive control measures in hospitals	8.3	Hegde et al. 2024 [23]
		Systematic review, CFR in Malaysia 1998-1999. Pig culling and transport ban; active surveillance for encephalitis cases; protective equipment for all persons who have exposure to pigs; education campaign	40	Hegde et al. 2024 [23]
		Systematic review, CFR in the Philippines 2014. Contact tracing implemented	53	Hegde et al. 2024 [23]
		Systematic review, CFR in Bangladesh 2001-2014	78	Hegde et al. 2024 [23]
		Systematic review, CFR in India 2001-2018	93	Hegde et al. 2024 [23]
		Systematic review, CFR in Bangladesh and Malaysia 1999-2016	61 (95% CI: 45.7-75.4)	Suman et al. 2024 [102]
RVF	Incubation period (d)	Systematic review of incubation periods up to 2011. Pooled estimate	Lognormal distribution with median of 4.0 (95% CI: 3.40-4.90)	Rudolph et al. 2014 [103]
		EpiLPS – Dataset collected from outbreak reports with reported exposure and symptom onset times	Semipar. with mean of 4.0 (95% CrI: 3.3-4.5)	Estimated
	Case fatality risk (%)	Systematic review, pooled estimate in Africa 1997-2020	27.5 (95% CI: 8.0-52.5)	Ebogo⬜Belobo et al. 2023 [104]
SARS-CoV-1	Basic reproduction number	Modelling study. Median estimate from study estimated using exponential doubling times of several epidemics in 2003. Without control measures. Data from Lipsitch et al. 2003 [105]	Mean: 2.9 (95% CI: 2.2-3.6)	Peak et al. 2017 [79]
		Modelling, R0 estimate at the start of the epidemic in Hong Kong (excluding superspreading events)	2.7 (95% C): 2.2 to 3.7)	Riley et al. 2003 [106]

		Modelling. Estimated R0 from outbreak at the National Taiwan University Hospital	Lognormal distribution with mean of 2.65	Chen et al. 2006 [107]
		Modelling, estimated R0 distribution for SARS	Median 1.1 (IQR: 0.43-2.41)	Chowell et al. 2004 [108]
		Modelling estimated R0 distribution Toronto 2003 (after implementing control measures). 77% of cases exposed in hospital setting	Median:0.58 (IQR: 0.24-1.18)	Chowell et al. 2004 [108]
		Modelling study, estimated R0 distribution Hong Kong 2003	Median: 1.1 (IQR: 0.44-2.29)	Chowell et al. 2004 [108]
		Modelling study, estimated R0 distribution Singapore 2003	Median: 1.17 (IQR: 0.47-2.47)	Chowell et al. 2004 [108]
	Dispersion parameter	Systematic review, range of k estimates	Range: 0.12 (90% CI: 0.08-0.42) to 0.20 (95% CI: 0.13-0.27)	Wang et al. 2021 [80]
	Serial interval (d)	Modelling study, estimated by analysing 205 probable cases reported in Singapore (after interventions were implemented)	Weibull distribution with mean of 8.4	Lipsitch et al. 2003 [105]
		Modelling study, estimated by analysing 205 probable cases reported in Singapore (before full-scale interventions)	Mean 10.0	Lipsitch et al. 2003 [105]
	Incubation period (d)	Systematic review, pooled analysis	Lognormal distribution with median of 4.0 (95% CI: 3.6-4.4)	Lessler et al. 2009 [40]
		EpiLPS – Dataset from Tsang et al. 2003 [109]	Semipar with mean of 4.70 (95% CrI: 3.90-5.50)	Estimated
	Latent period (d)	Modelling study referenced value. Assumed to be the same as the incubation period and a fixed value	6.5	Becker et al. 2005 [110]
		Modelling study. Assumed to be a fixed value.	6.81	Klinkenberg et al. 2006 [111]

		Modelling study assumed value. Average time of progression from latent infection to infectious	5.0	Lipsitch et al. 2003 [105]
	Infectious period (d)	Modelling study. Assumed value. Average duration of infectiousness	Mean: 5.0 (Range: 1-5)	Lipsitch et al. 2003 [105]
		Modelling study referenced value. Symptomatic period assumed to be infectious period	Gamma distribution with mean of 16.3	Lloyd-Smith et al. 2003 [112]
		Modelling study, estimated value	Gamma distribution with mean of 9.25	Fraser et al. 2004 [14]
		Modelling study referenced value. Effective infectious period	9.0	Becker et al. 2005 [110]
		Modelling study referenced value. Effective infectious period	3.87	Klinkenberg et al. 2006 [111]
	Case fatality risk (%)	Epidemiology review article	9.55 (95% CI: 8.94–10.2)	Hui et al. 2004 [113]
SARS-CoV-2 (Wild type) *	Basic reproduction number	Systematic review of R0 values from January 1 to August 31, 2020 (World)	Mean: 2.69 (95% CI: 2.40-2.98)	Ahammed et al. 2021 [114]
		Systematic review of R0 values from January 1 to August 31, 2020 (Asia)	Mean: 2.59 (95% CI: 2.19-2.94)	Ahammed et al. 2021 [114]
		Systematic review of R0 values from January 1 to August 31, 2020 (Europe)	Mean: 2.70 (95% CI: 2.26-3.13)	Ahammed et al. 2021 [114]
		Systematic review of R0 values from January 1 to August 31, 2020 (North America)	Mean: 3.69 (95% CI: 1.46-5.92)	Ahammed et al. 2021 [114]
		Systematic Review of transmission-Dynamic Models in Wuhan. R0 before the lockdown on the 23rd of January 2020	Median: 3.77 (IQR: 2.78-5.13)	Lin et al. 2020 [115]
		Systematic Review of transmission-Dynamic Models in Wuhan. R0 post lockdown	Median: 1.88 (IQR: 1.41-	Lin et al. 2020 [115]

		starting on the 23rd of January 2020	2.24)	
		Systematic review and meta-analysis of Epidemiologic, clinical, and laboratory findings. 1st December 2019 to 16th July 2020.	Mean: 3.32 (95% CI: 3.24-3.39)	Xie et al 2020 [116]
		Systematic Review and Meta-Analysis December 2019 up to March 2020	Mean: 2.99 (95% CI: 2.71-3.27)	Izadi et al. 2022 [117]
	Dispersion parameter	Systematic review of superspreading in a mix of countries from January to December 2020	Mean: 0.55 (95% CI: 0.30-0.79)	Du et al. 2022 [118]
	Serial interval (d)	Systematic review of serial intervals 2020-2023	Mean: 4.82 (95% CI: 4.50-5.14)	Xu et al. 2023 [19]
	Incubation period (d)	Systematic review of Incubation periods 2020-2023	Mean: 6.50 (95% CI: 5.88-7.12)	Xu et al. 2023 [19]
		EpiLPS - Using dataset from Backer et al. 2020 [119]	Lognormal distribution with mean of 4.40 (95% CrI: 4.0-4.80)	Estimated
		Modelling study, based on exposure information on COVID-19 cases in China	Weibull distribution with mean of 6.94	Xin et al. 2022 [120]
		Systematic Review of transmission-Dynamic Models in Wuhan. December 2019 and 21 February 2020	Median: 5.9 (IQR: 4.78-6.25)	Lin et al. 2020 [115]
	Latent period (d)	Modelling, estimated based on exposure information on COVID-19 cases in China	Gamma distribution with mean of 5.48 (95% CI: 5.06-5.86)	Xin et al. 2022 [116]
		Modelling, estimated based on exposure information on COVID-19 cases in China	Lognormal distribution with mean of 5.51	Xin et al. 2022 [116]
		Modelling, estimated based on exposure information on COVID-19 cases in China (symptomatic cases)	Gamma distribution with mean of 5.53 (95% CI: 5.09-5.99)	Xin et al. 2022 [116]

		Modelling, estimated based on exposure information on COVID-19 cases in China (asymptomatic cases)	Gamma distribution with mean of 5.24 (95% CI: 4.30-6.14)	Xin et al. 2022 [116]
	Infectious period (d)	Modelling, estimated from infectiousness profiles from a sample of 77 transmission pairs	Weibull distribution with mean of 9.3 (95% CI: 7.8-10)	He et al. 2020 [121]
		Modelling. Assumed that latent and the infectious period is approximately equal to the incubation period and the length of hospital stay.	12.53	Zhu et al. 2021 [122]
		Systematic Review of transmission-Dynamic Models in Wuhan December 2019 and 21 February 2020	Median: 9.94 (IQR: 3.93-13.5)	Lin et al. 2020 [115]
	Case fatality risk (%)	Systematic review of CFR's worldwide from January 1 to August 31, 2020	Mean: 2.67 (95% CI: 2.25-3.13)	Ahammed et al. 2021 [114]
		Systematic Review and Meta-Analysis December 2019 up to March 2020	Mean: 3.29	Izadi et al. 2020 [117]
		Systematic review and meta-analysis of Epidemiologic, clinical, and laboratory findings. 1st December 2019 to 16th July 2020	4.4	Xie et al 2020 [116]
		Systematic Review of transmission-Dynamic Models in Wuhan December 2019 and 21 February 2020	Median: 2.94 (IQR: 2.25-5.4)	Lin et al. 2020 [115]
	Infection fatality risk (%)	IFR from Japanese citizens who were evacuated from Wuhan on 29–31 January 2020	Range: 0.3-0.6	Nishiura et al. 2020 [123]
		Crude risk of death among all infected individuals in Wuhan city January-February, 2020	0.07 (95% CI: 0.05-0.09)	Mizumoto et al. 2020 [124]
		Estimating infection fatality ratio accounting for seroreversion. During the first wave of COVID-19 across different settings	0.49–2.53	Brazeau et al. 2022 [125]

SARS-CoV-2 (Alpha)	Basic reproduction number	Modelling, transmissibility model of Alpha variant in the United Kingdom fitted until the 24th of December 2020.	Alpha variant has a 43 to 90% higher reproduction number	Davies et al. 2021 [126]
	Serial interval (d)	Systematic review of serial intervals 2020-2023	Mean: 3.47 (95% CI: 2.52-4.41)	Xu et al. 2023 [19]
		Modelling. Estimated using household clusters from week 12/2021 until 22/2021 in Germany	Gamma distribution with mean of 4.5 (95% CI: 4.46-4.54)	Heiden and Buchholz 2022 [127]
	Incubation period (d)	Systematic review of Incubation periods 2020-2023	Mean: 4.92 (95% CI: 4.53-5.30)	Xu et al. 2023 [19]
	Latent period (d)	Epidemiological analysis. Assumed value as the ratio of mean duration of the latent and incubation periods	Median: (95% CI: 0.23 (0.04-0.50)	Hart et al. 2022 [128]
	Infectious period (d)	Epidemiological analysis. Assumed value of symptomatic infectious period	Gamma distribution with a mean of 3.5 (95% CI: 1.9-5.8)	Hart et al. 2022 [128]
	Case fatality risk (%)	Systematic review of CFR's worldwide from January 1, 2020, and March 31, 2023	2.62 (95% CI: 2.0-3.23)	Xia et al. 2024 [116]
	Infection fatality risk (%)	IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly partly vaccinated. Ages 20–39 years	0.03 (95% CI: 0.03-0.1)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly partly vaccinated. 40–59 years	0.37 (95% CI: 0.29-0.56)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly partly vaccinated. 60 + years	6.42 (95% CI: 4.69-7.44)	Zhang and Nishiura 2023 [129]
SARS-CoV-2 (Delta)	Basic reproduction number	Review of R0 from May to July 2021 using studies from China and the UK.	Mean: 5.08 (range: 3.20-8.0)	Liu and Rocklöv 2021 [130]

		Dispersion parameter	Modelling study. Calculated using 1,344 transmission pairs from the 11th of July to the 24th of July 2021 in South Korea. (Mask mandate, active case finding and immediately isolating laboratory-confirmed COVID-19 patients and exposed persons by using digital QR codes, 4-person limit for gatherings was implemented beginning July 19, 2021)	Negative binomial distribution with mean of 0.64 (95%CI: 0.57–0.72)	Ryu et al. 2022 [131]
			Modelling study. Calculated using 2,384 transmission pairs from the 25th of July to the 15th of August 2021 in South Korea. (Mask mandate, active case finding and immediately isolating laboratory-confirmed COVID-19 patients and exposed persons by using digital QR codes, 4-person limit for gatherings was implemented beginning July 19, 2021)	Negative binomial distribution with mean of 0.85 (95% CI: 0.75–0.98)	Ryu et al. 2022 [131]
			Modelling study. Using a likelihood-based estimating framework based on 126 observations from May to December 2021 in Guangdong, China. Under intense control measures	Negative binomial distribution with mean of 0.26 (95% CI: 0.16, 0.41)	Zhao et al. 2022 [132]
		Serial interval (d)	Systematic review of serial intervals 2020-2023	Mean: 3.59 (95% CI: 3.26-3.92)	Xu et al. 2023 [19]
			Modelling. Estimated value using household clusters from week 27/2021 until 49/2021 in Germany	Gamma distribution with mean of 4.19 (95% CI: 4.16-4.22)	Heiden and Buchholz 2022 [127]
		Incubation period (d)	Systematic review of Incubation periods 2020-2023	Mean: 4.63 (95% CI: 4.11-5.15)	Xu et al. 2023 [19]
			EpiLPS- Data set from Backer et al. 2022 [133]	Lognormal distribution with mean of 4.30 (95% CrI: 4.10-4.50)	Estimated
			Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China	Gamma distribution with mean of 5.04 (95% CI: 4.83-5.33)	Li et al. 2024 [134]

	Latent period (d)	Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China	Gamma distribution with mean of 4.40 (95% CI: 4.24-4.63)	Li et al. 2024 [134]
	Infectious period (d)	Viral shedding dynamics in fully vaccinated adults in the USA	Median: 6.0 (IQR: 5.0-8.0)	Garcia-Knight et a. 2022 [135]
	Case fatality risk (%)	Systematic review of CFR's worldwide from January 1, 2020, and March 31, 2023	2.01 (95% CI: 1.88-2.14)	Xia et al. 2024 [136]
		Systematic review of confirmed case-fatality risk. 18 January 2021 to December 2021	0.46 (95% CI: 0.2-0.73)	Yuan et al. 2023 [137]
	Infection fatality risk (%)	IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly was prioritised for vaccination and were fully vaccinated before the end of July 2021. Ages 20–39 years	0.01 (95% CI: 0.01-0.01)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly was prioritised for vaccination and were fully vaccinated before the end of July 2021. Ages 40–59 years	0.12 (95% CI: 0.1-0.16)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly was prioritised for vaccination and were fully vaccinated before the end of July 2021. Ages 60 + years	0.95 (95% CI: 0.7-1.1)	Zhang and Nishiura 2023 [129]
		Infection fatality risk in England at the end of the Delta period (November 2021)	0.11 (95% CI: 0.08-0.15)	Ward et al. 2024 [138]
SARS-CoV-2 (Omicron)	Basic reproduction number	Rapid review of R0 from November 2021 to December 2021 (Mix of countries)	Mean: 9.5 (IQR: 7.25-11.88)	Liu and Rocklöv 2022 [139]
	Effective reproduction number	Rapid review of RE from November 2021 to January 2022 (Mix of countries).	Mean: 3.4 (IQR: 0.88-9.40)	Liu and Rocklöv 2022 [139]
	Dispersion parameter	Modelling. 427 laboratory-confirmed cases from 25 November to 31 December 2021 in South Korea. (BA.1)	0.10 (95% CI: 0.08–0.13)	Guo et al. 2022 [140]

		Modelling. 67 epidemiologic linked cases from 2 to 21 January 2022 in Hong Kong (border control, physical distancing and contact tracing in place) (BA.1,2)	0.33 (95% CI: 0.17–0.62)	Guo et al. 2022 [140]
	Serial interval (d)	Systematic review of serial intervals 2020-2023	Mean: 3.21 (95% CI: 2.94-3.48)	Xu et al. 2023 [19]
	Incubation period (d)	Systematic review of Incubation periods. (BA.1)	Mean: 3.49 (95% CI: 3.13-4.86)	Xu et al. 2023 [19]
		Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China	Gamma distribution with mean of 3.41 (95% CI: 3.27-3.58)	Li et al. 2024 [134]
		Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China. (BA.1)	Gamma distribution with mean of 3.42 (95% CI: 3.00-3.89)	Li et al. 2024 [134]
		Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China. (BA.2)	Gamma distribution with mean of 3.39 (95% CI: 3.24-3.55)	Li et al. 2024 [134]
		EpiLPS - Dataset from Backer et al. 2022 [133]	Lognormal distribution with mean of 3.30 (95% CrI: 3.20-3.50)	Estimated
	Latent period (d)	Cross-sectional study in China, with 114 cases with COVID-19 Omicron variant BA.1.1 between January 2022 and February 2022	Gamma distribution with mean of 3.13 (95% CI: 2.82-3.48)	Xin et al. 2023 [120]
		Modelling, esitmated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China	Gamma distribution with mean of 2.58 (95% CI: 2.48-2.68)	Li et al. 2024 [134]
		Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China. (BA.1)	Gamma distribution with mean of 2.50 (95% CI: 2.27-2.76)	Li et al. 2024 [134]

		Modelling, estimated from infected individuals with clear transmission chains and infectors from epidemiological survey reports in China. (BA.2)	Gamma distribution with mean of 2.58 (95% CI: 2.48-2.69)	Li et al. 2024 [134]
	Infectious period (d)	Transmission period after symptom onset date (Spain). BA.1	Mean: 0.5 (IQR: -1.0-2.0)	Águila-Mejía et al. 2022 [141]
		Time to first negative viral culture (USA). (BA.2, 5, XBB)	Median: 4.0 (IQR: 3.0-4.0)	Edelstein et al. 2023 [142]
	Case fatality risk (%)	Systematic review of CFR's worldwide from January 1, 2020, and March 31, 2023	0.7 (95% CI: 0.67-0.73)	Xia et al. 2024 [136]
		Systematic review confirmed case-fatality risk. 18 January 2021 to December 2021	0.04 (95% CI: 0-0.61)	Yuan et al. 2023 [137]
		Systematic review, patient deaths/omicron patients. From 14/11/2021 to 07/03/2022	0.21	Ahmad et al. 2024 [143]
	Infection fatality risk (%)	IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly fully vaccinated. Ages 20–39 years	0 (95% CI: 0-0)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly fully vaccinated. Ages 40–59 years	0.02 (95% CI: 0.01-0.04)	Zhang and Nishiura 2023 [129]
		IFR for six epidemic waves in Osaka Japan from February 2020 to January 2022. Elderly fully vaccinated. Ages 60+ years	1.26 (95% CI: 0.9-2.54)	Zhang and Nishiura 2023 [129]
		Infection fatality risk in England at the end of the Omicron BA.1 and Omicron BA.2 period (April 2022)	0.06 (95% CI: 0.04-0.08)	Ward et al. 2024 [138]
ZIKV	Basic reproduction number	Review of R0 across global climate zones (1980-2018)	Mean: 3.02 (range: 0.16-9.40)	Liu et al. 2020 [144]
		Estimated R0 values for the outbreak in Miami 2016	Lognormal distribution with mean of 1.88 (95% CI: 1.53-2.32)	Liu et al. 2020 [144]

		Reproduction number for ZIKV sexual transmission	Median: 0.136 (95% CI: 0.009–0.521)	Gao et al. 2016 [145]
	Serial interval (d)	Systematic review, serial symptom onset interval in 15 couples via sexual transmission	Median: 12.0 (IQR:10.0-14.50)	Counotte et al 2018 [146]
	Incubation period (d)	Systematic review up to 2016. Pooled analysis	Lognormal distribution with median of 5.90 (95% CI: 4.40-7.60)	Lessler et al. 2016 [147]
		EpiLPS - Dataset from lessler et al. 2016 [147]	Semipar. with mean of 6.7 (95% CI: 5.8-7.6)	Estimated
	Latent period (d)	Modelling study. Referenced value. Assumed that human latent period is equivalent to the intrinsic incubation period and is a constant value	6.8	Agudelo et al. 2022 [148]
		Modelling study Referenced value. Assumed that latent period is equivalent to the incubation period	Gamma distribution with mean of 3.9	Kucharski et al. 2016 [149]
	Infectious period (d)	Systematic review up to 2016. Time period that Zika virus is detectable in blood	Mean: 9.9	Lessler et al. 2016 [147]
		Modelling study. Referenced value	Gamma distribution with mean of 5.0	Kucharski et al. 2016 [149]
		Systematic review, range of zika virus shedding in male genital tract	Range: 3-69	Moreira et al. 2017 [150]
	Case fatality risk (%)	Systematic review, CFR in the Americas up to 2018	Median: 0.02 (range: 0.002-0.324)	Cardona-Ospina et al. 2019 [151]
* SARS-CoV-2 wild type is defined as articles published regarding the first wave of COVID-19 prior to the emergence of the Alpha variant.				

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used Gemini 2.5 pro, within the Cursor editor in order to generate code used in this analysis. After using this tool/service, the

421   author(s) reviewed and edited the content as needed and take(s) full responsibility for the
422   content of the publication.
423

Acknowledgements

This work was supported by the ESCAPE project (101095619), co-funded by the European
Union. Views and opinions expressed are however those of the author(s) only and do not
necessarily reflect those of the European Union or European Health and Digital Executive
Agency (HADEA). Neither the European Union nor the granting authority can be held
responsible for them. This work was co-funded by UK Research and Innovation (UKRI)
under the UK government's Horizon Europe funding guarantee [grant number 10051037].
431
432
433
434

6. References

[1]   Rietveld J, Hobson T, Mani L, Avin S, Sundaram L. The UK's pandemic preparedness and early response to the COVID-19 pandemic. Glob Public Health 2024;19:2415499.

[2]   Rt Hon the Baroness Hallett DBE. UK Covid-19 Inquiry. House of Commons; 2024.

[3]   Looi M-K. What could the next pandemic be? BMJ 2023;381:909.

[4]   Adalja AA, Watson M, Toner ES, Cicero A, Inglesby TV. Characteristics of microbes most likely to cause pandemics and global catastrophes. Curr Top Microbiol Immunol 2019;424:1–20.

[5]   Prioritizing diseases for research and development in emergency contexts n.d. https://www.who.int/activities/prioritizing-diseases-for-research-and-development-in-emergency-contexts (accessed June 24, 2024).

[6]   Gressani O. EpiLPS: A Fast and Flexible Bayesian Tool for Estimating Epidemiological Parameters. 2021.

[7]   Gressani O, Torneri A, Hens N, Faes C. Flexible Bayesian estimation of incubation times. Am J Epidemiol 2025;194:490–501.

[8]   Gressani O, Wallinga J, Althaus CL, Hens N, Faes C. EpiLPS: A fast and flexible Bayesian tool for estimation of the time-varying reproduction number. PLoS Comput Biol 2022;18:e1010618.

[9]   Charniga K, Park SW, Akhmetzhanov AR, Cori A, Dushoff J, Funk S, et al. Best practices for estimating and reporting epidemiological delay distributions of infectious diseases. PLoS Comput Biol 2024;20:e1012520.

[10] Abbott S, Brand S, Pearson C, Funk S, Charniga K. primarycensored: Primary Event Censored Distributions 2025. https://doi.org/10.5281/zenodo.13632839.

[11] Cases of Crimean–Congo haemorrhagic fever infected in the EU/EEA, 2013–present. European Centre for Disease Prevention and Control 2021. https://www.ecdc.europa.eu/en/crimean-congo-haemorrhagic-fever/surveillance/cases-eu-since-2013 (accessed February 4, 2025).

[12] Strehl A, Ghosh J. Cluster ensembles --- A knowledge reuse framework for combining multiple partitions. J Mach Learn Res 2002;3:583–617.

[13] Rift valley fever n.d. https://www.who.int/health-topics/rift-valley-fever (accessed June 26, 2024).

[14] Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. Proc Natl Acad Sci U S A 2004;101:6146–51.

[15] Nash RK, Bhatia S, Morgenstern C, Doohan P, Jorgensen D, McCain K, et al. Ebola virus disease mathematical models and epidemiological parameters: a systematic

review. Lancet Infect Dis 2024;24:e762–73.

[16] Park J-E, Jung S, Kim A, Park J-E. MERS transmission and risk factors: a systematic
review. BMC Public Health 2018;18:574.

[17] Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the
reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic
review of the literature. BMC Infect Dis 2014;14:480.

[18] Muzembo BA, Kitahara K, Mitra D, Ntontolo NP, Ngatu NR, Ohno A, et al. The basic
reproduction number (R0) of ebola virus disease: A systematic review and meta-
analysis. Travel Med Infect Dis 2024;57:102685.

[19] Xu X, Wu Y, Kummer AG, Zhao Y, Hu Z, Wang Y, et al. Assessing changes in
incubation period, serial interval, and generation time of SARS-CoV-2 variants of
concern: a systematic review and meta-analysis. BMC Med 2023;21:374.

[20] Lai S, Qin Y, Cowling BJ, Ren X, Wardrop NA, Gilbert M, et al. Global epidemiology of
avian influenza A H5N1 virus infection in humans, 1997-2015: a systematic review of
individual case data. Lancet Infect Dis 2016;16:e108–18.

[21] DeWitt ME, Polk C, Williamson J, Shetty AK, Passaretti CL, McNeil CJ, et al. Global
monkeypox case hospitalisation rates: A rapid systematic review and meta-analysis.
EClinicalMedicine 2022;54:101710.

[22] Bunge EM, Hoet B, Chen L, Lienert F, Weidenthaler H, Baer LR, et al. The changing
epidemiology of human monkeypox-A potential threat? A systematic review. PLoS Negl
Trop Dis 2022;16:e0010141.

[23] Hegde ST, Lee KH, Styczynski A, Jones FK, Gomes I, Das P, et al. Potential for person-
to-person transmission of henipaviruses: A systematic review of the literature. J Infect
Dis 2024;229:733–42.

[24] Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-
19) infections. Int J Infect Dis 2020;93:284–6.

[25] Tindale LC, Stockdale JE, Coombe M, Garlock ES, Lau WYV, Saraswat M, et al.
Evidence for transmission of COVID-19 prior to symptom onset. Elife 2020;9.
https://doi.org/10.7554/eLife.57149.

[26] Casey-Bryars M, Griffin J, McAloon C, Byrne A, Madden J, Mc Evoy D, et al.
Presymptomatic transmission of SARS-CoV-2 infection: a secondary analysis using
published data. BMJ Open 2021;11:e041240.

[27] Ward T, Christie R, Paton RS, Cumming F, Overton CE. Transmission dynamics of
monkeypox in the United Kingdom: contact tracing study. BMJ 2022;379:e073153.

[28] Slifka MK, Gao L. Is presymptomatic spread a major contributor to COVID-19
transmission? Nat Med 2020;26:1531–3.

[29] Lo Iacono G, Cunningham AA, Fichet-Calvet E, Garry RF, Grant DS, Khan SH, et al.
Using modelling to disentangle the relative contributions of zoonotic and anthroponotic
transmission: the case of lassa fever. PLoS Negl Trop Dis 2015;9:e3398.

[30] Wolf T, Ellwanger R, Goetsch U, Wetzstein N, Gottschalk R. Fifty years of imported
Lassa fever: a systematic review of primary and secondary cases. J Travel Med
2020;27. https://doi.org/10.1093/jtm/taaa035.

[31] Fine PE, Jezek Z, Grab B, Dixon H. The transmission potential of monkeypox virus in
human populations. Int J Epidemiol 1988;17:643–50.

[32] Fraser C, Cummings DAT, Klinkenberg D, Burke DS, Ferguson NM. Influenza
transmission in households during the 1918 pandemic. Am J Epidemiol 2011;174:505–
14.

[33] Vink MA, Bootsma MCJ, Wallinga J. Serial intervals of respiratory infectious diseases: a
systematic review and analysis. Am J Epidemiol 2014;180:865–75.

[34] Nishiura H. Early efforts in modeling the incubation period of infectious diseases with an
acute course of illness. Emerg Themes Epidemiol 2007;4:2.

[35] Cori A, Valleron AJ, Carrat F, Scalia Tomba G, Thomas G, Boëlle PY. Estimating
influenza latency and infectious period durations using viral excretion data. Epidemics
2012;4:132–8.

[36] Carrat F, Luong J, Lao H, Sallé A-V, Lajaunie C, Wackernagel H. A "small-world-like"

model for comparing interventions aimed at preventing and controlling influenza
pandemics. BMC Med 2006;4:26.

[37] WHO. Pandemic Influenza Risk Management. 2017.

[38] Brugger J, Althaus CL. Transmission of and susceptibility to seasonal influenza in
Switzerland from 2003 to 2015. Epidemics 2020;30:100373.

[39] Tom BDM, Van Hoek AJ, Pebody R, McMenamin J, Robertson C, Catchpole M, et al.
Estimating time to onset of swine influenza symptoms after initial novel A(H1N1v) viral
infection. Epidemiol Infect 2011;139:1418–24.

[40] Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DAT. Incubation
periods of acute respiratory viral infections: a systematic review. Lancet Infect Dis
2009;9:291–300.

[41] Lessler J, Reich NG, Cummings DAT, New York City Department of Health and Mental
Hygiene Swine Influenza Investigation Team, Nair HP, Jordan HT, et al. Outbreak of
2009 pandemic influenza A (H1N1) at a New York City school. N Engl J Med
2009;361:2628–36.

[42] Tuite AR, Greer AL, Whelan M, Winter A-L, Lee B, Yan P, et al. Estimated
epidemiologic parameters and morbidity associated with pandemic H1N1 influenza.
CMAJ 2010;182:131–6.

[43] Van Kerkhove MD, Hirve S, Koukounari A, Mounts AW, H1N1pdm serology working
group. Estimating age-specific cumulative incidence for the 2009 influenza pandemic: a
meta-analysis of A(H1N1)pdm09 serological studies from 19 countries. Influenza Other
Respi Viruses 2013;7:872–86.

[44] Wong JY, Kelly H, Ip DKM, Wu JT, Leung GM, Cowling BJ. Case fatality risk of
influenza A (H1N1pdm09): a systematic review: A systematic review. Epidemiology
2013;24:830–41.

[45] Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, Leach S, et al. Time lines
of infection and disease in human influenza: a review of volunteer challenge studies. Am
J Epidemiol 2008;167:775–85.

[46] Elveback LR, Fox JP, Ackerman E, Langworthy A, Boyd M, Gatewood L. An influenza
simulation model for immunization studies. Am J Epidemiol 1976;103:152–65.

[47] Ferguson NM, Cummings DAT, Cauchemez S, Fraser C, Riley S, Meeyai A, et al.
Strategies for containing an emerging influenza pandemic in Southeast Asia. Nature
2005;437:209–14.

[48] Cauchemez S, Carrat F, Viboud C, Valleron AJ, Boëlle PY. A Bayesian MCMC
approach to study transmission of influenza: application to household longitudinal data.
Stat Med 2004;23:3469–87.

[49] Bettencourt LMA, Ribeiro RM. Real time bayesian estimation of the epidemic potential
of emerging infectious diseases. PLoS One 2008;3:e2185.

[50] Aditama TY, Samaan G, Kusriastuti R, Sampurno OD, Purba W, Misriyah, et al. Avian
influenza H5N1 transmission in households, Indonesia. PLoS One 2012;7:e29971.

[51] Yang Y, Halloran ME, Sugimoto JD, Longini IM Jr. Detecting human-to-human
transmission of avian influenza A (H5N1). Emerg Infect Dis 2007;13:1348–53.

[52] Ferguson NM, Fraser C, Donnelly CA, Ghani AC, Anderson RM. Public health. Public
health risk from the avian H5N1 influenza epidemic. Science 2004;304:968–9.

[53] Ward J, Lambert JW, Russell TW, Azam JM, Kucharski AJ, Funk S, et al. Estimates of
epidemiological parameters for H5N1 influenza in humans: a rapid review. medRxiv
2024:2024.12.11.24318702. https://doi.org/10.1101/2024.12.11.24318702.

[54] Beigel JH, Farrar J, Han AM, Hayden FG, Hyer R, de Jong MD, et al. Avian Influenza A
(H5N1) Infection in Humans. N Engl J Med 2005;353:1374–85.

[55] Huai Y, Xiang N, Zhou L, Feng L, Peng Z, Chapman RS, et al. Incubation period for
human cases of avian influenza A (H5N1) infection, China. Emerg Infect Dis
2008;14:1819–21.

[56] Cowling BJ, Jin L, Lau EHY, Liao Q, Wu P, Jiang H, et al. Comparative epidemiology of
human infections with avian influenza A H7N9 and H5N1 viruses in China: a population-
based study of laboratory-confirmed cases. Lancet 2013;382:129–37.

[57] Oner AF, Bay A, Arslan S, Akdeniz H, Sahin HA, Cesur Y, et al. Avian influenza A
(H5N1) infection in eastern Turkey in 2006. N Engl J Med 2006;355:2179–85.
[58] Van Kerkhove MD, Mumford E, Mounts AW, Bresee J, Ly S, Bridges CB, et al. Highly
pathogenic avian influenza (H5N1): pathways of exposure at the animal-human
interface, a systematic review. PLoS One 2011;6:e14582.
[59] Li FCK, Choi BCK, Sly T, Pak AWP. Evidence-based public health policy and practice:
Finding the real case-fatality rate of H5N1 avian influenza. Journal of Epidemiology and
Community Health (1979-) 2008;62:555–9.
[60] Bente DA, Forrester NL, Watts DM, McAuley AJ, Whitehouse CA, Bray M. Crimean-
Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and
genetic diversity. Antiviral Res 2013;100:159–89.
[61] Belhadi D, El Baied M, Mulier G, Malvy D, Mentré F, Laouénan C. The number of cases,
mortality and treatments of viral hemorrhagic fevers: A systematic review. PLoS Negl
Trop Dis 2022;16:e0010889.
[62] Belobo JTE, Kenmoe S, Kengne-Nde C, Emoh CPD, Bowo-Ngandji A, Tchatchouang S,
et al. Worldwide epidemiology of Crimean-Congo Hemorrhagic Fever Virus in humans,
ticks and other animal species, a systematic review and meta-analysis. PLoS Negl Trop
Dis 2021;15:e0009299.
[63] Perveen N, Khan G. Crimean-Congo hemorrhagic fever in the Arab world: A systematic
review. Front Vet Sci 2022;9:938601.
[64] Nasirian H. New aspects about Crimean-Congo hemorrhagic fever (CCHF) cases and
associated fatality trends: A global systematic review and meta-analysis. Comp
Immunol Microbiol Infect Dis 2020;69:101429.
[65] Wong ZSY, Bui CM, Chughtai AA, Macintyre CR. A systematic review of early modelling
studies of Ebola virus disease in West Africa. Epidemiol Infect 2017;145:1069–94.
[66] Velásquez GE, Aibana O, Ling EJ, Diakite I, Mooring EQ, Murray MB. Time From
Infection to Disease and Infectiousness for Ebola Virus Disease, a Systematic Review.
Clin Infect Dis 2015;61:1135–40.
[67] Khan SA, Imtiaz MA, Islam MM, Tanzin AZ, Islam A, Hassan MM. Major bat-borne
zoonotic viral epidemics in Asia and Africa: A systematic review and meta-analysis. Vet
Med Sci 2022;8:1787–801.
[68] Zhao S, Musa SS, Fu H, He D, Qin J. Large-scale Lassa fever outbreaks in Nigeria:
quantifying the association between disease reproduction number and local rainfall.
Epidemiol Infect 2020;148:e4.
[69] Doohan P, Jorgensen D, Naidoo TM, McCain K, Hicks JT, McCabe R, et al. Lassa fever
outbreaks, mathematical models, and disease parameters: a systematic review and
meta-analysis. Lancet Glob Health 2024;12:e1962–72.
[70] Akhmetzhanov AR, Asai Y, Nishiura H. Quantifying the seasonal drivers of transmission
for Lassa fever in Nigeria. Philos Trans R Soc Lond B Biol Sci 2019;374:20180268.
[71] Tuite AR, Watts AG, Kraemer MUG, Khan K, Bogoch II. Potential for Seasonal Lassa
Fever Case Exportation from Nigeria. Am J Trop Med Hyg 2019;100:647–51.
[72] Kenmoe S, Tchatchouang S, Ebogo-Belobo JT, Ka'e AC, Mahamat G, Guiamdjo Simo
RE, et al. Systematic review and meta-analysis of the epidemiology of Lassa virus in
humans, rodents and other mammals in sub-Saharan Africa. PLoS Negl Trop Dis
2020;14:e0008589.
[73] Dwalu E, Jetoh RW, Shobayo BI, Pewu I, Taweh F, Wilson-Sesay HW, et al. Trend of
Lassa fever cases and factors associated with mortality in Liberia, 2016 - 2021: a
secondary data analysis. Pan Afr Med J 2024;47:22.
[74] Qian GY, Edmunds WJ, Bausch DG, Jombart T. A mathematical model of Marburg virus
disease outbreaks and the potential role of vaccination in control. BMC Med
2023;21:439.
[75] Ajelli M, Merler S. Transmission potential and design of adequate control measures for
Marburg hemorrhagic fever. PLoS One 2012;7:e50948.
[76] Pavlin BI. Calculation of incubation period and serial interval from multiple outbreaks of
Marburg virus disease. BMC Res Notes 2014;7:906.

[77] Bettencourt LMA. An Ensemble Trajectory Method for Real-Time Modeling and
Prediction of Unfolding Epidemics: Analysis of the 2005 Marburg Fever Outbreak in
Angola. Mathematical and Statistical Estimation Approaches in Epidemiology 2009:143.
[78] Cuomo-Dannenburg G, McCain K, McCabe R, Unwin HJT, Doohan P, Nash RK, et al.
Marburg virus disease outbreaks, mathematical models, and disease parameters: a
systematic review. Lancet Infect Dis 2024;24:e307–17.
[79] Peak CM, Childs LM, Grad YH, Buckee CO. Comparing nonpharmaceutical
interventions for containing emerging epidemics. Proc Natl Acad Sci U S A
2017;114:4023–8.
[80] Wang J, Chen X, Guo Z, Zhao S, Huang Z, Zhuang Z, et al. Superspreading and
heterogeneity in transmission of SARS, MERS, and COVID-19: A systematic review.
Comput Struct Biotechnol J 2021;19:5039–46.
[81] Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DAT, et al. Hospital
outbreak of Middle East respiratory syndrome coronavirus. N Engl J Med
2013;369:407–16.
[82] Park SH, Kim Y-S, Jung Y, Choi SY, Cho N-H, Jeong HW, et al. Outbreaks of Middle
East Respiratory Syndrome in Two Hospitals Initiated by a Single Patient in Daejeon,
South Korea. Infect Chemother 2016;48:99–107.
[83] Cowling BJ, Park M, Fang VJ, Wu P, Leung GM, Wu JT. Preliminary epidemiological
assessment of MERS-CoV outbreak in South Korea, May to June 2015. Euro Surveill
2015;20:7–13.
[84] Cauchemez S, Fraser C, Van Kerkhove MD, Donnelly CA, Riley S, Rambaut A, et al.
Middle East respiratory syndrome coronavirus: quantification of the extent of the
epidemic, surveillance biases, and transmissibility. Lancet Infect Dis 2014;14:50–6.
[85] Lessler J, Rodriguez-Barraquer I, Cummings DAT, Garske T, Van Kerkhove M, Mills H,
et al. Estimating Potential Incidence of MERS-CoV Associated with Hajj Pilgrims to
Saudi Arabia, 2014. PLoS Curr 2014;6.
https://doi.org/10.1371/currents.outbreaks.c5c9c9abd636164a9b6fd4dbda974369.
[86] Okoli GN, Van Caeseele P, Askin N, Abou-Setta AM. A global systematic evidence
review with meta-analysis of the epidemiological characteristics of the 2022 Mpox
outbreaks. Infection 2024;52:901–21.
[87] Beer EM, Rao VB. A systematic review of the epidemiology of human monkeypox
outbreaks and implications for outbreak strategy. PLoS Negl Trop Dis
2019;13:e0007791.
[88] Blumberg S, Lloyd-Smith JO. Inference of R(0) and transmission heterogeneity from the
size distribution of stuttering chains. PLoS Comput Biol 2013;9:e1002993.
[89] Paredes MI, Ahmed N, Figgins M, Colizza V, Lemey P, McCrone JT, et al.
Underdetected dispersal and extensive local transmission drove the 2022 mpox
epidemic. Cell 2024;187:1374–86.e13.
[90] Okoli GN, Van Caeseele P, Askin N, Abou-Setta AM. Comparative evaluation of the
clinical presentation and epidemiology of the 2022 and previous Mpox outbreaks: a
rapid review and meta-analysis. Infect Dis 2023;55:490–508.
[91] Madewell ZJ, Charniga K, Masters NB, Asher J, Fahrenwald L, Still W, et al. Serial
Interval and Incubation Period Estimates of Monkeypox Virus Infection in 12
Jurisdictions, United States, May-August 2022. Emerg Infect Dis 2023;29:818–21.
[92] Chenchula S, Ghanta MK, Amerneni KC, Rajakarunakaran P, Chandra MB, Chavan M,
et al. A systematic review to identify novel clinical characteristics of monkeypox virus
infection and therapeutic and preventive strategies to combat the virus. Arch Virol
2023;168:195.
[93] Hatami H, Jamshidi P, Arbabi M, Safavi-Naini SAA, Farokh P, Izadi-Jorshari G, et al.
Demographic, Epidemiologic, and Clinical Characteristics of Human Monkeypox
Disease Pre- and Post-2022 Outbreaks: A Systematic Review and Meta-Analysis.
Biomedicines 2023;11. https://doi.org/10.3390/biomedicines11030957.
[94] Miura F, van Ewijk CE, Backer JA, Xiridou M, Franz E, Op de Coul E, et al. Estimated
incubation period for monkeypox cases confirmed in the Netherlands, May 2022. Euro

Surveill 2022;27:2200448.

[95] Asakura TR, Jung S-M, Murayama H, Ghaznavi C, Sakamoto H, Teshima A, et al.
Projecting international mpox spread in Asia: ongoing global health risk. medRxiv 2024.
https://doi.org/10.1101/2024.04.17.24305832.

[96] Endo A, Murayama H, Abbott S, Ratnayake R, Pearson CAB, Edmunds WJ, et al.
Heavy-tailed sexual contact networks and monkeypox epidemiology in the global
outbreak, 2022. Science 2022;378:90–4.

[97] Xiridou M, Miura F, Adam P, Op de Coul E, de Wit J, Wallinga J. The Fading of the
Mpox Outbreak Among Men Who Have Sex With Men: A Mathematical Modelling
Study. The Journal of Infectious Diseases 2024;230:e121–30.

[98] Benites-Zapata VA, Ulloque-Badaracco JR, Alarcon-Braga EA, Hernandez-Bustamante
EA, Mosquera-Rojas MD, Bonilla-Aldana DK, et al. Clinical features, hospitalisation and
deaths associated with monkeypox: a systematic review and meta-analysis. Ann Clin
Microbiol Antimicrob 2022;21:36.

[99] Nikolay B, Salje H, Hossain MJ, Khan AKMD, Sazzad HMS, Rahman M, et al.
Transmission of Nipah Virus - 14 Years of Investigations in Bangladesh. N Engl J Med
2019;380:1804–14.

[100]   Bradbury NV, Hart WS, Lovell-Read FA, Polonsky JA, Thompson RN. Exact
calculation of end-of-outbreak probabilities using contact tracing data. J R Soc Interface
2023;20:20230374.

[101]   Kenmoe S, Demanou M, Bigna JJ, Nde Kengne C, Fatawou Modiyinji A, Simo FBN,
et al. Case fatality rate and risk factors for Nipah virus encephalitis: A systematic review
and meta-analysis. J Clin Virol 2019;117:19–26.

[102]   Suman N, Khandelwal E, Chiluvuri P, Rami DS, Chansoria S, Jerry A, et al. NIPAH
virus encephalitis: Unveiling the epidemiology, risk factors, and clinical outcomes - A
systematic review and meta-analysis. J Pharm Bioallied Sci 2024;16:S102–5.

[103]   Rudolph KE, Lessler J, Moloney RM, Kmush B, Cummings DAT. Incubation periods
of mosquito-borne viral infections: a systematic review. Am J Trop Med Hyg
2014;90:882–91.

[104]   Ebogo-Belobo JT, Kenmoe S, Abanda NN, Bowo-Ngandji A, Mbaga DS, Magoudjou-
Pekam JN, et al. Contemporary epidemiological data of Rift Valley fever virus in
humans, mosquitoes and other animal species in Africa: A systematic review and meta-
analysis. Vet Med Sci 2023;9:2309–28.

[105]   Lipsitch M, Cohen T, Cooper B, Robins JM, Ma S, James L, et al. Transmission
dynamics and control of severe acute respiratory syndrome. Science 2003;300:1966–
70.

[106]   Riley S, Fraser C, Donnelly CA, Ghani AC, Abu-Raddad LJ, Hedley AJ, et al.
Transmission dynamics of the etiological agent of SARS in Hong Kong: impact of public
health interventions. Science 2003;300:1961–6.

[107]   Chen S-C, Chang C-F, Liao C-M. Predictive models of control strategies involved in
containing indoor airborne infections. Indoor Air 2006;16:469–81.

[108]   Chowell G, Castillo-Chavez C, Fenimore PW, Kribs-Zaleta CM, Arriola L, Hyman JM.
Model parameters and outbreak control for SARS. Emerg Infect Dis 2004;10:1258–63.

[109]   Tsang KW, Ho PL, Ooi GC, Yee WK, Wang T, Chan-Yeung M, et al. A cluster of
cases of severe acute respiratory syndrome in Hong Kong. N Engl J Med
2003;348:1977–85.

[110]   Becker NG, Glass K, Li Z, Aldis GK. Controlling emerging infectious diseases like
SARS. Math Biosci 2005;193:205–21.

[111]   Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in
emerging epidemics. PLoS One 2006;1:e12.

[112]   Lloyd-Smith JO, Galvani AP, Getz WM. Curtailing transmission of severe acute
respiratory syndrome within a community and its hospital. Proc Biol Sci 2003;270:1979–
89.

[113]   Hui DSC, Chan MCH, Wu AK, Ng PC. Severe acute respiratory syndrome (SARS):
epidemiology and clinical features. Postgrad Med J 2004;80:373–81.

[114] Ahammed T, Anjum A, Rahman MM, Haider N, Kock R, Uddin MJ. Estimation of
novel coronavirus (COVID-19) reproduction number and case fatality rate: A systematic
review and meta-analysis. Health Sci Rep 2021;4:e274.
[115] Lin Y-F, Duan Q, Zhou Y, Yuan T, Li P, Fitzpatrick T, et al. Spread and Impact of
COVID-19 in China: A Systematic Review and Synthesis of Predictions From
Transmission-Dynamic Models. Front Med 2020;7:321.
[116] Xie Y, Wang Z, Liao H, Marley G, Wu D, Tang W. Epidemiologic, clinical, and
laboratory findings of the COVID-19 in the current pandemic: systematic review and
meta-analysis. BMC Infect Dis 2020;20:640.
[117] Izadi N, Taherpour N, Mokhayeri Y, Sotoodeh Ghorbani S, Rahmani K, Hashemi
Nazari SS. Epidemiologic Parameters for COVID-19: A Systematic Review and Meta-
Analysis. Med J Islam Repub Iran 2022;36:155.
[118] Du Z, Wang C, Liu C, Bai Y, Pei S, Adam DC, et al. Systematic review and meta-
analyses of superspreading of SARS-CoV-2 infections. Transbound Emerg Dis
2022;69:e3007–14.
[119] Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus
(2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. Euro
Surveill 2020;25. https://doi.org/10.2807/1560-7917.ES.2020.25.5.2000062.
[120] Xin H, Li Y, Wu P, Li Z, Lau EHY, Qin Y, et al. Estimating the Latent Period of
Coronavirus Disease 2019 (COVID-19). Clin Infect Dis 2022;74:1678–81.
[121] He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral
shedding and transmissibility of COVID-19. Nat Med 2020;26:672–5.
[122] Zhu H, Li Y, Jin X, Huang J, Liu X, Qian Y, et al. Transmission dynamics and control
methodology of COVID-19: A modeling study. Appl Math Model 2021;89:1983–98.
[123] Nishiura H, Kobayashi T, Yang Y, Hayashi K, Miyama T, Kinoshita R, et al. The rate
of underascertainment of novel Coronavirus (2019-nCoV) infection: Estimation using
Japanese passengers data on evacuation flights. J Clin Med 2020;9:419.
[124] Mizumoto K, Kagaya K, Chowell G. Early epidemiological assessment of the
transmission potential and virulence of coronavirus disease 2019 (COVID-19) in Wuhan
City, China, January-February, 2020. BMC Med 2020;18:217.
[125] Brazeau NF, Verity R, Jenks S, Fu H, Whittaker C, Winskill P, et al. Estimating the
COVID-19 infection fatality ratio accounting for seroreversion using statistical modelling.
Commun Med (Lond) 2022;2:54.
[126] Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al.
Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England.
Science 2021;372. https://doi.org/10.1126/science.abg3055.
[127] An der Heiden M, Buchholz U. Serial interval in households infected with SARS-CoV-
2 variant B.1.1.529 (Omicron) is even shorter compared to Delta. Epidemiol Infect
2022;150:e165.
[128] Hart WS, Miller E, Andrews NJ, Waight P, Maini PK, Funk S, et al. Generation time of
the alpha and delta SARS-CoV-2 variants: an epidemiological analysis. Lancet Infect
Dis 2022;22:603–10.
[129] Zhang T, Nishiura H. Estimating infection fatality risk and ascertainment bias of
COVID-19 in Osaka, Japan from February 2020 to January 2022. Sci Rep
2023;13:5540.
[130] Liu Y, Rocklöv J. The reproductive number of the Delta variant of SARS-CoV-2 is far
higher compared to the ancestral SARS-CoV-2 virus. J Travel Med 2021;28.
https://doi.org/10.1093/jtm/taab124.
[131] Ryu S, Kim D, Lim J-S, Ali ST, Cowling BJ. Serial Interval and Transmission
Dynamics during SARS-CoV-2 Delta Variant Predominance, South Korea. Emerg Infect
Dis 2022;28:407–10.
[132] Zhao S, Guo Z, Chong MKC, He D, Wang MH. Superspreading potential of SARS-
CoV-2 Delta variants under intensive disease control measures in China. J Travel Med
2022;29. https://doi.org/10.1093/jtm/taac025.
[133] Backer JA, Eggink D, Andeweg SP, Veldhuijzen IK, van Maarseveen N, Vermaas K,

et al. Shorter serial intervals in SARS-CoV-2 cases with Omicron BA.1 variant compared with Delta variant, the Netherlands, 13 to 26 December 2021. Euro Surveill 2022;27:2200042.

[134] Li Y, Jiang X, Qiu Y, Gao F, Xin H, Li D, et al. Latent and incubation periods of Delta, BA.1, and BA.2 variant cases and associated factors: a cross-sectional study in China. BMC Infect Dis 2024;24:294.

[135] Garcia-Knight M, Anglin K, Tassetto M, Lu S, Zhang A, Goldberg SA, et al. Infectious viral shedding of SARS-CoV-2 Delta following vaccination: A longitudinal cohort study. PLoS Pathog 2022;18:e1010802.

[136] Xia Q, Yang Y, Wang F, Huang Z, Qiu W, Mao A. Case fatality rates of COVID 19 during epidemic periods of variants of concern: A meta-analysis by continents. Int J Infect Dis 2024;141:106950.

[137] Yuan Z, Shao Z, Ma L, Guo R. Clinical Severity of SARS-CoV-2 Variants during COVID-19 Vaccination: A Systematic Review and Meta-Analysis. Viruses 2023;15. https://doi.org/10.3390/v15101994.

[138] Ward T, Fyles M, Glaser A, Paton RS, Ferguson W, Overton CE. The real-time infection hospitalisation and fatality risk across the COVID-19 pandemic in England. Nat Commun 2024;15:4633.

[139] Liu Y, Rocklöv J. The effective reproductive number of the Omicron variant of SARS-CoV-2 is several times relative to Delta. J Travel Med 2022;29. https://doi.org/10.1093/jtm/taac037.

[140] Guo Z, Zhao S, Ryu S, Mok CKP, Hung CT, Chong KC, et al. Superspreading potential of infection seeded by the SARS-CoV-2 Omicron BA.1 variant in South Korea. J Infect 2022;85:e77–9.

[141] Del Águila-Mejía J, Wallmann R, Calvo-Montes J, Rodríguez-Lozano J, Valle-Madrazo T, Aginagalde-Llorente A. Secondary Attack Rate, Transmission and Incubation Periods, and Serial Interval of SARS-CoV-2 Omicron Variant, Spain. Emerg Infect Dis 2022;28:1224–8.

[142] Edelstein GE, Boucau J, Uddin R, Marino C, Liew MY, Barry M, et al. SARS-CoV-2 virologic rebound with nirmatrelvir-ritonavir therapy : An observational study: An observational study. Ann Intern Med 2023;176:1577–85.

[143] Ahmad SJ, Degiannis JR, Borucki J, Pouwels S, Rawaf DL, Lala A, et al. Fatality Rates After Infection With the Omicron Variant (B.1.1.529): How Deadly has it been? A Systematic Review and Meta-Analysis. J Acute Med 2024;14:51–60.

[144] Liu Y, Lillepold K, Semenza JC, Tozan Y, Quam MBM, Rocklöv J. Reviewing estimates of the basic reproduction number for dengue, Zika and chikungunya across global climate zones. Environ Res 2020;182:109114.

[145] Gao D, Lou Y, He D, Porco TC, Kuang Y, Chowell G, et al. Prevention and Control of Zika as a Mosquito-Borne and Sexually Transmitted Disease: A Mathematical Modeling Analysis. Sci Rep 2016;6:28070.

[146] Counotte MJ, Kim CR, Wang J, Bernstein K, Deal CD, Broutet NJN, et al. Sexual transmission of Zika virus and other flaviviruses: A living systematic review. PLoS Med 2018;15:e1002611.

[147] Lessler J, Ott CT, Carcelen AC, Konikoff JM, Williamson J, Bi Q, et al. Times to key events in Zika virus infection and implications for blood donation: a systematic review. Bull World Health Organ 2016;94:841–9.

[148] Agudelo S, Ventresca M. Modeling the spread of the Zika virus by sexual and mosquito transmission. PLoS One 2022;17:e0270127.

[149] Kucharski AJ, Funk S, Eggo RM, Mallet H-P, Edmunds WJ, Nilles EJ. Transmission Dynamics of Zika Virus in Island Populations: A Modelling Analysis of the 2013-14 French Polynesia Outbreak. PLoS Negl Trop Dis 2016;10:e0004726.

[150] Moreira J, Peixoto TM, Siqueira AM, Lamas CC. Sexually acquired Zika virus: a systematic review. Clin Microbiol Infect 2017;23:296–305.

[151] Cardona-Ospina JA, Henao-SanMartin V, Acevedo-Mendoza WF, Nasner-Posso KM, Martínez-Pulgarín DF, Restrepo-López A, et al. Fatal Zika virus infection in the

855  Americas: A systematic review. Int J Infect Dis 2019;88:49–59.