Landscape of Clinical Trials in Cancer Cachexia: Assessment of Trends From

1995-2024

Ana Regina Cabrera[1*], Kaitlyn Parker[1*], Deena B. Snoke[2], Bart Hammig[3], Nicholas P. Greene[1].

[1]Cachexia Research Laboratory, Exercise Science Research Center, Department of Health, Human Performance and

Recreation, University of Arkansas, Fayetteville AR

[2]Department of Medicine, University of Vermont College of Medicine, Burlington, VT, United States

[3]Public Health Program, Department of Health, Human Performance and Recreation, University of Arkansas,

Fayetteville AR

Corresponding Author:

Nicholas P. Greene

HPER 321E

University of Arkansas

155 Stadium Dr.

Fayetteville, AR 72701

Phone number: 479-575-6638

Fax: 479-575-2853

npgreene@uark.edu

* These authors contributed equally to this work

Abstract

Cancer cachexia (CC), a multifactorial syndrome characterized by unintentional weight loss, is a frequent complication of cancer that impacts patients' quality of life and survival. In this retrospective review, we evaluated the landscape of clinical trials (CTx) registered on ClinicalTrials.gov for the consideration of potential factors contributing to human heterogeneity in their design and analyses. Among CTx registered from 1995-2024, we observed a trend toward equality in including female participants, but lack of reporting of sex as a biological variable. Limited demographic diversity was observed with ~93% White population participation. There was a substantial divergence in the diagnostic criteria and a wide range of tools employed to measure CC. Lastly, few studies considered cancer type and stage as clinical variables. Overall, a substantial gap remains in our knowledge of CC in non-white and female populations. Ultimately, these underreported data across CC CTx complicate the comparison and interpretation of CTx results, both within understudied populations and in specific cancer types. The current evolution of knowledge and new methodologies used for CC assessment reinforce the need for a constant revision of the CC consensus definition and diagnosis criteria to align with current advances in our understanding of human heterogeneity in CC.

Background

Cancer is a critical global public health concern, with 9.7 million cancer-related deaths recorded worldwide
in 2022 (1). Among all cancer types, lung, gastrointestinal, and pancreatic cancers are together estimated to
comprise 38% of all cancer-related deaths in the United States (2). Cancer cachexia (CC), defined by Fearon et al. in
2011 as a multifactorial syndrome characterized by unintentional ongoing muscle loss which may occur with or
without fat loss that is irreversible by conventional nutritional support (3), is a frequent complication of cancer
which occurs at high rates in the cancer types mentioned above. CC is commonly clinically identified by decreased
body weight or changes in body composition, which inevitably leads to impaired physical function. Because reliable
body composition data are challenging, expensive, and impossible to retrospectively collect in a clinical setting,
diagnostic criteria for CC have focused on body weight loss, with patients losing more than 5% of body weight or
underweight patients (BMI<20 kg/m^2) losing more than 2% of body weight in the past six months meeting criteria
for CC (3). Clinical presentation of CC is a valuable metric to oncologists, as patients experiencing CC have
increased fatigue (4), muscle weakness (5), treatment-related toxicities (6, 7), and decreased survival (5).
Additionally, the frequency of CC is clinically significant, as its incidence increases in late-stage cancers (8), can be
experienced in up to 80% of patients depending on cancer type (3, 9, 10) and is the main cause of mortality in ~20-
40% of cancer-related deaths (11).

Despite its prevalence and impact, efforts to develop preventative and therapeutic treatments for CC,
including dietary supplements, exercise, and pharmaceutical drugs have had minimal success to date. The scientific
community has acknowledged that these challenges may be due to the widespread use of preclinical models that
may not faithfully recapitulate the human condition (12, 13), especially when extrapolated across cancers and other
biological variables. Additionally, characteristics of CC can vary significantly depending on cancer type, tumor
location, metabolism, mutational identity, and stage (7), which, due to the necessity of repeatability in disease
models, is often not well reflected in preclinical studies. This is further complicated by factors intrinsic to human
heterogeneity, which leads to significant variation in CC symptoms among individuals. One such factor that has
gained traction in recent years is the effect of biological sex on CC characteristics (14). Clinically, males appear to
exhibit well-described signs of CC earlier during cancer progression and undergo more significant declines in
muscle mass in comparison to females (14). The observation that males adhere to this more 'typical' pattern of CC
is partially due to the fact that historically, the majority of preclinical studies have utilized male mice or do not
address biological sex as an experimental variable, and thus our understanding of the manifestation of CC in females
is limited (15). Additionally, it is impossible to capture the complexities of human genetic, environmental, and
social factors such as race and ethnicity in preclinical models. However, recent patient data suggest that race and
ethnicity demographics do impact disease outcomes such as CC incidence and mortality (16-19), underscoring the
importance of studying diverse populations in clinical studies and trials to understand the genetic and environmental
factors, including health disparities, contributing to disease outcomes. Additionally, the breadth of diagnostic
measures reported across studies in human patients may contribute to inconsistencies in the literature that prevent
pooled or comparative analyses across trials. The aim of the present review was to conduct a comprehensive

assessment of CC clinical trials (CTx) between 1995-2024 and evaluate how these studies consider potential factors,

such as demographic characteristics, intervention type, cancer type, and reported measures, contributing to

heterogeneity of patient populations in their design, analyses, and interpretation. In doing so, we identify effective

measures used by these studies and areas where alternative methods and considerations should be employed to

address gaps in knowledge. Ultimately, the goal of this assessment is to aid in the advancement of future research

initiatives by the design of more comprehensive and effective CTx in CC.

Methods

This is a retrospective systematic review to analyze CTx investigating CC registered on ClinicalTrials.gov.

Eligibility Criteria

Inclusion criteria were interventional and observational CTx investigating CC in child and adult subjects,

both male and female, published on ClinicalTrials.gov. Trials were included if CC, cancer-induced sarcopenia, or

cancer-induced muscle wasting were the primary reason for the study. Completed trials, active trials, and terminated

trials were included. Exclusion criteria included trials that were unrelated to CC and trials measuring non-cancer

induced cachexia. No trials were excluded based on country, region, total enrollment, trial start or completion date.

Information Sources

The primary data source for this research was ClinicalTrials.gov, a comprehensive database providing

information about clinical research studies conducted throughout the world. For trials listed without study results,

PubMed was utilized to search for published manuscripts including results associated with them.

Search Strategy

This search was conducted through ClinicalTrials.gov from December 2023 to August 2024, regardless of

country and focusing on the condition/disease filter in the search generator. Terms related to CC, including "cancer

cachexia", "cachexia", "sarcopenia", and "muscle wasting" were searched one at a time. If "cachexia",

"sarcopenia", or "muscle wasting" was the primary search term in the condition filter, "cancer" was added as a

secondary search term in the other terms filter. No terms were added to the intervention/treatment filter or location

filter. All boxes under study status were selected. The "all" option was selected under the sex filter. All age ranges

were selected under the age filter. "Early Phase 1", "Phase 1", "Phase 2", "Phase 3", and "Phase 4" were selected

under the study phase filter. Both "Interventional" and "Observational" were selected under the study type filter.

"With results" and "Without results" were selected under the study results filter. No options were selected under the

study documents, funder type, and date range filters. No additional information was input into the "More ways to

search" section. When utilizing PubMed, the CTx official name followed by the study director's name were

searched.

Data Collection Process

Data from eligible CTx were extracted from ClinicalTrials.gov. Official study title, ClinicalTrials.gov ID

(NCT number), sponsor, study director, study start date, actual or estimated study completion date, study status,

study type, enrollment, intervention(s), participant demographics, primary and secondary outcome measures, the

type(s) of cancer included, CC definition used, and any listed publications were collected as reported on

ClinicalTrials.gov. The same data were extracted from trials that lacked results or other data on ClinicalTrials.gov

but were found in published manuscripts on PubMed. Published manuscripts were matched to CTx on

ClinicalTrials.gov by NCT number. Principal investigators and participation enrollment were used as secondary

confirmation to ensure the manuscript discussed the clinical trial of interest.

Data Synthesis

The extracted data was grouped based on a variety of measures, including start and completion date of the

trial, the treatment/intervention, type of cancer(s), CC definition used, and outcome measures. The list of CTx

included is in **Supplemental File 1**.

Figures and Tables

Figures were made with Biorender or R (v4.4.1). Tables and database were managed with Microsoft365

Word and Excel.

Results

The searches conducted in ClinicalTrials.gov generated 209 trials using "cancer cachexia"; "cachexia" +

"cancer" generated 226 results; "sarcopenia" + "cancer" generated 126 results; and "muscle wasting" + cancer"

generated 143 results. After removing duplicates, 491 individual trials remained. After title and abstract screening,

300 trials did not meet the inclusion criteria. Therefore, the remaining 191 trials were included in this analysis. Of

these, 106 trials had the status of complete, 10 of which had results published on ClinicalTrials.gov and 42 had

published results in manuscripts available from PubMed, resulting in 52 completed trials with results. The remaining

54 completed trials did not have results identified on either ClinicalTrials.gov or PubMed. In addition, 62 trials were

active, and 23 were terminated or withdrawn before completion at the conclusion of the search (**Supplemental File**

2). The reasons for termination are listed in **Supplemental File 3**, with none of the trials being terminated by safety

boards. Moreover, most of the CTx (116, 60.7%) did not have a phase stated, 15 (7.9%) were in phase 1, 31 (16.2%)

were in phase 2, 20 (10.5%) were in phase 3, 2 (1.0%) in phase 4, and 7 (3.7%) were in between two different

phases (**Supplemental File 4**).

Clinical Trial Demographics

We generated an overview of demographic composition of patients and how it was reported based on the

52 CTx with published findings. A summary of the reported demographics for age, biological sex, race, and

ethnicity is compiled in **Figure 1a**.

From the 52 CTx with results reported, 100% reported the ages of participants; of those, 50 trials (96.2%)

reported the mean or median age of patients, ranging from 11 to 75 years old. From these 52 CTx, eight trials

(15.4%) reported patient numbers by age categories (<18, 18-64, and >64). A total of 49 trials (94.2%) reported

biological sex of participants and only one from the CC-CTx completed with results (1.9%) reported clinical

outcomes by biological sex (NCT02148159). Among all trials with sex reporting, there were 3,103 females (40.4%)

and 4,573 males (59.6%). In addition, we calculated the percentage of female and male participants within each

study and the mean percentage had a similar distribution with 43.5% females and 56.5% males. However, we

observed different trends across years. We noted an average of 38.7% females and 61.3% males in CTx with results

reported from 2004 to 2014, while the average reported in CTx from 2015 to 2023 showed average distributions

closed to equality with 48.8% of female and 51.2% of male participants (**Figure 1b**). In addition, 17 trials (32.7%)

reported race of participants. Of these trials, all (100%) reported the participation of White patients. Fourteen trials

(82.4%) reported the participation of Black or African American patients, six trials (35.3%) included Asian patients,

two (11.8%) reported participation of Native Hawaiian/Pacific Islander individuals, one (5.9%) included Pardo

patients, 9 (52.9%) included an unknown or mixed category, and no trials included American Indian/Alaska Native

patients. Among all trials reporting race, there were 2,135 (92.6%) White patients, 77 (3.3%) Black or African

American patients, 21 (0.9%) Pardo, 12 (0.5%) Asian patients, 2 (0.1%) Native Hawaiian/Pacific Islander patients,

and 0 Native American/Alaskan Native patients or other races reported (**Figure 1c**). No trials reported clinical

outcomes by race in a subgroup analysis. In addition to race, only five trials (9.6%) reported the ethnicity of

participants. Among trials reporting ethnicity, 100% reported Hispanic/Latino patients making a 0.7% of the total

population included in the CTx completed with results. Besides Hispanic/Latino participants, no other specific

ethnicities were reported. No trials reported clinical outcomes by ethnicity in a subgroup analysis. Overall, CTx

include both sexes but fail to report sex as a biological variable. Additionally, there is a significant knowledge gap in

the study of CC among races and ethnicities other than the White population.

Cancer Cachexia Definition and Classification

Despite the existence of a consensus definition of cachexia, the application of the diagnostic criteria

remains only partially implemented in clinical practice. Therefore, we decided to classify the CTx according to the

diagnostic criteria used to define CC. Our analysis demonstrated that of all trials included in this study (n = 191),

61.8% defined how they classified CC (**Figure 2a**). A significant portion of the studies (64, representing 33.5%)

used the Fearon et al. consensus definition (3), stating a degree of loss and time frame for its occurrence. Studies that

used weight loss greater than 5% in the past six months as their criteria, or weight loss greater than 2% in

individuals already showing depletion with body mass index (BMI) <20 kg/m^2 or skeletal muscle mass loss were

classified as 'Fearon'. Studies that defined CC in a different way (54, 28.3%), were classified as 'Other'. Other

criteria included different consensus definitions such as the Asian Working Group for Sarcopenia (AWGS) (20)

European Society for Clinical Nutrition and Metabolism (ESPEN) (21), or European Working Group on Sarcopenia

in Older People (EWGSOP2) (22). Other definitions included variations in time statements (1-12 months) or no

statement regarding time frame, different body weight loss degrees (from 2-20%) or no statement about loss degree,

other BMI ranges, or definitions by caloric intake or anorexia. Some other research groups utilized scores and tools

such as the Edmonton Symptom Assessment Score (ESAS), Malnutrition Universal Screening Tool (MUST),

Eastern Cooperative Oncology Group (ECOG), Karnofsky Performance Index, among others. Lastly, 74 (38.2%)

CTx did not specify how they accounted for CC in their study. The distribution of the CC definitions used across

years (1995-2024) is represented in **Figure 2b**. Although the Fearon consensus definition was published in 2011, the

criteria mentioned above have been reported since 2005. It is noteworthy that despite having several criteria and a

consensus definition to diagnose CC, numerous studies fail to describe the diagnostic criteria used.

The current status of CTx stratified by their CC definition criteria and terms used for trial classification are

summarized in **Supplemental Files 5 and 6**. Of 52 CTx completed with results, 29 (55.8%) specified the use of the

Fearon et al. consensus definition. Fourteen trials (26.9%) used definitions other than the Fearon et al., with six of

those trials beginning after the consensus definition was established in 2011. Nine trials (17.3%) did not specify the

criteria used for how they defined CC. Moreover, five trials (9.6%) were classified under "sarcopenia", nine trials

(17.3%) were classified under "weight loss", four trials (7.7%) were classified under "muscle wasting" or "muscle

loss", and four trials (7.7%) were classified under "malnutrition" or "anorexia". Within the completed trials without

results**,** nine trials (16.6%) specified the use of the 2011 Fearon et al. consensus definition. Nineteen trials (35.2%)

used definitions other than the Fearon et al. definition, with 11 of those trials beginning after the consensus

definition was established in 2011. Twenty-six trials (48.2%) did not specify how they defined CC. From these trials

without results, six trials (11.1%) were classified under "sarcopenia", ten trials (18.5%) were classified under

"weight loss", six trials (11.1%) were classified under "muscle wasting" or "muscle loss", and three trials (5.5%)

were classified under "malnutrition" or "anorexia". For the CTx that were still active at the completion of the search,

24 (38.7%) specified the use of Fearon definition and eight trials (12.9%) used other definitions. Thirty (48.4%) did

not specify how they defined CC for their study. Among these active studies, 16 (25.8%) have been classified under

"sarcopenia", one trial (1.6%) under "weight loss", nine trials (14.5%) under "muscle wasting" or "muscle loss", and

one trial (1.6%) classified under "malnutrition" or "anorexia". From the withdrawn trials, two (8.7%) used Fearon

definition, 13 (56.5%) used other definitions, and eight (34.8%) did not specify their criteria to classify CC. Three

trials (13.0%) were classified under "weight loss", one trial (4.4%) was classified under "muscle wasting" or

"muscle loss", and one trial (4.4%) was classified under "malnutrition" or "anorexia".

Sample Size, Cancer Types, and Clinical Approaches Across Clinical Trials

We characterized study participants among CTx to determine the number of registered participants and the

cancer type and stage studied in each trial. Across all trials (n = 191), 72 (37.7%) enrolled between 1-50 patients, 49

trials (25.7%) enrolled between 51-100 patients, 51 (26.7%) enrolled between 101-500 patients, six (3.1%) enrolled

between 501-1000 patients, four (2.1%) enrolled more than 1000 patients, and nine trials (4.7%) did not specify the

total enrollment, most of which were from withdrawn trials. The distribution of patients' enrollment per clinical trial

status is shown in **Supplemental File 7**.

Analysis of the CTx by cancer type show that 50.3% of trials studied CC in a specific type of cancer, 8.9%

studied two cancer types, 5.2% studied three cancer types, 11.5% studied four or greater cancer types, and 24.1%

did not state the cancer type considered for their study (**Figure 3a**). Among these CTx, CC has been studied more

predominantly in lung and gastrointestinal related cancers (pancreatic, gastric, colorectal), either individually or in

combination with other cancer types. In addition, between 10-35% of the trials did not report a concurrent anti-

cancer treatment despite that it is standard of care (data not shown), and the majority did not report the cancer stages

represented in the study (146, representing 76.4%). Of those trials reporting patient cancer stage (n=45), 53.3%

investigated CC in advanced stage cancers (stage III-IV), while other stages were reported between 2-15% (**Figure**

3b). **Supplemental File 8** details the stratification of cancer type and stage per CTx status.

CTx studying the effects of CC employed either an observational (26.7%) or an interventional (73.3%)

study design. The most common intervention was pharmaceutical (33.5%, "drug" on the Figure), followed by

dietary supplements (14.7%), exercise (7.9%), and different multimodal approaches (9.4%). Multimodal

interventions included different combinations of dietary supplements, nutrition, exercise, drugs, counseling, and

psychological support. The interventions used by CTx status are detailed in **Supplemental File 9**. The frequency of

intervention types over time among CC-CTx can be visualized in **Figure 3c**. These trends indicate that drug

treatment interventions remain a popular approach over time. Additionally, a noticeable increase in the use of

exercise and dietary interventions is observed following the Fearon consensus definition in 2011.

Outcome Measures and Tools Used

To understand how CTx assessed CC, we summarized the evaluated parameters and the tools that have

been used to measure them. First, we looked for body weight, a parameter considered in most CC definitions, and

only 42.4% of studies included body weight as an outcome measured (**Figure 4a**). A total of 62.3% of studies report

the use of body composition across all trials, with 25.7% reporting muscle mass, 23.6% reporting lean body mass,

and 11.0% reporting fat mass, among others (**Figure 4b**). To assess body composition, computed tomography (CT)

was the most used technique (18.3%), followed by bioelectrical impedance analysis (BIA, representing 17.3%) and

dual energy x-ray absorptiometry (DXA, 12.6%). Physical function was an outcome measure in 50.3% of CTx,

including measures of muscle function or muscle strength (29.3%), physical activity levels (13.6%), and fatigue

(7.3%) (**Figure 4c**). To assess physical function, 26.2% of the trials used hand grip strength and 13.1% used a 6- or

10-minute walk test, with other methodologies used less frequently. Quality of life (QoL) was specified by 37.2% of

the trials (**Figure 4d**), and it was mostly evaluated by the Eastern Cooperative Oncology Group (ECOG)

Performance Status Scale (33.0%), the European Organization for the Research and Treatment of Cancer QoL

Questionnaire (EORTC QLQ, 20.9%), and by the Functional Assessment of Anorexia/Cachexia Treatment

(FAACT, 13.6%), among others. Nutrition and appetite were mentioned by 30.4% of the CTx and dietary recall or

food diaries were the methodologies most used (9.4%) (**Figure 4e**). Molecular measures were mentioned in 45.5%

of the studies, with ~45% of them stated in CTx studying CC in the last 5 years (**Figure 4f**). Of those studies

reporting molecular measures, most studies assessed blood samples (44.5%) while some reported use of muscle

biopsies (11.0%). Lastly, 21.5% of CTx specified the measure of safety and tolerability, and 19.4% reported survival

(**Figures 4g and 4h**). Outcomes measured and other tools that have been used to a lesser extent are detailed by

clinical trial status on the **Supplemental Files 10 and 11**.

Discussion

While significant progress has been made in the understanding of the pathogenesis of CC and in

establishing potential therapeutic interventions, CC significantly decreases quality of life and remains a main cause

of death for many cancer patients, while successful treatment strategies have not yet been identified. Through

evaluation of CC CTx, we looked to assess representation of aspects of human heterogeneity that are not well-

addressed in carefully controlled animal studies and assess changes in study approaches and improvements in these

metrics over time. Our findings suggest a large knowledge gap in how CC differs across patient populations and

how this impacts health outcomes. Expanding CTx to improve representation across patient populations may

improve progress toward effective treatments for CC.

Limited Diversity in Demographic Factors Reporting

It was not until 1993 that in the United States (U.S.), the National Institutes of Health (NIH) Revitalization

Act was passed requiring recruitment and inclusion of women and racial and ethnic minorities in clinical research

studies. In 2011, almost 20 years afterwards, the Institute of Medicine of the National Academies suggested

reporting results by biological sex be required for the publication of CTx results, as lack of reporting had

significantly slowed the advancement of women's health (23, 24). Recognizing sex-based biological differences is

essential for understanding mechanistic and pathophysiological divergence during the onset and progression of pre-

clinical studies of CC (14, 15). Therefore, the acknowledgment of women's inclusion in CTx alone is insufficient. In

2015, the NIH announced the requirement for sex to be considered as a biological variable in NIH-supported

preclinical and clinical studies. Herein, we identified that across all CC-CTx completed with results, 5.8% did not

report sex at all, while 59.6% of the total patient population was male. However, between 2015-2024, the percentage

of enrolled females increased to 48.4% of patients, reflecting increased equality of enrollment of males and females

in CTx over time suggesting enhanced equity. At present, significant advancements have been made in

understanding the pathogenesis of CC, but still very little is known about how biological sex as a variable affects

patient outcomes with CC. Despite improved equality of enrollment, trends are yet to come to fruition for the

consideration of biological sex as a variable in data analyses, as in twenty-nine years we only identify one CTx

reporting sex subgroup analysis. Our findings highlight that despite many positive changes in the conduct of clinical

studies, inequalities persist, likely contributing to suboptimal progress. Understanding sex as a biological variable in

clinical CC will ultimately increase the ability to accurately model the complications of the human condition, and in

turn, develop effective treatments improving health outcomes for both males and females with CC.

Approximately one-third of CTx completed with results reported race, while less than one-tenth reported

ethnicity. Of CTx reporting race and ethnicity, there was a lack of racial and ethnic diversity among enrolled

patients. Among CTx reporting race, 92.6% of all patients were White. In CTx reporting ethnicity, 100% focus

solely on Hispanic/Latino individuals, neglecting other ethnicities. This homogeneity fails to reflect the true

diversity of the general population and highlights the scarcity of information about CC in non-white individuals,

severely limiting the understanding of how race and ethnicity affect CC. Recent calls for applications from the

National Cancer Institute in the U.S. highlight the dire need to understand cancer burden in racial and ethnic

minorities represented in the U.S. population. Although contributions to this area of research are currently minimal,

there have been significant recent findings that provide evidence for possible genetic, environmental, and

socioeconomic differences across race and ethnicity warranting further studies. For instance, across all stages of

lung cancer, Black race strongly increased the odds of experiencing pretreatment weight loss (17) and in patients

with advanced lung cancer, Black race and Hispanic ethnicity presented with greater risk of developing CC (16).

The Florida Pancreas Collaborative has made great strides in understanding racial and ethnic differences in

pancreatic CC, reporting greater frequency of cachexia markers in Blacks with concurrent higher cancer incidence

and mortality and shorter survival time (18, 19). Additionally, Hispanic/Latino patients experienced increased

survival time compared to non-Hispanic White individuals and Black individuals (19). This suggests underlying

genetic, environmental, socioeconomic, and/or yet unknown factors predisposing patients to increased risk of

developing CC, which may differ beyond cancer type. Despite these observations, no CTx reported clinical

outcomes by race or ethnicity in a subgroup analysis. As with biological sex, race and ethnicity should be included

in subgroup analyses for clinical outcomes across CC studies; however, studies must be statistically powered for

these analyses and thus recruitment of a racially and ethnically diverse study population is critical.

Inconsistent Cancer Cachexia Definitions and Varied Diagnostic Criteria

Despite that a consensus CC definition was published in 2011 (3), we found considerable heterogeneity in

the definition and diagnostic criteria used across CTx, with only 33.5% of studies using the consensus definition

criteria to classify CC. Additionally, there have not been improvements in these metrics over time. Significant

variations in classification persist, with a significant portion (38.2%) of CTx not reporting the details of their

classification methods, which presents an enormous challenge in interpreting and comparing results across CTx.

Additionally, a significant portion of studies use description of symptoms to define CC, including sarcopenia,

muscle wasting/loss, anorexia, and malnutrition. The inconsistency in the terminology and diagnostic guidelines

may lead to misdiagnoses, absent medical records, and inaccurate reports of cachexia incidence, likely concealing

the visibility of the impact of CC on public health.

Consideration of Cancer Type Effects on Cancer Cachexia Patient Outcomes

We found that many CC-CTx have relatively small samples sizes, with most CTx in this review recruiting

100 participants or fewer. CTx on CC require many clinical assessments in patients who are often unwilling or

unable to contribute to research, and thus it can be challenging to recruit patients on a large scale. However, efforts

to recruit larger samples sizes could increase the ability to better understand CC in diverse populations and achieve

statistical power required for subgroup analyses. Additionally, almost one-third of CTx encompassed in the present

work included more than one cancer type in the study population. Recently, it has become more apparent that

different cancer types likely exhibit different patterns of tissue wasting and progression. This is clear not only in

clinical findings, but also in preclinical models of CC where different tumor types exhibit mechanistic differences

during disease development and progression (15). Thus, combining different cancer types without offering stratified

analyses likely conceals cancer type-specific changes. However, we report that 41.5% of completed CTx and 64.5%

of active CTx focused on a single cancer type, suggesting the evolving knowledge of varying clinical cancer

characteristics and treatments has informed the design of CTx to improve our understanding of cancer type-specific

CC outcomes. In addition, although CC is commonly diagnosed at advanced cancer stages, 76.4% of CTx included

in this review did not report a specific cancer stage. Moreover, 73.3% of the studies incorporated in this analysis

were interventional. Consequently, the lack of consideration for cancer stage as a clinical variable may skew the

results of the interventions, particularly by overlooking pre-cachectic states in which knowledge remains limited.

Growing Focus on Exercise Interventions in Cancer Cachexia Clinical Trials

The use of pharmaceutical approaches to treat or prevent CC remains the most popular intervention among

CC-CTx. However, we observed a growing focus on exercise interventions in the study of CC, either alone or in

combination with other treatment paradigms in recent years. Additionally, there was a relative increase in the

measure of physical function and quality of life, reinforcing the importance of maintaining muscle quality to

enhance well-being in patients with CC. Along with a growing understanding of the molecular mechanisms of tissue

wasting informed by preclinical studies, we observed an increase in the use of molecular measures in CTx. Broadly,

we observed wide variation in the outcome measures reported, even those non-invasive, affecting the ability to

analyze systematic reviews reinforcing clinical conclusions.

Body Weight is Underutilized as an Outcome in Cancer Cachexia Assessment

Although body weight and composition are fundamental measures of CC, we identified a surprisingly low

frequency of studies reporting body weight (42.4%) and body composition (62.3%). With more accurate tools for

body composition measurement, such as CT and MRI, recent studies have brought attention to body composition

changes across the cancer and treatment continuum within certain CC-associated cancer types (25, 26). Moreover,

body composition, reported by CT, is often a primary outcome of CC-CTx, and sexual dimorphisms in the

distribution of fat and muscle tissue in the thoracic and lumbar compartments must therefore be considered in the

interpretation of body composition assessment. We found that active CTx are more frequently using these

measurement tools, which will provide the field with a better understanding of body compositional changes during

disease progression. In light of these findings, consistency of diagnostic criteria and outcome reporting across future

CTx are essential in advancing our understanding of CC and improving clinical outcomes. To address this need, the

Cancer Cachexia Society has convened a CTx Advisory Board, a multidisciplinary group of CC experts to provide

transparent guidance on clinical trial design through a confidential and comprehensive review process.

Conclusions

Overall, a substantial gap persists in our understanding of CC in non-white and female populations. The

report of CC definition and transparency in diagnostic criteria, along with the absence of stratified variables such as

cancer type and biological sex are deficient, complicating the comparison and interpretation of CTx results. Lastly,

the arrival of new techniques and methodologies used for cachexia assessment reinforces the need for a regularly

revised and updated consensus definition and diagnosis criteria to align with new research findings and cutting-edge

technology.

Assessment of active, completed, and terminated CTx in this review reveals both the recent advancements

and remaining gaps in clinical CC research. Addressing the lack of subgroup analyses based on biological sex is

essential for comprehensive understanding of the differences in the development and progression of CC between

males and females. Similarly, investigating the diverse manifestations of CC among various racial and ethnic groups

is crucial for eliminating health disparities and developing treatment options that are effective in diverse

populations. Standardizing CC diagnostic criteria across future clinical trials can enhance the reliability and effective

comparability of trial findings. Additionally, small and heterogeneous study samples make drawing conclusions

challenging, emphasizing a need for larger recruitment and modified trials designs that minimize the combination of

cancer types, staging, and concurrent therapies within study populations. Current trends in intervention methods

mirror advancing knowledge of CC, with corresponding changes in outcome measures. However, significant gaps

and inconsistencies persist creating a dire need for continued improvement. By addressing these issues and

proposing more effective and standardized study designs, we aim to contribute to ongoing advancement of our

understanding of CC and ultimately improve patient outcomes.

Acknowledgements

The authors thank all the faculty, staff, and students of the Exercise Science Research Center at the University of
Arkansas.

Authors' contributions

ARC: drafting all the figures and final manuscript, reviewing, and editing. **KP:** creating the database, drafting the
manuscript, reviewing, and editing. **DBS:** creating figures, writing, reviewing, and editing. **BH**: conceptualization,
reviewing, and editing. **NPG:** conceptualization, resources, supervision, reviewing and editing. All authors reviewed
and approved the final manuscript.

Ethics approval and consent to participate

NA

Consent for publication

NA

Data availability

Correspondence and requests for materials should be addressed to Nicholas P. Greene.

Competing interests

The authors declare no conflict of interest.

Funding information

References

1.   Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global

cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36

cancers in 185 countries. CA: A Cancer Journal for Clinicians. 2024;74(3):229-63.

2.   Society AC. Cancer facts & figures 2023. Atlanta: American Cancer Society; 2023.

3.   Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition

and classification of cancer cachexia: an international consensus. Lancet Oncol.

2011;12(5):489-95.

4.   Stewart GD, Skipworth RJ, Fearon KC. Cancer cachexia and fatigue. Clin Med (Lond).

2006;6(2):140-3.

5.   Kilgour RD, Vigano A, Trutschnigg B, Lucar E, Borod M, Morais JA. Handgrip strength

predicts survival and is associated with markers of clinical and functional outcomes in advanced

cancer patients. Supportive Care in Cancer. 2013;21(12):3261-70.

6.   Andreyev HJ, Norman AR, Oates J, Cunningham D. Why do patients with weight loss

have a worse outcome when undergoing chemotherapy for gastrointestinal malignancies? Eur J

Cancer. 1998;34(4):503-9.

7.   Dewys WD, Begg C, Lavin PT, Band PR, Bennett JM, Bertino JR, et al. Prognostic effect

of weight loss prior to chemotherapy in cancer patients. Eastern cooperative oncology group.

Am J Med. 1980;69(4):491-7.

8.   Karuppannan M, Muthanna FMS, Mohd Fauzi F. Breaking Down Cachexia: A Narrative

Review on the Prevalence of Cachexia in Cancer Patients and Its Associated Risk Factors. Nutr

Cancer. 2024;76(5):404-18.

9.   Holmes S. Understanding cachexia in patients with cancer. Nursing Standard (through

2013). 2011;25(21):47-56; quiz 8.

10.   Vagnildhaug OM, Balstad TR, Almberg SS, Brunelli C, Knudsen AK, Kaasa S, et al. A

cross-sectional study examining the prevalence of cachexia and areas of unmet need in

patients with cancer. Supportive Care in Cancer. 2018;26(6):1871-80.

11.   Bruera E. Anorexia, cachexia, and nutrition. BMJ ens. 1997;315:1219-22.

12.   Baracos VE. Bridging the gap: are animal models consistent with clinical cancer

cachexia? Nat Rev Clin Oncol. 2018;15(4):197-8.

13.   Ballarò R, Costelli P, Penna F. Animal models for cancer cachexia. Curr Opin Support

Palliat Care. 2016;10(4):281-7.

14.   Zhong X, Zimmers TA. Sex Differences in Cancer Cachexia. Current Osteoporosis
Reports. 2020;18(6):646-54.

15.   Silva FMd, Cabrera AR, Greene NP. Exploring Heterogeneity: A Dive into Pre-Clinical
Models of Cancer Cachexia. American Journal of Physiology-Cell Physiology.0(0):null.

16.   Olaechea S, Liu A, Sarver B, Gilmore LA, Alvarez C, Yazdanbakhsh K, et al. Racial,
Ethnic, and Socioeconomic Characteristics Independently Predict for Cachexia Risk and
Associated Survival Outcomes in Stage IV NSCLC: A Brief Report. JTO Clinical and Research
Reports. 2023;4(4):100496.

17.   Lau SKM, Gannavarapu BS, Carter K, Gao A, Ahn C, Meyer JJ, et al. Impact of
Socioeconomic Status on Pretreatment Weight Loss and Survival in Non-Small-Cell Lung
Cancer. J Oncol Pract. 2018;14(4):e211-e20.

18.   Permuth JB, Clark Daly A, Jeong D, Choi JW, Cameron ME, Chen DT, et al. Racial and
ethnic disparities in a state-wide registry of patients with pancreatic cancer and an exploratory
investigation of cancer cachexia as a contributor to observed inequities. Cancer Med.
2019;8(6):3314-24.

19.   Permuth JB, Park MA, Chen D-T, Basinski T, Powers BD, Gwede CK, et al. Leveraging
real-world data to predict cancer cachexia stage, quality of life, and survival in a racially and
ethnically diverse multi-institutional cohort of treatment-naïve patients with pancreatic ductal
adenocarcinoma. Frontiers in Oncology. 2024;14.

20.   Chen L-K, Woo J, Assantachai P, Auyeung T-W, Chou M-Y, Iijima K, et al. Asian
Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and
Treatment. Journal of the American Medical Directors Association. 2020;21(3):300-7.e2.

21.   Arends J, Baracos V, Bertz H, Bozzetti F, Calder PC, Deutz NEP, et al. ESPEN expert
group recommendations for action against cancer-related malnutrition. Clin Nutr.
2017;36(5):1187-96.

22.   Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia:
revised European consensus on definition and diagnosis. Age Ageing. 2019;48(1):16-31.

23.   Lee E, Wen P. Gender and sex disparity in cancer trials. ESMO open. 2020;5:e000773.

24.   Institute of Medicine Board on Population H, Public Health P. The National Academies
Collection: Reports funded by National Institutes of Health. Sex-Specific Reporting of Scientific
Research: A Workshop Summary. Washington (DC): National Academies Press (US)

Copyright © 2012, National Academy of Sciences.; 2012.

25.   Al-Sawaf O, Weiss J, Skrzypski M, Lam JM, Karasaki T, Zambrana F, et al. Body
composition and lung cancer-associated cachexia in TRACERx. Nat Med. 2023;29(4):846-58.

26.   Babic A, Rosenthal MH, Sundaresan TK, Khalaf N, Lee V, Brais LK, et al. Adipose tissue

and skeletal muscle wasting precede clinical diagnosis of pancreatic cancer. Nat Commun.

2023;14(1):4317.

Figure Legends

Figure 1. Demographics reported in clinical trials completed with results published from 2004-2023. a)

Percentage of clinical trials including demographic data for age, biological sex, race, and ethnicity in the study

population. **b)** Percentage of females and males included in clinical trials with reported results across years by

clinical trial completion date (2004-2023). **c)** Percentage of White, Black, and Other races reported; others include

0.09% Native Hawaiian/Pacific Islander, 0.52% Asian, 0.91% Pardo, and 2.56% unknown (data not shown).

"Fearon" marker indicates the year of the publication of the cancer cachexia consensus definition. "Sex as

Biological Variable" indicates the year of the NIH expectation to consider sex as a biological variable in NIH-

supported preclinical and clinical studies.

Figure 2. Cancer cachexia diagnostic criteria used in trials from 1995-2024. a) Studies that used weight loss

greater than 5% in less than six months as their criteria, or weight loss greater than 2% in individuals already

showing depletion with body mass index (BMI) <20 kg/m^2 or skeletal muscle mass loss were classified as 'Fearon';

studies defining cachexia in a different way were classified as 'Other'. **b)** Distribution of the cachexia definitions

used across years.

Figure 3. Study population and approach. a) Cancer type and **b)** stages reported in clinical trials related to cancer

cachexia. **c)** Clinical trials approach. LOESS (locally weighted smoothing) regression analysis from

pharmacological (drugs), exercise, dietary, and multimodal interventions across years (1995-2024) investigated in

cancer cachexia clinical trials. "Fearon" marker indicates the year of the publication of the cancer cachexia

consensus definition.

Figure 4. Clinical trials outcomes across years (1995-2024). Number of trials and LOESS (locally weighted

smoothing) regression analysis from **a)** body weight, **b)** body composition, **c)** physical function, **d)** quality of life

(QoL), **e)** nutrition and appetite, **f)** molecular measures, **g)** safety and tolerability, and **h)** survival measured in cancer

cachexia clinical trials. "Fearon" marker indicates the year of the publication of the cancer cachexia consensus

definition.

Supplementary Information

Supplemental File 1. List of clinical trials included in this review. Identification number is the NCT code under

which the clinical trial is registered in ClinicalTrials.gov. The status is completed with (w/) results, completed

without (w/o) results, still active, or withdrawn (terminated) before completion.

Supplemental File 2. Summary of search results by terms and status of the clinical trials.

Supplemental File 3. This table list the reasons for withdrawn or anticipated termination before completion of

clinical trials studying cancer cachexia registered in ClinicalTrials.gov.

Supplemental File 4. Phases and status of clinical trials exploring cancer cachexia effects on ClinicalTrials.gov in

Phase1-4 or No Phase.

Supplemental File 5. Clinical trials registered on ClinicalTrials.gov that used Fearon's definition or any other

criteria to diagnose cancer cachexia within each status.

Supplemental File 6. Clinical trials stratified by clinical trial status studying cachexia but classified under other

terms different than "cachexia".

Supplemental File 7. Study population of clinical trials registered on ClinicalTrials.gov investigating cancer

cachexia.

Supplemental File 8. Cancer type and cancer stage of clinical trials registered on ClinicalTrials.gov investigating

cancer cachexia.

Supplemental File 9. Clinical trials approach and intervention utilized: drug testing, dietary supplement,

exercise/physical therapy, a combination of 2 or more interventions (multimodal), others, or none or non-stated

intervention of clinical trials registered on ClinicalTrials.gov investigating cancer cachexia.

Supplemental File 10. Outcome measures of cachexia in the clinical trials registered on ClinicalTrials.gov by

status: Body weight, body composition, physical function, quality of life, nutrition/appetite, molecular, safety and

tolerability, and survival.

Supplemental File 11. Measurement tools used to investigate cachexia by the clinical trials registered on

ClinicalTrials.gov by status.