Pharmacokinetic analysis of intermittent rapamycin administration in early-stage Alzheimer's Disease

Helen Annervik Wallgren[1,2], Miia Kivipelto[1,4,5,6], Pontus Plavén-Sigray[2,3*], Jonas E. Svensson[1,2*]

1. Theme Inflammation and Aging, Karolinska University Hospital, Stockholm, Sweden

2. Department of Clinical Neuroscience, Karolinska Institutet and Stockholm Health Care Services, Region Stockholm, Stockholm, Sweden.

3. Neurobiology Research Unit, Copenhagen University Hospital, Copenhagen, Denmark.

4. Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences, and Society, Karolinska Institutet, Stockholm, Sweden.

5. Ageing Epidemiology Research Unit (AGE), School of Public Health, Faculty of Medicine, Imperial College London, UK.

6. Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland.

*Contributed equally to this work

Corresponding author:

Jonas Svensson, Email: jonas.svensson@ki.se

Karolinska Institutet, Bioclinicum J4:14, Solnavägen 30, 171 64 Solna

Trial registration: ClinicalTrials.gov (NCT06022068) and EudraCT (2023–000127-36).

Keywords: rapamycin, sirolimus, mTOR inhibitor, Alzheimer´s disease, neurodegenerative disease, pharmacokinetics, intermittent dosing

Abstract

Background: rapamycin, an mTOR inhibitor used clinically for immunosuppression, shows promise for repurposing in age-related disorders including Alzheimer's disease (AD). While the pharmacokinetics of daily rapamycin are well-characterized in transplant populations, limited data exist on intermittent dosing regimens in patients with neurodegenerative conditions.

Method: this open-label pilot study investigated the pharmacokinetic properties of weekly oral rapamycin in 13 patients with early-stage AD. Participants received 7 mg weekly (11 patients) or reduced doses (2 mg and 4 mg; 2 patients) for 26 weeks. Blood concentrations were measured at four timepoints (pre-dose/Cmin, and 1-, 3-, and 48-hours post-dose) during week 13.

Results: moderate interindividual variability was observed across timepoints (coefficient of variation was 0.28-0.40), with the 48-hour sample showing the lowest variability (CoV = 0.28) and strongest correlation with Cmin from the previous dosing (r = 0.72). Estimate of terminal half-life (68.9 ± 13.6 hours) aligned with previous studies.

Conclusions: blood concentrations at Cmin were below immunosuppressive levels in all participants. Our findings suggest that weekly rapamycin administration in AD patients results in acceptable pharmacokinetic variability, supporting fixed-dose regimens in future trials. The 48-hour post-dose measurement appears optimal for monitoring blood concentrations. Additionally, our investigation into blood-brain barrier permeability revealed methodological challenges in directly measuring rapamycin in cerebrospinal fluid due to analytical sensitivity limitations. The foremost limitation of this study was the sparse blood sampling schedule, with Cmin collected from the previous dosing occasion which prevented a complete AUC-calculation.

1 Introduction

Rapamycin (also known as sirolimus), an mTOR inhibitor and immunosuppressive drug, has been in clinical use for nearly three decades in preventing rejection after organ transplantation (1). During the early 2000s, a series of experiments showed that rapamycin prolonged the lifespan in several model organisms, including yeast, C. elegans, and the fruit fly Drosophila melanogaster (2–4). In 2009 treatment with rapamycin was first shown to prolong lifespan of mice (5). These findings have since been replicated repeatedly (6,7) and subsequent animal research has shown a positive effect of rapamycin treatment on several age-related conditions, including neurodegenerative disease, such as in transgenic mouse models of Alzheimer's disease (8,9).

These promising pre-clinical results have led to calls for repurposing rapamycin, and structurally similar mTOR inhibitors known as "rapalogs", as treatments in a range of age-related diseases, with several studies already underway. Human clinical trials have been conducted to assess potential benefits of rapamycin and rapalogs in multiple age-related conditions, such as immunosenescence and decline in lean tissue mass (10–13) with more trials ongoing (14). Other trials have focused on neurodegenerative diseases, such as amyotrophic lateral sclerosis (15), multiple system atrophy (16), multiple sclerosis (17), and Alzheimer's disease (clinicaltrial.gov ID: NCT04629495).

When rapamycin is used for immunosuppression, therapeutic drug monitoring (TDM) using whole blood trough values is recommended (18). The reason for this is a relatively narrow therapeutic window, together with an substantial observed interindividual variation in bioavailability and elimination (19). It is unclear whether the interindividual variation presents the same challenge in new non-immunosuppressive indications where other dosing regimens have been used (11,13,17). Therefore, the question of whether TDM is necessary or if fixed dosing would be sufficient remains to be determined.

For treatment in new indications where immunosuppression is not desired, it has been suggested to administer rapamycin at lower doses but also using an intermittent dosing regimen. These proposals are based on rapamycin's specific binding profile to subtypes of the mTOR complex, where it has been hypothesized that intermittent dosing could reduce side effects while maintaining treatment efficacy (14). Whereas the pharmacokinetics of daily dosing of rapamycin is well described in healthy populations and in patients following organ transplantation, less is known about the pharmacokinetic properties in other groups such as patients with neurodegenerative disease. Additionally, human data on rapamycin's pharmacokinetics under intermittent dosing remain scarce, as does evidence regarding its ability to cross the blood-brain barrier (BBB).

Here we investigate the pharmacokinetic properties of oral tablet rapamycin administered once weekly to patients with early Alzheimer's disease. We aim to:

1. Provide data on the interindividual variability in blood concentration in order to guide decisions on whether to use a fixed dose or TDM in future rapamycin-trials applying intermittent dosing regimens.
2. Assess the optimal time point for blood concentration monitoring during intermittent dosing.
3. Provide data on rapamycin's pharmacokinetic properties during long-term intermittent treatment, including BBB permeability.

Methods

Clinical trial design

The study is part of a single-center, open-label, one-arm pilot clinical trial where treatment with rapamycin was evaluated in patients with early-stage Alzheimer's disease (20). The study was conducted in accordance with the Declaration of Helsinki and the International Conference on

Harmonisation for Good Clinical Practice (ICH GCP) and was approved by the Swedish Medical

Products Agency (5.1–2023-8283), and the Swedish Ethical Review Authority (2023–03075-02).

Before study start, the trial was registered at ClinicalTrials.gov (NCT06022068) and EudraCT

(2023–000127-36). Written informed consent was collected from all participants and their

designated study partners before initiating any study procedures.

The study was conducted as a collaboration between Karolinska Institutet and the Karolinska

University Hospital, Theme Inflammation and Aging. Patients followed at the Solna Memory clinic

were eligible for inclusion. A full description of inclusion and exclusion criteria has been

published elsewhere (20). In short, patients between 50-80 years, diagnosed with amyloid-

positive Alzheimer's disease, either with mild cognitive impairment (MCI) or early-stage

Alzheimer's dementia (not above stage 4 according to the National Institute of Aging-Alzheimer's

Association 2018 criteria) (21), and with a normal or clinically acceptable medical history and

physical examination were eligible to participate in the trial. All study participants underwent

routine laboratory testing and brain imaging with MRI prior to the first dose of the study drug.

Throughout the study, participants were continuously monitored for safety and adverse events

through regular clinical follow-up visits, including safety blood tests.

Dosing regimen and sampling procedures

Participants were administered a weekly dose of rapamycin for 26 weeks, starting with an initial

dose of 3 mg, which, if well tolerated, was increased to the target dose of 7 mg in the second

week. Participants were prescribed 1 mg tablets of rapamycin (Tablet Rapamune®) which they

collected at the hospital pharmacy. A study nurse dispensed the drug into a pill organizer which

participants were asked to bring to clinical check-ups where it was verified that the tablets were

taken according to the study protocol.

At the mid-trial follow-up visit (week 13), blood samples for sirolimus concentration were

collected at four time points over 48 hours. Blood concentration tests were taken just prior to the

weekly dose (C_{min}), and at 1-, 3- and 48-hours post-dose. The 1h sample was chosen with the aim

of capturing C_{max} based on estimates from prior studies (22,23). Participants were fasting at the

time for the pre-dose test due to simultaneous sampling of routine laboratory variables. Following

the blood draw the participant was instructed to swallow the weekly dose with a glass of water

and received immediately thereafter a light meal. The pre-dose, 1- and 3-hour blood samples

were collected by a study nurse as part of the study visit. The participants were given the option

to select a caregiver closer to their home for the 48-hour blood test. For assessment of BBB

permeability of the study drug, a lumbar puncture for collection of a cerebrospinal fluid (CSF)

sample was performed before initiation of the study treatment and again within 28 days of the last

dose.

Sirolimus whole blood concentrations were analyzed at Karolinska University Hospital

Laboratory, Huddinge, Medical Unit Clinical Pharmacology, a unit which routinely performs

clinical sirolimus concentration analyses. Analysis was performed on a Thermo Fisher (Thermo

Scientific, Waltham, MA, USA) liquid chromatography–tandem mass spectrometry (LC-MS/MS)

machine. All blood samples were collected, stored and transported as per clinical routine

outlined in written instruction supplied by the laboratory.

Blood brain-barrier-permeability analysis

At steady state the CSF concentration of a drug that passes freely over the BBB can be expected

to be similar to the free concentration in blood (24). Given that the free fraction of rapamycin is

only about 0.17% of whole blood (25), it is uncertain whether standard laboratory methods can

reliably quantify it in CSF. The lower limit of quantification (LLOQ) for rapamycin in whole blood

using LC-MS/MS—the gold standard method—typically ranges from 0.5 to 1 ng/mL (26), which is

more than an order of magnitude higher than expected CSF concentrations at therapeutic doses.

To determine whether lower CSF concentrations of rapamycin can be quantified, we conducted

a method assessment as part of this pharmacokinetic study. Rapamycin (purity > 98%, Thermo

Scientific Chemicals) was used to prepare calibration standards in blank CSF (i.e., CSF from

individuals not treated with rapamycin). The study was performed at Drug Discovery and

Development Platform, SciLifeLab in Uppsala, using a triple quadrupole mass spectrometer

(Xevo™ TQ-S micro, Waters Corp.), with the aim of determining the LLOQ of rapamycin in CSF.

Further methodological details are provided in the Supplementary Materials.

Pharmacokinetic modeling and statistical analysis

The interindividual variation in blood concentration of rapamycin was assessed through

calculating the standard deviation (SD) and the coefficient of variation (CoV) at each of the

sampling times separately (C_{min}, 1h, 3h, and 48h tests). Since the C_{min} sample was collected

from a separate dosing occasion a complete area under the curve (AUC) of the blood

concentration was not possible to calculate. The concentration data collected post dosing was

performed at 1h, 3h and 48h, allowing for calculation of AUC_{0-48h}. However, given the relatively

sparse sampling scheme with only one sample after the distribution phase, together with the

substantially higher concentrations in the two early samples compared to the 48h sample, the

observed variation in AUC_{0-48h} values will be mainly influenced by the variation in the 3h sample.

Consequently, this makes it not a valid measure of the true intraindividual variation in AUC.

In order to estimate terminal half-life (terminal t½), more than one concentration sample

collected after the distribution phase (i.e., during the elimination phase) of the drug is needed. To

overcome this, we made the assumption that C_{min} is stable between doses and hence used C_{min}

from the previous dosing occasion to provide an additional time point (denoted 168h below). To

underscore that the terminal half-life calculation was based on blood concentration values from

two separate dosing occasions this parameter is denoted as "*pseudo-*" terminal t½ below, and it

was calculated as:

$pseudo\text{-}terminal\ t½\ = \frac{0,693}{\lambda}$

where λ denotes the terminal elimination rate constant, calculated as the negative slope of the

logarithmed concentration versus time for the two last time points (i.e., 48h and 168h).

In order to assess the association between the different post-dosing samples, and the stability

over time, Pearson's correlation coefficient was calculated between C_{min} from the previous

dosing occasion and concentration estimates at the sampling times post-dose i.e., at 1h, 3h, and

48h. The following heuristics were used to describe the strength of association between

variables: a correlation estimate between 0–0.19 was described as negligible, 0.2–0.39 as weak,

0.40–0.59 as moderate, 0.6–0.79 as strong and 0.8–1 as very strong (25).

Pharmacokinetic calculations were performed using Microsoft Excel (Microsoft Corporation,

Redmond, WA, USA). Data visualization and additional statistical analyses were conducted using

R software (version 4.2.2, R Foundation for Statistical Computing, Vienna, Austria).

Results

14 patients, eight women and six men with an average age of 60.9 ± 4.2 (mean ± SD) years were

included in the study. Six patients had a diagnosis of MCI and eight of dementia, with mean time

since diagnosis of 6.6 ± 4.2 months and Montreal Cognitive Assessment (MoCa) score of 23.8 ± 3

(maximum 30) at inclusion. The participants' mean weight was 75.5 ± 16 kg with body surface

area of 1.9 ± 0.2 m^2.

One participant (male) did not tolerate the study drug due to nausea and was excluded from the

trial two weeks after initiating the treatment (i.e., before sampling of blood concentration).

Consequently, the results are based on the remaining 13 participants, who all completed the 26-

week treatment. Two participants (both female) received a lower dose of rapamycin (2 and 4 mg

respectively). The reason for the reduced dose was side effects in one participant (vertigo

following initial dose) and safety reasons (chronically elevated aspartate transaminase) for the

second participant. For both participants the treatment was well tolerated at these doses. Three

participants did not perform the 3h sample, one of whom also provided blood sample at 24h

instead of 48h. All participants were on a stable dose of a cholinesterase inhibitor (minimum of 4

weeks of target dose before initiation of study drug). No serious adverse events were observed

during the trial.

Average concentration values from participants taking 7 mg rapamycin (11 out of 13) at the

different timepoints are reported in Table 1, together with the results from the two participants

taking reduced doses. For participants receiving full dose C_{min} was 1.17 ± 0.44 ng/mL (CoV =

0.38), and 1h after dosing 26.73 ± 10.74 ng/mL (CoV = 0.4).

Figure 1. *Rapamycin whole blood concentration before intake of weekly dose (C_{min}) and at three*

different time points following dosing (1, 3 and 48 hours post dose, with one participant tested at

24h instead of 48h post dose). Data from participants on a weekly dose of 7 mg in gray, with the

participants on 2 and 4 mg doses shown in yellow and blue respectively.

Table 1. Rapamycin whole blood concentration.

Time point	mean ±SD (ng/mL)	CoV
Cmin (~168h)		
7mg	1,17±0,44	0.38
4mg	0.58	NA
2mg	NA	NA
1 -hour		
7mg	26,7±10,7	0.4
4mg	14	NA
2mg	9.5	NA
3-hour		
7mg	23,5±8,85	0.38
4mg	13	NA
2mg	7.2	NA
48-hour		
7mg	4,11±1,15	0.28
4mg	1.7	NA
2mg	1.3	NA

Rapamycin whole blood concentration for participants taking a weekly rapamycin dose of 7 mg (n = 11), 4 mg (n = 1) and 2 mg (n = 1). CoV, Coefficient of Variation; NA, not available, SD, Standard Deviation

165 Correlation analysis of blood concentrations at different time points for participants taking 7 mg

166 per week are reported in Figure 2. The C_{min} value had the strongest correlation to the 48h sample

167 (r = 0.72) and the weakest correlation to the 1h sample (r = 0.076). A very strong correlation was

168 seen between the 1h and 3 h sample (r = 0.87) and between the 3h and 48h sample (r = 0.80).

Figure 2. *Correlation estimates for rapamycin whole blood concentration before intake of weekly dose (Cmin) and at three different time points following dosing (1, 3 and 48 hours post dose). Only participants taking 7mg per week is included in this analysis.*

For participants receiving 7 mg, pseudo-terminal t½ was 68.85 ± 13.64 hours (CoV 0.20). The participant missing the 48-hour sample was not included in this calculation. The participant receiving a reduced dose of 4 mg showed a pseudo-terminal t½ of 77.3 hours and for the participant receiving 2 mg we could not estimate pseudo-terminal t½ due to a non-quantifiable blood concentration at Cmin.

For method development of rapamycin quantification in CSF, an LLOQ of 20 ng/mL was achieved.

Sample pre-concentration did not significantly enhance sensitivity, yielding an LLOQ of 10–20

ng/mL. See Supplementary Material for further details.

Discussion

The purpose of the study was to characterize the pharmacokinetics of rapamycin when given

weekly to patients with early Alzheimer's disease. The drug is of high interest in the field of aging

research, with multiple ongoing studies evaluating its safety and efficacy for new indications (14).

However, limited data on the pharmacokinetic properties for long-term, intermittently

administered rapamycin exist, with only a few studies evaluating the topic (26,27). In our study,

blood concentrations showed moderate interindividual variance, with CoV ranging from 0.28 to

0.40 across timepoints. The 48h sample demonstrated the lowest variance (CoV = 0.28) and the

strongest correlation with C_{min} from the previous dosing (r = 0.72), suggesting 48h as the time

point of choice for control of blood concentration based on this series of samples.

When rapamycin is used for immunosuppression following organ transplantation, TDM using

C_{min}/trough values is recommended based on two key factors: the drug's narrow therapeutic

window in this indication (5-15 ng/mL) and a substantial observed interindividual

pharmacokinetic variability (19). At blood concentrations below 5 ng/mL, immunosuppression

may be insufficient, while levels exceeding 15 ng/mL increase the risk of adverse effects and

toxicity. For age-related disorders where immunosuppression is not desired, intermittent and/or

lower dosing of rapamycin and rapalogs is typically employed (10–13), requiring reassessment of

these factors. In our study, the 11 patients receiving weekly 7 mg doses, blood concentrations at

48h averaged 4.11 ± 1.15 ng/mL (range: 2.7–6.6 ng/mL, CoV = 0.28), with only two participants

exceeding 5 ng/mL. Notably, no participants exceeded 5 ng/mL at C_{min}. While earlier time points

showed moderately higher variability (CoV = 0.38–0.40), this still remains lower than that

observed in transplant patients on rapamycin treatment (28) as well as with specific commonly

used drugs such as certain beta blockers (29) and antidepressants (30).

Although the minimum effective concentration for indications other than immunosuppression

remains unknown, our findings demonstrate that weekly dosing avoids concentration levels that

could be of high safety-concern, while exhibiting relatively low pharmacokinetic variability,

supporting a fixed-dose approach in future trials.

No recommendations exist for the optimal timing of testing the blood concentration of rapamycin

when administered as a weekly dose. Potential sampling points include C_{max}, C_{min}, or at some

earlier time during the elimination phase (i.e., between the end of the distribution phase and

C_{min}). In daily dosing of rapamycin, C_{min} has been shown to have an excellent correlation to AUC

(23,28). In our data, based on a weekly dosing scheme of 7 mg rapamycin, values of C_{min} are low

(0.75–2 ng/mL), approaching the LLOQ of 0.5 ng/mL at our hospital lab. Given a lower dose or the

use of an analysis method with higher LLOQ, it is likely that some samples would not be able to

be quantified, as demonstrated by our 2 mg dose participant's unmeasurable C_{min}. Time to peak

concentration (T_{max}) for orally administered rapamycin is reported at approximately 1h, although

some studies indicate considerable interindividual variation, with T_{max} ranging from 0,5 to 3h (19).

In our data, 11 participants showed the highest concentration at 1h, however, two participants

peaked at the 3h sample. This variation complicates reliable C_{max} capture with single-point

sampling, reflected by the fact that the highest variation (CoV = 0.40) in our data was at the 1h

timepoint. Both C_{min} and 3h samples showed CoV of 0.38, while the 48h sample demonstrated

lowest variation (CoV = 0.28). Taken together, our results suggest that sampling early during the

elimination phase is the best option when rapamycin is administered as a weekly dose.

In our data, the long terminal half-life (pseudo-terminal t½) of 68.85 ± 13.64 hours is well in line

with that observed in multiple previous studies evaluating pharmacokinetic properties after a

single dose rapamycin in healthy volunteers, as well as after multiple doses in transplant

recipients (22,23,31). The two participants treated with reduced doses (2 and 4 mg) showed

consistently lower blood concentrations across all time points, with the 2 mg dose producing the

lowest results, suggesting dose proportionality for rapamycin administered once per week within

this dose range.

The extent to which rapamycin passes the BBB is unclear. The drugs high molecular weight (914.2

Da) and its status as an efflux pump substrate suggest limited permeability. One study of

rapamycin in patients with amyotrophic lateral sclerosis showed no quantifiable levels of

sirolimus in CSF using LC-MS/MS (15). To assess whether LC-MS/MS could reliably quantify

rapamycin in CSF, we developed a method achieving an LLOQ of 20 ng/mL. While further

optimization might lower this threshold, reaching the single-digit pg/mL range - necessary to

definitively rule out the presence of a pharmacologically relevant concentration - was deemed

unfeasible. Given this, we elected not to perform any analysis of rapamycin concentration on the

CSF data collected from the study participants in our trial. This constrained LLOQ should also be

taken into consideration when interpreting the published dataset where no rapamycin could be

detected in CSF of patients on the drug (15). While some clinical data suggest BBB passage—

such as rapamycin's efficacy in treating cerebral manifestations of tuberous sclerosis complex

in humans (33)—the question of its extent of brain delivery remains open. Given the expected low

free concentration in CSF, one option for future research is to perform post-mortem analysis of

brain tissue from rapamycin-fed laboratory animals. Alternatively, labelling rapamycin with a

radioactive nuclide and perform micro-dosing experiments using positron emission tomography

(PET) could provide further insights into its brain exposure.

The foremost limitation of this study was the sparse blood sampling schedule, with c_{min}

collected from the previous dosing occasion. This prevented a complete AUC estimation and

limited the analysis in two ways: 1) we could not evaluate interindividual variation in overall drug

exposure but had to restrain the analysis to the separate sampling time-points; 2) more than one

concentration sample collected after the distribution phase of the drug is needed to perform a

full modelling of the drug's pharmacokinetic parameters, such as t1/2 . To overcome this, we

assumed stable C_{min} between doses and used the previous dosing occasion's C_{min} as an

additional timepoint, which allowed for estimates of pseudo-terminal $t_{1/2}$.

In conclusion, our data show that weekly rapamycin administration in patients with early

Alzheimer's disease results in moderate interindividual variability in blood concentration,

suggesting that a fixed dosing regimen may be appropriate for future trials using a similar design.

Of the four time points assessed (C_{min}, 1h-, 3h-, and 48h-hours post dose) the 48h sample

showed the lowest interindividual variability and strongest correlation to C_{min} from the previous

dose, supporting it as the optimal sampling time for monitoring blood concentration.

List of abbreviations

AUC	Area under the curve
BBB	Blood-brain barrier
CoV	Coefficient of variation
CSF	Cerebrospinal fluid
LC-MS/MS	Liquid chromatography–tandem mass spectrometry
LLOQ	Lower limit of quantification
MCI	Mild cognitive impairment
SD	Standard deviation
TDM	Therapeutic drug monitoring

Declarations

Ethics approval and consent to participation

The study was approved by the Swedish Medical Products Agency (5.1–2023-8283), and the

Swedish Ethical Review Authority (2023–03075-02). Written informed consent was collected

from all participants and their designated study partners before initiating any study procedures.

Availability of data and materials

The datasets used and analyzed during the current study are available from the

corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This study was supported by a Longevity Impetus grant from the Norn Group, Åhlén Stiftelsen, Demensfonden, The Swedish Society of Medicine (SLS), Loo and Hans Osterman Stiftelse, Stiftelsen för Åldersjukdomar Karolinska Institutet, Stiftelsen för Gamla Tjänarinnor, Tore Nilssons Stiftelse för Medicinsk Forskning, Åke Wibergs stiftelse (M24-0117), Swedish Brain Foundation (PD2024-0444), and Magnus Bergvall stiftelse.

Authors' contributions

JS, PPS and MK conceived of and designed the study. PPS and MK directed and supervised conduction of the study. JS and HA performed the data collection and drafted the manuscript. JS, HA and PPS performed data analysis. All authors contributed to the interpretation of the results, critically reviewed the manuscript, and approved the final manuscript.

Acknowledgements

We wish to thank Associate Professor Staffan Rosenborg, MD, PhD at the Clinical Pharmacology Unit, Karolinska University Hospital Huddinge for valuable discussions on the blood sirolimus analysis method, and Associate Professor Anna Matton, PhD at the Karolinska Institute, for providing blank CSF. The authors would like to acknowledge support of the SciLifeLab Drug Discovery and Development Platform, Sweden.

References

1. Mohamed MA, Elkhateeb WA, Daba GM. Rapamycin golden jubilee and still the miraculous drug: a potent immunosuppressant, antitumor, rejuvenative agent, and potential contributor in COVID-19 treatment. Bioresour Bioprocess. 13 juni 2022;9(1):65.

2. Alvers AL, Wood MS, Hu D, Kaywell AC, Dunn, Jr. WA, Aris JP. Autophagy is required for extension of yeast chronological life span by rapamycin. Autophagy. 16 augusti 2009;5(6):847–9.

3. Robida-Stubbs S, Glover-Cutter K, Lamming DW, Mizunuma M, Narasimhan SD, Neumann-Haefelin E, m.fl. TOR Signaling and Rapamycin Influence Longevity by Regulating SKN-1/Nrf and DAF-16/FoxO. Cell Metab. maj 2012;15(5):713–24.

4. Bjedov I, Toivonen JM, Kerr F, Slack C, Jacobson J, Foley A, m.fl. Mechanisms of Life Span Extension by Rapamycin in the Fruit Fly Drosophila melanogaster. Cell Metab. januari 2010;11(1):35–46.

5. Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, m.fl. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. Nature. juli 2009;460(7253):392–5.

6. Bitto A, Ito TK, Pineda VV, LeTexier NJ, Huang HZ, Sutlief E, m.fl. Transient rapamycin treatment can increase lifespan and healthspan in middle-aged mice. eLife. 23 augusti 2016;5:e16351.

7. Miller RA, Harrison DE, Astle CM, Baur JA, Boyd AR, De Cabo R, m.fl. Rapamycin, But Not Resveratrol or Simvastatin, Extends Life Span of Genetically Heterogeneous Mice. J Gerontol Ser A. februari 2011;66A(2):191–201.

8. Spilman P, Podlutskaya N, Hart MJ, Debnath J, Gorostiza O, Bredesen D, m.fl. Inhibition of mTOR by Rapamycin Abolishes Cognitive Deficits and Reduces Amyloid-β Levels in a Mouse Model of Alzheimer's Disease. Ferrari PF, redaktör. PLoS ONE. 01 april 2010;5(4):e9979.

9. Lin AL, Zheng W, Halloran JJ, Burbank RR, Hussong SA, Hart MJ, m.fl. Chronic Rapamycin Restores Brain Vascular Integrity and Function Through NO Synthase Activation and Improves Memory in Symptomatic Mice Modeling Alzheimer's Disease. J Cereb Blood Flow Metab. september 2013;33(9):1412–21.

10. Mannick JB, Giudice GD, Lattanzi M, Valiante NM, Praestgaard J, Huang B, m.fl. mTOR inhibition improves immune function in the elderly.

11. Kraig E, Linehan LA, Liang H, Romo TQ, Liu Q, Wu Y, m.fl. A randomized control trial to establish the feasibility and safety of rapamycin treatment in an older human cohort: Immunological, physical performance, and cognitive effects. Exp Gerontol. maj 2018;105:53–69.

12. Mannick JB, Teo G, Bernardo P, Quinn D, Russell K, Klickstein L, m.fl. Targeting the biology of ageing with mTOR inhibitors to improve immune function in older adults: phase 2b and phase 3 randomised trials. Lancet Healthy Longev. maj 2021;2(5):e250–62.

13. Harinath G, Lee V, Nyquist A, Moel M, Morgan SL, Isman A, m.fl. Safety and efficacy of rapamycin on healthspan metrics after one year: PEARL Trial Results [Internet]. Geriatric Medicine; 2024 [citerad 05 februari 2025]. Tillgänglig vid: http://medrxiv.org/lookup/doi/10.1101/2024.08.21.24312372

14.   Konopka AR, Lamming DW, RAP PAC Investigators, Grasso BA, Marrah RC, Decker S, m.fl. Blazing a trail for the clinical use of rapamycin as a geroprotecTOR. GeroScience. 06 oktober 2023;45(5):2769–83.

15.   Mandrioli J, D'Amico R, Zucchi E, De Biasi S, Banchelli F, Martinelli I, m.fl. Randomized, double-blind, placebo-controlled trial of rapamycin in amyotrophic lateral sclerosis. Nat Commun. 17 augusti 2023;14(1):4970.

16.   Palma J, Martinez J, Millar Vernetti P, Ma T, Perez MA, Zhong J, m.fl. MTOR Inhibition with Sirolimus in Multiple System Atrophy: A Randomized, Double-Blind, Placebo-Controlled Futility Trial and 1-Year Biomarker Longitudinal Analysis. Mov Disord. april 2022;37(4):778–89.

17.   Bagherpour B, Salehi M, Jafari R, Bagheri A, Kiani-Esfahani A, Edalati M, m.fl. Promising effect of rapamycin on multiple sclerosis. Mult Scler Relat Disord. november 2018;26:40–5.

18.   U.S. Food and Drug Administration (FDA). 2025. Summary of Product Characteristics for Rapamune. [Internet]. [citerad 07 februari 2025]. Tillgänglig vid: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/021083s069s070,021110s087s088l bl.pdf#page=43

19.   Mahalati K, Kahan BD. Clinical Pharmacokinetics of Sirolimus: Clin Pharmacokinet. 2001;40(8):573–85.

20.   Svensson JE, Bolin M, Thor D, Williams PA, Brautaset R, Carlsson M, m.fl. Evaluating the effect of rapamycin treatment in Alzheimer's disease and aging using in vivo imaging: the ERAP phase IIa clinical study protocol. BMC Neurol. 04 april 2024;24(1):111.

21.   Jack CR, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, m.fl. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. Alzheimers Dement. april 2018;14(4):535–62.

22.   Brattstr??m C, S??we J, Jansson B, L??nnebo A, Nordin J, Zimmerman JJ, m.fl. Pharmacokinetics and Safety of Single Oral Doses of Sirolimus (Rapamycin) in Healthy Male Volunteers: Ther Drug Monit. oktober 2000;22(5):537–44.

23.   Zimmerman JJ, Kahan BD. Pharmacokinetics of Sirolimus in Stable Renal Transplant Patients after Multiple Oral Dose Administration. J Clin Pharmacol. maj 1997;37(5):405–15.

24.   Lin J. CSF as a Surrogate for Assessing CNS Exposure: An Industrial Perspective. Curr Drug Metab. 01 januari 2008;9(1):46–59.

25.   Swinscow. TDV Statistics at Square One. Ninth Edition. Vol. 1997. BMJ Publishing Group;

26.   Harinath G, Lee V, Nyquist A, Moel M, Wouters M, Hagemeier J, m.fl. The bioavailability and blood levels of low-dose rapamycin for longevity in real-world cohorts of normative aging individuals. GeroScience [Internet]. 28 januari 2025 [citerad 05 februari 2025]; Tillgänglig vid: https://link.springer.com/10.1007/s11357-025-01532-w

27.   Cohen EEW, Wu K, Hartford C, Kocherginsky M, Eaton KN, Zha Y, m.fl. Phase I Studies of Sirolimus Alone or in Combination with Pharmacokinetic Modulators in Advanced Cancer Patients. Clin Cancer Res. 01 september 2012;18(17):4785–93.

28.   Kahan B, Napoli K, Kelly P, Podbielski J, Hussein I, Urbauer D, m.fl. Therapeutic drug monitoring of sirolimus: correlations with efficacy and toxicity. Clin Transplant. april 2000;14(2):97–109.

29.   Ågesen FN, Weeke PE, Tfelt-Hansen P, Tfelt-Hansen J, for ESCAPE-NET. Pharmacokinetic variability of beta-adrenergic blocking agents used in cardiology. Pharmacol Res Perspect. augusti 2019;7(4):e00496.

30.   Reis M, Chermá MD, Carlsson B, Bengtsson F. Therapeutic Drug Monitoring of Escitalopram in an Outpatient Setting. Ther Drug Monit. december 2007;29(6):758–66.

31.   Leung LY, Lim HK, Abell MW, Zimmerman JJ. Pharmacokinetics and Metabolic Disposition of Sirolimus in Healthy Male Volunteers After a Single Oral Dose. Ther Drug Monit. februari 2006;28(1):51–61.