1   **Word count (excluding abstract, references, and Tables): 3514**
2   **Abstract word count: 229**
3   **Number of Tables and Figures: 1 Table and 4 Figures**
4   **Supplementary material: 12 Tables and 8 Figures**

Cross-disorder comparison of Brain Structures among 4,836 Individuals with Mental Disorders and Controls utilizing Danish population-based Clinical MRI Scans

Authors: Stefano Cerri, PhD[1,2], Vardan Nersesjan, MD, PhD[1], Kiril Vadimovic Klein, MSc[2],
Enric Cristòbal Cóppulo, MSc[1], Sebastian Nørgaard Llambias, MSc[2], Mostafa Mehdipour
Ghazi, PhD[2], Mads Nielsen, PhD[2, *], Michael Eriksen Benros, MD, PhD[1,3, *]

Affiliations:
1: Copenhagen Research Centre for Biological and Precision Psychiatry, Mental Health
Centre Copenhagen, Copenhagen University Hospital, Hellerup, Denmark.
2: Department of Computer Science, University of Copenhagen, Copenhagen, Denmark
3: Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of
Copenhagen, Copenhagen, Denmark
* shared last authorship

Correspondence:
- Stefano Cerri, PhD; stefano.cerri@regionh.dk,
Gentofte Hospitalsvej 19, st., 2900, Hellerup, Denmark, +45 50345029
- Professor Michael E. Benros, MD, PhD; benros@dadlnet.dk,
Gentofte Hospitalsvej 19, st., 2900, Hellerup, Denmark, +45 26255239

Keywords: mental disorders, brain morphology, observational studies, electronic health
records, clinical cohort

ABSTRACT:

Large-scale mega-analyses of worldwide combined Magnetic Resonance Imaging (MRI) studies have demonstrated brain differences between individuals with mental disorders and controls. However, the potential of large-scale observational studies using population-based clinical MRI data remains unexplored. We analyzed clinical MRI data from 23,545 patients in the Eastern half of Denmark (Capital Region of Denmark and Region Zealand). 2,774 patients with mental disorders and 2,062 non-psychiatric controls fulfilled our inclusion and exclusion criteria. Patients with mental disorders exhibited smaller thalamic ($d=-0.298$) and amygdala volumes ($d=-0.250$), with larger ventricles ($d=0.272$), and thinner insula ($d=-0.177$), all $p<0.0001$. Analysis across all ROIs revealed a widespread pattern of thinner cortex ($d=-0.180$), especially in the temporal pole ($d=-0.234$) and superior frontal ($d=-0.212$) regions, and increased extracerebral cerebrospinal fluid ($d=0.264$). For volumetric measurements, findings were consistent across different inclusion and exclusion criteria but varied for cortical thickness measurements. Utilizing this currently largest population-based MRI cohort for mental disorders, we demonstrate that clinical MRI scans can detect brain structural differences among patients with mental disorders in real-world clinical settings, aiding in the stratification of patients without mental disorders. Cross-disorder analyses reveal shared neuroanatomical changes, including globally smaller brain volumes and thinner cortex. Integrating large-scale clinical MRI data with electronic health records holds promise for improved patient stratification and tracking of disease progression for future longitudinal cross-disorder studies, bridging real-world MRI data with clinical trajectories for further biological subgrouping.

INTRODUCTION

Prior brain imaging studies aiming to identify brain patterns in mental disorders have often been constrained by small sample sizes with publication bias towards positive findings, highlighting the critical need for larger datasets to ensure reproducibility.[1,2] Nevertheless, Magnetic Resonance Imaging (MRI) scans have proven effective in detecting subtle brain changes across various mental disorders in large-scale studies.[3]

To produce more reliable results, the pioneering consortium ENIGMA (Enhancing NeuroImaging Genetics through Meta-Analysis)[4] pools neuroimaging data from multiple research centers worldwide to increase statistical power and generalize findings across different mental disorders. Despite their studies being the largest to date,[3, 5-10] heterogeneity in imaging protocols, MRI processing, inclusion and exclusion criteria for patients and controls regarding diagnoses, treatments, and demographics across centers might introduce noise and hinder the generalizability of their findings. Additionally, the lack of comprehensive patient history information, in connection to the ENIGMA MRI data, constrains their ability to correlate brain findings with patient clinical data. Most importantly, as the ENIGMA consortia data is the largest dataset to date, validation of their findings in independent large-scale datasets is strongly needed.

Recent advances in automatic tools for extracting cortical and subcortical information from MRI scans have enabled the analysis of large-scale clinical MRI data,[11-13] available in quantities far exceeding typical research study samples. Clinical MRI data, often accompanied by electronic health records (EHRs), allows for better patient stratification and more precise inclusion and exclusion criteria based on complete patient histories. Furthermore, as recent advances have shown shared clinical and genetic features across the spectrum of mental disorders,[14,15] cross-disorder investigations are increasingly conducted within psychiatric research.[16-21]

Given these advancements, our study has two primary objectives: First, to assess whether prior findings based on high-quality MRI data can be replicated with routinely acquired clinical MRI scans. Second, to investigate the influence of inclusion and exclusion criteria based on medication and patient history on brain changes associated with mental disorders, both cross-disorder and within diagnostic subgroups. By addressing these objectives, we aim to highlight the potential of large-scale clinical MRI data for advancing the understanding of the neurobiological mechanisms underlying mental disorders and exploring the utility of current real-world MRI in contributing to diagnostic biomarkers.

METHODS

We conducted a retrospective cohort study utilizing all clinical MRI scans from the Eastern half of Denmark (the Capital Region of Denmark and Region Zealand) acquired during the year 2019, totaling 313,927 MRI images obtained during 28,129 MRI visits on 23,545 unique patients. We first defined our cases with mental disorders and non-psychiatric control using complete EHRs available for all patients, including demographics, diagnoses, and medications. Secondly, anatomical brain regions of interest were defined based on prior large-scale studies and meta-analyses comparing mental disorder patients with controls, to investigate the reproducibility of previous findings. Lastly, we explored the robustness and dynamics of these differences by adding stricter exclusion criteria for cases and non-psychiatric controls. We obtained legal and ethical approvals for the proposed study from the

98 Data Protection Agency and the Ethics Committee of the Capital Region of Denmark (R-
99 22002033, and P-2020-101).

Study Participants

Individuals ≥18 years of age with mental disorders were included as cases using the following
ICD-10 codes: any Mental Disorders (F00-F99), and specifically Dementia (F00-F03 and
G30), Substance Use Disorder (F10-19), Schizophrenia Spectrum Disorder (F20-29),
Depression (F32-F33), and Anxiety Disorder (F40-F48). For patients with multiple of these
diagnoses, a hierarchy was applied—Dementia, Schizophrenia Spectrum Disorder,
Depression, Anxiety Disorder, and Substance Use Disorder—based on decreasing ICD-10
code values. Substance Use Disorder was placed last to avoid overriding other psychiatric
diagnoses and isolate the effect of substance abuse. Individuals had to be diagnosed with a
mental disorder within three months of the MRI scan. This threshold was determined through
sensitivity analysis (**Supplementary Figure S1**). Controls were ≥18 years of age and without
mental or neurological disorders. These basic inclusion and exclusion criteria were termed
"Population A", similar to the minimum criteria to be included in ENIGMA consortium
studies. "Population B" additionally included brain medications as exclusion criteria for non-
psychiatric controls. Lastly, further exclusions were based on neurological medications and
neurological comorbidities for cases ("Population C"). Details for each criteria are provided
in **Figure 1**, and specific medication and diagnosis codes are listed in **Supplementary
Methods**. Subgroup analyses included only diagnoses with ≥50 individuals meeting
Population C criteria, to minimize false-positive findings and overestimation of effect sizes.[1,2]
Cross-disorder analyses included all mental disorders (F00–F99, G30) to investigate
neuroanatomical patterns shared across the diagnostic spectrum.

MRI Processing

The analysis focused solely on T1-weighted scans due to their superior contrasts for
anatomical structures. We estimated T1-weighted sequences from all available MRI scans,
resulting in 80,624 T1-weighted scans (**Supplementary Methods**). Each T1-weighted
DICOM sequence was converted to NIfTI using dcm2niix.[22] We processed each T1w NIfTI
scan with recon-all-clinical[11-13,23,24] to obtain volumetric and cortical thickness estimates.
Recon-all-clinical extracts the same cortical and subcortical metrics as the standard
FreeSurfer recon-all pipeline[25] used in ENIGMA studies (**Supplementary Figure S2**), with
the addition of extracerebral cerebrospinal fluid volume, and it is optimized for lower-quality,
heterogeneous clinical MRI scans. Measurements were averaged bilaterally for regions of
interest (ROIs).

All scans were subjected to automated quality control, explicitly designed to remove clear
segmentation failures and extreme morphometric outliers while preserving biologically
meaningful variability. Low-quality segmentations, as predicted by recon-all-clinical Dice[26]
estimates, were automatically discarded.[11] Additional quality control involved removing
extreme outliers based on interquartile ranges (below or above 2 for each ROI,
Supplementary Methods). Our quality control steps and inclusion and exclusion criteria
allowed us to detect and remove any potential MRI scans affected by severe motion, poor
tissue contrast, acquisition-related artifacts, or structural brain abnormalities. Thresholds were
computed across the entire population to avoid defining 'normal' variation based on any
single diagnostic group and to reduce the risk of selectively excluding illness-related
morphometric variation. For patients with multiple MRI scans passing quality control,
inclusion, and exclusion criteria, the best-predicted scan from recon-all-clinical was used.
When performing a sensitivity analysis on slice thickness, we first applied the high- and low-

resolution threshold before selecting the best-predicted scan per patient, thereby increasing
our sample size and overlap between the two groups. All MRI processing steps are illustrated
in **Figure 1** and detailed in **Supplementary Methods**.

Outcomes of interest

We aimed to replicate known differences between mental disorder patients and non-
psychiatric controls, and dementia patients and non-psychiatric controls, based on ROIs
identified in previous studies examining regions affected across disorders (10 ROIs: Lateral
Ventricles, Thalamus, Amygdala, Hippocampus, Cortical Thickness, Pars Opercularis
Thickness, Medial Orbital Frontal Thickness, Rostral Anterior Cingulate Thickness, Superior
Temporal Thickness, and Insula Thickness),[3,5-10,16,27] and well-known studies for dementia (4
ROIs: Lateral Ventricles, Hippocampus, Cortical Thickness, Entorhinal Thickness).[28-30] For
mental disorder subgroups, we identified the following ROIs from previous ENIGMA
studies: Substance Use Disorder (7 ROIs: Amygdala, Hippocampus, Insula Thickness, Middle
Temporal Thickness, Paracentral Thickness, Precentral Thickness, and Supramarginal
Thickness),[9] Schizophrenia Spectrum Disorder (12 ROIs: Hippocampus, Amygdala,
Thalamus, Nucleus Accumbens, Pallidum, Lateral Ventricles, Cortical Thickness, Fusiform
Thickness, Inferior Temporal Thickness, Middle Temporal Thickness, Superior Temporal
Thickness, and Insula Thickness),[5,6] Depression (8 ROIs: Hippocampus, Amygdala, Lateral
Ventricles, Posterior Cingulate Thickness, Fusiform Thickness, Insula Thickness, Medial
Orbitofrontal Thickness, Rostral Anterior Cingulate Thickness),[7,8] and Anxiety Disorder (4
ROIs: Putamen, Pallidum, Hippocampus, and Amygdala).[10]

Statistical Analyses

We reported group differences in brain volumes and cortical thickness between cases with
mental disorders and non-psychiatric controls for each ROI with multiple linear regression
models using statsmodels.[31] A binary variable, "Diagnosis" (0=non-psychiatric controls,
1=patients), was the predictor of interest. We included sex, age, intracranial volume (ICV)
obtained with recon-all-clinical, and site as covariates in the model. Site was defined as the
unique combination of scanner and performing hospital; sites without both diagnostic groups
were excluded to ensure within-site identifiability of diagnosis effects. The full model is then
described as: ROI ~ Diagnosis + Age + Sex +[ICV] + Site, where ICV was included for
volumetric measures only. To ease comparisons with prior ENIGMA studies,[5-10] effect size
estimates were calculated using Cohen's d[32] computed from the t-statistic of the "Diagnosis"
variable from the regression models (Equation 10).[23] The 95% confidence intervals were
derived using Equation 15,[33] utilizing the asymptotic standard error from Equation 17.[33] For
each comparison, we reported uncorrected two-tailed p-values for the t-statistics of the
"diagnosis" binary variable from the linear regression model, accompanied by an "*" in case
of significance after multiple comparisons correction for each patient group (Bonferroni,
α=0.05).[34] The number of hypotheses for Bonferroni correction corresponds to the number of
ROIs used in each patient group comparison. Differences between our statistical model and
those used in prior ENIGMA studies are summarized in Supplementary Table 9. The site
distribution for each comparison is provided in Supplementary Tables 10–16.

RESULTS

In the Capital Region of Denmark, a total of 313,927 MRI images were conducted in 28,129
MRI visits during the year 2019 on 23,545 unique patients. Out of these, a total of 4,836
subjects (2,776 cases and 2,062 non-psychiatric controls) from 36 sites fulfilled the inclusion

and exclusion criteria under the most inclusive criteria (Population A), followed by 4,146
subjects from 35 sites for Population B (2,764 cases and 1,382 non-psychiatric controls), and
2,161 subjects from 33 sites for Population C (785 cases and 1,376 non-psychiatric controls)
(**Table 1** and **Figure 1**). Most MRI scans were acquired with Philips and Siemens scanners
using 3D acquisitions, with magnetic fields of either 1.5T or 3T and slice thicknesses ranging
between 1 and 6 mm, with peaks around 1-3 mm. There were no substantial differences in
MRI scanner manufacturers, acquisition protocols, or slice thicknesses across criteria
(**Supplementary Figure S3**).

MRI differences among individuals with Mental Disorders compared to non-psychiatric
controls

Results for Population A, B, and C are presented in **Figure 2A-C**, with numerical values in
Supplementary Table S17, **Table S18**, and **Table S19**. Under Population C, all comparisons
between our clinical MRI data and prior mega-analyses[3,5-10,16,27] were significant, except for
the Rostral Anterior Cingulate after correction for multiple comparisons. Mental disorder
patients showed smaller volumes in the thalamus (d=-0.298), amygdala (d=-0.250), and
hippocampus (d=-0.145), larger lateral ventricular volume (d=0.272), and a general pattern of
thinner cortex (d=-0.180), particularly in the insula (d=-0.177).

For cortical thickness, most comparisons under Population A and B showed no significant
differences between mental disorder patients and non-psychiatric controls, with a greater
pattern of thinner cortex in unmedicated patients and those without brain comorbidities
(Population C d=-0.180 vs. Population B d=-0.044). Consistent findings were observed across
criteria for volumetric measures, despite Population C having approximately 3.5 times fewer
cases and 1.5 times fewer non-psychiatric controls than Population A.

Dementia patients exhibited large effect sizes: smaller hippocampal volume (d=-0.632), larger
lateral ventricles (d=0.757), and a general pattern of thinner cortex (d=-0.648), especially in
the entorhinal cortex (d=-0.690). To ensure that our findings for mental disorders were not
biased by large effect sizes in the dementia subgroup, we performed analysis excluding
dementia patients and reassessed Population C (**Figure 2C**). Effect sizes were smaller but
consistent with previous findings and all comparisons, except for the hippocampus and
Medial Orbitofrontal, were still significant after correction for multiple comparisons.

We identified additional brain differences not reported as main findings in prior ENIGMA
studies[3,5-10,16,27] under Population C (**Figure 3**). After correcting for multiple comparisons
across all ROIs, we identified 27 significant differences, highlighting a general pattern of
thinner cortex across most cortical areas (cortical thickness d=-0.180, temporal pole d=-0.234,
superior frontal d=-0.212, and middle temporal d=-0.191,)), smaller deep gray matter
structures (thalamus d=-0.298, amygdala d=-0.250, pallidum d=-0.192, and nucleus
accumbens d=-0.180), and larger ventricular volumes (inferior lateral ventricles d=0.351, 3rd
ventricle d=0.313 and lateral ventricles d=0.272). We also reported higher amounts of
extracerebral cerebrospinal fluid (d=0.264). These differences were also found for Population
A (**Supplementary Figure S4**) and B (**Supplementary Figure S5**), although to a smaller
extent, especially regarding thinner cortex (21 and 22 significant differences, respectively).

Comparison between mental disorder subgroups and non-psychiatric controls

For Population B, statistically significant brain differences between mental disorder
subgroups and controls were found for 19 of 32 ROIs; and after correction for multiple
comparisons, 15 of 32 remained significant (**Figure 4**). Population A and C yielded 14 and 9

significant comparisons, respectively (**Supplementary Figure S6** and **Supplementary**
Figure S7). Effect sizes were generally consistent but smaller compared to ENIGMA studies,
and most non-significant differences are related to subcortical thickness areas. For the
Substance Use Disorder subgroup, we observed larger effect sizes for the amygdala,
hippocampus, and insula than previously reported.[9] Schizophrenia Spectrum Disorder
subgroup comparisons revealed consistent patterns of thinner cortex and smaller deep gray
matter volumes compared with ENIGMA studies,[5,6] with only opposite findings for the
hippocampus, lateral ventricles, and pallidum. For the Depression subgroup, effect sizes were
similar for most comparisons, although mostly not significant, while we noticed a large effect
for lateral ventricles compared to ENIGMA.[7,8] For Anxiety Disorders, we only reported a
negative effect for the pallidum in contrast with ENIGMA.[10]

Effect of Slice Thickness

We assessed the influence of MRI slice thickness on our findings by separating scans into
high-resolution (<1.5 mm) and low-resolution (≥1.5 mm) groups (approximately 40% of
cases and non-psychiatric controls overlapped between high- and low-resolution groups).
Results for Population C are shown in **Figure 2D**. Volumetric comparisons were consistent,
while slight differences were observed for cortical thickness comparisons. Similar results
were found across criteria for volumetric comparisons, with larger differences in cortical
thickness comparisons (**Supplementary Figure S8**).

DISCUSSION

To our knowledge, this is the first large-scale population-based neuroimaging study using
real-world clinical MRI scans to investigate brain changes in mental disorders. We
successfully replicated many previously reported findings from meta-analyses of brain
differences between patients with mental disorders and non-psychiatric controls. Results in
patients with dementia were included as a reference, confirming that the observed differences
in other mental disorders were not primarily driven by dementia. However, when examining
mental disorder subgroups, we observed smaller effect sizes than those reported in previous
meta-analyses. This discrepancy likely stems from differences in the definition of our control
group, which included patients undergoing MRI scans for possible brain-related conditions,
highlighting future avenues for predicting changes in a more challenging control cohort than
typically used in research studies relying on healthy controls. This study is also the largest to
date that investigates brain changes associated with mental disorders as a whole, rather than
focusing on specific subgroups, as seen in previous ENIGMA studies.[3] Our findings reveal a
general pattern of thinner cortex and smaller deep gray matter volumes among patients with
mental disorders broadly consistent with the ENIGMA cross-disorder studies[16-21], which
report overlapping patterns of structural abnormalities across several psychiatric conditions—
particularly schizophrenia, bipolar disorder, and major depression—although disorder-
specific differences also exist and effect sizes vary considerably across the diagnostic
spectrum. These results suggest that clinical MRI data, supplemented with EHR information,
can identify biomarkers for mental disorders and improve patient diagnosis and trajectories.

Our exclusion criteria, based on brain medications and comorbidities from EHRs, influenced
our findings. We noted consistent effect sizes for volumetric measurements when applying
stricter exclusion criteria, even though our sample sizes decreased 1.5-fold for cases and 3.5-
fold for non-psychiatric controls. We also observed a more pronounced pattern of thinner
cortex in unmedicated patients and those without brain comorbidities compared to the overall
patient group. This may reflect selection biases in clinical MRI referrals, whereby

unmedicated patients or those without detectable brain comorbidities who nevertheless
undergo neuroimaging may represent individuals with more acute or atypical psychiatric
presentations, whereas patients with brain comorbidities may be scanned for reasons unrelated
to psychiatric disease severity. Furthermore, due to the cross-sectional design, we cannot
disentangle medication effects from illness severity or disease duration. Additionally, we
found that effect sizes for the Substance Use disorders group were substantially larger for
Population C than those reported in a previous ENIGMA study,[9] as well as our Population A
and B, suggesting that different inclusion and exclusion criteria might play an even bigger
role, thus warranting further investigation.

When examining mental disorder subgroups, the effect sizes were generally in the same
direction as in the ENIGMA studies but often not significant after correcting for multiple
comparisons. This might be due to the larger sample sizes in the combined ENIGMA studies,
which translate into more significant comparisons, even with small effect sizes. In addition,
many ENIGMA cohorts include patients with long-standing illness, whereas our inclusion
criteria required a psychiatric diagnosis before or within three months of MRI acquisition.
Because MRI is typically performed close to diagnosis in clinical practice, our cohort likely
captures individuals at earlier stages of mental disorder disease trajectory with limited
cumulative exposure to illness-related processes and treatments. Additionally, our findings
suggest that cortical thickness changes are more challenging to detect in clinical MRI scans
due to the subtlety of these changes and the lower resolution and quality of clinical images.
However, comparing the effect sizes in our study with those reported in ENIGMA studies is
complex. Discrepancies in inclusion and exclusion criteria, processing tools, modeling
choices, and MRI scan quality between the ENIGMA and our studies likely impact the
findings. We used Population B to closely match those in ENIGMA studies. However, several
centers in prior ENIGMA studies employed less strict protocols, especially for cases, that are
more similar to our less strict criteria (Population A). Indeed, under our strictest criteria
(Population C), we observed the least significant comparisons with ENIGMA studies.

The main strength of this study is the use of a large-scale population-based MRI dataset from
real-world clinical settings. The integration of EHRs allows for rigorous and flexible
inclusion and exclusion criteria, enabling a better understanding of the impact of medication
and comorbidities on brain structure, a better match with prior studies' inclusion and
exclusion criteria, and enhancing the ability to track disease progression and correlate brain
changes with patient histories.

Limitations include our real-world data with clinical MRI scans being typically of lower
quality than those used in research studies, including scans acquired with contrast agents,
which may affect tissue contrast and thus segmentation quality. Our findings suggest that
differences in effect sizes are only to a small degree influenced by the slice thickness of the
scans, particularly for cortical thickness measurements. Furthermore, we could not use
established automatic methods like FreeSurfer,[25] FSL,[35] and SPM,[36] as these tools cannot yet
handle the low-resolution MRI scans in our clinical cohort. Instead, we employed recently
developed automatic tools[10-12] to obtain cortical and subcortical measurements, which might
require further validation. Additionally, we did not perform visual quality control of the
segmentations, which is a limitation, as even in ENIGMA studies visual inspection is used to
confirm segmentation quality. However, visual quality control is often infeasible in large-
scale clinical datasets. Although site effects were modeled by scanner and hospital in our
linear model, we did not apply more complex harmonization techniques like ComBat[37]
because several sites included very small sample sizes, which can yield unstable estimates. A
prior ENIGMA study[38] has shown that effect sizes can vary depending on site-effect

modeling strategies, highlighting the importance of evaluating harmonization approaches
when sufficient site-specific data are available.

Our cohort is the largest population-based cohort, yet smaller than the combined worldwide
cohorts in prior ENIGMA mega-analyses studies. However, our study is based on MRI scans
acquired in a single year (2019) only, thus, there is substantial potential in future studies to
mitigate the subtle nature of subcortical differences, where more data is needed to validate our
findings further. Moreover, we did not age- and sex-match controls in this study due to the
limited sample size, but we adjusted for these covariates in our analysis. Our sample size was
also yet insufficient for studying brain changes in other mental disorders such as bipolar
disorder, attention-deficit/hyperactivity disorder, and autism spectrum disorder. Additionally,
future work may investigate how factors such as duration of illness, age at onset, cumulative
medication exposure, and sociodemographic factors may influence cortical thickness and
volumetric changes in mental disorders.

Clinical MRI scans can effectively detect cortical and subcortical differences among patients
with mental disorders and also distinguish them from other patients without mental disorders.
When examining mental disorder subgroups, we observed consistent, though smaller, effect
sizes compared to previous ENIGMA mega-analyses studies using high-quality MRI data.
These differences are likely due to our control group comprising patients undergoing MRI
scans for potential clinical diagnoses. In contrast, the studies in the ENIGMA mega-analyses
compare to healthy controls, which makes prediction easier, whereas, in a real-world clinical
context, the separation from other individuals undergoing examination is of importance for
specificity.

Our findings reveal shared brain differences across mental disorders. Factors such as slice
thickness, comorbidities, and medication can affect the detection of structural changes.
Population selection should reflect the study objective: real-world clinical samples
(Population A) capture brain patterns representative of routine care but introduce greater
heterogeneity, whereas strictly defined samples (Population C) minimize variability and
improve specificity, at the cost of broader generalizability. We recommend standardizing
inclusion/exclusion criteria, incorporating EHR-based comorbidity data, and leveraging
routine clinical MRI scans—including lower-resolution images—to enable large-scale
biomarker discovery while accounting for scan quality. These strategies can support more
precise investigations of neurobiological mechanisms and advance diagnostic and therapeutic
approaches.

ACKNOWLEDGEMENTS

363　Stefano Cerri has received funding from the Lundbeck Foundation (R449-2023-1512).
364　Mostafa Mehdipour Ghazi has received funding from the Lundbeck Foundation (R400-2022-
365　617). Mads Nielsen is supported by the Pioneer Centre for AI, Danish National Research
366　Foundation, grant number P1. Michael Eriksen Benros has received funding from the
367　Lundbeck Foundation (R278-2018-1411 and R383-2022-285).

References:

1. Ioannidis, J. P. (2011). Excess significance bias in the literature on brain volume abnormalities. Archives of general psychiatry, 68(8):773–780.

2. Button, K. S., Ioannidis, J. P., Mokrysz, C., et al. (2013). Power failure: why small sample size undermines the reliability of neuroscience. Nature reviews neuroscience, 14(5):365–376.

3. Thompson, P. M., Jahanshad, N., Ching, C. R., et al. (2020). ENIGMA and global neuroscience: A decade of large-scale studies of the brain in health and disease across more than 40 countries. Translational psychiatry, 10(1):100.

4. Thompson, P. M., Stein, J. L., Medland, S. E., et al. (2014). The ENIGMA Consortium: large-scale collaborative analyses of neuroimaging and genetic data. Brain imaging and behavior, 8:153–182.

5. Van Erp, T. G., Hibar, D. P., Rasmussen, J. M., et al. (2016). Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. Molecular psychiatry, 21(4):547–553.

6. Van Erp, T. G., Walton, E., Hibar, D. P., et al. (2018). Cortical brain abnormalities in 4474 individuals with schizophrenia and 5098 control subjects via the enhancing neuro imaging genetics through meta analysis (ENIGMA) consortium. Biological psychiatry, 84(9):644–654.

7. Schmaal, L., Veltman, D. J., van Erp, T. G., et al. (2016). Subcortical brain alterations in major depressive disorder: findings from the ENIGMA Major Depressive Disorder working group. Molecular psychiatry, 21(6):806–812.

8. Schmaal, L., Hibar, D., Sämann, P. G., et al. (2017). Cortical abnormalities in adults and adolescents with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working Group. Molecular psychiatry, 22(6):900– 909.

9. Mackey, S., Allgaier, N., Chaarani, B., et al. (2019). Mega-analysis of gray matter volume in substance dependence: general and substance-specific regional effects. American Journal of Psychiatry, 176(2):119–128.

10. Groenewold, N. A., Bas-Hoogendam, J. M., Amod, A. R., et al. (2023). Volume of subcortical brain regions in social anxiety disorder: mega-analytic results from 37 samples in the ENIGMA-Anxiety Working Group. Molecular psychiatry, 28(3):1079–1089.

11. Billot, B., Magdamo, C., Cheng, Y., Arnold, S. E., Das, S., and Iglesias, J. E. (2023). Robust machine learning segmentation for large-scale analysis of heterogeneous clinical brain MRI datasets. Proceedings of the National Academy of Sciences, 120(9):e2216399120.

12. Gopinath, K., Greve, D. N., Das, S., Arnold, S., Magdamo, C., and Iglesias, J. E. (2023). Cortical analysis of heterogeneous clinical brain MRI scans for large-scale neuroimaging studies. In International Conference on Medical Image Computing and Computer-Assisted Intervention, pages 35–45. Springer.

13. Gopinath, K., Greve, D. N., Magdamo, C., et. al. (2025). "Recon-all-clinical": Cortical surface reconstruction and analysis of heterogeneous clinical brain MRI. Medical Image Analysis, 103608.

14. Cai N, Verhulst B, Andreassen OA, et al. (2025). Assessment and ascertainment in
psychiatric molecular genetics: challenges and opportunities for cross-disorder research. Mol
Psychiatry Apr;30(4):1627-1638.

15. Schork AJ, Won H, Appadurai V, et al. (2019). A genome-wide association study of
shared risk across psychiatric disorders implicates gene regulation during fetal
neurodevelopment. Nat Neurosci. 22(3):353-361.

16. Hettwer, M., Larivière, S., Park, B., et al. (2022). Coordinated cortical thickness
alterations across six neurodevelopmental and psychiatric disorders. Nature communications,
13(1):6851.

17. Opel N, Goltermann J, Hermesdorf M, et al. (2020). Cross-Disorder Analysis of Brain
Structural Abnormalities in Six Major Psychiatric Disorders: A Secondary Analysis of Mega-
and Meta-analytical Findings From the ENIGMA Consortium. Biol Psychiatry.

18. Cheon, E. J., Bearden, C. E., Sun, D., et al. (2022). Cross disorder comparisons of brain
structure in schizophrenia, bipolar disorder, major depressive disorder, and 22q11. 2 deletion
syndrome: A review of ENIGMA findings. Psychiatry and Clinical Neurosciences, 76(5),
140-161.

19. Boedhoe, P. S., van Rooij, D., Hoogman, M., et al. (2020). Subcortical brain volume,
regional cortical thickness, and cortical surface area across disorders: findings from the
ENIGMA ADHD, ASD, and OCD working groups. American Journal of Psychiatry, 177(9),
834-843.

20. Patel, Y., Parker, N., Shin, J., et al. (2021). Virtual histology of cortical thickness and
shared neurobiology in 6 psychiatric disorders. JAMA psychiatry, 78(1), 47-63.

21. Patel, Y., Shin, J., Abé, C., et al. (2022). Virtual ontogeny of cortical growth preceding
mental illness. Biological psychiatry, 92(4), 299-313.

22. Li, X., Morgan, P. S., Ashburner, J., Smith, J., and Rorden, C. (2016). The first step for
neuroimaging data analysis: DICOM to NIfTI conversion. Journal of neuroscience methods,
264:47–56.

23. Billot, B., Greve, D. N., Puonti, O., et al. (2023). SynthSeg: Segmentation of brain MRI
scans of any contrast and resolution without retraining. Medical image analysis, 86:102789.

24. Iglesias, J. E., Billot, B., Balbastre, Y., et al. (2023). SynthSR: A public AI tool to turn
heterogeneous clinical brain scans into high-resolution T1-weighted images for 3D
morphometry. Science advances, 9(5):eadd3607.

25. Fischl, B. (2012). FreeSurfer. Neuroimage, 62(2):774–781.

26. Dice, L. R. (1945). Measures of the amount of ecologic association between species.
Ecology, 26(3), 297-302.

27. Goodkind, M., Eickhoff, S. B., Oathes, D. J., et al. (2015). Identification of a common
neurobiological substrate for mental illness. JAMA psychiatry, 72(4):305–315.

28. Killiany, R. J., Gomez-Isla, T., Moss, M., et al. (2000). Use of structural magnetic
resonance imaging to predict who will get Alzheimer's disease. Annals of Neurology: Official

Journal of the American Neurological Association and the Child Neurology Society,
47(4):430–439.

29. Dickerson, B. C., Goncharova, I., Sullivan, M., et al. (2001). MRI-derived entorhinal and
hippocampal atrophy in incipient and very mild Alzheimer's disease. Neurobiology of aging,
22(5):747–754.

30. Apostolova, L. G., Green, A. E., Babakchanian, S., et al. (2012). Hippocampal atrophy
and ventricular enlargement in normal aging, mild cognitive impairment (MCI), and
Alzheimer Disease. Alzheimer Disease & Associated Disorders, 26(1):17–27.

31. Seabold, S. and Perktold, J. (2010). Statsmodels: econometric and statistical modeling
with python. SciPy, 7:1.

32. Cohen, J. (1992). Quantitative methods in psychology: A power primer. Psychol. Bull.,
112:1155–1159.

33. Nakagawa, S. and Cuthill, I. C. (2007). Effect size, confidence interval and statistical
significance: a practical guide for biologists. Biological reviews, 82(4):591–605.

34. Bonferroni, C. (1936). Teoria statistica delle classi e calcolo delle probabilita.
Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze, 8:3–
62.

35. Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., and Smith, S. M.
(2012). FSL. Neuroimage, 62(2):782–790.

36. Ashburner, J. and Friston, K. J. (2005). Unified segmentation. neuroimage, 26(3):839–
851.

37. Fortin, J. P., Cullen, N., Sheline, Y. I., et al. (2018). Harmonization of cortical thickness
measurements across scanners and sites. Neuroimage, 167, 104-120.

38. Sun, D., Rakesh, G., Haswell, C. C., et al. (2022). A comparison of methods to harmonize
cortical thickness measurements across scanners and sites. Neuroimage, 261, 119509.
472

Figure legend

Figure 1: Illustration of the MRI processing pipeline, selected inclusion and exclusion criteria, and analysis. More details are provided in Supplementary Methods.

Figure 2: Comparisons between mental disorders and Non-Psychiatric Controls. Panel A: Patients with mental disorders vs. non-psychiatric controls across inclusion and exclusion criteria. Panel B: Patients with dementia vs. non-psychiatric controls across inclusion and exclusion criteria. Panel C: Patients with mental disorders vs. non-psychiatric controls, excluding dementia patients from the cases group across inclusion and exclusion criteria. Panel D: Impact of slice thickness on differences between patients with mental disorders and non-psychiatric controls for Population C. Results for Population A and B can be found in Supplementary Figure S8. Numerical values for all the figures can be found in Supplementary Table S17 to Table S22.

Figure 3: Mental Disorders vs. Non-Psychiatric Controls. Comparisons for all ROI for Population C. Results for Population A and B can be found in Supplementary Figure S4 and Figure S5. Numerical values for all the figures can be found in Supplementary Table S23 to Table S25.

Figure 4: Mental Disorders Subgroups vs. Non-Psychiatric Controls. Comparisons with ENIGMA studies for Population B. R: Right, L: Left, and B: Bilateral. Results for Population A and C can be found in Supplementary Figure S6 and Supplementary Figure S7. Numerical values for all the figures can be found in Supplementary Table S26 to Table S28.

Table 1: Demographic of the clinical MRI dataset of individuals with mental disorders and non-psychiatric controls.

Group	Population A				Population B				Population C			
	N	Sites	Sex [male/female] [female %]	Age [mean ± std] [min, max]	N	Sites	Sex [male/female] [female %]	Age [mean ± std] [min, max]	N	Sites	Sex [male/female] [female %]	Age [mean ± std] [min, max]
Mental Disorders (any)	2774	36	[1589/1185] [43%]	[52.65±18.45] [18, 95]	2764	35	[1584/1180] [43%]	[52.60±18.46] [18, 95]	785	33	[458/327] [42%]	[45.56±18.25] [18, 92]
Non-Psychiatric Controls	2062	36	[1217/845] [41%]	[51.37±17.74] [18, 98]	1382	35	[797/585] [42%]	[48.11±16.93] [18, 98]	1376	33	[793/583] [42%]	[48.11±16.94] [18, 98]
Mental Disorders without Dementia	2606	36	[1509/1097] [42%]	[51.28±18.01] [18, 95]	2596	35	[1504/1092] [42%]	[51.22±18.01] [18, 95]	751	33	[438/313] [42%]	[44.30±17.53] [18, 89]
Non-Psychiatric Controls	2062	36	[1217/845] [41%]	[51.37±17.74] [18, 98]	1382	35	[797/585] [42%]	[48.11±16.93] [18, 98]	1376	33	[793/583] [42%]	[48.11±16.94] [18, 98]
Dementia	262	26	[138/124] [47%]	[75.09±9.82] [34, 93]	262	26	[138/124] [47%]	[75.09±9.82] [34, 93]	60	16	[35/25] [42%]	[74.18±9.09] [46, 92]
Non-Psychiatric Controls	1818	26	[1067/751] [41%]	[51.33±17.80] [18, 98]	1194	26	[689/505] [42%]	[47.75±16.96] [18, 98]	1048	16	[619/429] [41%]	[48.04±16.84] [18, 98]
Substance Use Disorder	574	33	[254/320] [56%]	[58.58±15.06] [19, 90]	569	32	[252/317] [56%]	[58.50±15.10] [19, 90]	119	20	[46/73] [61%]	[53.77±14.75] [19, 78]
Non-Psychiatric Controls	2005	33	[1183/822] [41%]	[51.40±17.74] [18, 98]	1335	32	[772/563] [42%]	[48.12±16.93] [18, 98]	1201	20	[702/499] [42%]	[47.82±16.88] [18, 98]
Schizophrenia Spectrum Disorder	435	28	[213/222] [51%]	[41.29±17.53] [18, 88]	435	28	[213/222] [51%]	[41.29±17.53] [18, 88]	214	23	[102/112] [52%]	[36.28±15.97] [18, 78]
Non-Psychiatric Controls	1987	28	[1169/818] [41%]	[51.33±17.74] [18, 98]	1326	28	[762/564] [43%]	[48.04±16.94] [18, 98]	1269	23	[734/535] [42%]	[48.00±16.86] [18, 98]
Depression	676	33	[459/217] [32%]	[53.28±17.00] [18, 91]	674	32	[458/216] [32%]	[53.27±17.03] [18, 91]	179	28	[127/52] [29%]	[48.44±16.57] [18, 85]
Non-Psychiatric Controls	2047	33	[1208/839] [41%]	[51.39±17.73] [18, 98]	1372	32	[792/580] [42%]	[48.15±16.92] [18, 98]	1335	28	[774/561] [42%]	[48.04±16.88] [18, 98]
Anxiety Disorder	524	31	[365/159] [30%]	[48.06±16.05] [18, 87]	524	31	[365/159] [30%]	[48.06±16.05] [18, 87]	155	24	[117/38] [25%]	[43.17±14.88] [18, 84]
Non-Psychiatric Controls	2024	31	[1193/831] [41%]	[51.39±17.76] [18, 98]	1350	31	[775/575] [43%]	[48.08±16.96] [18, 98]	1289	24	[743/546] [42%]	[48.14±16.85] [18, 98]

Population A: Non-Psychiatric Controls: age ≥18, No Mental Disorders, No Neurological Disorder. Cases: age ≥18, Mental Disorders.
Population B: Non-Psychiatric Controls: Population A + No Brain Medications. Cases: Same as Population A.
Population C: Non-Psychiatric Controls: Same as Population B. Cases: Population A + No Neurological Medications, No Neurological Comorbidities.

Figure 1: Illustration of the MRI processing pipeline, with selected inclusion and exclusion criteria, and analysis.

Figure 2: Mental disorders vs. Non-Psychiatric Controls comparisons.

504 **Figure 3: Mental Disorders vs. Non-Psychiatric Controls. Comparisons for all ROI for Population C.**

Figure 4: Mental Disorders Subgroups vs. Non-Psychiatric Controls. Comparisons with

ENIGMA studies for Population B.