Brief interactive lifestyle preventive education in the primary care clinic: protocol for a randomized clinical trial

March 18, 2025

Dylan Arroyo[1, *], Daniel Keyes[1, 2, 3¶], David Steinberger[1,], Ghada Abdulshafi[1,&,] Ali Jafri[1, &],

Hemica Hasan[1, &], Alice Nassar[1,], Anukul Karn[1], Preanka Dhanoa[1,] Rachel Nkrumah[1,] Batoul

Dabajeh[1]

[1] Department of Graduate Medical Education, Trinity Health-Livonia Hospital, Livonia, Michigan, United States

[2] Michigan State University COM Osteopathic Medical Specialties, Lansing, Michigan, United States

[3] University of Michigan, School of Medicine, Ann Arbor, Michigan, United States

Corresponding author: Daniel Keyes, MD, MPH

Email: Daniel.Keyes@trinity-health,org

¶ Conceptualization, Methodology, Project administration, Writing – original draft, Writing – review & editing: All authors contributed equally to this work.

* Methodology, Project administration, Writing – original draft, Writing – review & editing: All authors contributed equally to this work.

& Project administration, Writing – review & editing

Abstract

Introduction

It is widely accepted that prevention is far more impactful than curative medicine concerning overall health efficacy. The primary care setting faces the challenges of increasingly important time constraints and conflicting priorities; however, prevention is a priority for improving health outcomes. In a previous pilot study, we investigated the impact of a simple instructional video on patients' willingness to change in the emergency department setting. The initial intervention was passive, requiring subjects to watch a video without interaction. The current study evaluates the practicality and impact of bringing a brief *interactive* educational video intervention to patients at their primary care clinic visit.

Materials and methods

This is a randomized, non-blinded prospective study of education for adult patients presenting to the primary care clinic immediately following their primary care appointment.

Patients will be randomized to receive the interactive intervention video during this encounter with the practitioner in the clinic. The video is interactive because it asks the patients to answer simple questions about the content in the video to keep them engaged. For instance, if the participant indicates that they do not smoke, the video will "skip" the smoking cessation education and move directly to the other prevention domains. Afterwards, all patients will be presented with a questionnaire about their readiness and confidence to initiate lifestyle changes (transtheoretical stage of change measured on a "readiness ruler") and additional survey instruments. Most of these questions will come from existing validated scales, including the Gillespie & Lenz behavior modification tool, the Pittsburgh Sleep Quality Index, and select questions from the US HCAHPS Patient Satisfaction Survey (Appendix B). They will be asked for access to their medical record and their willingness to be contacted for a follow-up survey.

Outcome measures

The study's primary outcome will be the results of the Lifestyle Readiness to Change and Confidence to Change questionnaires. Secondary outcomes will include satisfaction with the primary care clinic visit using select questions from HCAHPS (Appendix B) and whether they intend to change any specific lifestyle behavior (dichotomous, yes/no) and, if yes, to specify which one(s), seeking medical care outside of the clinic visit as determined by a review of the patient's medical record at intervals of 30 days and six months. HIV will not be evaluated outside of its inclusion in the Charlson Comorbidity Index (CCI). A follow-up survey at approximately 3-12 months will evaluate new diagnoses related to chest pain and ask about the individual's lifestyle changes.

Statistical Summary Plan

64   Logistical regression analysis will be conducted for the primary outcome variable,
65 controlling for the Charlson Comorbidity Index, sex, age, and other predictive variables. We
66 estimate an approximate sample size of 350 patients.
67
68 **Discussion**
69   The Lifewise Primary Care trial is the first evaluation of comprehensive preventive
70 education using video to minimize interference with the practitioner's workflow.
71
72 **Trial Registration**
73   The study was approved by Trinity Health at Ann Arbor IRB (IRB number: E-24-1117).
74 Trial was prospectively registered in the Clinicaltrials.gov database (registration number:
75 NCT06730737).

Introduction

Preventive measures are critical in avoiding and limiting the severity of diseases. It is
intuitive and widely accepted that preventing disease is much more potent than treating
morbidities once they have developed. It has been estimated that health behaviors contribute as
much as four-fifths of health outcomes, which can be attributed to behavioral and social factors,
with clinical care only contributing a small proportion of influence [1,2]. This implies that the
clinical encounter should always promote behavior changes, emphasizing prevention. However,
contemporary healthcare is characterized by scheduling efficiencies emphasizing shorter times
spent in direct contact between the provider and the patient. Direct observations of primary care
practices have noted that prevention is under-emphasized [3]. Various reasons have been given
for the lack of time for lifestyle change counseling. It is thus implicit that new methods are
needed to educate patients about self-care and prevention that require less of a clinician's time.

This project will evaluate the impact of a video-based educational intervention
administered in the primary care clinic following the practitioner's first encounter with the
patient. It will also endeavor to examine the feasibility and efficiency of introducing a patient
and the accompanying family to the importance of adopting changes in lifestyle behaviors.

Receptivity to health behavior changes has been extensively investigated. The
transtheoretical model is a highly-used measurement of receptivity to change and may be used to
optimize intervention [4]. This model relates that there are six stages people fall into when
adopting a more healthful lifestyle. The first stage is *pre-contemplation,* in which people have no
plans to take action to change their behavior in the near future. This may be due to a need for
more information or previous unsuccessful attempts to change. *Contemplation* occurs when
people want to change their behavior within the next six months. They may be more aware of the
benefits and costs, but this awareness can create trepidation and cause people to remain in this
stage for extended periods. The third stage is *preparation*, defined as an individual's readiness to
change their behavior within the next month. They have a plan and are ready for an action-
oriented program. The following stages include *action,* where changes are initiated within the
last six months; *maintenance,* where people continue positive behaviors and work to prevent
relapse; and *termination,* where people have no temptation to return to their old behavior. The
transtheoretical model is often applied to preventive behavioral change. For example, it was
reported that current smokers typically fall into a spread of 40% pre-contemplation, 40%
contemplation, and 20% preparation [4]. This research suggests that when faced with the
potential for serious adverse health outcomes, patients may be able to accelerate through some
early stages, such as moving from pre-contemplation to contemplation or onto the preparation
stage.
There have been previous attempts to implement prevention and education in the primary
care clinic. These attempts include written after-visit summaries (AVS) shared with the patient.

However, it has been reported that the AVS is not the most effective tool, considering that only
82.8% of patients recalled receiving an AVS, and 67.4% of patients consulted their AVS [5]. The
information provided in the AVS is typically particular to the reason for that visit and includes
contact information for a follow-up appointment. Another study measured whether patients given
the AVS have better or worse glaucoma medication recall and revealed that 51.2% did not recall
receiving any AVS. Those who recalled receiving one paradoxically had *lower* medication recall
scores (remembering the name or color of the bottle or cap, treatment eye(s), and dosing
regimen) than those who did not [6]. These results show that the intended goal of printed AVS,
which is to enhance patient care, fails to be consistently achieved.

Moreover, there have been previous attempts at smoking cessation education in the
primary care clinic. One example of these studies consisted of a training program focused on
motivational interviewing to reduce smoking among adolescent patients. Follow-up assessments
were conducted at 1, 3, and 6 months post-interview [7]. It was reported that abstinence rates at
the 6-month follow-up were significantly higher in the motivational interview group compared to
the standard advice control group [7].

Innovative approaches are needed to help patients absorb and retain the information
required to properly ensure a healthy future. Showing a video discussing ways to enhance health
while the patient is in the PCP office could be that medium. In one study, patients who watched
videos while they were waiting to be seen resulted in a higher proportion of patients rated their
stay as "excellent" or "very good" (65%) in comparison to those who were not shown a video
(58.1%) [8]. Those who received the video discharge instructions tested better on a post-test
survey than the standard instructions group [9]. Another smaller study showed that after viewing
an educational video on myocardial infarction, participants in the group shown the video had
statistically significant improvement in the results of their knowledge assessment compared to
the group not shown the video. The intervention group had a mean number of correct responses
of 27 out of 37 questions as opposed to 20 in the control [10]. This supports using a video format
as a potentially practical approach for preventive education.

Objectives

141 Primary objective: This study will evaluate the utility of presenting patients attending a
142 primary care clinic with a focused video that will expose them to evidence-based education
143 based on four of the core "pillars of preventive lifestyle health" promulgated by the American
144 College of Lifestyle Medicine [11], including successful strategies for diet, exercise, sleep
145 hygiene, and smoking cessation.
146

Materials and methods

This will be a randomized, controlled, unblinded, two-armed superiority clinical trial.
The study will be conducted according to the Standard Protocol Items: Recommendations for

Interventional Trials (SPIRIT) Checklist. The study was approved by Trinity Health at Ann
Arbor IRB (IRB number: E-24-1117). Trial was prospectively registered in the Clinicaltrials.gov
database (registration number: NCT06730737).
The study's primary outcome will be the results from the Lifestyle Readiness to Change
and Confidence to Change questionnaire, which uses the *readiness ruler* (see Appendix B) [12].
The intervention will include four primary prevention domains promulgated by the American
College of Lifestyle Medicine: diet, exercise, sleep hygiene, and smoking cessation.
Secondary aims include:
1) To evaluate the ability to implement a preventive education strategy by health
practitioners in the primary care clinic setting
2) To assess the impact of age on willingness to change lifestyle behaviors
3) To determine those receiving the video concerning patient satisfaction utilizing select
HCAHPS questions, compared to the control group

Eligibility

Inclusion criteria:
● All adults, including all ethnicities and genders, between the ages of 18 and 80 who come
into the primary care clinic will be recruited to participate in the study.
Exclusion criteria:
● Non-English-speaking patient
● Unable or unwilling to consent to the study
● Unable or unwilling to hear a video on a smartphone or computer tablet
● The patient is in hospice care
● Patients with advanced dementia, incapable of answering the survey questions in the
opinion of the person administering the survey
○ Note: research associates will consult the patient's primary practitioner or the
175   physician members of the research team for any questions regarding dementia
176 Note:
177   ● There are no relevant concomitant care and interventions that are permitted or prohibited
178   during the trial.

Participant Enrollment and Screening

Adult patients aged 18 to 80 will be recruited by a study team member or a research
associate before their appointment or upon arrival at an Academic Internal Medicine ambulatory
clinic. Please see the flow diagram for the randomization process (Figure 1 and 2).
The randomization module within REDCap will be used to allocate the video
intervention. Those randomized for the intervention will be asked if they are interested and

willing to watch a video about the clinic visit and their health. Those who agree to watch will
then interact with a video featuring lifestyle changes relating to diet, exercise, sleeping, and
smoking and why they are important for one's overall health. The media is interactive because it
contains questions for the user to answer at different points within the video sequence. The
control group will not watch the video.
The study includes two surveys: the first will be administered in the clinic, and the
second will be a follow-up survey administered upon phone contact with consenting patients.
Target follow-up is planned for 3-12 months following the initial visit.
Whether receiving the video intervention or not, all participants will have provided verbal
consent to answer the questionnaires (see the included script at the heading of the initial survey
instrument (Appendix B). They will then fill out the questionnaire at the end of the video either
alone, with the assistance of an accompanying relative, or with the assistance of a research team
member to determine their stage of the transtheoretical model. A study identification number will
be placed on the completed survey. The ID number and the patient's identifying information will
be noted in REDCap. Any patient who declines to participate will not have their patient health
information (PHI) included in the study. The REDCap file containing the patients' personally
identifiable information will be kept on password-protected servers using the institution's
REDCap.

203 Study procedures

Research associates and/or research team members will screen for eligible patients
using a screening form based on the inclusion/exclusion criteria mentioned above.
Results will be analyzed at two stages: first, with the results of the questionnaires
administered as part of the initial visit, and second, after concluding the 3-12 month follow-up.
The consent, educational video, and questionnaire will all be conducted while the patient is in the
waiting room, after the clinician (physician or advanced practice provider) has completed the
initial patient encounter, or after the provider contact has been completed.
The patients will fill out the composite questionnaire. This questionnaire is designed with
the following criteria: prioritize using validated tools capable of self-administration. The
Readiness-to-Change questionnaire evaluates patient readiness to implement ten lifestyle
behaviors based on a 5-point rating scale. A rating of 5 represents having no interest in the
suggested lifestyle change, which is consistent with the precontemplation stage of the
transtheoretical model. A rating of 4 means an interest in implementing the lifestyle change in
the next few months (contemplation stage of the TTM). A rating of 3 represents having the plan
to implement the lifestyle change within the next month (preparation stage of the TTM). A rating
of 2 represents having already started the lifestyle change within the past few months (action
stage of TTM). A rating of 1 means having maintained lifestyle behavior changes for over 6
months (maintenance stage of TTM). The Confidence-to-Change questionnaire evaluates patient
confidence in implementing the same lifestyle behaviors as the Readiness-to-Change

questionnaire. However, the Confidence-to-Change questionnaire is based on a 3-point rating

scale. Ratings are as follows: 3-very confident, 2-somewhat confident, and 1-not very confident

[12]. The Confidence-to-Change questionnaire addresses any barriers or hesitancy that may

impact patient confidence in implementing better lifestyle behaviors. Additional questions will

include measures of satisfaction taken from the HCAHPS, specifically addressing their

practitioner and visit (Appendix B).

Sometime after the visit (target: 3-12 months later), they will receive a phone follow-up

call, asking them if they have adopted any lifestyle changes since watching the video, among

other inquiries (Appendix C). We will use patients' responses at this later stage (follow-up phase)

to help determine how sustainable the preventive education intervention has been.

Randomized patients who participate in the study will receive the video intervention.

In a previous pilot study, we investigated the impact of a simple instructional video on patients'

willingness to change in the emergency department setting [13]. The initial intervention was

passive, requiring subjects to watch a video without interaction. The current study evaluates the

practicality and impact of bringing a brief *interactive* educational video intervention to patients

attending the primary care clinic visit.

All patients, including those randomized to "no video learning," will be asked to

complete a questionnaire. The questions will be administered orally or self-administered for

those willing to participate using the tablet interface. Responses will be recorded using REDCap

electronic data capture software and the related platform.

It is estimated that the sample will include a total of 350 participants. Due to the

minimal risks in this study, a data monitoring committee is not necessary. An interim analysis

will be conducted after 250 patients have been enrolled. It is anticipated that the total data

collection will take six months. For those patients who agree to receive a call-back, contact will

be attempted during the 3 to 12-month follow-up period. A research associate, a resident, or an

attending physician will complete the follow-up by telephone or email correspondence with the

subject. Up to 3 follow-up attempts will be made. Available follow-up data from the interactive

electronic health record (EHR) will be obtained on all patients. The estimated time to view the

video is approximately 10-15 minutes. The time required for responses to the survey component

is estimated at 10-12 minutes.

The EHR will be used to determine several variables: age, race, ethnicity, sex and

gender, BMI, and the Charlson Comorbidity Index (CCI). Michigan Data Analytics will assist in

generating the patient's CCI from ICD-10 codes. Information from the patient's chart will only

be obtained from those who provide informed consent.

The research team will conduct training for the research as per protocol. This training

will include informed consent, knowledge of the inclusion and exclusion criteria, using the EPIC

EHR, and practice administering the study. The study will be performed using a convenience

sample when team members are available to facilitate the study procedures.

Upon completion of the study, the results will be analyzed and presented at regional

and national scientific meetings and for publication. The protocol will be submitted for

publication prior to the initiation of the trial. Authorship of the resulting publication will follow
the guidelines of the International Committee of Medical Journal Editors (ICMJE).No
professional writers will be used for publication. No individual patient identifiers will be
revealed during the abstract presentation or publication of the results. The data will be
maintained in a password-protected electronic file on a hospital server using REDCap, a secure
electronic data capture system.
269
270 Note that all videos will use the interactive format. Patients will not be video recorded.

Data Analysis and Statistical Considerations

The study team members will collect demographic data upon registration to the patient's
clinic. Patients from ages 18 to 80 will be included in the study. Following informed consent,
patients will be randomized to either the intervention or control arms of the study. Only the
interventional arm will receive a video link or QR code or start on the video using a provided
computer tablet or the patient's phone for the streaming content, which provides the focused
lifestyle preventive education. The control patients will not receive a video and will go directly
to the follow-up questionnaire. Hence, both groups will receive the follow-up survey by research
team members.

The Demographic data includes age, sex, race, and insurance status. The patient's BMI,
Charlson Comorbidity Index, smoking status, answers to the social influences of health
questionnaires, phone number, and frequency of their ED visits will be documented and taken
from the patient's EHR. Their zip code will also be collected for geographic and socioeconomic
status analysis.

The study's primary outcome is the readiness and confidence to change scores as a
function of video exposure, measured on a continuous "readiness ruler" or using a dichotomous
categorization of ages 18-39 and 40-80 [14].

Secondary outcomes will include readiness to change in specific domains, including
exercise, smoking, diet, sleep hygiene, living an overall healthy lifestyle, satisfaction with the
hospital, and satisfaction with the practitioner dichotomized between those who watched the
video and those who did not. Part of this data will be obtained directly from the patient's medical
record.

We will collaborate with Michigan Data Analytics, which will provide large-scale data
abstraction to assist with obtaining the Charlson Comorbidity Index (CCI) for participating
subjects.

The remaining data will be collected using the Readiness-to-Change and Confidence-to-
Change Lifestyle Questionnaires.

Statistical Considerations

Demographic characteristics will be analyzed using basic descriptive statistics. Logistic
Regression Analysis will be used to evaluate the major outcomes of the study based on
predictive variables included in the data collection fields (Appendix A). Binary (if dichotomous)
or multivariate (if the outcome is multiple or continuous, such as the readiness ruler) logistic
regression will be used. Statistical significance will be defined utilizing an $\square$ of $p < 0.05$ and/or
95% confidence intervals. Data analysis will be performed by intention-to-treat utilizing multiple
imputation techniques for missing data analysis. The reasons for missing data will be reported.

Data Confidentiality and Storage

307   All team members will be trained in HIPAA and safe data handling practices. To
308 maintain patient confidentiality and privacy, the data will be maintained on a password-protected
309 secure server and using the secure, password-protected REDCap platform.

Data and Safety Monitoring Plan

The principal investigator (PI) is responsible for this project's conduct and shall ensure
that all work and services described, or incidental to those described, are conducted following
accepted standards of medical practice. The PI and co-investigators will maintain all master
subject lists, and all data collected will be maintained in a password-protected secure server and
reviewed at regular intervals (see below) to ensure adherence to the investigational procedures
outlined in the protocol.

The approved research team members will obtain information regarding the patient's
lifestyle and demographic habits. Data will be accepted from primary care clinic records and
charts. Each data abstractor will use a pre-formulated data collection tool to enhance the
uniformity of information. Statistical analysis for inter-observer variability will be conducted on
a subset of duplicate charts to ensure the quality of the data abstracted and used for the study.

Meetings will be held at regular intervals to discuss data management, general IRB
compliance, and other study aspects and to analyze the collected data. The PI will evaluate data
quality and ensure that data evaluation is performed frequently. All data will be documented and
retained in the study files.

Regulatory Requirements

327 Informed Consent

All patients will be consented. Informed consent will be obtained before the randomization of
subjects. If they approve, the research team member will administer the informed consent form
attached to this protocol.

Those who approve to watch the interactive video will be given a QR code to watch on their
smartphone or a tablet with a link to the video, which the patient or the research team member may
start.

334 Participant Privacy

Subject confidentiality will be maintained using a randomly generated participant identification
number/code. The participant identification number will be stored with the patient's identifying
information, such as name and room number, for verification purposes. Identifying information
will be kept in a password-protected, secure REDCap file on a password-protected server. Such
information will only be accessible to the research team trained in research ethics (CITI training)
and HIPAA compliance.
341
342
343 Using personally identifiable data is necessary to determine eligibility, abstract chart data, and
344 follow up with patients. All patient information collected will be stored in a password-protected
345 secure server using REDCap and only accessed when necessary by authorized research
346 associates.

347 HIPAA

All parties involved shall maintain the confidentiality of PHI to ensure compliance with
privacy regulations, including the Health Insurance Portability and Accountability Act of
1996 (HIPAA). The study team will consist of only personnel trained in research ethics (CITI
training) and HIPAA compliance.

Patient eligibility will be determined by screening in Epic EHR. We will request a waiver of

HIPAA authorization for this portion of the study and access to the patient's medical record.

354 Unanticipated Problems and Adverse Events

This project is considered to be minimal risk, such as loss of protected health information and
discomfort of answering questions. Any unanticipated problems will be promptly reported to the

PI, project's research coordinator and the IRB. If necessary, the project will be discontinued until
resolution in consultation with Trinity Health, Ann Arbor IRB. This trial involves preventive
health education and does not involve any interventions that carry a risk of physical harm.
Therefore, provisions for ancillary and post-trial care related to trial participation, as well as
compensation for harm, are not applicable.

362 Conflicts of Interests

363
364 No conflicts of interest have been identified, and the authors wish to acknowledge no known
365 conflicts of interest for this study. Note that Dylan Arroyo received funding as a Clinical
366 Research and Innovation Scholarship Program (CRISP) Scholar from the University of
367 Michigan-Dearborn. This research is not being sponsored or funded by any other organization.

368 Limitations and Assumptions

369 Since this is planned as a single-institution study, this will limit generalizability.

12   REFERENCES

1. Park H, Roubal AM, Jovaag A, Gennuso KP, Catlin BB. Relative Contributions of a Set of Health Factors to Selected Health Outcomes. Am J Prev Med. 2015 Dec;49(6):961-9. doi: 10.1016/j.amepre.2015.07.016. PMID: 26590942.

2. Hood CM, Gennuso KP, Swain GR, Catlin, BB. County Health Rankings: Relationships Between Determinant Factors and Health Outcomes. Am J Prev Med. 2016 Feb;50(2):129-35. doi: 10.1016/j.amepre.2015.08.024. PMID: 26526164.

3. Stange KC, Flocke SA, Goodwin MA, Kelly RB, Zyzanski SJ. Direct observation of rates of preventive service delivery in community family practice. Prev Med. 2000 Aug;31(2 Pt 1):167-76. doi: 10.1006/pmed.2000.0700. PMID: 10938218.

4. Prochaska JO, Velicer WF. The transtheoretical model of health behavior change. Am J Health Promot. 1997 Sep-Oct;12(1):38-48. doi: 10.4278/0890-1171-12.1.38. PMID: 10170434.

5. Pathak S, Summerville G, Kaplan CP, Nouri SS, Karliner LS. Patient-Reported Use of the After Visit Summary in a Primary Care Internal Medicine Practice. J Patient Exp. 2020 Oct;7(5):703-707. doi: 10.1177/2374373519879286. Epub 2019 Oct 4. PMID: 33294604; PMCID: PMC7705830.

6. Hirabayashi KJ, Pomerantz M, Radell JE, Chadha N, Thomas S, Serle JB. The Efficacy of the After-visit Summary in Medication Recall Among Glaucoma Patients. J Glaucoma. 2020 Jul;29(7):529-535. doi: 10.1097/IJG.0000000000001518. PMID: 32332333.

7. Colby SM, Monti PM, O'Leary Tevyaw T, Barnett NP, Spirito A, Rohsenow DJ, Riggs S, Lewander W. Brief motivational intervention for adolescent smokers in medical settings. Addict Behav. 2005 Jun;30(5):865-74. doi: 10.1016/j.addbeh.2004.10.001. Epub 2004 Nov 10. PMID: 15893085.

8. Papa L, Seaberg DC, Rees E, Ferguson K, Stair R, Goldfeder B, Meurer D. Does a waiting room video about what to expect during an emergency department visit improve patient satisfaction? CJEM. 2008 Jul;10(4):347-54. doi: 10.1017/s1481803500010356. PMID: 18652727.

9. Wood EB, Harrison G, Trickey A, Friesen MA, Stinson S, Rovelli E, McReynolds S, Presgrave K. Evidence-Based Practice: Video-Discharge Instructions in the Pediatric Emergency Department. J Emerg Nurs. 2017 Jul;43(4):316-321. doi: 10.1016/j.jen.2016.11.003. Epub 2017 Mar 28. PMID: 28359707.

10. Dhawan N, Saeed O, Gupta V, Desai R, Ku M, Bhoi S, Verma S. Utilizing video on myocardial infarction as a health educational intervention in patient waiting areas of the developing world: A study at the emergency department of a major tertiary care hospital in India. Int Arch Med. 2008 Jul 29;1(1):14. doi: 10.1186/1755-7682-1-14. PMID: 18662408; PMCID: PMC2547095.

11. Lifestyle medicine certification. (2024). Accessed: September 20, 2024:
https://lifestylemedicine.org/overview/#:~:text=Most%20physicians%20and%20health%20p
rofessionals,six%20pillars%20of%2
12. Gillespie, N. D., & Lenz, T. L. (2011). Implementation of a tool to modify behavior in a
chronic disease management program. *Advances in preventive medicine*, *2011*, 215842.
https://doi.org/10.4061/2011/21584.
13. Keyes D C, Arroyo D L, Abdulshafi G W, et al. (November 30, 2024) Seizing the Teachable
Moment: A Pilot Study of Preventive Lifestyle Education in the Emergency Department.
Cureus 16(11): e74825. doi:10.7759/cureus.74825
14. Prochaska JO, DiClemente CC. Stages of change in the modification of problem behaviors.
Prog Behav Modif. 1992;28:183-218. PMID: 1620663.

Appendices

Appendix A: Data collection fields

REDCap will be used to collect all data

- Medical Record Number
- Age
- Sex
- Gender
- Race
- Ethnicity
- Smoking Status and use of tobacco products
- BMI
- Charlson Comorbidity Index
- Insurance Status
- Social Influencers of Health Questionnaire (if available)
- Patient zip code
- Insurance status
- Follow-up questionnaire items: see page 21, Appendix C for questions included during telephonic and EHR follow-up.
- If the patient is living in an institution or living independently
- HIV will not be evaluated outside of its inclusion in the aggregate Charlson Comorbidity Index (CCI), which will prevent any possibility of calculating HIV status indirectly).

Appendix B: Primary Care Clinic Questionnaires and Script

Hello! My name is (Research assistant/associate name), and I am a research associate here at the clinic. You (or a family member) came to our primary clinic today. To participate in our study, you will be asked to complete a brief questionnaire about your lifestyle habits, such as sleep, smoking, diet, and exercise. We hope that your participation will help us do a better job of giving patients instructions before they leave the clinic. Would that be okay?

DO NOT PROCEED IF INFORMED CONSENT IS NOT OBTAINED.

If informed consent is obtained, proceed with randomization of video or no video.
If the patient is randomized to video, tell the patient: "Your provider (optional - give the name of provider here) would like you to watch a video that will teach you about how to live a healthy lifestyle. Would you like to watch it now?"

457 NOTE: The following two questions will be administered using REDCap data collection tool. .
458 For each prompt, the participant will be asked about their readiness score and then about their
459 confidence level immediately after, using the corresponding visual analog scale (VAS (- i.e., in
460 parallel).

On the scale below, enter how you feel right now about the lifestyle behavior

1	2	3	4	5
No interest in this lifestyle behavior at this time	Thinking about implementing this lifestyle behavior sometime in the next few months	Plan to implement this lifestyle behavior sometime in the next month	Recently (within the previous 6 months) began this lifestyle behavior	Have been maintaining this lifestyle behavior for 6+ months

461
462
463
Source: Gillespie and Lenz 2011

Readiness-to-Change Lifestyle Questionnaire

Name: _______________________________

Date: _______________________________

____

Rating   Readiness-to-Change (USE "READINESS RULER" ABO VE

____

Rating	Readiness-to-Change (USE "READINESS RULER" ABOVE
5	No interest in this lifestyle behavior at this time
4	Thinking about implementing this lifestyle behavior sometime in the next few months
3	Plan to implement this lifestyle behavior sometime in the next month
2	Recently (within the previous 6 months) began this lifestyle behavior
1	Have been maintaining this lifestyle behavior for 6+ months

Rating	Lifestyle behavior
______	Purposeful exercise at least 5 times per week
______	Incorporate "extra" physical activity throughout the daily routine ("extra" = taking stairs rather than the elevator, walking pets, parking further away in the parking lot, etc.)
______	Participate in sporting activity at least 1-2 times weekly (e.g., golf, volleyball, basketball, tennis)
______	Eat 5 or more servings of fruits and vegetables daily
______	Consistently choose foods with whole grains
______	Choose foods high in fiber
______	Lose or maintain body weight
______	Avoid smoking or tobacco use
______	Implementing strategies to help with lifestyle modifications
______	Live an overall healthy lifestyle

1	2	3
Not very confident	Somewhat confident	Very confident

Confidence-to-Change Lifestyle Questionnaire

Name: _________________________________

Date: _________________________________

Rating	Confidence-to-Change

501	3	Very confident
502	2	Somewhat confident
503	1	Not very confident

504 ____

505   Rating   Lifestyle behavior

506 ____

507   ______   Purposeful exercise at least 5 times per week

508   ______   Incorporate "extra" physical activity throughout the daily routine ("extra"
509   = taking stairs rather than the elevator, walking pets, parking further away
510   in the parking lot, etc.)

511   ______   Participate in sporting activity at least 1-2 times weekly (e.g., golf,
512   volleyball, basketball, tennis)

______   Eat 5 or more servings of fruits and vegetables daily

______   Consistently choose foods with whole grains

______   Choose foods high in fiber

______   Lose or maintain body weight

______   Avoid smoking or tobacco use

______   Implementing strategies to help with lifestyle modifications

______   Live an overall healthy lifestyle

Continue immediately to the next set of questions (PSQI)

The Pittsburgh Sleep Quality Index (PSQI)

1. When do you usually go to bed? ______________
2. How long (in minutes) has it taken you to fall asleep each night? ________________
3. When have you usually gotten up in the morning? ______________
4. How many hours of actual sleep do you get at night? (This may be different than the
number of hours you spend in bed) ____________________

5. During the past month, how often have you had trouble	Not during the past month (0)	Less than once a week (1)	Once or twice a week (2)	Three or more times a week (3)

sleeping because you…				
a. Cannot get to sleep within 30 minutes				
b. Wake up in the middle of the night or early morning				
c. Have to get up to use the bathroom				
d. Cannot breathe comfortably				
e. Cough or snore loudly				
f. Feel too cold				
g. Feel to hot				
h. Have bad dreams				
i. Have pain				
j. Other reason(s), please describe, including how often you have had trouble sleeping				

because of this reason(s):				
7. During the past month, how often have you taken medicine (prescribed or "over the counter") to help you sleep?				
7. During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity?				
8. During the past month, how much of a problem has it been for you to keep up enthusiasm to get things done?				
	Very good (0)	Fairly good (1)	Fairly bad (2)	Very bad (3)
9. how would you rate your sleep quality over the past month?				

*After each participant completes the survey, the research associate scores the Pittsburgh Sleep
Quality Index questionnaire and enters the score in REDCap.

Total Sleep Index Score: _______ (Range 0 to 21)

Do you currently smoke? ☐ Yes ☐ no Did you previously smoke? ☐ Yes ☐ no

If yes to either, how many packs per day (PPD) and how many years?

PPD: ______ Years: ______ (Maximum = 99)

Do you plan to make any lifestyle changes as a result of today's visit? If so, what kind? Check all
that apply.

- No planned changes
- Initiating a change in my diet
- Initiating or increasing my exercise in a safe manner
- Improving my sleep habits and/or hours of daily sleep
- Stop smoking
- Other ____

545 **Select questions from the US Governmental HCAHPS Patient Satisfaction Survey**

546 During this hospital stay, how often did doctors treat you with courtesy and respect?

- Never
- Sometimes
- Usually
- Always

551 During this hospital stay, how often did doctors listen carefully to you?

- Never
- Sometimes
- Usually
- Always

556 During this hospital stay, how often did doctors explain things in a way you could understand?

- Never
- Sometimes
- Usually
- Always

561 Using any number from 0 to 10, where 0 is the worst hospital possible, and 10 is the best hospital
562 possible, what number would you use to rate this hospital during your stay?

- 0   Worst hospital possible
- 1
- 2
- 3
- 4
- 5
- 6
- 7

● 8
● 9
● 10   Best hospital possible

Would you recommend this hospital to your friends and family

● Definitely no
● Probably no
● Probably yes
● Definitely yes

When I left the hospital, I had a good understanding of the things I was responsible for in
managing my health.

● Strongly disagree
● Disagree
● Agree
● Strongly agree

In general, how would you rate your overall health?

● Excellent
● Very good
● Good
● Fair
● Poor

What is the highest grade or level of school that you have completed?

● 8th grade or less
● Some high school but did not graduate
● Good
● High school graduate or GED
● Some college or 2-year degree
● 4-year college graduate
● More than a 4-year college degree

May we contact you for a follow-up after this visit? If so, what number? _____________________
600
601 How to to make this more effective in helping you make changes to prevent diseases in your
602 life? (If randomized to the video, ask how the video can be improved)
603 ____
604 ____

605 # Appendix C: <u>Follow-up</u> Survey and Script
606
Hello! Is this (Patient Name)?

This is (Insert research associate name); I'm part of the research team. You consented at your
primary care clinic visit on (insert clinic visit date) that we could call you at this number for a
follow-up survey regarding your lifestyle behaviors. Are you free to answer our survey of a few
questions?

If no:
That is okay! When would be a good time for us to call you back for a follow-up?
Date/Time: _______________

If yes:
Have you adopted any of the lifestyle changes discussed in the video shown to you in the clinic
regarding diet, sleep, exercise, or smoking?
If so, what changes have you adopted?

Those are all the questions! Thank you so much for your time and have a great day!

Follow-up Patient Interview - Prior to interview:

List diagnoses from the last two office visits:

____

____

____

Is chest pain included as one of the above diagnoses? ❏ No ❏ Yes

❏ Don't Know

Interview: Please select the party providing the information for this interview. Check all
that apply (may include more than one person present during the interview). **Note:** only
patients willing and able to participate in the original interview will complete the survey.
However, some patients may request another person to answer the questionnaires. Please
indicate here who provided answers to the questionnaires:

❏ Patient ❏ Spouse/Significant Other ❏ Sibling ❏ Child ❏ Other Family Member ❏

Medical Care Provider ❏ Other: _______________________________

1. Did the patient receive follow-up care after their index PCP visit?

❏ No ❏ Yes ❏ Don't Know

2. Is the patient doing well?

❏ No ❏ Yes ❏ Don't Know

3. Have you adopted any lifestyle changes, including diet, exercise, sleep, or smoking?

❏ No ❏ Yes ❏ Don't Know

Specify here which changes:
____
____

Note any special considerations here, e.g. patient in hospital now, at a follow-
up appointment during the time of call, the patient is deceased:

____

Appendix D: Educational Videos Used for Prevention and Script

If informed consent is obtained, proceed with randomization of video or no video.

If the patient is randomized to video, tell the patient: "Your provider (optional - give the name of

provider here) would like you to watch a video that will teach you how to live a healthy lifestyle.

Would you like to watch it now?"

Link for Video Playlist:

https://bit.ly/Lifestyle_Medicine_Prevention

	STUDY PERIOD				
	Enrollment	Allocation	Post-allocation		Close-out
TIMEPOINT	$-t_1$	0	t_1 (time 0)	t_2 (3-12 months)	t_x
ENROLMENT:					
Eligibility screen	X				
Informed consent	X				
HIPAA Waiver	X				
Randomization/ Allocation		X			
INTERVENTIONS:					
Watch Video			X		
Standard-of-care			X		
ASSESSMENTS:					
*Baseline Assessment**	X				
*Primary Outcomes***			X		X
*Secondary outcomes****			X		X
Follow-up Survey				X	X

Fig 1. Schedule of enrollment, interventions, and assessments
*Sociodemographic, clinical, and lifestyle characteristics
**Readiness/Confidence to Change questionnaires
***Satisfaction with the primary care clinic visit and others

Figure

Fig 2. Flowchart for the study process.
Note that all videos will use the interactive format. Patients will not be video recorded.

Figure