Brief interactive lifestyle preventive medicine video education in the primary care clinic: protocol for a randomized clinical trial

December 22, 2025

Dylan Arroyo[1,2¶], Daniel Keyes[1,3,4*¶&], Ghadah W. Abdulshafi[1,2¶], Ali Jafri[1], Hemica Hasan[1],

David Steinberger[1], Batoul Dabajeh[1], Alice Nassar[1], Anukul Karn[1], Rachel Nkrumah[1], Preanka

Dhanoa[1], Kerby Shedden[5&]

[1] Department of Graduate Medical Education, Trinity Health Livonia Hospital, Livonia,

Michigan, United States of America

[2] Clinical Research and Innovation Scholarship Program (CRISP), University of Michigan-

Dearborn, Dearborn, Michigan, United States of America

[3] Michigan State University COM Osteopathic Medical Specialties, Lansing, Michigan, United

States of America

[4] University of Michigan, School of Medicine, Ann Arbor, Michigan, United States of America

[5] Department of Statistics, University of Michigan, Ann Arbor, Michigan, United States of

America

*Corresponding author: Daniel Keyes, MD, MPH

Email: Daniel.Keyes@trinity-health.org

22 [¶]These authors contributed equally to this work.

23 [&]These authors also contributed equally to this work.

Abstract

Introduction

Preventive counseling improves long-term health, but primary care clinicians face time

constraints limiting patient education. In a prior emergency department pilot study, a brief

passive video trended towards increased patients' willingness to change health behaviors. The

current trial, conducted at the primary care clinic, evaluates the feasibility and impact of a short,

interactive prevention video delivered during a primary care office visit.

Methods and analysis

We will conduct a prospective, randomized, non-blinded clinical trial among adult patients

presenting to a primary care clinic. Immediately after their appointment, participants will be

randomized to receive either the interactive prevention video with a practitioner or usual care.

The video maintains engagement through simple in-video questions and adapts content based on

patient responses (e.g., skipping smoking cessation for non-smokers). After the encounter, all

participants will complete surveys assessing readiness and confidence to initiate lifestyle change

including a transtheoretical "readiness ruler." Survey items will be drawn from validated

instruments, including the Gillespie & Lenz Readiness and Confidence to Participate in Lifestyle

Activities Surveys, the Pittsburgh Sleep Quality Index (PSQI), and select questions from the U.S.

Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) Survey.

Patients will be asked for medical record access and permission for follow-up contact.

Outcomes

The primary outcome is index-visit Lifestyle Readiness and Confidence to Change scores.

Secondary outcomes include satisfaction with the clinic visit, intention to change specific

lifestyle behaviors, and healthcare use outside the clinic at 30 days and 6 months, as assessed via

record review. A 3-12-month follow-up survey will assess self-reported lifestyle changes and

new diagnoses.

Statistical plan and sample size

Analyses will follow intention-to-treat. Ordinal logistic regression will compare primary

outcomes between arms, adjusting for Charlson Comorbidity Index, age, sex, gender, and

prespecified predictors. Logistic regression will evaluate binary follow-up outcomes. Complete-

case analyses will be used, and differential attrition will be reported. We plan to enroll

approximately 350 patients.

Trial registration

ClinicalTrials.gov NCT06730737.

Introduction

Preventive measures are critical in avoiding and limiting the severity of diseases. It is

intuitive and widely accepted that preventing disease is much more potent than treating

morbidities once they have developed. It has been estimated that health behaviors contribute as

much as four-fifths of health outcomes, which can be attributed to behavioral and social factors,

with clinical care only contributing a small proportion of influence [1,2]. This implies that the

clinical encounter should always promote behavior changes, emphasizing prevention. However,

contemporary healthcare is characterized by scheduling efficiencies emphasizing shorter times

spent in direct contact between the provider and the patient. Direct observations of primary care

practices have noted that prevention is under-emphasized [3]. Various reasons have been given

for the lack of time for lifestyle change counseling. It is thus implicit that new methods are

needed to educate patients about self-care and prevention that require less of a clinician's time.

This project will evaluate the impact of a video-based educational intervention

administered in the primary care clinic following the practitioner's first encounter with the

patient. It will also endeavor to examine the feasibility and efficiency of introducing patients and

their accompanying family members to the importance of adopting changes in lifestyle

behaviors.

Receptivity to health behavior changes has been extensively investigated. The

transtheoretical model is a highly used measurement of receptivity to change and may be used to

optimize intervention [4]. This model relates that there are six stages people fall into when

adopting a more healthful lifestyle. The first stage is *pre-contemplation,* in which people have no

plans to take action to change their behavior in the near future. This may be due to a need for

more information or previous unsuccessful attempts to change. *Contemplation* occurs when

people want to change their behavior within the next six months. They may be more aware of the

benefits and costs, but this awareness can create trepidation and cause people to remain in this

stage for extended periods. The third stage is *preparation*, defined as an individual's readiness to

change their behavior within the next month. They have a plan and are ready for an action-

oriented program. The following stages include *action,* where changes are initiated within the

last six months; *maintenance,* where people continue positive behaviors and work to prevent

relapse; and *termination,* where people have no temptation to return to their old behavior. The

transtheoretical model is often applied to preventive behavioral change. For example, it was

reported that current smokers typically fall into a spread of 40% pre-contemplation, 40%

contemplation, and 20% preparation [4]. This research suggests that when faced with the

potential for serious adverse health outcomes, patients may be able to accelerate through some

early stages, such as moving from pre-contemplation to contemplation or onto the preparation

stage.

There have been previous attempts to implement prevention and education in the primary

care clinic. These attempts include written after-visit summaries (AVS) shared with the patient.

However, it has been reported that the AVS is not the most effective tool, considering that only

82.8% of patients recalled receiving an AVS, and 67.4% of patients consulted their AVS [5]. The

information provided in the AVS is typically specific to the reason for that visit and includes

contact information for a follow-up appointment. Another study measured whether patients given

the AVS have better or worse glaucoma medication recall and revealed that 51.2% did not recall

receiving any AVS. Those who recalled receiving one paradoxically had *lower* medication recall

scores (remembering the name or color of the bottle or cap, treatment eye(s), and dosing

regimen) than those who did not [6]. These results show that the intended goal of printed AVS,

which is to enhance patient care, fails to be consistently achieved.

Moreover, there have been previous attempts at smoking cessation education in the

primary care clinic. One example of these studies consisted of a training program focused on

motivational interviewing to reduce smoking among adolescent patients. Follow-up assessments

were conducted at 1, 3, and 6 months post-interview [7]. It was reported that abstinence rates at

the 6-month follow-up were significantly higher in the motivational interview group compared to

the standard advice control group [7].

Innovative approaches are needed to help patients absorb and retain the information

required to properly ensure a healthy future. Showing a video on ways to enhance health while

the patient is in the primary care provider (PCP) office could be an effective medium. In one

study, patients who watched videos while they were waiting to be seen resulted in a higher

proportion of patients rated their stay as "excellent" or "very good" (65%) in comparison to

those who were not shown a video (58.1%) [8]. Those who received the video discharge

instructions tested better on a post-test survey than the standard instructions group [9]. Another

smaller study showed that after viewing an educational video on myocardial infarction,

participants in the group shown the video had statistically significant improvement in the results

of their knowledge assessment compared to the group not shown the video. The intervention

group had a mean number of correct responses of 27 out of 37 questions as opposed to 20 in the

control [10]. This supports the use of video as a practical approach to preventive education.

Objectives

Primary objective: This study will evaluate the utility of presenting patients attending a

primary care clinic with a focused video that will expose them to evidence-based education

based on four of the core "pillars of preventive lifestyle health" promulgated by the American

College of Lifestyle Medicine [11], including successful strategies for diet, exercise, sleep

hygiene, and smoking cessation.

Study hypotheses

We hypothesize that patients receiving the interactive video intervention will exhibit

greater readiness and confidence to change lifestyle behaviors compared to patients in the control

group, as measured post-intervention.

Null hypothesis (H_0): There is no difference in readiness and confidence to change

lifestyle behaviors between the intervention group and the control group.

Alternative hypothesis (H_1): The intervention group will demonstrate significantly greater

readiness and confidence to change lifestyle behaviors than the control group.

Materials and methods

141   This will be a randomized, controlled, unblinded, two-armed superiority clinical trial.

142 The study will be conducted according to the Standard Protocol Items: Recommendations for

143 Interventional Trials (SPIRIT) Checklist. The study was approved by Trinity Health at Ann

144 Arbor Institutional Review Board (IRB number: E-24-1117). The trial was prospectively

145 registered in the Clinicaltrials.gov database (registration number: NCT06730737).

146   The study's primary outcome will be the results from the Lifestyle Readiness to Change

147 and Confidence to Change questionnaires, both of which use a *readiness ruler* [12]. The

148 intervention will include four primary prevention domains promulgated by the American College

149 of Lifestyle Medicine: diet, exercise, sleep hygiene, and smoking cessation.

150 Secondary aims include:

151   1) To evaluate the ability to implement a preventive education strategy in the primary care

152    clinic setting

153   2) To assess the impact of age on willingness to change lifestyle behaviors

154   3) To determine those receiving the video concerning patient satisfaction, utilizing select

155    Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) Survey

156    questions, compared to the control group

Eligibility

159 Inclusion criteria:

160  • All adults, including all ethnicities and genders, between the ages of 18 and 80 who come

161   into the primary care clinic will be recruited to participate in the study.

162 Exclusion criteria:

163  • Non-English-speaking patient

164  • Unable or unwilling to consent to the study

165  • Unable or unwilling to hear a video on a smartphone or computer tablet

166  • The patient is in hospice care

● Patients with advanced dementia, incapable of answering the survey questions in the

opinion of the person administering the survey

○ Note: research associates will consult the patient's primary practitioner or the

physician members of the research team for any questions regarding dementia

Note:

● There are no relevant concomitant care and interventions that are permitted or prohibited

during the trial.

Participant enrollment and screening

Adult patients aged 18 to 80 will be recruited by a study team member or a research

associate before their appointment or upon arrival at an Academic Internal Medicine ambulatory

clinic. Please see the flow diagram for the randomization process and the table for the study

process (Figure 1 and Table 1).

Fig 1. Flowchart for the Study Process. Note that all videos will use the interactive format.

Patients will not be video recorded.

Table 1. Schedule of Enrollment, Interventions, and Assessments.

	STUDY PERIOD				
	Enrollment	Allocation	Post-allocation		Close-out
TIMEPOINT	$-t_1$	0	t_1 (time 0)	t_2 (3-12 months)	t_x

ENROLLMENT:					
Eligibility Screen	X				
Informed Consent	X				
HIPAA Waiver	X				
Randomization/Allocation		X			
INTERVENTIONS:					
Watch Video			X		
Standard-of-care			X		
ASSESSMENTS:					
Baseline Assessment*	X				
Primary Outcomes**			X		X
Secondary Outcomes***			X		X
Follow-up Survey				X	X

*Sociodemographic, clinical, and lifestyle characteristics

**Readiness/Confidence to Change questionnaires

***Satisfaction with the primary care clinic visit and others

The randomization module within Research Electronic Data Capture (REDCap) will be

used to allocate the video intervention. Those randomized for the intervention will be asked if

they are interested and willing to watch a video about the clinic visit and their health. Those who

agree to watch will then interact with a video featuring lifestyle changes relating to diet, exercise,

sleeping, and smoking and why they are important for one's overall health. The media is

interactive because it contains questions for the user to answer at different points within the

video sequence. The control group will not watch the video.

The study includes two surveys: the first will be administered in the clinic, and the

second will be a follow-up survey administered upon phone contact with consenting patients.

Target follow-up is planned for 3-12 months following the initial visit.

Whether receiving the video intervention or not, all participants will have provided verbal

consent to answer the questionnaires. They will then fill out the questionnaire at the end of the

video either alone, with the assistance of an accompanying relative, or with the assistance of a

research team member to determine their stage of the transtheoretical model. A study

identification number will be placed on the completed survey. The ID number and the patient's

identifying information will be noted in REDCap. Any patient who declines to participate will

not have their patient health information (PHI) included in the study. The REDCap file

containing the patients' personally identifiable information will be kept on password-protected

servers using the institution's REDCap.

Study procedures

Research associates and/or research team members will screen for eligible patients using

a screening form based on the inclusion/exclusion criteria mentioned above. Results will be

analyzed at two stages: first, with the results of the questionnaires administered as part of the

initial visit, and second, after concluding the 3-12 month follow-up survey. The consent,

educational video, and questionnaire will all be conducted while the patient is in the waiting

room, after the clinician (physician or advanced practice provider) has completed the initial

patient encounter, or after the provider contact has been completed.

The patients will fill out the composite questionnaire. This questionnaire is designed with

the following criteria: prioritize using validated tools capable of self-administration. The

Readiness-to-Change questionnaire evaluates patient readiness to implement ten lifestyle

behaviors based on a 5-point rating scale. A rating of 5 represents having no interest in the

suggested lifestyle change, which is consistent with the precontemplation stage of the

transtheoretical model (TTM). A rating of 4 means an interest in implementing the lifestyle

change in the next few months (contemplation stage of the TTM). A rating of 3 represents

having the plan to implement the lifestyle change within the next month (preparation stage of the

TTM). A rating of 2 represents having already started the lifestyle change within the past few

months (action stage of TTM). A rating of 1 means having maintained lifestyle behavior changes

for over 6 months (maintenance stage of TTM). The Confidence-to-Change questionnaire

evaluates patient confidence in implementing the same lifestyle behaviors as the Readiness-to-

Change questionnaire. However, the Confidence-to-Change questionnaire is based on a 3-point

rating scale. Ratings are as follows: 3-very confident, 2-somewhat confident, and 1-not very

confident [12]. The Confidence-to-Change questionnaire addresses any barriers or hesitancy that

may impact patient confidence in implementing better lifestyle behaviors. Additional questions

will include measures of satisfaction taken from the HCAHPS, specifically addressing their

practitioner and visit.

Sometime after the visit (target: 3-12 months later), they will receive a phone follow-up

call, asking them if they have adopted any lifestyle changes since watching the video, among

other inquiries. We will use patients' responses at this later stage (follow-up phase) to help

determine how sustainable the preventive education intervention has been.

Randomized patients who participate in the study will receive the video intervention. In a

previous pilot study, we investigated the impact of a simple instructional video on patients'

willingness to change in the emergency department setting [13]. The initial intervention was

passive, requiring subjects to watch a video without interaction. The current study evaluates the

practicality and impact of bringing a brief *interactive* educational video intervention to patients

attending the primary care clinic visit.

All patients, including those randomized to "no video learning," will be asked to

complete a questionnaire. The questions will be administered orally or self-administered for

those willing to participate using the tablet interface. Responses will be recorded using REDCap

electronic data capture software and the related platform.

A sample size of approximately 350 was chosen based on feasibility and the expectation

of sufficient power to detect moderate differences in readiness and confidence scores, guided by

prior pilot study results. Due to the minimal risks in this study, a data monitoring committee is

not considered necessary. An interim analysis will be conducted after 250 patients have been

enrolled. It is anticipated that the total data collection will take six months. For those patients

who agree to receive a call-back, contact will be attempted during the 3 to 12-month follow-up

period. A research associate, a resident, or an attending physician will complete the follow-up by

telephone or email correspondence with the subject. Up to 3 follow-up attempts will be made.

Available follow-up data from the interactive electronic health record (EHR) will be obtained on

all patients. The estimated time to view and interact with the video is approximately 10-15

minutes. The time required for responses to the survey component is estimated at 10-12 minutes.

The EHR will be used to determine several variables: age, race, ethnicity, sex and gender,

BMI, and the Charlson Comorbidity Index (CCI). Michigan Data Analytics will assist in

generating the patient's CCI from ICD-10 codes. Information from the patient's chart will only

be obtained from those who provide informed consent.

The research team training will include informed consent, knowledge of the inclusion and

exclusion criteria, using the EPIC EHR, and practice administering the study. The study will be

performed using a convenience sample when team members are available to facilitate the study

procedures.

Upon completion of the study, the results will be analyzed and presented at regional and

national scientific meetings and for publication. The protocol will be submitted for publication

prior to the initiation of the trial. Authorship of the resulting publication will follow the

guidelines of the International Committee of Medical Journal Editors (ICMJE). No professional

writers will be used for publication. No individual patient identifiers will be revealed during the

abstract presentation or publication of the results. The data will be maintained in a password-

protected electronic file on a hospital server using REDCap, a secure electronic data capture

system.

Statistical methods

The study team members will collect demographic data upon registration to the patient's

clinic. Patients from ages 18 to 80 will be included in the study. After informed consent, patients

will be randomized to either the intervention or the control arm of the study. Only the

interventional arm will interact with the video using a provided computer tablet which focuses on

lifestyle preventive education. Control patients will not receive a video and will go directly to the

follow-up questionnaire. Hence, both groups will receive the follow-up survey from research

team members.

The Demographic data includes age, sex, gender, race, and insurance status. The patient's

BMI, Charlson Comorbidity Index, smoking status, responses to the social influences of health

questionnaires, phone number, and frequency of ED visits will be documented from the patient's

EHR. Their zip code will also be collected for geographic and socioeconomic status analysis.

The study's primary outcome is the readiness and confidence to change scores as a

function of video exposure, measured on a continuous "readiness ruler." [14].

Secondary outcomes will include readiness to change in specific domains, including

exercise, smoking, diet, sleep hygiene, living an overall healthy lifestyle, satisfaction with the

hospital, and satisfaction with the practitioner, dichotomized between those who watched the

video and those who did not. Part of this data will be obtained directly from the patient's medical

record.

We will collaborate with Michigan Data Analytics, which will provide large-scale data

abstraction to assist with obtaining the Charlson Comorbidity Index (CCI) for participating

subjects. The remaining data will be collected using the Readiness-to-Change and Confidence-

to-Change Lifestyle Questionnaires, the Pittsburgh Sleep Quality Index (PSQI), and U.S.

Governmental HCAHPS Patient Satisfaction Survey.

Descriptive statistics will be used to summarize baseline characteristics by study arm,

including means and standard deviations for continuous variables and proportions for categorical

variables. All analyses will follow the intention-to-treat principle.

Outcomes collected at the clinic visit (e.g., readiness to change and confidence to change)

will be compared between intervention and control groups using ordinal logistic regression,

appropriate for ordinal outcome variables. As these measurements are taken after randomization

and intervention (where applicable), no pre-intervention baseline measures will be available.

Follow-up outcomes, including receipt of additional care, adoption of lifestyle changes,

and overall well-being, are recorded as binary or categorical variables (e.g., Yes/No/Don't

Know) and will be analyzed using logistic regression models. Clinic visit scores may be included

as covariates to assess their association with follow-up outcomes.

A complete-case approach will be used for all primary and secondary outcome analyses.

Differences in loss to follow-up between study arms will be reported, and characteristics of

completers and non-completers will be compared to assess potential attrition bias. Multiple

imputation is not planned due to the structure of data collection involving a single follow-up time

point.

Separate regression models will be used to analyze outcomes collected at the clinic visit

and follow-up. Given the design of the study, which includes a single post-intervention follow-up

and no pre-intervention baseline, repeated measures modeling will not be used in the primary

analysis. Secondary analyses may include generalized estimating equations or other repeated

measures approaches to assess the consistency of findings.

Data confidentiality and storage

325   All team members will be trained in HIPAA and safe data handling practices. To

326 maintain patient confidentiality and privacy, the data will be maintained on a password-protected

327 secure server and using the secure, password-protected REDCap platform.

329 Data and safety monitoring plan

The principal investigator (PI) is responsible for this project's conduct and shall ensure
that all work and services described, or incidental to those described, are conducted following
accepted standards of medical practice. The PI and co-investigators will maintain all master
subject lists, and all data collected will be maintained in a password-protected secure server and
reviewed at regular intervals to ensure adherence to the investigational procedures outlined in the
protocol.

The approved research team members will obtain information regarding the patient's
lifestyle and demographic habits. Data will be accepted from primary care clinic records and
charts. Each data abstractor will use a pre-formulated data collection tool to enhance the
uniformity of information. Statistical analysis for inter-observer variability will be conducted on
a subset of duplicate charts to ensure the quality of the data abstracted and used for the study.

Meetings will be held at regular intervals to discuss data management, general IRB
compliance, and other study aspects and to analyze the collected data. The PI will evaluate data
quality and ensure that data evaluation is performed frequently. All data will be documented and
retained in the study files.

346 Regulatory requirements

347 Informed consent

All patients will be consented. Informed consent will be obtained before the randomization

of subjects. If they approve, the research team member will administer the informed consent form

prior to randomization.

Those who approve to watch the interactive video will be given a QR code to watch on

their smartphone or a tablet with a link to the video, which the patient or the research team member

may start.

Participant privacy

Subject confidentiality will be maintained using a randomly generated participant

identification number/code. The participant identification number will be stored with the

patient's identifying information, such as name and room number, for verification purposes.

Identifying information will be kept in a password-protected, secure REDCap file on a password-

protected server. Such information will only be accessible to the research team trained in

research ethics (CITI training) and HIPAA compliance.

Using personally identifiable data is necessary to determine eligibility, abstract chart

data, and follow up with patients. All patient information collected will be stored in a password-

protected secure server using REDCap and only accessed when necessary by authorized research

associates.

HIPAA

All parties involved shall maintain the confidentiality of PHI to ensure compliance

with privacy regulations, including the Health Insurance Portability and Accountability Act

of 1996 (HIPAA). The study team will consist of only personnel trained in research ethics

(CITI training) and HIPAA compliance.

Patient eligibility will be determined by screening in Epic EHR. We will request a

waiver of HIPAA authorization for this portion of the study and access to the patient's

medical record.

Unanticipated problems and adverse events

This project is considered to be minimal risk, such as loss of protected health information

and discomfort of answering questions. Any unanticipated problems will be promptly reported to

the PI, project's research coordinator, and the IRB. If necessary, the project will be discontinued

until resolution in consultation with Trinity Health, Ann Arbor IRB. This trial involves

preventive health education and does not involve any interventions that carry a risk of physical

harm. Therefore, provisions for ancillary and post-trial care related to trial participation, as well

as compensation for harm, are not applicable.

Conflicts of interests

The authors wish to acknowledge no known conflicts of interest for this study. Note that

Dylan Arroyo and Ghadah Abdulshafi received funding as Clinical Research and Innovation

Scholarship Program (CRISP) Scholars from the University of Michigan-Dearborn, however, no

specific funding was received for this project. The clinical scholar funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript. This

research is not being sponsored or funded by any other organization.

Limitations and assumptions

390   Since this is planned as a single-institution study, this will limit generalizability.

Data availability

This manuscript describes a study protocol. No data has been generated or analyzed at

this stage. Upon completion of the study, all de-identified individual-level data underlying the

findings will be made publicly available in a recognized public repository in accordance with

Trinity Health IRB policies and participant consent provisions.

Study materials such as the survey instruments (including the Readiness-to-Change and

Confidence-to-Change questionnaires, Pittsburgh Sleep Quality Index [PSQI], and HCAHPS

items) and the video are available from the corresponding author upon reasonable request and

will be made publicly available upon peer-reviewed publication.

References

1. Park H, Roubal AM, Jovaag A, Gennuso KP, Catlin BB. Relative Contributions of a Set

of Health Factors to Selected Health Outcomes. Am J Prev Med. 2015 Dec;49(6):961-9. doi:

10.1016/j.amepre.2015.07.016. PMID: 26590942.

2. Hood CM, Gennuso KP, Swain GR, Catlin BB. County Health Rankings: Relationships

Between Determinant Factors and Health Outcomes. Am J Prev Med. 2016 Feb;50(2):129-

35. doi: 10.1016/j.amepre.2015.08.024. Epub 2015 Oct 31. PMID: 26526164.

3. Stange KC, Flocke SA, Goodwin MA, Kelly RB, Zyzanski SJ. Direct observation of rates

of preventive service delivery in community family practice. Prev Med. 2000 Aug;31(2 Pt

1):167-76. doi: 10.1006/pmed.2000.0700. PMID: 10938218.

4. Prochaska JO, Velicer WF. The transtheoretical model of health behavior change. Am J

Health Promot. 1997 Sep-Oct;12(1):38-48. doi: 10.4278/0890-1171-12.1.38. PMID:

10170434.

5. Pathak S, Summerville G, Kaplan CP, Nouri SS, Karliner LS. Patient-Reported Use of

the After Visit Summary in a Primary Care Internal Medicine Practice. J Patient Exp. 2020

Oct;7(5):703-707. doi: 10.1177/2374373519879286. Epub 2019 Oct 4. PMID: 33294604;

PMCID: PMC7705830.

6. Hirabayashi KJ, Pomerantz M, Radell JE, Chadha N, Thomas S, Serle JB. The Efficacy

of the After-visit Summary in Medication Recall Among Glaucoma Patients. J Glaucoma.

2020 Jul;29(7):529-535. doi: 10.1097/IJG.0000000000001518. PMID: 32332333.

7. Colby SM, Monti PM, O'Leary Tevyaw T, Barnett NP, Spirito A, Rohsenow DJ, Riggs S,

Lewander W. Brief motivational intervention for adolescent smokers in medical settings.

Addict Behav. 2005 Jun;30(5):865-74. doi: 10.1016/j.addbeh.2004.10.001. Epub 2004 Nov

10. PMID: 15893085.

8. Papa L, Seaberg DC, Rees E, Ferguson K, Stair R, Goldfeder B, Meurer D. Does a

waiting room video about what to expect during an emergency department visit improve

patient satisfaction? CJEM. 2008 Jul;10(4):347-54. doi: 10.1017/s1481803500010356.

PMID: 18652727.

9. Wood EB, Harrison G, Trickey A, Friesen MA, Stinson S, Rovelli E, McReynolds S,

Presgrave K. Evidence-Based Practice: Video-Discharge Instructions in the Pediatric

Emergency Department. J Emerg Nurs. 2017 Jul;43(4):316-321. doi:

10.1016/j.jen.2016.11.003. Epub 2017 Mar 28. PMID: 28359707.

10. Dhawan N, Saeed O, Gupta V, Desai R, Ku M, Bhoi S, Verma S. Utilizing video on

myocardial infarction as a health educational intervention in patient waiting areas of the

developing world: A study at the emergency department of a major tertiary care hospital in

India. Int Arch Med. 2008 Jul 29;1(1):14. doi: 10.1186/1755-7682-1-14. PMID: 18662408;

PMCID: PMC2547095.

11. American College of Lifestyle Medicine. Lifestyle medicine certification. 2024 [cited

2024 Sep 20] Available

from:https://lifestylemedicine.org/overview/#:~:text=Most%20physicians%20and%20health

%20professionals,six%20pillars%20of%2

12. Gillespie ND, Lenz TL. Implementation of a tool to modify behavior in a chronic disease

management program. Adv Prev Med. 2011;2011:215842. doi: 10.4061/2011/215842. Epub

2011 Mar 8. PMID: 21991432; PMCID: PMC3169365.

13. Keyes DC, Arroyo DL, Abdulshafi GW, Dabajeh B, Polgar J, Sakr R, Hasan S. Seizing

the Teachable Moment: A Pilot Study of Preventive Lifestyle Education in the Emergency

Department. Cureus. 2024 Nov 30;16(11):e74825. doi: 10.7759/cureus.74825. PMID:

39737259; PMCID: PMC11684556.

14. Prochaska JO, DiClemente CC. Stages of change in the modification of problem

behaviors. Prog Behav Modif. 1992;28:183-218. PMID: 1620663.