Caribbean Women Face Higher Obesity and Diabetes Amid Socioeconomic Struggles – A Cross-

Sectional Study

Authors: Cesar Barrabi, Amanda Fowler

Affiliation: Western Atlantic University School of Medicine

Abstract

This cross-sectional study examined gender disparities in health outcomes across 30 Caribbean nations,

focusing on the intersection of noncommunicable diseases (NCDs), reproductive health, and

socioeconomic inequality. Using publicly available data from the Pan American Health Organization,

United Nations Development Programme, and International Diabetes Federation, we first identified

significantly higher obesity prevalence (43.88% vs. 29.77%) and a greater proportion of diabetes-related

deaths among women (13.21% vs. 10.17%) in the Caribbean compared to men. We then compared

regional outcomes to high-income neighbors in North America, selecting the U.S. and Canada as

reference countries. The Caribbean showed substantially higher maternal mortality (9.39 vs. 1.41 per

10,000 live births) and infant mortality (11.58 vs. 5 per 1,000 live births), as well as lower Inequality-

adjusted Human Development Index scores (0.556 vs. 0.839). Socioeconomic disparities were also

pronounced, with higher rates of food insecurity (41.71% vs. 8.15%) and female unemployment (10.67%

vs. 4.45%) in the Caribbean. Together, these findings reveal how interlocking health and social

vulnerabilities disproportionately affect women in the region. Addressing these disparities will require

coordinated policy action that extends beyond healthcare access to target the structural determinants

driving gendered health inequities.

Introduction

Gender health disparities are a well-documented global issue, with women often facing worse outcomes in chronic diseases, mental health, and access to healthcare (1). In the Caribbean, these disparities are particularly stark, with women facing higher rates of obesity and diabetes and poorer reproductive health outcomes, as demonstrated in regional mortality assessments and global risk analyses (2, 3). While high-income countries like the United States of America (US) and Canada (CAN) also struggle with noncommunicable diseases (NCDs) (4), Caribbean nations such as the Bahamas and Barbados experience these challenges within more fragile healthcare systems and greater socioeconomic inequalities (5). Although lifestyle factors like poor nutrition and inactivity are widely recognized contributors (6), the deeper socio-economic influences on NCDs among Caribbean women remain underexplored.

Despite its relative wealth, the Bahamas exhibits high obesity rates and diabetes-related mortality, consistent with global trends showing disproportionate NCD burdens in middle-income nations (7). Caribbean women are 60% more likely to have diabetes and twice as likely to be obese as men, underscoring gendered health disparities [4]. Recent studies indicate that obesity rates among Bahamian women continue to rise, contributing to downstream chronic disease burdens including diabetes and kidney disease (8, 9). Targeted policy interventions are needed to expand healthcare access, address food insecurity, and reduce socioeconomic inequality, particularly among young adults and women (10). While North America offers extensive literature on NCDs, research specifically examining these complex relationships in Caribbean women is scarce.

Beyond chronic diseases, reproductive health poses additional risks. Caribbean women face higher maternal mortality rates, early menstruation, and late pregnancies, which contribute to breast cancer risks and poor reproductive outcomes (11, 12). The stigma around conditions like cervical cancer further limits healthcare access, creating a cycle of late diagnoses and higher mortality(13). These observed disparities

suggest that comprehensive approaches integrating medical treatment and addressing socioeconomic

barriers may be beneficial, consistent with findings from prior studies.

This study contributes new regional evidence on how structural inequities—spanning healthcare access,

socioeconomic vulnerability, and gender-specific risks—shape women's health across the Caribbean. By

integrating chronic disease, reproductive health, and policy data, our findings underscore the urgent need

for coordinated public health strategies that address the multifaceted drivers of health inequality. Regional

responses must move beyond behavioral interventions to include structural reforms that prioritize equity,

resilience, and long-term health system investment.

Materials & methods

Ethical Considerations

This study used publicly available, aggregated data from international organizations and did not involve

human subjects. Therefore, formal ethical approval was not required.

Study Design and Data Collection

This cross-sectional study analyzed gender disparities in health outcomes across 30 Caribbean nations

using publicly available data from the Pan American Health Organization (PAHO), United Nations

Development Programme (UNDP), and International Diabetes Federation (IDF) (14-16). The analysis

included metrics on noncommunicable diseases (NCDs), reproductive health outcomes, and

socioeconomic indicators, comparing these to benchmarks to NA. The dataset focused on 2019 data,

which represented the most recent year with comprehensive health and economic information. These data

sources were selected for their coverage, credibility, and relevance, particularly given the limited

availability of peer-reviewed literature addressing Caribbean health outcomes.

The primary outcomes assessed were obesity and diabetes prevalence, maternal and adolescent health

metrics, and socioeconomic disparities, including Human Development Index (HDI) scores, food

insecurity, and unemployment rates. Additional variables related to lifestyle, healthcare access, and

reproductive health were included to provide broader context.

In a separate analysis, we assessed national-level policy implementation related to NCD prevention using

data from the 2022 PAHO ENLACE interactive scorecard. This tool evaluates country-level progress on

commitments outlined in the 2011 UN Political Declaration and the 2014 UN Outcome Document on

NCDs. Four policy indicators were selected: (1) national evidence-based clinical guidelines for NCD

management, (2) policies or campaigns promoting physical activity, (3) implementation of the

International Code of Marketing of Breast-milk Substitutes, and (4) restrictions on the marketing of

unhealthy foods and beverages to children. Each indicator was classified by PAHO as fully implemented

(green), partially implemented (yellow), not implemented (red), or unknown (grey).

We focused on 2022 data as it was the most recent round available; no data were collected in 2019, and

previous years used different indicator definitions, limiting comparability. Countries were grouped into

two regions: North America (n = 2; Canada and the United States) and the Caribbean (n = 13; Antigua and

Barbuda, Bahamas, Barbados, Belize, Dominica, Grenada, Guyana, Jamaica, Saint Kitts and Nevis, Saint

Lucia, Saint Vincent and the Grenadines, Suriname, and Trinidad and Tobago).

Data Analysis

All data processing and visualization, including scatterplots and trend lines, were conducted using

Microsoft Excel. Pearson correlation, Principal Component Analysis (PCA), Bootstrapping, and

hierarchical clustering were implemented in Python.

For gender-wise comparisons, Student's t-tests were used to compare male-to-female differences within

Caribbean nations for each metric. Welch's t-tests were used for country-level comparisons, grouping the

United States and Canada as one group (North America or NA) and the 30 Caribbean nations as the other

group (Caribbean or CAR), with a significance cutoff of $p < 0.05$ (17). This test was chosen due to

account for unequal sample sizes. Data for gestational diabetes rates for the Caribbean regions were

retrieved from the International Diabetes Federation (IDF, 2021), using regional data. A Geographic

Information System (GIS) map of the Caribbean was generated in QGIS 3.34 T using GADM shapefiles,

with mapping performed by Fiver professional Ayesha Suraweera (18, 19). No statistics were performed

in QGIS, this is simply reporting values obtained from IDF.

Data imputation was performed exclusively for the Pearson correlation and Principal Component Analysis

(PCA) analyses. Countries with less than 75% of data for each variable were removed, and the remaining

missing data was imputed using Predictive Mean Matching within the Multiple Imputation by Chained

Equations framework. This approach maintains the original data distribution by matching predicted

values to the closest observed values, ensuring realistic imputation. After data imputation, Pearson

correlation coefficients were calculated to examine associations between gestational diabetes,

malnutrition rates, and socio-economic indicators (20).

PCA was performed using the imputed data to reduce the dimensionality of the dataset with socio-

economic and health indicators across the 30 countries (21). Variables with high collinearity were

removed based on Pearson correlations before PCA, and additional variables were hand-selected based on

relevance to women's health metrics, socio-economic indicators, and other health-related metrics. A scree

plot confirmed that the first two principal components explained the majority of the variance (sFigure1A).

Ward's hierarchical clustering was applied to the PCA scores to group countries based on their socio-

economic and health profiles (sFigure2), while the elbow method was applied to select the number of

clusters (sFigure1B). Bootstrapping was performed with 1,000 iterations to assess the stability of the PCA

results, achieving a 95% confidence interval.

Results

The Caribbean Shows Distinct Gendered Mortality Patterns, with Women Facing Higher Diabetes

Mortality

We first examined 2019 PAHO data across 30 Caribbean nations to evaluate gender-specific health

outcomes. External mortality rates were significantly higher in men (110.68 per 100,000) compared to

women (31.01; $p < 0.001$), while NCD mortality was higher in women (83.18%) than in men (75.33%; p

< 0.01). CD mortality showed no significant gender difference (Figure 1A).

To explore which conditions contributed to the observed NCD disparity, we assessed specific causes of

death. Mortality from ischemic heart disease, respiratory diseases, cirrhosis, and stomach cancer was

consistently higher in men than in women (all $p < 0.05$; Figure 1B).

We next examined diabetes-related mortality, given its known burden in the region. Women had a higher

proportion of diabetes-related deaths than men (13.21% vs. 10.17%; $p < 0.001$). When comparing regions,

the Caribbean reported a substantially higher proportion of diabetes-related mortality (62.22%) than NA

(11.65%; $p < 0.001$), highlighting a striking regional disparity (Figure 1C).

To assess potential contributors to this pattern, we analyzed lifestyle metrics. Women in the Caribbean

reported significantly higher rates of insufficient physical activity than men (67.79% vs. 53.47%; $p <$

0.001), and obesity prevalence was also higher among women (43.88% vs. 29.77%; $p < 0.001$; Figure

1D).

Together, these findings reveal distinct gendered health patterns across the Caribbean. While men

experience higher mortality from several leading NCD causes, women carry a disproportionate burden of

diabetes-related mortality, likely shaped by both behavioral and structural risk factors.

The Caribbean Faces Greater Socioeconomic and Health Challenges Compared to the USA/Canada

We then aimed to expand on the observed gendered health disparities, and assessed socioeconomic

conditions in the Caribbean relative to NA. The region had significantly lower Human Development

Index (HDI) scores (0.76 vs. 0.93; $p < 0.001$) and Inequality-Adjusted HDI (0.56 vs. 0.84; $p < 0.01$),

reflecting more limited development and deeper structural inequality (Figure 2A).

Food insecurity was also considerably higher in the Caribbean (41.71%) compared to NA (8.15%; $p <$

0.001), despite no significant regional differences in overall inflation rates. Rates in Haiti (18.7), Aruba

(4.3), and Barbados (4.1) were the highest recorded, suggesting that inflation alone does not explain the

region's more severe access challenges (Figure 2B).

We next explored mortality from external causes. Homicide mortality was significantly higher in the

Caribbean (7.31) than in NA (1.55; $p < 0.01$), while suicide mortality showed no significant difference

(6.1 vs. 3.87), though elevated rates in Guyana (17.0) and Suriname (11.8) contributed to regional

variability (Figure 2C).

To examine economic instability through a gendered lens, we compared unemployment rates. Women's

unemployment was significantly higher in the Caribbean (10.67%) than in NA (4.45%; $p < 0.01$), whereas

men's unemployment showed no regional difference (Figure 2D). Prior research has linked

unemployment to greater risk of diabetes and chronic disease, particularly among women, suggesting that

these disparities may exacerbate existing health inequities (23).

Together, these findings highlight how lower human development, limited food access, and gendered

employment gaps distinguish the Caribbean's socioeconomic landscape from its northern neighbors—and

may help explain the region's persistent health challenges.

Reproductive and Early-Life Health Indicators Underscore Deep Regional and Gender Inequities

To build on the observed health and socioeconomic disparities, we examined reproductive and early-life

health outcomes in the Caribbean compared to NA. Maternal mortality was significantly higher in the

Caribbean (9.39 per 10,000 live births) than in NA (1.41; $p < 0.001$), and infant mortality showed a

similar trend, with rates more than twice as high in the Caribbean (11.58 vs. 5 per 1,000 live births; $p <$

0.001; Figure 3A).

We next assessed adolescent health metrics to identify early-life contributors to regional disparities.

Adolescent fertility rates were higher in the Caribbean (34.26 per 10,000) and trended toward significance

when compared to NA (11.3; $p = 0.05006$). Adolescent mortality was significantly elevated in the

Caribbean (26.38 vs. 13.7 per 10,000; $p < 0.01$), highlighting additional risk in this age group (Figure

3B).

To contextualize these disparities, we examined broader demographic and health system indicators. Life

expectancy, median age, and total health spending did not differ significantly between regions. However,

out-of-pocket health expenditures were more than twice as high in the Caribbean (32.5%) relative to NA

(13.4%; $p < 0.01$), indicating a heavier financial burden on individuals seeking care (Figure 3C).

We also explored gestational diabetes mellitus (GDM) prevalence as a marker of maternal health

vulnerability. Both the Caribbean and North America reported a GDM prevalence of 20.7%, placing the

region behind South-East Asia (25.9%) but ahead of South and Central America (15.8%; Figure 3D).

These rates signal an emerging challenge for maternal health across the hemisphere.

Taken together, these findings underscore the Caribbean's reproductive and adolescent health challenges,

compounded by higher financial barriers to care. These gaps may contribute to the persistence of gender

and regional inequities in health outcomes.

Diabetes and Obesity Outcomes Strongly Correlate with Socioeconomic Development and

Healthcare Access in Caribbean Populations

We examined how socioeconomic conditions relate to population health by conducting Pearson correlation analyses across Caribbean nations with at least 75% data availability, with results visualized in a correlation heatmap (Figure 5A). Missing values were imputed specifically for these analyses. A complete correlation matrix is provided in the supplementary materials.

Diabetes mortality showed a strong inverse correlation with inequality-adjusted development, with higher IHDI values associated with lower age-adjusted diabetes deaths (r = -0.884). Adult obesity prevalence was also negatively associated with healthcare system strength, as measured by the Universal Health Coverage (UHC) index (r = -0.841). These findings align with previous reports demonstrating that countries with lower HDI tend to have higher diabetes-related mortality (22).

Economic development was moderately correlated with multiple reproductive and adolescent health outcomes. Higher GDP per capita was associated with increased low birth weight prevalence (r = 0.610) and lower adolescent fertility (r = -0.710), suggesting complex links between economic status, maternal health, and early-life reproductive patterns (23).

Finally, adolescent mortality (per 1,000 females aged 15–19) showed a moderate positive correlation with preterm birth rates (r = 0.583), indicating that adverse birth outcomes may contribute to elevated mortality in this age group (24). These findings point to the layered relationship between development, healthcare systems, and health outcomes in Caribbean populations, particularly among women and adolescents.

Principal Component Analysis Reveals Three Distinct Health and Socioeconomic Clusters Across the Caribbean

To identify underlying patterns in health and socioeconomic indicators, we performed Principal Component Analysis (PCA) across countries with sufficient data availability. Two principal components explained the majority of variance in the dataset (Figure 5B). PC1 accounted for 49.0% of the variance and reflected a gradient from stronger socioeconomic conditions and lower NCD mortality to higher

adolescent fertility and mortality. Positive contributors included adolescent fertility and adolescent mortality, while negative contributors included GDP per capita, inequality-adjusted HDI, and female NCD mortality. PC2 explained an additional 13.2% of the variance and captured variation in healthcare access and lifestyle-related factors, driven by out-of-pocket expenditure and insufficient physical activity.

Hierarchical clustering applied to PCA scores revealed three country groups with distinct profiles. The first cluster—Canada, the United States, and Cuba—had the most favorable outcomes, including the lowest adolescent mortality (1.41 per 1,000) and fertility rates (24.6 births per 1,000), and the highest life expectancy (81.9 years). The second cluster—Antigua and Barbuda, Barbados, Grenada, Saint Lucia, and Trinidad and Tobago—showed intermediate outcomes, with adolescent mortality of 1.74 per 1,000, adolescent fertility of 35.6 per 1,000, and a life expectancy of 77.7 years.

The third cluster, composed of the Bahamas, Belize, Dominican Republic, Guyana, Haiti, Jamaica, Saint Vincent and the Grenadines, and Suriname, demonstrated the greatest vulnerability. This group had the highest adolescent mortality (3.70 per 1,000), adolescent fertility (54.0 per 1,000), and the lowest life expectancy (73.7 years). It also reported the highest maternal mortality (123.1 per 100,000), with Haiti as an outlier at 128.4 per 100,000.

Together, these findings highlight how adolescent and maternal health disparities cluster with broader socioeconomic instability, reinforcing the need for region-specific interventions that address both health outcomes and structural inequality.

Caribbean and North American Countries Differ in Policy Implementation for NCD Prevention
"National action on NCD prevention varies widely across the Americas, as shown by comparisons of four key policy indicators from the 2022 PAHO ENLACE NCD scorecard. This analysis included North America (NA: United States and Canada) and 13 countries in the Caribbean (CAR: Antigua and Barbuda,

Bahamas, Barbados, Belize, Dominica, Grenada, Guyana, Jamaica, Saint Kitts and Nevis, Saint Lucia,

Saint Vincent and the Grenadines, Suriname, and Trinidad and Tobago) (Figure 6).

We began by examining the presence of national NCD guidelines, which were fully implemented in both

NA countries. In CAR, full implementation was reported in Saint Lucia, Saint Vincent and the

Grenadines, and Trinidad and Tobago. Six countries—including Belize, Barbados, Bahamas, Guyana,

Jamaica, and Suriname—had partially implemented guidelines, while the remaining four had taken no

action. This mixed picture suggests that although some Caribbean countries are progressing toward

national NCD care standards, full adoption remains limited.

We then assessed physical activity awareness policies, which were fully implemented in both NA

countries. In CAR, Barbados, Belize, and Saint Lucia had achieved full implementation, while Suriname

and Trinidad and Tobago reported partial implementation. The remaining eight countries, including

Jamaica, Bahamas, and Antigua and Barbuda, had no such policies in place. The gap in physical activity

promotion across much of the region may reflect both resource limitations and the absence of coordinated

national strategies.

Regulation of breastmilk substitutes showed minimal uptake in both regions. Neither NA country had

implemented policies aligned with the International Code of Marketing of Breast-milk Substitutes.

Among CAR countries, only Guyana had fully implemented this policy. All others, including Saint Lucia,

Jamaica, and Saint Vincent and the Grenadines, had not adopted any regulatory measures. Despite

growing global consensus on the importance of protecting breastfeeding, this remains a neglected policy

area across most of the Caribbean.

Finally, marketing restrictions to children revealed the sharpest contrast. Canada and the United States

had both fully implemented these protections. In contrast, all 13 CAR countries reported no

implementation. This includes countries like Barbados and Belize that have otherwise made progress in

other areas. The complete lack of child-targeted marketing regulation across the Caribbean raises

concerns about commercial influences on childhood diet and long-term NCD risk.

Overall, NA demonstrated strong and consistent policy implementation across the four indicators, while

CAR countries showed significant variation and persistent gaps. Even among relatively stronger

performers like Saint Lucia, Belize, and Barbados, implementation remains incomplete—particularly in

areas related to infant and child health.

Discussion

This study examined gender disparities in health outcomes across the Caribbean, focusing on how

socioeconomic conditions, healthcare access, and systemic inequalities disproportionately affect women's

health. The analysis highlighted the intersection of chronic diseases, reproductive health disparities, and

economic stressors, with a particular focus on the Bahamas due to its notable obesity rates, limited

healthcare access, and elevated mortality metrics. Building upon prior research, this study provides novel

insights into the complex relationship between gender, health outcomes, and socioeconomic factors in the

region (2, 3, 25).

Our findings revealed significant gender disparities in NCDs, including obesity and diabetes. Women in

the Bahamas exhibited higher obesity rates (54.1%) than men (38.3%), a disparity that aligns with broader

NCD trends observed across the Caribbean (2, 3). Previous research has shown that socioeconomic

inequalities contribute to adverse health outcomes in the region, including elevated risks for preterm birth

and other preventable conditions. These systemic inequities often interact with healthcare access

challenges—such as high out-of-pocket costs and limited preventive services—which delay diagnosis and

treatment for chronic diseases (26, 27). Our Pearson correlations findings identified a strong negative

correlation between diabetes mortality and IHDI ($r = -0.884$), which underscores how broader

socioeconomic development is associated with improved management of chronic diseases like diabetes.

Additionally, PCA analysis further demonstrated that adolescent health metrics distinctly segregate
Caribbean nations from higher-income countries, highlighting persistent healthcare access gaps (Figure
5B-D).

Reproductive health disparities also emerged as a defining element of gendered health inequality in the
Caribbean. Maternal mortality rates in the region (9.39 per 10,000 live births ± 5.22) were substantially
higher than those in North America, and adolescent fertility rates (34.26 per 10,000 ± 19.49) also
exceeded those observed in high-income countries ($p < 0.001$ and $p = 0.05$, respectively). These patterns
echo previous findings linking structural inequity and gender-based health disparities to weaker
reproductive health systems and limited access to prenatal care (28, 29). Our analysis further revealed a
negative correlation between adolescent fertility and GDP per capita ($r = -0.710$), reinforcing prior
evidence that economic hardship is closely tied to early childbearing and long-term reproductive risk (30,
31). These associations support the growing recognition that reproductive and chronic disease outcomes
are not separate phenomena, but rather interlinked manifestations of broader gendered and socioeconomic
inequities in health systems across the Caribbean.

Socioeconomic conditions—including high income inequality and food insecurity—continue to shape
these health outcomes. Despite having the highest GDP per capita in the region, the Bahamas remains one
of the most unequal societies, as indicated by its IHDI and GINI coefficients. This contradiction helps
explain its simultaneous burden of economic wealth and elevated obesity prevalence, which we identified
as the highest in the Caribbean. These patterns suggest that economic growth alone is insufficient to
reduce chronic disease risk without equity-focused health and social policies.

While social conditions remain central, emerging evidence also points to biological contributors. Genetic
studies have identified African ancestry-linked variants, such as ZRANB3, that impair insulin secretion
and increase diabetes risk (32), reinforcing the critical role of the insulin secretory pathway, where defects
can directly contribute to the development of diabetes (33, 34).

Similar disparities have been observed across other chronic conditions. For instance, studies show that

cataract burden in Caribbean small island developing states (SIDS) disproportionately affects

socioeconomically disadvantaged populations (35). Economic constraints often push families toward low-

cost, ultra-processed foods, a pattern that has been extensively documented in food insecurity research,

particularly in countries like Haiti (29, 36). These outcomes reinforce longstanding evidence that social

and economic inequality—not individual behavior—is the dominant force shaping population health (29,

37, 38).

The Caribbean's economic dependence on tourism further compounds these structural vulnerabilities.

While tourism boosts national GDP, it often generates low-wage, seasonal employment with limited job

security or healthcare benefits (39). This economic structure mirrors prior findings on how gender-based

and economic vulnerabilities shape health outcomes in the Caribbean (40, 41). As a result, the apparent

economic growth tied to tourism has not translated into improved or equitable health outcomes across the

region.

Recent studies of breast cancer and other chronic disease risks in the Caribbean show that economic

growth has not translated into equitable health outcomes, reinforcing how persistent social and economic

divides continue to widen regional health disparities (41). These findings reinforce the need for economic

diversification and gender-responsive health financing strategies that strengthen long-term health

resilience.

Our PCA-derived clusters offer further insight into how structural inequality manifests across the region.

Cuba's clustering alongside high-income countries like the U.S. and Canada is striking, given its much

lower GDP ($103.43 billion vs. $1.74 trillion for Canada and $21.54 trillion for the U.S. in 2019) and

ongoing economic constraints (42). Cuba's strong public health indicators may reflect its decades-long

investment in preventive care and universal health coverage, though concerns persist regarding the

sustainability of these gains and the reliability of reported data amid deepening resource limitations (43).

In contrast, countries in the red cluster—such as the Bahamas, Belize, Dominican Republic, Guyana,

Haiti, Jamaica, Saint Vincent and the Grenadines, and Suriname—exhibited consistent patterns of

socioeconomic vulnerability, under-resourced health systems, and disproportionately high gender-related

disparities in NCD outcomes (5). Meanwhile, countries in the green cluster—such as Antigua and

Barbuda, Barbados, Grenada, Saint Lucia, and Trinidad and Tobago—tended to maintain stronger health

metrics, likely supported by more stable governance structures and sustained public health investment.

These findings highlight how structural stability and policy prioritization can buffer against broader

regional inequities, while countries facing overlapping stressors may require more intensive, targeted

interventions.

The wide gaps in policy implementation observed in this study underscore the urgent need for

coordinated, equity-driven public health reform across the Caribbean. While countries in North America

have achieved broad adoption of foundational NCD prevention policies, most Caribbean countries remain

at an early stage, particularly in the areas of child-targeted marketing, breastfeeding protections, and

physical activity promotion (9). These findings align with prior work showing that NCD risk in Caribbean

youth is rising rapidly in the absence of structural policy safeguards, especially for obesity and diet-

related conditions (10).

The complete absence of marketing restrictions to children across all 13 Caribbean countries included in

this analysis is especially concerning given the region's documented challenges with ultra-processed food

exposure, school-based advertising, and limited regulatory enforcement (9). Similarly, breastmilk

substitute regulations—a critical tool for protecting infant nutrition—were implemented in only one

country, reflecting a neglected policy space despite long-standing WHO recommendations. These gaps

are not only technical oversights but manifestations of deeper systemic inequities that leave Caribbean

women and children particularly vulnerable to preventable NCDs (5, 11). Several structural constraints

likely contribute to this pattern. Past studies have highlighted barriers including inconsistent health

financing, limited legislative infrastructure, and poor integration of NCD services into primary care

systems, particularly following environmental and economic shocks (3). For small island states, the

capacity to draft, enforce, and monitor public health legislation often remains limited—further reinforcing

dependence on donor priorities and regional agencies for guidance and technical support.

Improving policy adoption will require more than regional consensus statements; it will demand clear

political commitment, cross-country collaboration, and investment in health system readiness. Models

like the PAHO ENLACE scorecard can provide useful benchmarking, but implementation support must

go beyond measurement. Prior research has shown that achieving population-level impact requires

aligning fiscal policies, regulatory tools, and public education strategies—particularly when tackling

upstream drivers of NCDs such as poverty, food insecurity, and gender inequity (28, 41).

Efforts to advance universal healthcare coverage, reduce out-of-pocket costs, and embed preventive

services into routine care are particularly promising for women's health outcomes in vulnerable settings

(30). Strengthening regional surveillance systems and enabling shared policy templates across ministries

of health could also lower the administrative burden of reform. Ultimately, closing the implementation

gap will require structural investment in public health governance that centers equity and resilience—

ensuring that NCD prevention is not only planned, but enacted.

This study has several limitations. Its cross-sectional design limits causal inference, allowing only for the

identification of associations between economic conditions, healthcare access, and health outcomes. The

analysis relied on publicly available datasets from PAHO, UNDP, and IDF due to the limited availability

of peer-reviewed data on Caribbean health metrics, which may introduce inconsistencies in data quality

across countries (14-16, 19, 40). Data imputation was used sparingly and only for the Pearson correlation

and PCA analyses to preserve sample size; while necessary, this may have introduced minor bias.

Additionally, data collection disruptions during the COVID-19 pandemic limited the availability of recent

socioeconomic indicators, particularly for smaller Caribbean states.

In conclusion, this study highlights how structural inequities in healthcare systems, economic

vulnerability, and social conditions continue to disproportionately shape women's health outcomes across

the Caribbean. Targeted interventions, stronger regional surveillance, and policy reforms that prioritize

equity will be critical to narrowing these gaps and improving long-term health outcomes for vulnerable

populations.

Competing Interests Statement

The authors declare no competing interests related to this study.

Funding Statement

This research received no external funding. The study was conducted with independent resources

provided by Dr. Cesar Barrabi.

Author Contributions

C.B. conceptualized the study, designed the methodology, collected and analyzed the data, interpreted the

results, and wrote the manuscript. A.F. contributed to data interpretation and conducted the literature

review. All authors reviewed and approved the final version of the manuscript for submission.

Data Availability Statement

This study utilized publicly available data from multiple sources. Health indicators, including NCD

prevalence and mortality rates, were obtained from the Pan American Health Organization (PAHO)

(https://opendata.paho.org/en/core-indicators/core-indicators-dashboard). Socioeconomic metrics, such as

the Inequality-Adjusted Human Development Index (IHDI) and employment rates, were sourced from the

United Nations Development Programme (UNDP) (https://hdr.undp.org/data-center/human-development-

index#/indicies/HDI). Diabetes-related data, including gestational diabetes prevalence, were retrieved

from the International Diabetes Federation (IDF) (https://diabetesatlas.org/data/en/). All datasets are

publicly accessible, and further details on data retrieval can be provided upon request.

References

1.　　Hay K, McDougal L, Percival V, Henry S, Klugman J, Wurie H, et al. Disrupting gender norms in
health systems: making the case for change. Lancet. 2019;393(10190):2535-49.
2.　　Guzman-Vilca WC, Carrillo-Larco RM. Mortality attributable to type 2 diabetes mellitus in Latin
America and the Caribbean: a comparative risk assessment analysis. BMJ Open Diabetes Res Care.
2022;10(1).
3.　　Hassan S, Nguyen M, Buchanan M, Grimshaw A, Adams OP, Hassell T, et al. Management Of
Chronic Noncommunicable Diseases After Natural Disasters In The Caribbean: A Scoping Review.
Health Aff (Millwood). 2020;39(12):2136-43.
4.　　Control CfD, Prevention. Diabetes prevalence and undiagnosed rates in the USA. 2022.
5.　　Sobers-Grannum N MM, Nielsen A, Guell C, Samuels TA, et al. . Female Gender Is a Social
Determinant of Diabetes in the Caribbean: A Systematic Review and Meta-Analysis. PLOS ONE.
2015;10(5): e0126799.
6.　　Galicia-Garcia U, Benito-Vicente A, Jebari S, Larrea-Sebal A, Siddiqi H, Uribe KB, et al.
Pathophysiology of Type 2 Diabetes Mellitus. Int J Mol Sci. 2020;21(17).
7.　　(NCD-RisC) NRFC. Worldwide trends in underweight and obesity from 1990 to 2022: a pooled
analysis of 3663 population-representative studies with 222 million children, adolescents, and adults.
Lancet. 2024;Mar 16;403(10431):1027-1050.
8.　　Wang S, Rexrode KM, Florio AA, Rich-Edwards JW, Chavarro JE. Maternal Mortality in the
United States: Trends and Opportunities for Prevention. Annu Rev Med. 2023;74:199-216.
9.　　Vega-Salas MJ MC, Nunes R, Hidalgo-Arestegui A, Curi-Quinto K, Penny ME, Cueto S,
Lovegrove JA, Sánchez A, Vimaleswaran KS. School environments and obesity: a systematic review of
interventions and policies among school-age students in Latin America and the Caribbean. Int J Obes
(Lond). 2023;47(1):5-16.
10.　　Savage MJ DN, Hennis PJ, James RM, Grazette N, Ferguson TS, McFarlane S, Armstrong H,
Cox H, Varley I. Tackling non-communicable disease risk in young adults across the Caribbean: a call to
action. BMJ Glob Health. 2025;10(2):e017010.
11.　　Rocío Sáenz GN, Ingrid Gómez-Duarte, Karol Rojas, Arachu Castro, Edson Serván-Mori.
Persistent inequities in maternal mortality in Latin America and the Caribbean, 1990-2019. Int J Equity
Health. 2019;May 10;23(1):96.
12.　　Kristy Samaroo AH, Lyronne K Olivier, Jameel Ali. Breast Cancer in the Caribbean. Cureus.
2021;13(8):e17042.
13.　　Thomas-Purcell K, Bailey A, Sealy DA, Song G, Ashing KT. Cervical Cancer Stigma Among
Caribbean Population: A Descriptive Paper. Int J Public Health. 2024;69:1606725.
14.　　International Diabetes Federation. Brussels, Belgium . IDF Diabetes Atlas (2021) 10thedn.
Brussels, Belgium Retrieved: 2/15/25 URL: http://www.diabetesatlas.org.

15. United Nations Development Programme (UNDP). (2025). Human Development Report 2025:
Human Development Index (HDI) and Inequality-adjusted Human Development Index (IHDI). Retrieved
March 8, 2025, from https://hdr.undp.org/.
16. Pan American Health Organization/World Health Organization. Core Indicators Portal. Region of
the Americas. Washington D.C. Retrieved: 02/15/2025 URL:https://opendata.paho.org/en/core-indicators
[Internet].
17. Fu Q, Hoijtink H, Moerbeek M. Sample-size determination for the Bayesian t test and Welch's
test using the approximate adjusted fractional Bayes factor. Behav Res Methods. 2021;53(1):139-52.
18. Open Source Geospatial Foundation. QGIS Geographic Information System (2024). Version 3.34.
Retrieved: 2/20/25 URL: https://qgis.org.
19. GADM. Database of Global Administrative Areas (2022) Retrieved Shapefiles: 2/20/25
URL:https://gadm.org/data.html.
20. Lee JH, Huber JC, Jr. Evaluation of Multiple Imputation with Large Proportions of Missing Data:
How Much Is Too Much? Iran J Public Health. 2021;50(7):1372-80.
21. Jake Lever MK, Naomi Altman. Principal component analysis. Nature methods. 2017;14:641-2.
22. Balooch Hasankhani M, Mirzaei H, Karamoozian A. Global trend analysis of diabetes mellitus
incidence, mortality, and mortality-to-incidence ratio from 1990 to 2019. Sci Rep. 2023;13(1):21908.
23. Grbic D, Supic ZT, Todorovic J, Nesic D, Karic S, Jurisic A, et al. Factors associated with low
birth weight in low-income populations in the Western Balkans: insights from the multiple indicator
cluster survey. Front Public Health. 2024;12:1394060.
24. Perez MJ, Chang JJ, Temming LA, Carter EB, Lopez JD, Tuuli MG, et al. Driving Factors of
Preterm Birth Risk in Adolescents. AJP Rep. 2020;10(3):e247-e52.
25. Roberts C, et al. Economic and health disparities in the Caribbean. Public Health Journal.
2018;9(2):33-40.
26. Philip McHale KF, Andy Pennington, Daniela K Schlüter, Ben Barr, David Taylor-Robinson.
How do socioeconomic inequalities and preterm birth interact to modify health and education outcomes?
A narrative systematic review. BMJ Open. 2025;15(1):e084147.
27. Reichenberger V, Corona AP, Ramos VD, Shakespeare T, Hameed S, Penn-Kekana L, et al.
Access to primary healthcare services for adults with disabilities in Latin America and the Caribbean: a
review and meta-synthesis of qualitative studies. Disabil Rehabil. 2024;46(25):6011-20.
28. Sherrie L Kelly TW, Dominic Delport, Debra ten Brink, Rowan Martin-Hughes, Caroline SE
Homer, Jennifer Butler, Olanike Adedeji, Davide De Beni, Federica Maurizio, Howard S Friedman,
Doretta Di Marco, Federico Tobar, Maria Pilar de la Corte Molina, Andre S Richards, Nick Scott Health
and economic benefits of achieving contraceptive and maternal health targets in Small Island Developing
States in the Pacific and Caribbean. BMJ Glob Health. 2023;8(2):e010018.
29. Susan Salmond CD. Time to Reflect and Take Action on Health Disparities and Health Inequities.
Orthop Nurs. 2022;41(2):64–85.
30. Alex P Sanchez-Covarrubias MJC, Priscilla Barreto-Coehlo, Cheryl Alexis, Jameel Ali,
Alexandra Diaz-Barbe, Raleigh Butler, Saida Bowe, DuVaughn Curling, Vincent DeGennaro, Leah V
Dodds, Hedda Dyer, Darron Halliday, Patricia Jeudin, Dwight Lowe, Kristy Samaroo, Gillian Wharfe,
Matthew Schlumbrecht, Isildinha M Reis, Judith Hurley, Sophia George. Reproductive Risk Factor
Patterns in Caribbean Women With Breast Cancer Across 4 Generations. JAMA Netw Open.
2024;7(10):e2438091.
31. Sarah Neal CH, Venkatraman Chandra-Mouli, Sonja Caffe, Alma Virginia Camacho. Trends in
adolescent first births in five countries in Latin America and the Caribbean: disaggregated data from
demographic and health surveys. Reprod Health. 2018;15:146.
32. Adeyemo AA, Zaghloul NA, Chen G, Doumatey AP, Leitch CC, Hostelley TL, et al. ZRANB3 is
an African-specific type 2 diabetes locus associated with beta-cell mass and insulin response. Nat
Commun. 2019;10(1):3195.
33. Barrabi C, Zhang K, Liu M, Chen X. Pancreatic beta cell ER export in health and diabetes. Front
Endocrinol (Lausanne). 2023;14:1155779.

34.   Zhu R, Li X, Xu J, Barrabi C, Kekulandara D, Woods J, et al. Defective endoplasmic reticulum
export causes proinsulin misfolding in pancreatic beta cells. Mol Cell Endocrinol. 2019;493:110470.
35.   Patel R, et al. The Impact of Sociodemographic and Socioeconomic Factors on the Burden of
Cataract in Small Island Developing States (SIDS) in the Caribbean from 1990 to 2016. Ophthalmic
Epidemiol. 2020;27(2):132-140.
36.   Rehana Rasul VR, Rodney Sufra, Lily D Yan, Inddy Joseph, Nour Mourra, Shalom Sabwa, Marie
M Deschamps, Daniel W Fitzgerald, Jean W Pape, Denis Nash, Margaret L McNairy. Extreme Food
Insecurity and Malnutrition in Haiti: Findings from a Population-Based Cohort in Port-au-Prince, Haiti.
Nutrients. 2022;14(22):4854.
37.   Sisitha Jayasinghe MAF, Andrew P Hills. Social justice equity in healthy living medicine - An
international perspective. Prog Cardiovasc Dis. 2022;71:64-68.
38.   Emily Brown Weida PP, Falguni Patel, Mariana Chilton. Financial health as a measurable social
determinant of health. PLOS ONE. 2020;15(5):e0233359.
39.   Evans KS, Brown. Socioeconomic vulnerabilities in Caribbean women. Global Health Action.
2021;13(2):38-44.
40.   World   Bank.   World   Development   Indicators   (2025).   Retrieved: 2/15/25   URL:
https://data.worldbank.org/indicator.
41.   Francis Poitier RK, Bassey Ebenso. Impact of sustained health policy and population-level
interventions on reducing the prevalence of obesity in the Caribbean region: A qualitative study from The
Bahamas. Front Public Health. 2022;10:926672.
42.   Keck CW, Reed GA. The curious case of Cuba. Am J Public Health. 2012;102(8):e13-22.
43.   Salicrup LA, Ordunez P, Engelgau MM. Hypertension control activities in Latin America and the
Caribbean: opportunities for late-stage (T4) translation research. Rev Panam Salud Publica. 2018;42:e22.

Figure Legends

Figure 1: Gender-Based Disparities in Mortality and Health Indicators Across the Caribbean All

data were obtained from PAHO using 2019 data. Welch's t-tests were performed to compare men and

women across all variables. (A) External causes mortality rate (age-adjusted per 100,000 population) and

the proportion of deaths from noncommunicable diseases (NCDs) and communicable diseases (CDs) (%)

by gender. (B) Age-adjusted mortality rates (per 100,000 population) for ischemic heart disease,

respiratory diseases, cirrhosis, and stomach cancer by gender. (C) Diabetes mortality rate (per 100,000

population) and proportional diabetes mortality (%) comparing USA/Canada and the Caribbean by

gender. (D) Prevalence of insufficient physical activity and obesity (%) by gender.

Figure 2: The Caribbean Faces Greater Socioeconomic and Health Challenges Compared to the

USA/Canada HDI and IHDI data were obtained from UNDP for 2019, while all other data were from

PAHO for 2019. Welch's t-tests were used to compare the USA/Canada (US/CAN) with 30 Caribbean

nations across all variables. (A) Human Development Index (HDI) and Inequality-adjusted HDI (IHDI)

scores comparing US/CAN and CAR regions. (B) Food insecurity prevalence (%) and inflation (%) in

US/CAN and CAR regions. (C) Homicide and suicide mortality rates (per 100,000 population) by region.

(D) Unemployment rates (%) for men and women in US/CAN and CAR regions.

Figure 3: The Caribbean Exhibits Greater Health and Economic Vulnerabilities Compared to the

USA/Canada Welch's t-tests were used to compare the USA/Canada (US/CAN) with 30 Caribbean

nations across all variables. (A) Maternal mortality ratio (per 1,000,000 live births) and infant mortality

rate (per 1,000 live births) comparing US/CAN and CAR regions. (B) Adolescent fertility rate (per 1,000

population) and adolescent mortality rate (per 100,000 population) by region. (C) Life expectancy at birth

and median age (in years) for US/CAN and CAR regions. (D) Current health expenditure as a percentage

of GDP and out-of-pocket expenditure as a percentage of current health expenditure (%) comparing

US/CAN and CAR regions.

Figure 4: Prevalence of Gestational Diabetes Mellitus (GDM) by Region, 2021

All data were obtained from the International Diabetes Federation (IDF) for the year 2021. The dataset

represents the prevalence of hyperglycemia in pregnancy (age 20-49) across different regions. The map

was created in QGIS 3.34 T using shapefiles from GADM to visualize the regional prevalence of GDM.

Darker shades indicate higher prevalence rates, highlighting disparities in gestational diabetes across the

regions. South-East Asia: 25.9%, North America and Caribbean: 20.7%, South and Central America:

15.8%, Europe: 15%, Middle East and North Africa: 14.1%, Western Pacific: 14%, Africa: 13%.

Figure 5: Correlation of Health and Demographic Variables and Distinct Regional Health Profiles

in the Caribbean Only data with at least 75% availability were included for Pearson correlation and

Principal Component Analysis (PCA). (A) illustrates the relationships between health and demographic

variables across countries with at least 75% data availability. Positive correlations are shown in red, and

negative correlations in blue, with stronger correlations indicated by more intense colors. (B) PCA

identified three distinct clusters of countries. Principal Component 1 (PC1) reflects socioeconomic status

and health outcomes, while Principal Component 2 (PC2) represents healthcare access and lifestyle-related health risks. Clusters highlight the USA, Canada, and Cuba with favorable health outcomes, intermediate Caribbean nations with moderate health and socioeconomic challenges, and more vulnerable Caribbean nations with higher adolescent mortality and fertility rates.

Figure 6: Implementation of NCD-Related Policies Across the Caribbean and North America Data are from the 2022 PAHO ENLACE interactive NCD scorecard. Each bar represents the number of countries classified by implementation status across four key policy areas: national NCD guidelines, physical activity awareness, regulation of breastmilk substitutes, and marketing restrictions to children. Green indicates full implementation, yellow partial implementation, red no implementation, and grey unknown status. Panel A shows results for North America (n = 2; United States and Canada), both of which fully implemented three of the four policies, with no action taken on breastmilk substitute regulation. Panel B displays results for the Caribbean (n = 13; Antigua and Barbuda, Bahamas, Barbados, Belize, Dominica, Grenada, Guyana, Jamaica, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, Suriname, and Trinidad and Tobago), where implementation was more variable. Most CAR countries had not implemented policies on marketing to children or breastmilk substitutes, while a subset had made progress on NCD guidelines and physical activity awareness.

Supplemental Figure Legends

Figure S1: Principal Component Analysis (PCA) and Hierarchical Clustering Selection for Caribbean Nations Only data with at least 75% availability were included for Principal Component Analysis (PCA) and clustering. (A) The scree plot displays the eigenvalues for each principal component, showing that the first two components explain the majority of variance in the dataset. (B) The elbow plot illustrates the total within-cluster sum of squares, identifying the optimal number of clusters where

additional clusters provide minimal variance reduction. These analyses were conducted on imputed data

to extract key socio-economic and health patterns across 30 Caribbean nations.

Figure S2: Hierarchical Clustering of Caribbean Nations Based on Socioeconomic and Health

Indicators Only data with at least 75% availability were included for hierarchical clustering. (A) The

dendrogram illustrates Ward's hierarchical clustering applied to PCA scores, grouping Caribbean nations

based on shared socio-economic and health characteristics. The y-axis represents Euclidean distance, with

shorter distances indicating greater similarity. (B) The clustering analysis identified three primary country

groups: one comprising nations with favorable health and economic indicators, another with intermediate

Caribbean nations facing moderate health and socioeconomic challenges, and a third with more

vulnerable nations experiencing higher adolescent mortality and fertility rates.

HYPERGLYCAEMIA IN PREGNANCY (20-49 Y)
PREVALENCE OF GESTATIONAL DIABETES MELLITUS
(GDM), % FROM IDF

Legend

Prevalence of
Hyperglycemia in
Pregnancy (20-49y).

13.0 - 14.0	15.0 - 16.0
14.0 - 15.0	16.0 - 17.0
	17.0 - 18.0
	18.0 - 19.0
19.0 - 20.0	23.0 - 24.0
20.0 - 21.0	24.0 - 25.0
21.0 - 22.0	25.0 - 26.0
22.0 - 23.0	

0   250   500 km

A

B

A

B

A

B

A

Hierarchical Clustering Dendrogram