Caribbean Women Face Higher Obesity and Diabetes Amid Socioeconomic Struggles – A Cross-

Sectional Study

Authors: Cesar Barrabi, Amanda Fowler

Affiliation: Western Atlantic University School of Medicine

Abstract

This cross-sectional study examined gender disparities in health outcomes across 30 Caribbean nations,

focusing on the intersection of non-communicable diseases, reproductive health, and socioeconomic

factors. Using publicly available data from the Pan American Health Organization, United Nations

Development Programme, and International Diabetes Federation, we first identified significantly higher

rates of obesity (43.88% vs. 29.77%) and a greater proportion of diabetes-related deaths among women

(13.21% vs. 10.17%) in the Caribbean compared to men. We next aimed to assess how the Caribbean

compared to the Global North, selecting the U.S. and Canada as representative high-income countries

within the same geographic region. Our analysis revealed substantially higher maternal mortality (1.41 vs.

9.39 per 10,000 live births) and infant mortality rates (5 vs. 11.58 per 1,000 live births) in the Caribbean

than in the U.S. and Canada. The Caribbean also had significantly lower Inequality-adjusted Human

Development Index scores (0.839 vs. 0.556), along with higher rates of food insecurity (8.15% vs.

41.71%), female unemployment (4.45% vs. 10.67%), and other socioeconomic challenges, highlighting

systemic barriers that disproportionately affect women. These findings demonstrate how intertwined

medical and socioeconomic factors potentially promote gender and regional health disparities in the

Caribbean, emphasizing the critical need for policies that not only improve healthcare access but also

address broader social and economic inequities to promote lasting health equity.

Introduction

Gender health disparities are a well-documented global issue, with women often facing worse outcomes

in chronic diseases, mental health, and access to healthcare (1). In the Caribbean, these disparities are

particularly stark, with women facing higher rates of obesity and diabetes and poorer reproductive health

outcomes, as demonstrated in regional mortality assessments and global risk analyses (2, 3). While high-

income countries like the U.S. and Canada also struggle with NCDs (4), Caribbean nations such as the

Bahamas and Barbados experience these challenges within more fragile healthcare systems and greater

socioeconomic inequalities (5). Although lifestyle factors like poor nutrition and inactivity are widely

recognized contributors (6), the deeper socio-economic influences on NCDs among Caribbean women

remain underexplored.

Despite its relative wealth, the Bahamas exhibits high obesity rates and diabetes-related mortality,

consistent with global trends showing disproportionate NCD burdens in middle-income nations (7).

Caribbean women are 60% more likely to have diabetes and twice as likely to be obese as men,

underscoring gendered health disparities [4]. Recent studies indicate that obesity rates among Bahamian

women continue to rise, contributing to downstream chronic disease burdens including diabetes and

kidney disease (8, 9). Targeted policy interventions are needed to expand healthcare access, address food

insecurity, and reduce socioeconomic inequality, particularly among young adults and women (10). While

the Global North offers extensive literature on NCDs, research specifically examining these complex

relationships in Caribbean women is scarce.

Beyond chronic diseases, reproductive health poses additional risks. Caribbean women face higher

maternal mortality rates, early menstruation, and late pregnancies, which contribute to breast cancer risks

and poor reproductive outcomes (11, 12). The stigma around conditions like cervical cancer further limits

healthcare access, creating a cycle of late diagnoses and higher mortality(13). These observed disparities

suggest that comprehensive approaches integrating medical treatment and addressing socioeconomic

barriers may be beneficial, consistent with findings from prior studies.

This study aims to fill these research gaps by examining gender disparities in NCDs and reproductive

health across 30 Caribbean nations compared to the U.S. and Canada. By focusing on the intersection of

healthcare access, socio-economic conditions, and gender-specific health challenges, this research offers

critical insights into the underlying factors driving health inequities in Caribbean women.

Materials & methods

Ethical Considerations

This study used publicly available, aggregated data from international organizations and did not involve

human subjects. Therefore, formal ethical approval was not required.

Study Design and Data Collection

This cross-sectional study analyzed gender disparities in health outcomes across 30 Caribbean nations

using publicly available data from the Pan American Health Organization (PAHO), United Nations

Development Programme (UNDP), and International Diabetes Federation (IDF) (14-16). The analysis

included metrics on non-communicable diseases (NCDs), reproductive health outcomes, and socio-

economic indicators, comparing these to benchmarks from the U.S. and Canada. The dataset focused on

2019 data, as this was the most recent year with complete health/economic health information. These data

sources were selected due to their comprehensive coverage, credibility, and the limited availability of

peer-reviewed literature addressing health outcomes specifically across Caribbean nations.

The primary outcomes assessed were obesity and diabetes prevalence, maternal and adolescent health

metrics, and socio-economic disparities, including Human Development Index (HDI) scores, food

insecurity, and unemployment rates. Additional variables related to lifestyle factors, healthcare access,

and reproductive health were included to provide a broader context to gender-specific health outcomes in

the Caribbean.

Data Analysis

All data processing and visualization, including scatterplots and trend lines, were conducted using

Microsoft Excel. Pearson correlation, Principal Component Analysis (PCA), Bootstrapping, and

hierarchical clustering were implemented in Python.

For gender-wise comparisons, Student's t-tests were used to compare male-to-female differences within

Caribbean nations for each metric. Welch's t-tests were used for country-level comparisons, grouping the

United States and Canada as one group (Global North) and the 30 Caribbean nations as the other group

(Global South), with a significance cutoff of $p < 0.05$ (17). This test was chosen due to account for

unequal sample sizes. Data for gestational diabetes rates for the Caribbean regions were retrieved from

the International Diabetes Federation (IDF, 2021), using regional data. A Geographic Information System

(GIS) map of the Caribbean was generated in QGIS 3.34 T using GADM shapefiles, with mapping

performed by Fiver professional Ayesha Suraweera (18, 19). No statistics were performed in QGIS, this is

simply reporting values obtained from IDF.

Data imputation was performed exclusively for the Pearson correlation and Principal Component Analysis

(PCA) analyses. Countries with less than 75% of data for each variable were removed, and the remaining

missing data was imputed using Predictive Mean Matching within the Multiple Imputation by Chained

Equations framework. This approach maintains the original data distribution by matching predicted

values to the closest observed values, ensuring realistic imputation. After data imputation, Pearson

correlation coefficients were calculated to examine associations between gestational diabetes,

malnutrition rates, and socio-economic indicators (20).

PCA was performed using the imputed data to reduce the dimensionality of the dataset with socio-

economic and health indicators across the 30 countries (21). Variables with high collinearity were

removed based on Pearson correlations before PCA, and additional variables were hand-selected based on

relevance to women's health metrics, socio-economic indicators, and other health-related metrics. A scree

plot confirmed that the first two principal components explained the majority of the variance. Ward's

hierarchical clustering was applied to the PCA scores to group countries based on their socio-economic

and health profiles, while the elbow method was applied to select the number of clusters. Bootstrapping

was performed with 1,000 iterations to assess the stability of the PCA results, achieving a 95% confidence

interval.

<u>Results</u>

We first examined 2019 PAHO metrics across 30 Caribbean nations to compare gender-specific health

outcomes. For clarity, all metrics are reported per 100,000 population unless specified. External mortality

rates were significantly higher in men (110.68) compared to women (31.01; $p < 0.001$). In contrast, non-

communicable disease (NCD) mortality rates were higher in women (83.18) than men (75.33; $p < 0.01$),

while communicable disease mortality showed no significant gender difference.

Since women had higher overall NCD mortality rates, we analyzed specific NCD categories to determine

which conditions contributed most to this disparity. Surprisingly, mortality rates for ischemic heart

disease, respiratory diseases, cirrhosis, and stomach cancer were all significantly higher in men than

women (all $p < 0.05$).

To identify the specific conditions contributing to higher mortality in women, we further examined

diabetes-related mortality. The proportion of diabetes-related mortality was higher in women (13.21%)

compared to men (10.17%; $p < 0.001$). Regionally, the Caribbean (men and women, metric from IDF)

exhibited a substantially higher proportion of diabetes-related mortality (62.22%) compared to the

USA/Canada (11.65%; $p < 0.001$), underscoring the significant impact of diabetes in the Caribbean.

To explore potential contributors to higher diabetes-related mortality in women, we analyzed lifestyle-related metrics such as physical activity and obesity rates. Women in the Caribbean had significantly higher rates of self-reported insufficient physical activity (67.79%) compared to men (53.47%; $p < 0.001$). Obesity rates also showed a pronounced gender gap, with 43.88% of women being obese compared to 29.77% of men ($p < 0.001$). These findings suggest that lifestyle factors are associated with gender disparities in diabetes-related mortality.

Overall, these findings highlight significant gender differences in health outcomes in the Caribbean. While women had higher rates of non-communicable diseases, obesity, and insufficient physical activity, men exhibited higher mortality rates from specific causes, including ischemic heart disease, respiratory diseases, cirrhosis, and stomach cancer.

The Caribbean Faces Greater Socioeconomic and Health Challenges Compared to the USA/Canada

To build on the observed gender disparities in health outcomes, we next examined socioeconomic and health challenges in the Caribbean compared to the USA/Canada. The analysis revealed that the Caribbean had significantly lower HDI and Inequality-Adjusted HDI (IHDI) scores, reflecting greater socioeconomic challenges and inequalities. HDI was lower in the Caribbean (0.76) compared to the USA/Canada (0.93; $p < 0.001$), and the IHDI followed a similar trend (0.56 vs. 0.84; $p < 0.01$).

To assess the broader impact of these socioeconomic disparities, we analyzed resource access and economic stability. Food insecurity was markedly higher in the Caribbean (41.71%) compared to the USA/Canada (8.15%; $p < 0.001$), while inflation rates did not differ significantly between regions. Interestingly, Haiti (18.7), Aruba (4.3) and Barbados (4.1) had the highest rates of inflation. This finding suggests that despite similar economic inflation, the Caribbean continues to face substantial challenges in accessing resources, which could contribute to health disparities.

Recognizing the well-documented issues of crime and violence in the Caribbean (22), we also examined

mortality rates related to external causes. Homicide mortality rates were significantly higher in the

Caribbean (7.31) compared to the USA/Canada (1.55; $p < 0.01$). Although suicide mortality rates

appeared higher in the USA/Canada (6.1) than in the Caribbean (3.87), this difference was not statistically

significant. Notably, high rates in Guyana (17) and Suriname (11.8) influenced the Caribbean average,

highlighting the variability within the region.

To further investigate socioeconomic factors influencing health outcomes, we analyzed unemployment

rates for potential gender-specific challenges. Women's unemployment was significantly higher in the

Caribbean (10.67%) compared to the USA/Canada (4.45%; $p < 0.01$), while men's unemployment rates

showed no significant regional difference. These results emphasize disproportionate socioeconomic

challenges faced by Caribbean women, a concern given that previous meta-analyses have shown

unemployment to be associated with increased risk of diabetes and related health issues (23).

Overall, these findings indicate that the Caribbean's socioeconomic environment—characterized by lower

human development, higher food insecurity, and gender-specific unemployment disparities—may play a

critical role in shaping health outcomes, particularly among women.

Reproductive Health Disparities Highlight Broader Economic and Health Vulnerabilities in the

Caribbean

To build on our findings of socioeconomic and mortality disparities, we examined reproductive and health

vulnerabilities in the Caribbean compared to the USA/Canada. Maternal mortality rates were significantly

higher in the Caribbean (9.39 per 10,000 births) compared to the USA/Canada (1.41; $p < 0.001$). Infant

mortality rates were also more than double in the Caribbean (11.58 per thousand) compared to the

USA/Canada (5; $p < 0.001$).

Since maternal and infant health are closely linked to broader health outcomes, we also assessed

adolescent health metrics to identify potential early-life contributors to gender disparities in health.

Adolescent fertility rates per 10,000 women in the Caribbean (34.26) is trending toward significance

compared to the USA/Canada (11.3; p = 0.05006). Adolescent mortality rates were significantly higher in

the Caribbean (26.38 per 10,000) than in the USA/Canada (13.7; p < 0.01).

To assess potential broader health and economic impacts, we analyzed life expectancy, median age, and

healthcare spending. While no significant differences were observed in these metrics, out-of-pocket

healthcare expenditures were significantly higher in the Caribbean (32.5% of Current health expenditure)

compared to the USA/Canada (13.4%; p < 0.01). This disparity suggests a greater financial burden on

individuals in the region, potentially limiting access to healthcare services.

We further explored reproductive health disparities by analyzing the prevalence of hyperglycemia in

pregnancy, including gestational diabetes mellitus (GDM). The Caribbean and North America reported a

GDM prevalence of 20.7%, following South-East Asia (25.9%) and ahead of South and Central America

(15.8%). These findings emphasize the need for targeted strategies to address reproductive and maternal

health challenges, particularly in regions with high GDM prevalence and broader reproductive health

disparities.

Overall, these results indicate that the Caribbean faces significant reproductive and economic challenges,

with particularly high maternal and adolescent health vulnerabilities and financial barriers to healthcare.

These issues may contribute to the persistence of health disparities and hinder progress toward equitable

healthcare outcomes.

Diabetes and Obesity Outcomes Strongly Correlate with Socioeconomic Development and

Healthcare Access in Caribbean Populations

To further understand the relationships between health and demographic variables, we performed Pearson

correlation analysis, visualized in a heatmap (Figure 5A). These analyses included only countries with at

least 75% data availability, with missing data imputed specifically for these analyses using the Multiple

Imputation by Chained Equations framework. The heatmap illustrates the strength of correlations, with

darker colors indicating stronger associations. A detailed correlation table is available in the

supplementary materials.

Key findings highlight the complex interplay between socioeconomic factors, healthcare systems, and

women's health outcomes. Higher human development appears to be associated with better health

outcomes, as evidenced by the strong negative correlation between diabetes mortality (age-adjusted) and

the IHDI ($r = -0.884$). Similarly, stronger healthcare systems may help mitigate obesity rates, with a

negative correlation observed between adult obesity prevalence and the Universal Health Coverage

(UHC) index ($r = -0.841$). These findings are supported by previous literature that found countries with

lower HDI values had higher age-standardized death rates due to type 2 diabetes mellitus, indicating a

negative correlation between socioeconomic development and diabetes mortality (24).

We next examined the relationship between economic factors and birth complications. A moderate

positive correlation was found between GDP per capita (PPP-adjusted) and low birth weight prevalence (r

$= 0.610$), suggesting that economic factors may play a role in maternal health outcomes. Furthermore,

higher GDP per capita was moderately associated with lower adolescent fertility rates ($r = -0.710$),

indicating that economic stability may contribute to delayed childbirth among adolescents (25).

The analysis also revealed significant insights into adolescent health. The adolescent mortality rate (per

1,000 females aged 15-19) showed a moderate positive correlation with preterm birth rates ($r = 0.583$),

suggesting that adverse early-life health conditions could contribute to higher mortality in this age group

(26).

To understand how countries segregate based on socioeconomic, NCD, and reproductive health metrics,

we conducted Principal Component Analysis (PCA), which identified two primary components

explaining the variance in our dataset (Figure 5B). Principal Component 1 (PC1) explained 49.0% of the

variance and represented a gradient from higher socioeconomic status and health outcomes to higher

adolescent mortality and fertility rates. It was defined by high positive contributions from adolescent

mortality and fertility rates and strong negative contributions from deaths from noncommunicable

diseases (female), GDP per capita, and the inequality-adjusted HDI. Principal Component 2 (PC2)

explained 13.2% of the variance and was influenced by out-of-pocket healthcare expenditure, insufficient

physical activity, and a mix of health and socioeconomic metrics, representing variability in healthcare

access and lifestyle-related health factors. Hierarchical clustering based on PCA scores identified three

distinct clusters with specific health and socioeconomic profiles.

The first cluster (Canada, United States, Cuba) showed favorable health outcomes and stronger

socioeconomic conditions, with the lowest adolescent mortality (1.41 per 1,000) and fertility rates (24.6

births per 1,000 women aged 15-19) and the highest life expectancy (81.9 years). The second cluster

(Antigua and Barbuda, Barbados, Grenada, Saint Lucia, Trinidad and Tobago) exhibited intermediate

socioeconomic and health challenges, with moderate adolescent mortality (1.74 per 1,000) and fertility

rates (35.6 births per 1,000), and a life expectancy of 77.7 years. The third cluster (Bahamas, Belize,

Dominican Republic, Guyana, Haiti, Jamaica, Saint Vincent and the Grenadines, Suriname) faced more

severe socioeconomic challenges, with higher adolescent mortality (3.70 per 1,000) and fertility rates

(54.0 births per 1,000) and the lowest life expectancy (73.7 years). This cluster also exhibited the highest

maternal mortality ratio (123.1 per 100,000 live births), with Haiti as an outlier, showing extreme

vulnerabilities including a maternal mortality ratio of 128.4 per 100,000. These clusters demonstrate how

socioeconomic and adolescent health metrics intersect, emphasizing that women's health disparities in the

Caribbean are deeply embedded within broader regional vulnerabilities.

Discussion

This study examined gender disparities in health outcomes across the Caribbean, focusing on how socioeconomic conditions, healthcare access, and systemic inequalities disproportionately affect women's health. The analysis highlighted the intersection of chronic diseases, reproductive health disparities, and economic stressors, with a particular focus on the Bahamas due to its notable obesity rates, limited healthcare access, and elevated mortality metrics. Building upon prior research, this study provides novel insights into the complex relationship between gender, health outcomes, and socioeconomic factors in the region (2, 3, 27).

Our findings revealed significant gender disparities in non-communicable diseases (NCDs), including obesity and diabetes. Women in the Bahamas exhibited higher obesity rates (54.1%) than men (38.3%), a disparity that aligns with broader NCD trends observed across the Caribbean (2, 3). Previous research has shown that socioeconomic inequalities contribute to adverse health outcomes in the region, including elevated risks for preterm birth and other preventable conditions. These systemic inequities often interact with healthcare access challenges—such as high out-of-pocket costs and limited preventive services— which delay diagnosis and treatment for chronic diseases (28, 29). Our Pearson correlations findings identified a strong negative correlation between diabetes mortality and IHDI (r = -0.884), which underscores how broader socioeconomic development is associated with improved management of chronic diseases like diabetes. Additionally, PCA analysis further demonstrated that adolescent health metrics distinctly segregate Caribbean nations from higher-income countries, highlighting persistent healthcare access gaps (Figure 5B-D).

Reproductive health disparities also emerged as a critical issue. The Caribbean exhibited higher maternal mortality (9.39 per 10,000 lb ± 5.22) and adolescent fertility rates (34.26 per 10,000 ± 19.49) compared to the USA/Canada, with rates significantly higher (p < 0.001 and p = 0.05, respectively). Previous studies have shown that gender-based health disparities are influenced by policy-level inequities, and that reproductive health outcomes are often intertwined with broader chronic disease patterns (30, 31). Our

finding of a strong negative correlation between adolescent fertility and GDP per capita (r = -0.710) aligns

with previous research linking early reproductive patterns to long-term chronic disease risks and

highlighting the role of economic inequality in shaping health outcomes (32, 33).

Socioeconomic factors, such as income inequality and food insecurity, underpin many of these health

inequities. Our previous study found that despite its high GDP per capita, the Bahamas has stark income

inequality, evidenced by high IHDI and GINI coefficients. Additionally, the Bahamas had the highest

obesity prevalence in the Caribbean, underscoring the country's heightened vulnerability to metabolic

disease. While socioeconomic factors play a major role, emerging research suggests individuals of

African descent may also carry genetic variants that impair insulin secretion. One such variant, identified

in the gene *ZRANB3*, has been linked to impaired insulin secretion (34), reinforcing the critical role of the

insulin secretory pathway, where defects can directly contribute to the development of diabetes (35, 36).

Similar patterns of socioeconomic vulnerability have been observed in other chronic conditions; for

example, cataract burden in Caribbean SIDS has been shown to disproportionately affect populations

facing systemic disadvantage (37). Economic constraints often force households toward inexpensive,

ultra-processed foods, a pattern reflected in regional studies of food insecurity and malnutrition, such as

those conducted in Haiti (31, 38). These conditions are shaped by broader systemic inequities, with

literature emphasizing that social and economic disadvantage—not individual behavior—primarily drives

health outcomes (31, 39, 40).

The Caribbean's dependence on tourism as an economic driver exacerbates these disparities, reflecting

broader patterns of systemic inequality and financial vulnerability identified in prior research (41, 42).

While tourism boosts GDP, it offers primarily seasonal, low-wage employment without robust healthcare

benefits, a structure that mirrors prior findings on how economic and gender-based vulnerabilities shape

health outcomes in the Caribbean (33, 43). Recent analyses of breast cancer risk factors in the Caribbean

suggest that economic growth has not translated into equitable health outcomes, highlighting how

underlying social and economic divides may widen health disparities (37).

The PCA-derived clusters highlight significant regional disparities in health and socioeconomic metrics

among Caribbean nations, offering insights into how broader systemic factors shape women's health

outcomes. Cuba's placement alongside high-income nations like the U.S. and Canada is surprising given

its much lower GDP ($103.43 billion vs. $1.74 trillion for Canada and $21.54 trillion for the U.S. in

2019) and ongoing political and economic challenges (44). While Cuba's strong emphasis on preventive

healthcare and universal coverage may partially explain its favorable health indicators, questions remain

about the sustainability of these outcomes amidst economic hardships and potential discrepancies in

reported data (45). The green cluster (Antigua and Barbuda, Barbados, Grenada, Saint Lucia, Trinidad

and Tobago) aligns with literature showing that these nations, despite moderate resources, maintain

relatively strong health metrics through targeted public health programs and comparatively stable

economies (46). Conversely, the red cluster (Bahamas, Belize, Dominican Republic, Guyana, Haiti,

Jamaica, Saint Vincent and the Grenadines, Suriname) is characterized by more severe socioeconomic

challenges and gender-related disparities in chronic disease, consistent with literature showing that

women in the Caribbean face disproportionate NCD burdens linked to systemic inequities (5). These

findings illustrate how stronger economies and healthcare systems can mitigate disparities, while

vulnerable nations may require targeted interventions to address entrenched inequalities effectively.

Policy interventions targeting healthcare access and socioeconomic inequality could potentially mitigate

these disparities, as suggested by previous research on preterm birth outcomes and access barriers in the

Caribbean and Latin America (28, 29). Strategies such as universal healthcare coverage, reduced out-of-

pocket costs, and targeted economic support programs could significantly improve women's health

outcomes in vulnerable Caribbean populations (41). Implementing universal healthcare coverage could

address health inequities, as suggested by the observed correlation between stronger healthcare systems

(UHC index) and lower obesity rates ($r = -0.841$), reinforcing the potential impact of reducing out-of-

pocket costs and enhancing access to preventive care. Addressing extreme food insecurity through

broader economic and public health strategies is also essential, particularly in countries where

malnutrition reflects deeper structural instability (38).

This study has several limitations. The cross-sectional design inherently restricts the ability to draw causal

inferences between economic and healthcare factors and health outcomes, allowing only for the

identification of associations. The reliance on publicly available datasets from PAHO, UNDP, and IDF

was necessary due to the scarcity of peer-reviewed literature on Caribbean health outcomes (47).

However, these aggregated sources may introduce variability in data quality and collection methods

across different countries. Data imputation was applied exclusively to the Pearson correlation and PCA

analyses, representing a small proportion of the dataset. This approach maintained data integrity by

preserving original data distributions, though it may have introduced minor biases in these specific

analyses. Lastly, the COVID-19 pandemic disrupted data collection, limiting the availability of recent

socioeconomic indicators, particularly in smaller Caribbean nations (42). These limitations should be

considered when interpreting the findings and their broader applicability.

In conclusion, this study highlights how systemic inequities in economic policies, healthcare access, and

social determinants of health disproportionately affect women's health outcomes in the Caribbean.

Addressing these disparities through targeted interventions and systemic changes is critical to advancing

health equity and improving long-term outcomes for vulnerable populations.

Competing Interests Statement

The authors declare no competing interests related to this study.

Funding Statement

This research received no external funding. The study was conducted with independent resources

provided by Dr. Cesar Barrabi.

Author Contributions

335 C.B. conceptualized the study, designed the methodology, collected and analyzed the data, interpreted the

336 results, and wrote the manuscript. A.F. contributed to data interpretation and conducted the literature

337 review. All authors reviewed and approved the final version of the manuscript for submission.

Data Availability Statement

This study utilized publicly available data from multiple sources. Health indicators, including non-

communicable disease prevalence and mortality rates, were obtained from the Pan American Health

Organization (PAHO) (https://opendata.paho.org/en/core-indicators/core-indicators-dashboard).

Socioeconomic metrics, such as the Inequality-Adjusted Human Development Index (IHDI) and

employment rates, were sourced from the United Nations Development Programme (UNDP)

(https://hdr.undp.org/data-center/human-development-index#/indicies/HDI). Diabetes-related data,

including gestational diabetes prevalence, were retrieved from the International Diabetes Federation (IDF)

(https://diabetesatlas.org/data/en/). All datasets are publicly accessible, and further details on data retrieval

can be provided upon request.

References

1.   Hay K, McDougal L, Percival V, Henry S, Klugman J, Wurie H, et al. Disrupting gender
norms in health systems: making the case for change. Lancet. 2019;393(10190):2535-49.
2.   Guzman-Vilca WC, Carrillo-Larco RM. Mortality attributable to type 2 diabetes mellitus in
Latin America and the Caribbean: a comparative risk assessment analysis. BMJ Open Diabetes
Res Care. 2022;10(1).

3. Hassan S, Nguyen M, Buchanan M, Grimshaw A, Adams OP, Hassell T, et al.
Management Of Chronic Noncommunicable Diseases After Natural Disasters In The Caribbean:
A Scoping Review. Health Aff (Millwood). 2020;39(12):2136-43.
4. Control CfD, Prevention. Diabetes prevalence and undiagnosed rates in the USA. 2022.
5. Sobers-Grannum N MM, Nielsen A, Guell C, Samuels TA, et al. . Female Gender Is a
Social Determinant of Diabetes in the Caribbean: A Systematic Review and Meta-Analysis.
PLOS ONE. 2015;10(5): e0126799.
6. Galicia-Garcia U, Benito-Vicente A, Jebari S, Larrea-Sebal A, Siddiqi H, Uribe KB, et al.
Pathophysiology of Type 2 Diabetes Mellitus. Int J Mol Sci. 2020;21(17).
7. (NCD-RisC) NRFC. Worldwide trends in underweight and obesity from 1990 to 2022: a
pooled analysis of 3663 population-representative studies with 222 million children,
adolescents, and adults. Lancet. 2024;Mar 16;403(10431):1027-1050.
8. Wang S, Rexrode KM, Florio AA, Rich-Edwards JW, Chavarro JE. Maternal Mortality in
the United States: Trends and Opportunities for Prevention. Annu Rev Med. 2023;74:199-216.
9. Vega-Salas MJ MC, Nunes R, Hidalgo-Arestegui A, Curi-Quinto K, Penny ME, Cueto S,
Lovegrove JA, Sánchez A, Vimaleswaran KS. School environments and obesity: a systematic
review of interventions and policies among school-age students in Latin America and the
Caribbean. Int J Obes (Lond). 2023;47(1):5-16.
10. Savage MJ DN, Hennis PJ, James RM, Grazette N, Ferguson TS, McFarlane S,
Armstrong H, Cox H, Varley I. Tackling non-communicable disease risk in young adults across
the Caribbean: a call to action. BMJ Glob Health. 2025;10(2):e017010.
11. Rocío Sáenz GN, Ingrid Gómez-Duarte, Karol Rojas, Arachu Castro, Edson Serván-
Mori. Persistent inequities in maternal mortality in Latin America and the Caribbean, 1990-2019.
Int J Equity Health. 2019;May 10;23(1):96.
12. Kristy Samaroo AH, Lyronne K Olivier, Jameel Ali. Breast Cancer in the Caribbean.
Cureus. 2021;13(8):e17042.
13. Thomas-Purcell K, Bailey A, Sealy DA, Song G, Ashing KT. Cervical Cancer Stigma
Among Caribbean Population: A Descriptive Paper. Int J Public Health. 2024;69:1606725.
14. International Diabetes Federation. Brussels, Belgium . IDF Diabetes Atlas (2021)
10thedn. Brussels, Belgium Retrieved: 2/15/25 URL: http://www.diabetesatlas.org.
15. United Nations Development Programme (UNDP). (2025). Human Development Report
2025: Human Development Index (HDI) and Inequality-adjusted Human Development Index
(IHDI). Retrieved March 8, 2025, from https://hdr.undp.org/.
16. Pan American Health Organization/World Health Organization. Core Indicators Portal.
Region of the Americas. Washington D.C. Retrieved: 02/15/2025
URL:https://opendata.paho.org/en/core-indicators [Internet].
17. Fu Q, Hoijtink H, Moerbeek M. Sample-size determination for the Bayesian t test and
Welch's test using the approximate adjusted fractional Bayes factor. Behav Res Methods.
2021;53(1):139-52.
18. Open Source Geospatial Foundation. QGIS Geographic Information System (2024).
Version 3.34. Retrieved: 2/20/25 URL: https://qgis.org.
19. GADM. Database of Global Administrative Areas (2022) Retrieved Shapefiles: 2/20/25
URL:https://gadm.org/data.html.
20. Lee JH, Huber JC, Jr. Evaluation of Multiple Imputation with Large Proportions of
Missing Data: How Much Is Too Much? Iran J Public Health. 2021;50(7):1372-80.
21. Jake Lever MK, Naomi Altman. Principal component analysis. Nature methods.
2017;14:641-2.
22. Maharaj S, Seepersad R, Sooknanan J, Anderson S, Franco D, Ramdass A, et al. A
public health approach to reducing violence within the CARICOM region. Front Public Health.
2024;12:1344387.

23. Varanka-Ruuska T, Rautio N, Lehtiniemi H, Miettunen J, Keinanen-Kiukaanniemi S,
Sebert S, et al. The association of unemployment with glucose metabolism: a systematic review
and meta-analysis. Int J Public Health. 2018;63(4):435-46.
24. Balooch Hasankhani M, Mirzaei H, Karamoozian A. Global trend analysis of diabetes
mellitus incidence, mortality, and mortality-to-incidence ratio from 1990 to 2019. Sci Rep.
2023;13(1):21908.
25. Grbic D, Supic ZT, Todorovic J, Nesic D, Karic S, Jurisic A, et al. Factors associated with
low birth weight in low-income populations in the Western Balkans: insights from the multiple
indicator cluster survey. Front Public Health. 2024;12:1394060.
26. Perez MJ, Chang JJ, Temming LA, Carter EB, Lopez JD, Tuuli MG, et al. Driving Factors
of Preterm Birth Risk in Adolescents. AJP Rep. 2020;10(3):e247-e52.
27. Roberts C, et al. Economic and health disparities in the Caribbean. Public Health
Journal. 2018;9(2):33-40.
28. Philip McHale KF, Andy Pennington, Daniela K Schlüter, Ben Barr, David Taylor-
Robinson. How do socioeconomic inequalities and preterm birth interact to modify health and
education outcomes? A narrative systematic review. BMJ Open. 2025;15(1):e084147.
29. Reichenberger V, Corona AP, Ramos VD, Shakespeare T, Hameed S, Penn-Kekana L,
et al. Access to primary healthcare services for adults with disabilities in Latin America and the
Caribbean: a review and meta-synthesis of qualitative studies. Disabil Rehabil.
2024;46(25):6011-20.
30. Sherrie L Kelly TW, Dominic Delport, Debra ten Brink, Rowan Martin-Hughes, Caroline
SE Homer, Jennifer Butler, Olanike Adedeji, Davide De Beni, Federica Maurizio, Howard S
Friedman, Doretta Di Marco, Federico Tobar, Maria Pilar de la Corte Molina, Andre S Richards,
Nick Scott Health and economic benefits of achieving contraceptive and maternal health targets
in Small Island Developing States in the Pacific and Caribbean. BMJ Glob Health.
2023;8(2):e010018.
31. Susan Salmond CD. Time to Reflect and Take Action on Health Disparities and Health
Inequities. Orthop Nurs. 2022;41(2):64–85.
32. Alex P Sanchez-Covarrubias MJC, Priscilla Barreto-Coehlo, Cheryl Alexis, Jameel Ali,
Alexandra Diaz-Barbe, Raleigh Butler, Saida Bowe, DuVaughn Curling, Vincent DeGennaro,
Leah V Dodds, Hedda Dyer, Darron Halliday, Patricia Jeudin, Dwight Lowe, Kristy Samaroo,
Gillian Wharfe, Matthew Schlumbrecht, Isildinha M Reis, Judith Hurley, Sophia George.
Reproductive Risk Factor Patterns in Caribbean Women With Breast Cancer Across 4
Generations. JAMA Netw Open. 2024;7(10):e2438091.
33. Sarah Neal CH, Venkatraman Chandra-Mouli, Sonja Caffe, Alma Virginia Camacho.
Trends in adolescent first births in five countries in Latin America and the Caribbean:
disaggregated data from demographic and health surveys. Reprod Health. 2018;15:146.
34. Adeyemo AA, Zaghloul NA, Chen G, Doumatey AP, Leitch CC, Hostelley TL, et al.
ZRANB3 is an African-specific type 2 diabetes locus associated with beta-cell mass and insulin
response. Nat Commun. 2019;10(1):3195.
35. Barrabi C, Zhang K, Liu M, Chen X. Pancreatic beta cell ER export in health and
diabetes. Front Endocrinol (Lausanne). 2023;14:1155779.
36. Zhu R, Li X, Xu J, Barrabi C, Kekulandara D, Woods J, et al. Defective endoplasmic
reticulum export causes proinsulin misfolding in pancreatic beta cells. Mol Cell Endocrinol.
2019;493:110470.
37. Patel R, et al. The Impact of Sociodemographic and Socioeconomic Factors on the
Burden of Cataract in Small Island Developing States (SIDS) in the Caribbean from 1990 to
2016. Ophthalmic Epidemiol. 2020;27(2):132-140.
38. Rehana Rasul VR, Rodney Sufra, Lily D Yan, Inddy Joseph, Nour Mourra, Shalom
Sabwa, Marie M Deschamps, Daniel W Fitzgerald, Jean W Pape, Denis Nash, Margaret L

McNairy. Extreme Food Insecurity and Malnutrition in Haiti: Findings from a Population-Based Cohort in Port-au-Prince, Haiti. Nutrients. 2022;14(22):4854.

39.   Sisitha Jayasinghe MAF, Andrew P Hills. Social justice equity in healthy living medicine - An international perspective. Prog Cardiovasc Dis. 2022;71:64-68.

40.   Emily Brown Weida PP, Falguni Patel, Mariana Chilton. Financial health as a measurable social determinant of health. PLOS ONE. 2020;15(5):e0233359.

41.   Evans KS, Brown. Socioeconomic vulnerabilities in Caribbean women. Global Health Action. 2021;13(2):38-44.

42.   Peterson RR. Over the Caribbean Top: Community Well-Being and Over-Tourism in Small Island Tourism Economies. Int J Community Wellbeing. 2022;1–38.

43.   Francis Poitier RK, Bassey Ebenso. Impact of sustained health policy and population-level interventions on reducing the prevalence of obesity in the Caribbean region: A qualitative study from The Bahamas. Front Public Health. 2022;10:926672.

44.   World Bank. World Development Indicators (2025). Retrieved: 2/15/25 URL: https://data.worldbank.org/indicator.

45.   Keck CW, Reed GA. The curious case of Cuba. Am J Public Health. 2012;102(8):e13-22.

46.   Salicrup LA, Ordunez P, Engelgau MM. Hypertension control activities in Latin America and the Caribbean: opportunities for late-stage (T4) translation research. Rev Panam Salud Publica. 2018;42:e22.

47.   Singh KP, Taylor. Methodological considerations in cross-sectional studies. Research Methodology Journal. 2023;8(1):73-81.

Figure Legends

Figure 1: Gender-Based Disparities in Mortality and Health Indicators Across the Caribbean All data were obtained from PAHO using 2019 data. Welch's t-tests were performed to compare men and women across all variables. (A) External causes mortality rate (age-adjusted per 100,000 population) and the proportion of deaths from noncommunicable diseases and communicable diseases (%) by gender. (B) Age-adjusted mortality rates (per 100,000 population) for ischemic heart disease, respiratory diseases, cirrhosis, and stomach cancer by gender. (C) Diabetes mortality rate (per 100,000 population) and proportional diabetes mortality (%) comparing USA/Canada and the Caribbean by gender. (D) Prevalence of insufficient physical activity and obesity (%) by gender.

Figure 2: The Caribbean Faces Greater Socioeconomic and Health Challenges Compared to the USA/Canada HDI and IHDI data were obtained from UNDP for 2019, while all other data were from PAHO for 2019. Welch's t-tests were used to compare the USA/Canada (US/CAN) with 30 Caribbean

nations across all variables. (A) Human Development Index (HDI) and Inequality-adjusted HDI (IHDI)

scores comparing US/CAN and CAR regions. (B) Food insecurity prevalence (%) and inflation (%) in

US/CAN and CAR regions. (C) Homicide and suicide mortality rates (per 100,000 population) by region.

(D) Unemployment rates (%) for men and women in US/CAN and CAR regions.

Figure 3: The Caribbean Exhibits Greater Health and Economic Vulnerabilities Compared to the

USA/Canada Welch's t-tests were used to compare the USA/Canada (US/CAN) with 30 Caribbean

nations across all variables. (A) Maternal mortality ratio (per 1,000,000 live births) and infant mortality

rate (per 1,000 live births) comparing US/CAN and CAR regions. (B) Adolescent fertility rate (per 1,000

population) and adolescent mortality rate (per 100,000 population) by region. (C) Life expectancy at birth

and median age (in years) for US/CAN and CAR regions. (D) Current health expenditure as a percentage

of GDP and out-of-pocket expenditure as a percentage of current health expenditure (%) comparing

US/CAN and CAR regions.

Figure   4:   Prevalence   of   Gestational   Diabetes   Mellitus   (GDM)   by   Region,   2021

All data were obtained from the International Diabetes Federation (IDF) for the year 2021. The dataset

represents the prevalence of hyperglycemia in pregnancy (age 20-49) across different regions. The map

was created in QGIS 3.34 T using shapefiles from GADM to visualize the regional prevalence of GDM.

Darker shades indicate higher prevalence rates, highlighting disparities in gestational diabetes across the

regions. South-East Asia: 25.9%, North America and Caribbean: 20.7%, South and Central America:

15.8%, Europe: 15%, Middle East and North Africa: 14.1%, Western Pacific: 14%, Africa: 13%.

Figure 5: Correlation of Health and Demographic Variables and Distinct Regional Health Profiles

in the Caribbean Only data with at least 75% availability were included for Pearson correlation and

Principal Component Analysis (PCA). (A) illustrates the relationships between health and demographic

variables across countries with at least 75% data availability. Positive correlations are shown in red, and

negative correlations in blue, with stronger correlations indicated by more intense colors. (B) PCA

identified three distinct clusters of countries. Principal Component 1 (PC1) reflects socioeconomic status

and health outcomes, while Principal Component 2 (PC2) represents healthcare access and lifestyle-

related health risks. Clusters highlight the USA, Canada, and Cuba with favorable health outcomes,

intermediate Caribbean nations with moderate health and socioeconomic challenges, and more vulnerable

Caribbean nations with higher adolescent mortality and fertility rates.

Supplemental Figure Legends

Figure S1: Principal Component Analysis (PCA) and Hierarchical Clustering Selection for

Caribbean Nations Only data with at least 75% availability were included for Principal Component

Analysis (PCA) and clustering. (A) The scree plot displays the eigenvalues for each principal component,

showing that the first two components explain the majority of variance in the dataset. (B) The elbow plot

illustrates the total within-cluster sum of squares, identifying the optimal number of clusters where

additional clusters provide minimal variance reduction. These analyses were conducted on imputed data

to extract key socio-economic and health patterns across 30 Caribbean nations.

Figure S2: Hierarchical Clustering of Caribbean Nations Based on Socioeconomic and Health

Indicators Only data with at least 75% availability were included for hierarchical clustering. (A) The

dendrogram illustrates Ward's hierarchical clustering applied to PCA scores, grouping Caribbean nations

based on shared socio-economic and health characteristics. The y-axis represents Euclidean distance, with

shorter distances indicating greater similarity. (B) The clustering analysis identified three primary country

groups: one comprising nations with favorable health and economic indicators, another with intermediate

Caribbean nations facing moderate health and socioeconomic challenges, and a third with more

vulnerable nations experiencing higher adolescent mortality and fertility rates.

HYPERGLYCAEMIA IN PREGNANCY (20-49 Y)
PREVALENCE OF GESTATIONAL DIABETES MELLITUS
(GDM), % FROM IDF

0   250   500 km

Legend

Prevalence of
Hyperglycemia in
Pregnancy (20-49y).

13.0 - 14.0	15.0 - 16.0	19.0 - 20.0	23.0 - 24.0
14.0 - 15.0	16.0 - 17.0	20.0 - 21.0	24.0 - 25.0
	17.0 - 18.0	21.0 - 22.0	25.0 - 26.0
	18.0 - 19.0	22.0 - 23.0	

A

B