Network-based Molecular Constraints on *in vivo* Synaptic Density Alterations in Schizophrenia

Sidhant Chopra[1-3], Patrick D. Worhunsky[4], Mika Naganawa[5], Xi-Han Zhang[1], Ashlea Segal[6], Loïc Labache[9], Edwina Orchard[1,7], Vanessa Cropley[2,3], Stephen Wood[2,3,8], Gustavo A. Angarita[4], Kelly Cosgrove[4], David Matuskey[4,5], Nabeel B. Nabulsi[5], Yiyun Huang[5], Richard E. Carson[5], Irina Esterlis[1,4,5], Patrick D. Skosnik[4], Deepak C. D'Souza[4], Avram J. Holmes[9]*****, Rajiv Radhakrishnan[4,5]*

1. Department of Psychology, Yale University, New Haven, CT, USA
2. Orygen, Parkville, Melbourne, VIC, Australia
3. Centre for Youth Mental Health, The University of Melbourne, Melbourne, VIC, Australia
4. Department of Psychiatry, Yale University, New Haven, CT, USA
5. Department of Radiology and Biomedical Imaging, Yale University, New Haven, CT, USA
6. Department of Neuroscience, Yale University, New Haven, CT, USA
7. Ann S. Bowers Women's Brain Health Initiative, University of California Santa Barbara, CA, USA
8. School of Psychology, University of Birmingham, UK
9. Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, NJ, USA

*These authors have contributed equally to this work.

Corresponding Author: Sidhant Chopra (sidhantchopra4@gmail.com; sidhant.chopra@orygen.org)

Abstract

Converging neuroimaging, genetic, and post-mortem evidence highlights the fundamental role of synaptic density reductions in schizophrenia pathogenesis. However, the brain-wide spatial pattern of these alterations and the mechanisms underlying this patterning remain to be established. Here, using $[^{11}C]$UCB-J radiotracer positron emission tomography (PET) imaging in individuals with schizophrenia (n=29) and healthy controls (n=93), we find a prominent and widespread pattern of lower synaptic density (0.58 < Cohen's D < 1.47; p_{FWE}<0.05) in patients. The left hemisphere is substantially more impacted than the right (Cohen's D = 1.14; p < .001), with frontal, temporal, cingulate, thalamic, striatal and hippocampal areas particularly affected. Synaptic density alterations were not spatially aligned with gray matter alterations indexed using anatomical Magnetic Resonance Imaging. Lower synaptic density in the left hemisphere is associated with higher normative concentrations of $GABA_{A/BZ}$, $5HT_{2A}$, mGluR5 and $5HT_{1B}$ (r_{cca} = .68; p = .022). Simulation-based network diffusion models identified regions that may represent the initial sources of pathology, nominating left inferior frontal areas (p_{FWE} < .05) as potential foci from which synaptic pathology initiates and then propagates to structurally connected and molecularly similar areas. Overall, our findings provide *in vivo* evidence for widespread synaptic density deficits in schizophrenia that are left-lateralized, independent of gray matter alterations, aligned to specific neurochemical systems, and suggest that such synaptic pathology may propagate in a pattern consistent with axonal networks.

64 **Introduction**
65

Post-mortem, genetic and neuroimaging studies have converged to implicate loss of synapses as a
fundamental pathophysiological process in schizophrenia[1-4]. The advent of positron emission tomography
(PET) radioligands with high specificity for synaptic vesicle proteins[5] have made it possible to examine
presynaptic synaptic terminal density *in vivo*[6], with preliminary studies in schizophrenia[7-9] demonstrating
reductions in specific cortical and subcortical areas. However, the brain-wide spatial patterning of synaptic
pathology, and the mechanisms shaping this pattern, remain poorly understood.
72

Research in neurological conditions shows that the brains vulnerability to pathology is not spatially random
but instead reflects intrinsic properties of large-scale systems[10], cellular composition[11], and receptor
distributions[12]. Similar principles may apply in schizophrenia, where gray matter morphological deficits
preferentially impact higher-order association networks and align with normative receptor expression
profiles. Moreover, several lines of evidence suggest that brain pathology in schizophrenia is constrained
by macroscale axonal networks. For instance, grey matter reductions in patients correlate with the
microstructure of adjacent white matter[13], across spatially distributed regions[14,15], with normative
connectome organization[16,17], and with reductions in structurally connected and functionally coupled
regions[18-20]. Moreover, multiple longitudinal and modelling-based studies have now demonstrated that
pathology may spread from an initial site to axonally connected territories[20-23]. However, widely used MRI-
based morphometric measures predominantly index a composite of tissue properties such as intracortical
myelin, neurite packing, glial density, iron and water content and vascular effects, rather than synaptic
terminals. Thus, direct *in vivo* measures are required to test whether the systems-level features confer
vulnerability and shape the spatial organization of synaptic pathology in schizophrenia.
87

Here, we address this gap by mapping the *in vivo* brain-wide spatial pattern of synaptic density alterations
using [11C]UCB-J PET in a large single-site cohort of 122 individuals, including 29 with schizophrenia. We
demonstrate a prominent, widespread pattern of reduced presynaptic terminal density in schizophrenia, with
stronger effects in the left hemisphere and in frontal, temporal, cingulate, thalamic, striatal, and
hippocampal regions. These reductions were spatially independent of grey matter volume alterations
measured with structural MRI and closely aligned with normative distributions of specific neurotransmitter
and transporter systems. Network diffusion models further nominated left prefrontal regions as potential
sources of pathology, from which synaptic deficits may propagate to axonally connected and molecularly
similar territories. Together, these findings provide the first i*n vivo* evidence that presynaptic terminal
deficits in schizophrenia are widespread, left-lateralized, and shaped by both molecular architecture and
macroscale network organization.
99

100 **Methods**
101 *Sample characteristics*
102

A total of 122 individuals, including 29 individuals diagnosed with schizophrenia were included in this
study. Individuals with schizophrenia were recruited via clinician referrals, paper, and web advertisements.
The majority of individuals with schizophrenia were recruited using criteria that excluded based on the
following criteria: regular exposure to drugs of abuse (except nicotine and caffeine) within the past 3 months
based on history or positive urine drug screen, lifetime substance use disorder (except for nicotine and

caffeine), weekly alcohol consumption exceeding National Institute on Alcohol Abuse and Alcoholism
(NIAAA) guidelines (4 or more drinks on any single day and/or 21 or more drinks per week for males),
history of significant head injury resulting in unconsciousness, unstable medical condition, significant prior
exposure to radiation, metal in the body which would exclude MRI scan or claustrophobia. A subset of
individuals with schizophrenia were recruited using a less restrictive criteria, where substance use was not
an exclusion. We repeated our primary analyses after excluding subjects with a history of, or current,
substance abuse (n=7; SFig1). Data from 13 of the included individuals with schizophrenia have been part
of a previously published study[7]. The healthy control subjects were included from studies performed at the
Yale PET Center using the [^{11}C]UCB-J PET to investigate brain synaptic density across a variety of
neuropsychiatric conditions. Healthy control individuals were excluded for a current and/or lifetime
diagnosis of a psychiatric disorder, current or past serious medical or neurological illness (including a
history of head injury with loss of consciousness), metal in body which would result in MRI
contraindication, or a history of substance abuse or dependence. Demographic and sample characteristics
are provided in Table 1. All study protocols were approved by the Yale University Human Investigation
Committee and Radiation Safety Committee. All participants provided written informed consent prior to
participation.

124
PET and MR imaging acquisition and processing
[^{11}C]UCB-J was synthesized using established methods[24]. All [^{11}C]UCB-J PET measurements were
conducted on the HRRT (Siemens Medical Solutions), which acquires 207 slices (1.2-mm slice separation)
with a reconstructed image resolution of ~3 mm. Before every [^{11}C]UCB-J injection, a transmission scan
was performed for attenuation correction. Dynamic PET data (frames: 6×0.5 min, 3×1 min, 2×2 min, and
16×5 min) were acquired and reconstructed using the MOLAR algorithm[25]. Event-by-event motion
correction was performed using Polaris Vicra optical tracking (NDI Systems, Waterloo, Canada) with
reflectors mounted on the subject's head via a swim cap[26]. All gray matter voxels were corrected for partial
volume effects using the Müller-Gärtner algorithm[27]. Then, the simplified reference tissue model 2
(SRTM2)[28,29] with the centrum semiovale as the reference region, was applied using 60 mins of dynamic
data to generate parametric images of binding potential (BP_{ND}) [6,7,30,31]. Mean voxel-level BP_{ND} maps for
patients and controls are provided in SFig. 2.

137
A high-resolution T1-weighted structural MPRAGE scan (TR/ TE = 2530/3.34, flip angle = 7°, in-plane
resolution = 0.98 × 0.98 mm, matrix size = 256 × 256, slice thickness = 1 mm, slices = 176) was acquired
on a 3T MAGNETOM Prisma scanner (Siemens, Erlangen, Germany). Registration of parametric PET
images to Montreal Neurological Institute (MNI152) standard space was performed using SPM12
(Wellcome Trust center for Neuroimaging, London, UK)[32]. For each subject, PET images were motion-
corrected by registering each frame to a summed image (0–10 min post-injection). A linear registration
aligned the summed PET image to the T1-weighted MR image, followed by a nonlinear registration to the
MNI template. The combined transformations were then applied to the parametric PET image. The
standardized parametric PET images were smoothed with a 4mm full width half maximum Gaussian
kernel then masked using an enhanced gray-matter tissue prior map thresholded at .25[33].

148
149 ***Mapping brain-wide synaptic density alterations***
150 General liner models examining lower BP_{ND} in individuals with schizophrenia compared to controls,
151 adjusting for age and sex, were fit at each grey matter voxel. Non-parametric voxel-level family-wise error

152 (FWE) correction was implemented using CAT12[34], using 5000 permutations, with significance assessed
153 at $p_{FWE} < 0.05$ (Fig1A).
154
To examine hemispheric asymmetry in brain-wide synaptic pathology, t-statistic map which indexed
differences in synaptic density between patients and controls, were then parcellated using a functional
previously validated homotopic atlas into 300 discrete cortical[35] and 32 subcortical[36] areas of approximately
equal size. Hemispheric asymmetry was quantified by computing right-left differences in the test-statistic
across each homotopic parcel (negative values indicating leftward asymmetry). A one-sample t-test was
used to test for systematic asymmetry. To generate a non-parametric null distribution, left and right values
were randomly swapped within each homotopic pair (10,000 permutations), and significance was assessed
as the proportion of permuted mean differences exceeding the observed mean difference.
163
Association between grey-matter volume and synaptic density alterations
Voxel-based morphometry (VBM), implemented in the CAT12 toolbox[34], was used to index brain-wise
grey matter volume alteration between individuals with schizophrenia and controls within the current
sample, as well as four previously published independent samples[20] which included individuals with
medication-naive first-episode psychosis[37] ($N_{case/control}$=59/27), early psychosis[38] ($N_{case/control}$=121/57), and
two cohorts of established schizophrenia[39,40] ($N_{case/control}$=66/72; $N_{case/control}$=70/62). The VBM procedure
used for each sample has been previously described in detail[20]. The resulting brain-wide voxel-level t-
statistic map from each study, which indexed differences in gray matter volume between patients and
controls, were then parcellated using the same 332 region atlas described above. The volume alteration of
a region was estimated as the mean test-statistic (comparing patients and controls) of all voxels
corresponding to that region. The same parcellation procedure was followed for synaptic density alteration
map from the current sample (Fig 1B), allowing for direct comparison between synaptic and gray-matter
volume alteration maps. Product-moment correlations (r) were used to examine spatial associations between
synaptic density alteration and gray matter alterations within the current sample and four independent
cohorts (Fig 1C). Statistical significance of associations between spatial maps was assessed using a
benchmark null model that preserved spatial autocorrelation, allowing us to evaluate whether the observed
findings were specific to the empirically observed alteration maps or were a generic property of the intrinsic
spatial structure of these maps. The procedure used a spin-test to rotate cortical region-level t-values from
the synaptic density alteration map 10,000 times[41,42]. For brain-wide analyses, the rotation was applied to
one hemisphere and then mirrored for the other hemisphere. For hemisphere-specific analyses, the rotations
were computed independently for each hemisphere. Sub-cortical t-values were randomly permuted without
replacement. All *p*-values were quantified as the fraction of null correlation values exceeding the observed
correlation, considering both positive and negative tails of the null distribution (two-tailed). Statistical
significance was assessed at $p < .05$ and inference was FWE corrected using Bonferroni correction (Fig 1C).
188
Quantifying network, neurochemical and cellular enrichment of *synaptic density alterations*
Network-level enrichment. To examine whether synaptic density alterations were preferentially enriched
within specific large-scale canonical functional networks and cytoarchitectonic territories of each
hemisphere, we used two brain system classifications: i) intrinsic functional networks based on the 17
cortical Yeo[43,44] and ii) cytoarchitectonic types of laminar differentiation based on the canonical von
Economo and Koskinas atlas [41,45,46]. Synaptic pathology was estimated averaging the t-statistic across
regions belonging to each functional network or cytoarchitectonic subdivision (Fig. 2A-B). Statistical

significance of observed systems-level synaptic pathology was compared to mean values derived using the
spin-test with 10,000 permutations, as previously described. Statistical inference was assessed at p<.05
(two-tailed) and FWE corrected using Bonferroni correction across 17 networks. To quantify the effect size
for each network, we computed z-scores by comparing the observed test-statistic to the null distribution[18].
200
Neurochemical and cellular associations. To examine whether synaptic density alterations were
preferentially enriched with normative neurochemical or cellular densities, we used two sources of data.
For regional normative neurochemical densities, we used 18 different group-consensus PET binding maps
from healthy population collated by the *neuromaps* toolbox[47]. Following previous work, for tracers that had
multiple samples, a sample-size weighted average map was computed[48]. For cases where two different
tracers targeted the same receptor/transporter, the one with the larger sample size was retained. Further
details regarding each PET map are provided elsewhere[47]. Each PET map was masked and parcellated into
the same 332 region atlas as the synaptic density alteration maps. Regional transcriptomic cell-type
concentrations were deconvolved from whole brain bulk tissue transcriptions provided by the Allens
Human Brain Atlas (AHBA)[49], with 24 different cell-types being imputed using recently published single-
nucleus RNA-sequencing data[50] (see Zhang et al[51] for detailed overview). Briefly, the transcription
signatures identifying each class of cell in the AHBA bulk samples were derived from cortical single-
nucleus RNA sequencing (snRNA-seq) data of eight cortical areas. This included 24 cellular classes with
distinct laminar specialization, developmental origin, morphology, spiking pattern, and broad projection
targets[52]. These cells include 9 GABAergic inhibitory interneurons (PAX6, SNCG, VIP, LAMP5, LAMP5
LHX6, Chandelier, PVALB, SST CHODL, SST), 9 glutamatergic excitatory neurons (L2/3 IT, L4 IT, L5
IT, L6 IT, L5 ET, L5/6 NP, L6 CT, L6b, L6 IT Car3), and 6 non-neuronal cells (Astro, Endo, VLMC, Oligo,
OPC, Micro/PVM). This procedure resulted in cell density estimates for each of the 24 cell-types at each
cortical region using the same atlas as the synaptic density alteration maps. Due to data availability, non-
cortical areas were excluded for analyses that used these cellular data.
Given that regional neurochemical and cellular profiles are multivariate, we examined enrichment
between each of these two features and synaptic density alterations using robust non-parametric canonical
correlation analyses[53]. Two canonical correlation analyses models were computed, examining the
associations between regional synaptic density alterations and receptor (Fig. 2C) or cell densities (SFig5).
Statistical significance of the observed canonical correlations (r_{cca}) compared to null correlations derived
using the spin-test[41,42] with 10,000 permutations. Statistical inference was assessed at p<.05 (two-tailed).
To examine which cell or receptor/transporter maps contributed most to synaptic density alterations, we
quantified canonical loadings by computing product moment correlations between the 18 PET or 24 cell
input maps and corresponding canonical variate. To assess statistical significance of canonical loadings, we
used robust bootstrapping to estimate the standard error for each input map (10,000 bootstraps), computed
z-scores by dividing the correlation by the standard error and used these z-scores to compute two-tailed p-
values[54]. Maps showing reliable loadings on the canonical variate were those that survived a Bonferroni
FWE-correction across input features ($p_{FWE} < 0.05$; two-tailed; Fig 2D).
234
235 **Network diffusion model**
236
237 We used a network diffusion model to test whether the pattern of lower synaptic density is consistent with
238 a diffusion process that spreads through the brain and the extent to which certain brain regions may act as
239 sources, or initiation sites, of pathological spread (Fig4)[55]. The network diffusion model simulates the

240   dynamic spread of pathology between the nodes of a weighted network via a process of diffusion (Fig4A),
241   defined as

$$f(t) = e^{-\alpha H t} f_0 \, ,$$

where t is the model diffusion time, which has arbitrary units (a.u.), and $f(t)$ is a vector characterizing the
amount of diffusion in each region at time t. The model was constrained by a weighted structural
connectivity matrix, as the base of diffusion modelling. Here, the structural connectivity matrix was derived
using tractography applied to high-resolution diffusion-weighted imaging data the Human Connectome
Project[56] (details of the procedure provided in the Supplement). The strength of the diffusion process is
controlled by a constant (α) and H is the Laplacian of the weighted base matrix. f_0 represents the initial
distribution of pathology. We repeatedly initialized the model using each of the 332 regions as the starting
seed, such that the initial state was set to 1 for the seed region, and 0 for all other regions. At each
initialization, using a constant of $\alpha = 1$, the network diffusion model was used to estimate the diffusion at
all other regions at time $t = 0$ to 50. In this way, we were able to determine whether a diffusion process
seeded from each region resulted in a spatial distribution of synaptic density loss that matched the
empirically observed patterns.

Consistent with prior work[20,55,57], model performance was evaluated as the product-moment correlation
between the predicted diffusion and observed volume abnormalities at each time step and for each seed,
with the maximum correlation across all time steps. The seed region was excluded when correlating
predicted and observed volume abnormalities to ensure that our analysis was not influenced by large
synaptic abnormalities in the seeds. The performance of the model in capturing the empirical maps of
synaptic density abnormalities was compared to its performance in capturing surrogate maps generated
using the two benchmark models derived using either the spin-test applied to the empirical synaptic
density alteration map, or a network rewiring algorithm applied to the empirical structural connectivity
matrix. Seed regions were considered significant if they were assigned a $p_{FWE} < .05$ for both benchmark
null models. Further details about benchmark null models used to evaluate the network diffusion model
can be found in the Supplement.

Results

Widespread synaptic density abnormalities in schizophrenia

We find prominent and widespread pattern of lower synaptic density in individuals with schizophrenia (Fig.
1A), compared to controls, with frontal, temporal, cingulate, thalamic, striatal, hippocampal and cerebellar
areas surviving stringent voxel-level family-wise error (FWE) correction ($0.58 <$ Cohens $D < 1.47$; $p_{FWE} <$
0.05, Fig. 1A). An analysis of hemispheric asymmetry indicated that the left hemisphere was substantially
more impacted than the right hemisphere ($D = 1.14$, p < 0.001). We repeated this analyses after adjusting
for antipsychotic medication exposure (SFig. 1A), after excluding individuals with a history of substance
abuse (SFig. 1B), and after removing partial volume correction (SFig. 1C), finding a highly similar and
significantly correlated pattern of lower synaptic density. Mean voxel-level BP_{ND} maps for patients and
controls are provided in SFig. 2).

284
285
286   *Table 1. Sample Demographics*

	Schizophrenia (N=29)	Control (N=93)	*p*-value
Age, mean years (SD)	43.21 (14.66)	45.40 (16.83)	0.501
Females, N (%)	4 (13.8%)	37 (39.8%)	0.018
Mass dose, mean in ug/kg (SD)	0.02 (0.02)	0.02 (0.02)	0.647
Activity at time of injection, mean MBq (SD)	235.53 (201.00)	194.61 (125.28)	0.332
Antipsychotic exposure, mean CPZ eqs. (SD)	2693.34 (2738.91)	-	-
PANSS Total, mean (SD)	68.39 (13.51)	-	-
PANSS Positive, mean (SD)	17.46 (5.26)	-	-
PANSS Negative, mean (SD)	16.64 (5.06)	-	-
PANSS General, mean (SD)	33.93 (6.88)	-	-

287   *CPZ eqs. = lifetime exposure to antipsychotics as chlorpromazine (CPZ) equivalents (eqs.). Listed p-values*
288   *correspond to two-sample t-tests, except for gender-differences, where they correspond to a X^2 proportion*
289   *test. PANSS= Positive and Negative Syndrome Scale.*
290
291
292   ***Synaptic density abnormalities in schizophrenia do not overlap with grey matter abnormities***
293
The brain-wide spatial patterns of synaptic and volumetric alteration were not correlated within the current
($r = 0.03$; $p = .0496$), nor across four independent samples (Fig. 1C). This was not because the current
sample was neuroanatomically distinctive, since there were significant correlations between the volume
alteration maps of the current sample and those from other datasets ($0.33 < r < 0.41$; $p_{FWE} < 0.05$),
except for the first episode psychosis dataset ($r = .-0.02$; $p = 0.81$). This is consistent a reduced
magnitude of grey matter alterations reported in first episode psychosis, compared to chronic schizophrenia.
These results remained consistent when examining each hemisphere independently (SFig. 3A), when
deformation-based, rather than voxel-based, morphometry was used to measure gray matter volume
differences between groups (SFig. 3B) and, when partial volume correction was not implemented for the
synaptic density maps (SFig. 3C).
304
305   Overall, these results demonstrate prominent and widespread lowering of synaptic density in schizophrenia,
306   which is left lateralized and spatially independent of commonly reported MRI-derived gray matter volume
307   alterations across different stages of illness.
308
309
310

311
312
***Fig. 1. Widespread and left-lateralized lower synaptic density in schizophrenia.* a)** Voxel-level effect-size
map of difference in binding potential (non-displaceable) between individuals with schizophrenia (SCZ)
and healthy controls (HC), adjusting for age and sex. Test statistics (t-values) were converted to
standardized effect-sizes (Cohen's *D*), with the black outline indicated voxel-level family-wise error
corrected (FWE) significance at p<0.05. **b)** Effect-size map parcellated using the 300 homotopic cortical
regions[35] and subcortical Tian 32 region[36] atlases, with the mean value of voxels assigned to each region.
Bottom bar displays mean asymmetry index across all regions, with the p-value representing overall
significant leftward laterality in synaptic loss. **c)** Thick black outline highlights spatial correlation
(Pearson's r) between parcellated synaptic density alterations (SYN) and gray matter volume (GMV) within
the current sample (top left box), as well as four independent samples including antipsychotic-naive first-
episode psychosis (FEP), early psychosis (EP) and two established schizophrenia (SCZ) cohorts. No
association between synaptic density alterations and gray matter alterations were found. All other boxes
represent correlations between GMV maps between the current and independent samples. Asterisks (*)
indicate statistical significance (*p* < .05) completed to spatial autocorrelation preserving null models, with
a circle indicating survival against FWE correction.

328
329 *Network & molecular enrichment of synaptic pathology*
330
Brain abnormalities in psychiatric and neurological disorders preferentially affect specific systems such as
macroscale functional networks, distinct cytoarchitectonic regions and neurotransmitter systems[11,58,59]. This
spatial organization of pathology suggests that the functional and molecular properties of distributed brain
regions may confer differential vulnerability to synaptic pathology. Supporting this conjecture, in the right
hemisphere, we find evidence for preferential accumulation of synaptic pathology within aspects of the
functional frontoparietal/cognitive control network (FPN), in particular, dorsal precuneus and posterior
cingulate areas (FPN C subdivision; $Z = 3.02;$ $p < .001;$ Fig. 2A). We also find preliminary evidence that
while the temporoparietal network is preferentially impacted in the left hemisphere ($Z = 1.70;$ $p = 0.047;$
Fig. 2B), it is relatively spared in the right (Z=-1.72; p = 0.025; Fig. 2A), suggestive of functional network-
level asymmetry in synaptic pathology. No significant enrichment was observed for any cytoarchitectonic
types of laminar differentiation.
342
We found a significant association between the spatial pattern of synaptic alterations and neurotransmitter
concentrations within the left hemisphere ($r_{cca} = 0.68;$ $p = 0.022$; Fig. 2C). When examining the
canonical loadings of each receptor/transporter, we found that regions with typically higher concentrations
of $GABA_{A/BZ}$, $5HT_{2A}$, mGluR5 and $5HT_{1B}$, were reliably ($p_{FWE} < 0.05;$ Fig. 2D) associated with lower
synaptic density in individuals with schizophrenia. No significant neurotransmitter association was detected
in the right hemisphere and no significant cell type associations were detected in either hemisphere.
349
350
351
352

Functional network enrichment

a. Left hemisphere

b. Right hemisphere

+ z-score: More synaptic abnormality than expected   * p<.05
- z-score: Less synaptic abnormality than expected   ⊛ p<.05 FWE

Receptors and transporter associations

c. Left hemisphere CCA

$r = .68; p = .022$

d. CCA loadings

Negative   Positive

* p<.05
⊛ p<.05 FWE

Fig. 2. Network and neurochemical correlates of synaptic pathology in schizophrenia. a-b) Enrichment of synaptic pathology within left (a) and right (b) hemisphere functional networks. Bars plot each networks corresponding z-scores relative to the null distribution generated by the spatial permuting procedure. A positive z score indicates greater synaptic loss than expected, whereas a negative z-score indicates less synaptic loss than expected. Asterisks (*) indicate statistical significance ($p < 0.05$), with a circle indicating survival against family-wise error (FWE) correction. **c)** Significant canonical correlation between regional synaptic deficits and normative receptor/transporter density in the left hemisphere. The synaptic variate (y-axis) in this case is the regional synaptic deficit/effect size, whereas the neurotransmitter variate is a normalized weighted combination of regional receptor/transporter densities. The non-parametric *p*-value was computed using the spatial permuting procedure. **d)** Canonical loading (Pearson correlation between normative receptor/transporter density data and neurotransmitter variate) corresponding to (c), where the loading of each receptor/transporter map onto the significant neurochemical variate are plotted. positive loadings indicate that lower synaptic density is associated with higher levels, whereas negative loadings indicate that lower synaptic density is associated with lower levels of the specific molecular system. Asterisks (*) indicate statistical significance using bootstrapping (p<0.05), with a circle indicating survival of FWE correction. TPN: temporoparietal network, DN: default network, FPN: frontoparietal network (sometimes referred to as cognitive control network), LN: limbic network, VAN: salience/ventral attention network, DAN: dorsal attention network, SMN: somatomotor network, VIS: visual network.

373
374 ***Synaptic alterations may be constrained by white-matter pathways***
375
To understand the spatiotemporal dynamics of synaptic loss in schizophrenia and to identify regions from
where pathology may initiate, we used a network diffusion model[55] to directly test whether synaptic loss
propagates through the brain in a way consistent with a process of diffusion between axonally connected
regions (Fig. 3A). The performance (r_{max}) for each brain region is shown in Fig. 3B. Two regions, both in
the left hemisphere showed the best performance and surviving FWE correction (p_{FWE} < .05) across both
null models: frontal operculum/anterior insula (r_{max} = 0.54; Fig. 3C) and the middle temporal gyrus
(r_{max} = 0.53; Fig. 3C). Observed and predicted synaptic pathology using the best performing seed region
(left frontal operculum/anterior insula) are shown in Fig 4D. These findings suggest that synaptic pathology
may initiate in left hemisphere inferior frontal and middle temporal regions and spread to structurally
connected regions.
386
Given the previous association found between neurotransmitter systems and synaptic alteration, we
examined an exploratory network diffusion model where the interregional connectivity matrix was
weighted by receptor/transporter similarity between regions (see Supplement for details). In this way, the
model preferentially diffused pathology not only between regions that were structurally connected, but
between those that also share a similar neurotransmitter composition. In this model, the left frontal
operculum/anterior insula areas again showed the best performance survived FWE correction (p_{FWE} < .05)
across both null models, suggesting that synaptic pathology may propagate between structurally connected
and molecularly similar areas.
395
396

a. Network Diffusion Model

b. Regional performance c. Significant regions d. Significant region performance

398
399
Fig 3. Network diffusion model of synaptic pathology. (a) To understand how global connectivity and
local neurochemical vulnerabilities shape synaptic pathology, we simulated a spreading process using a
network diffusion model. Using each of the 332 parcellated regions as a seed, we retained the maximum
correlation between the simulated and observed synaptic abnormalities. We then compared
maximum r values (r_{max}) for each region to a distribution of maximum r values from two benchmark null
models accounting for spatial autocorrelations in the synaptic pathology maps and basic topological
properties of the connectivity matrix (see Methods). **(b)** Model performance (r_{max}) for each region. **(c)** Seed
regions (purple) with significantly ($p_{FWE} < .05$) greater r_{max} than both null models **(d)** Observed and
network diffusion model predicted synaptic deficit, using the best performing seed (left frontal
operculum/anterior insula).

Discussion

We show a brain-wide pattern of lower *in vivo* synaptic density in individuals with schizophrenia, compared to a large sample of healthy participants. This pattern is left lateralized and spatially independent of commonly reported MRI-derived gray matter volume alterations across different stages of psychotic illness. Functionally defined frontoparietal/control areas were enriched for synaptic pathology in the right hemisphere, whereas the spatial pattern of lower synaptic density left hemisphere is strongly coupled to the normative neurochemical composition, with areas rich in GABA$_{A/BZ}$, 5HT$_{2A}$, mGluR5 and 5HT$_{1B}$ receptors most strongly affected. Using simulation-based network diffusion models, left inferior frontal/anterior insula areas were identified as initial sources from which synaptic pathology initiates and spreads to axonally connected and neurochemically similar areas.

Widespread lower synaptic density in schizophrenias

The quantification of synaptic density within the living brain used [^{11}C]UCB-J, a PET tracer that targets the Synaptic Vesicle Protein 2A (SV2A)[6,60] which is localized and monodispersed on vesicular membranes within all chemical pre-synaptic boutons [61-63]. This tracer has highly specific binding to SV2A[24], good test-retest reliability[6,64], and displacement within gray matter using a drug with specific binding to SV2A[6]. In primates, regional SV2A levels are strongly correlated (r >.95) with synaptophysin[6], a validated marker of *ex vivo* presynaptic density. Moreover, [^{11}C]UCB-J uptake recapitulates anatomically inferred patterns of synaptic density[6], has been shown to be sensitive to synaptic pathology across multiple neurodegenerative conditions[6,65-67], and is specific to synaptic loss measured using electron microscopy, rather than reductions in protein levels[68]. Nonetheless, it is possible that the alterations detected here represent reduced SV2A or vesicular organelles in schizophrenia, in the absence of changes in terminal numbers. For instance, initial evidence suggested that a polymorphism in the SV2A gene may increase risk for schizophrenia, though this has not been replicated in recent large-scale genome-wise association studies[69]. Furthermore, while sustained vesicular depletion occurs as a secondary effect in neurological disorders initiated by illness-linked proteins[70-72], limited post-mortem evidence in schizophrenia actually suggests increased numbers and clustering of synaptic vesicles[73,74]—findings that may result from *ex vivo* artifacts[75,76]. Given the sensitivity of [^{11}C]UCB-J to detecting synaptic terminal loss in other neurological conditions, and the strong converging genetic and post-mortem evidence demonstrating lower presynaptic terminals and dendritic spines in schizophrenia, it is likely that lower levels of [^{11}C]UCB-J binding detected here in schizophrenia are likely a result of lower numbers of synaptic terminals.

We find that areas of gray matter in individuals with schizophrenia, especially thalamic, striatal, hippocampal, frontal, temporal, cingulate, cerebellar and occipital regions, show lower levels of synaptic density compared to controls, with medium to large effect sizes ($0.58 < d < 1.47$). These findings align with previous region-of-interest studies examining *in vivo* synaptic density in early psychosis and schizophrenia populations, which have reported lower tracer binding in frontal, anterior cingulate, hippocampal, temporal, occipital and striatal regions[7-9,77,78], using both measures of specific binding to SV2A, such as that used in the current study, and measures that additionally index non-specific binding. Extending this prior work, our results suggest that synaptic alterations are not only isolated to these regions but are far more widespread and left lateralized in chronic schizophrenia than previously reported. Our

finding are also consistent with large-scale neuroanatomical studies in schizophrenia showing subtle left
lateralized cortical thinning driven by regions of the language network [79].

Patients with schizophrenia often exhibit alterations in cortical thickness and gray matter volume[80-85], which
can progressively worsen in some individuals[37,86,87]. Given the limited evidence from post-mortem studies
for a reduction in the number of neurons, it has been theorized that the substrate underlying these MRI
abnormities are alterations in synaptic, dendritic, and axonal organization[88]. However, there is limited direct
evidence linking MRI signals to neurite changes in psychotic illnesses. Alternatively, MRI signal
differences between patients and controls may reflect differences in properties related to myelination, soma,
glia, vasculature, or extracellular space. The lack of neurobiological understanding behind MRI signal
alterations in psychotic illness have led others to suggest that artefactual alterations between cases and
controls in hydration, cholesterol levels, physical and mental activity, and stress[89-91] may be causes of the
observed volumetric changes. Our finding that the pattern of synaptic and gray matter alterations does not
spatially correlate within the same or independent samples across different stages of illness suggests that a
lowering of presynaptic terminals is not the primary substrate underlying gray matter MRI abnormalities in
psychotic illness. This finding is also consistent with prior work demonstrating that age-related loss of
synaptic density has a different spatial pattern to age-related MRI-derived gray matter volume loss[92].
Although, future work should examine association with a broader range of MRI-derived gray matter metrics
such as cortical thickness and surface area.

Local molecular vulnerability and global axonal connectivity constrain synaptic abnormalities

In many neurological illnesses, pathological processes selectively target or initiate from specific genetic,
cellular[93,94], neurochemical[95,96], and functional systems[10,97]. Here, we find evidence for the preferential
accumulation of synaptic pathology within functionally defined control systems, specifically dorsal
precuneus and posterior cingulate areas. This is consistent with a large body of work demonstrating that
higher-order transmodal association areas are a primary site of pathology in psychosis[98,99]. Our findings
also indicate a strong spatial correspondence of left hemisphere synaptic pathology with normative receptor
profiles. Specifically, regions typically rich in $GABA_{A/BZ}$, $5HT_{2A}$, mGluR5 and $5HT_{1B,}$ receptors, show
vulnerability to synaptic pathology. All three neurotransmiter systems have been shown to be disrupted or
track symptom severity in schizophrenia[100-102], suggesting that synapses associated with these systems are
vulnerable to pathology. This is also consistent with genetic findings showing that enrichment of common
variant associations is restricted to genes that are expressed in both excitatory and inhibitory neurons[103,104]
and in genes encoding proteins involved in general synaptic features. Notably, we did not find associations
with dopamine receptors or transporters, despite finding lower synaptic density in striatal regions, likely
because our brain-wide spatial analysis is dominated by variation in cortical patterns that overlaps poorly
with the predominantly striatal distribution of dopamine maps. Moreover, dopaminergic abnormalities in
schizophrenia are thought to reflect presynaptic functional dysregulation rather than regional
receptor/synapse loss.

We also did not find that densities of transcriptomically-defined cell types were predictive of the spatial
pattern of synaptic pathology. Many post-mortem and genetic studies have shown layer specific cell
abnormalities, commonly in frontal supragranular cells[105,106] and particularly in cortical layer three[107-109].

The current spatial resolution of PET does not allow for layer specific inference, therefore a spatial
correspondence with layer specific cell types may become apparent with improving resolutions.

We used a network diffusion model to understand the spatiotemporal dynamics underlying the local
molecular and brain-wide connectivity constraints. We find that left frontal operculum/anterior insula and
middle temporal cortices are a putative initial source of synaptic density loss in schizophrenia.
Neuroanatomical alterations in these regions are some of the earliest neural alterations reported in the initial
stages of illness[110-112]. Our exploratory analysis which parametrized the network diffusion model using a
neurotransmitter similarity matrix again nominated left frontal operculum/anterior insula areas as putative
source regions. This is consistent with recent large-scale studies that have subtyped schizophrenia based on
the trajectory of gray matter changes, which found that the most common subtype is characterized by
initiation of pathology in inferior fronto-insular areas[21,113]. Our results are in line with a spreading process
in which pathological processes preferentially propagate via axonal connections to molecularly similar
areas. The precise mechanism driving this process remains unclear. Trans-neuronal spreading processes
have often been seen in prion-like spread of misfolded proteins, but limited evidence exists for visible
deposits of aggregated pathological proteins in psychotic illness (although see[114-116]). However, although
speculative, subtle changes in protein homeostasis[117] may interact with regional neurotransmitter
levels[118,119] and spread to synaptically connected distal brain region[120,121]. Alternatively, and given the
commonly reported finding of functional brain alterations in psychotic disorders, dysfunction of one region
may trigger abnormal functional activity in connected sites that, over time, may trigger synaptic changes as
a result of aberrant neurotransmission or a loss of trophic support[122]. Indeed, trans-neuronal propagation of
excitatory-inhibitory imbalance, with consequent excitotoxic stress, could plausibly contribute to the
reduced synaptic density observed in schizophrenia[123].

Limitations

Our analysis of synaptic density depended on group-level summary values and may not directly reflect
synaptic differences for individual patients. While the extent of interpatient heterogeneity in synaptic
pathology is currently unknown, subsequent work could use larger samples with synaptic density imaging
to derive patient-specific measures, such as those obtained through normative modelling[124,125]. The cross-
sectional nature of this study precludes inference about the temporal onset of synaptic pathology, and
further longitudinal studies will be able to clarify whether these alterations occur during neurodevelopment,
at transition to illness or at later stages. An important consideration here is that the cell and receptor density
maps used in the current study were derived from healthy samples and therefore represent normative
cytoarchitecture and chemoarchitecture. Future data aggregation efforts examining cell-type and receptor
maps from patient populations may provide additional evidence regarding synaptic vulnerabilities in
schizophrenia. Our approach to characterize structural connectivity is also limited by the accuracy of
diffusion MRI[126]. While the processing procedures we applied enhanced the biological validity of our
structural connectivity measures as much as possible[127], further developments in non-invasive connectivity
mapping and tractography will be required to allow more precise mechanistic inferences about constrains
imposed by axonal pathways.
545

546 Overall, our findings provide *in vivo* evidence for widespread presynaptic terminal deficits in schizophrenia
547 that are left-lateralized, independent of gray matter alterations, aligned to specific neurochemical systems,
548 and suggest that such synaptic pathology may propagate in a way consistent with axonal networks.

Acknowledgements
SC is supported by the University of Melbourne McKenzie Fellowship and the Brain Behavior Research
Foundation. IE (Nancy Taylor Foundation and VA NCPTSD), DM (NINDS R01NS124819)). AS is
supported by the Yale Wu Tsai Postdoctoral Fellowship). VC is supported by an Australian NHMRC
Investigator Grant (1177370) and a University of Melbourne Dame Kate Campbell Fellowship. This study
was supported in part by the National Institute on Drug Abuse (NIDA) grant R01 DA052454-03
(GAA), U54 AA027989 (KC), Dana Foundation David Mahoney program (RR), CTSA Grant Number
UL1 TR001863 from the National Center for Advancing Translational Science (NCATS) (RR), National
Center for Homelessness among Veterans (36C24820Q1276) (RR), and National Institute for Mental
Health (NIMH) grant R01MH120080 (AJH). We also thank Dr. Jack Tsai (NCHAV) for administrative
support.
601

602 **Declaration of Interests**
603 The authors declare no competing interests.
604

References

Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502-508 (2022). https://doi.org/10.1038/s41586-022-04434-5

Osimo, E. F., Beck, K., Reis Marques, T. & Howes, O. D. Synaptic loss in schizophrenia: a meta-analysis and systematic review of synaptic protein and mRNA measures. *Molecular Psychiatry* **24**, 549-561 (2019). https://doi.org/10.1038/s41380-018-0041-5

Howes, O. D. & Onwordi, E. C. The synaptic hypothesis of schizophrenia version III: a master mechanism. *Molecular Psychiatry* **28**, 1843-1856 (2023).

Fish, K. N., Sweet, R. A., MacDonald, M. L. & Lewis, D. A. Regional Specificity of Cortical Layer 3 Dendritic Spine Deficits in Schizophrenia. *JAMA Psychiatry* (2025). https://doi.org/10.1001/jamapsychiatry.2025.2221

Husain, M. O. *et al.* A systematic review and meta-analysis of neuroimaging studies examining synaptic density in individuals with psychotic spectrum disorders. *BMC psychiatry* **24**, 460 (2024).

Finnema, S. J. *et al.* Imaging synaptic density in the living human brain. *Science translational medicine* **8**, 348ra396-348ra396 (2016).

Radhakrishnan, R. *et al.* In vivo evidence of lower synaptic vesicle density in schizophrenia. *Molecular Psychiatry* (2021). https://doi.org/10.1038/s41380-021-01184-0

Onwordi, E. C. *et al.* Synaptic density marker SV2A is reduced in schizophrenia patients and unaffected by antipsychotics in rats. *Nature Communications* **11**, 246 (2020). https://doi.org/10.1038/s41467-019-14122-0

Yoon, J. H. *et al.* Reductions in synaptic marker SV2A in early-course Schizophrenia. *Journal of Psychiatric Research* (2023).

Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L. & Greicius, M. D. Neurodegenerative diseases target large-scale human brain networks. *Neuron* **62**, 42-52 (2009).

Pak, V. *et al.* Distinctive whole-brain cell types predict tissue damage patterns in thirteen neurodegenerative conditions. *Elife* **12**, RP89368 (2024).

Wiesman, A. I. *et al.* Neurochemical organization of cortical proteinopathy and neurophysiology along the Alzheimer's disease continuum. *Alzheimer's & Dementia* (2024).

Di Biase, M. A. *et al.* Linking Cortical and Connectional Pathology in Schizophrenia. *Schizophr Bull* **45**, 911-923 (2019). https://doi.org/10.1093/schbul/sby121

Wannan, C. M. J. *et al.* Evidence for Network-Based Cortical Thickness Reductions in Schizophrenia. *American Journal of Psychiatry*, appi.ajp.2019.18040380 (2019). https://doi.org/10.1176/appi.ajp.2019.18040380

Cauda, F. *et al.* The morphometric co-atrophy networking of schizophrenia, autistic and obsessive spectrum disorders. *Hum Brain Mapp* **39**, 1898-1928 (2018). https://doi.org/10.1002/hbm.23952

Hettwer, M. *et al.* Coordinated cortical thickness alterations across six neurodevelopmental and psychiatric disorders. *Nature Communications* **13**, 1-14 (2022).

Cauda, F. *et al.* Brain structural alterations are distributed following functional, anatomic and genetic connectivity. *Brain* **141**, 3211-3232 (2018).

Shafiei, G. *et al.* Spatial patterning of tissue volume loss in schizophrenia reflects brain network architecture. *Biological psychiatry* **87**, 727-735 (2020).

Georgiadis, F. *et al.* Connectome architecture shapes large-scale cortical alterations in schizophrenia: a worldwide ENIGMA study. *Molecular Psychiatry*, 1-13 (2024).

Chopra, S. *et al.* Network-Based Spreading of Gray Matter Changes Across Different Stages of Psychosis. *JAMA Psychiatry* (2023). https://doi.org/10.1001/jamapsychiatry.2023.3293

21 Jiang, Y. *et al.* Neuroimaging biomarkers define neurophysiological subtypes with distinct
trajectories in schizophrenia. *Nature Mental Health* **1**, 186-199 (2023).
22 Jiang, Y. *et al.* Neuroimaging epicenters as potential sites of onset of the neuroanatomical pathology
in schizophrenia. *Science Advances* **10**, eadk6063 (2024).
https://doi.org/doi:10.1126/sciadv.adk6063
23 Jiang, Y. *et al.* Gray matter volume heterogeneity by stage, site of origin and pathophysiology in
schizophrenia. *Nature Mental Health* **3**, 803-813 (2025). https://doi.org/10.1038/s44220-025-
00449-9
24 Nabulsi, N. B. *et al.* Synthesis and preclinical evaluation of 11C-UCB-J as a PET tracer for imaging
the synaptic vesicle glycoprotein 2A in the brain. *Journal of Nuclear Medicine* **57**, 777-784 (2016).
25 Carson, R. E., Barker, W. C., Liow, J.-S. & Johnson, C. A. in *2003 IEEE Nuclear Science*
Symposium. Conference Record (IEEE Cat. No. 03CH37515). 3281-3285 (IEEE).
26 Jin, X. *et al.* List-mode reconstruction for the Biograph mCT with physics modeling and event-by-
event motion correction. *Physics in Medicine & Biology* **58**, 5567 (2013).
27 Müller-Gärtner, H. W. *et al.* Measurement of radiotracer concentration in brain gray matter using
positron emission tomography: MRI-based correction for partial volume effects. *Journal of*
Cerebral Blood Flow & Metabolism **12**, 571-583 (1992).
28 Naganawa, M. *et al.* (Soc Nuclear Med, 2022).
29 Wu, Y. & Carson, R. E. Noise reduction in the simplified reference tissue model for neuroreceptor
functional imaging. *Journal of Cerebral Blood Flow & Metabolism* **22**, 1440-1452 (2002).
30 Koole, M. *et al.* Quantifying SV2A density and drug occupancy in the human brain using [11 C]
UCB-J PET imaging and subcortical white matter as reference tissue. *European journal of nuclear*
medicine and molecular imaging **46**, 396-406 (2019).
31 Rossano, S. *et al.* Assessment of a white matter reference region for 11C-UCB-J PET quantification.
Journal of Cerebral Blood Flow & Metabolism **40**, 1890-1901 (2020).
32 Ashburner, J. *et al.* SPM12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK* **2464**
(2014).
33 Lorio, S. *et al.* New tissue priors for improved automated classification of subcortical brain
structures on MRI. *Neuroimage* **130**, 157-166 (2016).
34 Gaser, C. *et al.* CAT–A computational anatomy toolbox for the analysis of structural MRI data.
biorxiv, 2022.2006. 2011.495736 (2022).
35 Yan, X. *et al.* Homotopic local-global parcellation of the human cerebral cortex from resting-state
functional connectivity. *NeuroImage* **273**, 120010 (2023).
36 Tian, Y., Margulies, D. S., Breakspear, M. & Zalesky, A. Topographic organization of the human
subcortex unveiled with functional connectivity gradients. *Nature Neuroscience* **23**, 1421-1432
(2020). https://doi.org/10.1038/s41593-020-00711-6
37 Chopra, S. *et al.* Differentiating the effect of antipsychotic medication and illness on brain volume
reductions in first-episode psychosis: A Longitudinal, Randomised, Triple-blind, Placebo-
controlled MRI Study. *Neuropsychopharmacology* **46**, 1494-1501 (2021).
38 Lewandowski, K. E., Bouix, S., Ongur, D. & Shenton, M. E. Neuroprogression across the early
course of psychosis. *Journal of psychiatry and brain science* **5** (2020).
39 Bustillo, J. R. *et al.* Glutamatergic and neuronal dysfunction in gray and white matter: a
spectroscopic imaging study in a large schizophrenia sample. *Schizophrenia bulletin* **43**, 611-619
(2017).
40 Çetin, M. S. *et al.* Thalamus and posterior temporal lobe show greater inter-network connectivity
at rest and across sensory paradigms in schizophrenia. *Neuroimage* **97**, 117-126 (2014).
https://doi.org/10.1016/j.neuroimage.2014.04.009
41 Markello, R. D. & Misic, B. Comparing spatial null models for brain maps. *NeuroImage* **236**,
118052 (2021).
42 Alexander-Bloch, A. F. *et al.* On testing for spatial correspondence between maps of human brain
structure and function. *Neuroimage* **178**, 540-551 (2018).

43   Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional
connectivity. *Journal of neurophysiology* (2011).
44   Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
Functional Connectivity MRI. *Cerebral Cortex (New York, N.Y.: 1991)* **28**, 3095-3114 (2018).
https://doi.org/10.1093/cercor/bhx179
45   von Economo, C. F. & Koskinas, G. N. *Die cytoarchitektonik der hirnrinde des erwachsenen*
menschen. (J. Springer, 1925).
46   Scholtens, L. H., de Reus, M. A., de Lange, S. C., Schmidt, R. & van den Heuvel, M. P. An MRI
Von   Economo   –   Koskinas   atlas.   *NeuroImage*   **170**,   249-256   (2018).
https://doi.org/https://doi.org/10.1016/j.neuroimage.2016.12.069
47   Markello, R. D. *et al.* Neuromaps: structural and functional interpretation of brain maps. *Nature*
Methods **19**, 1472-1479 (2022).
48   Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional organization
of   the   human   neocortex.   *bioRxiv*,   2021.2010.2028.466336   (2022).
https://doi.org/10.1101/2021.10.28.466336
49   Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain
transcriptome. *Nature* **489**, 391-399 (2012).
50   Jorstad, N. L. *et al.* Transcriptomic cytoarchitecture reveals principles of human neocortex
organization. *Science* **382**, eadf6812 (2023).
51   Zhang, X.-H. *et al.* The Cellular Underpinnings of the Human Cortical Connectome. *bioRxiv* (2023).
52   Jorstad, N. L. *et al.* Comparative transcriptomics reveals human-specific cortical features. *Science*
382, eade9516 (2023).
53   Winkler, A. M., Renaud, O., Smith, S. M. & Nichols, T. E. Permutation inference for canonical
correlation   analysis.   *NeuroImage*   **220**,   117065   (2020).
https://doi.org/10.1016/j.neuroimage.2020.117065
54   Zimmermann, J., Griffiths, J. D. & McIntosh, A. R. Unique Mapping of Structural and Functional
Connectivity   on   Cognition.   *Journal   of   Neuroscience*   **38**,   9658-9667   (2018).
https://doi.org/10.1523/JNEUROSCI.0900-18.2018
55   Raj, A., Kuceyeski, A. & Weiner, M. A network diffusion model of disease progression in dementia.
Neuron **73**, 1204-1215 (2012). https://doi.org/10.1016/j.neuron.2011.12.040
56   Van Essen, D. C. *et al.* The WU-Minn human connectome project: an overview. *Neuroimage* **80**,
62-79 (2013).
57   Raj, A. *et al.* Network diffusion model of progression predicts longitudinal patterns of atrophy and
metabolism in Alzheimer's disease. *Cell reports* **10**, 359-369 (2015).
58   Hansen, J. Y. *et al.* Local molecular and global connectomic contributions to cross-disorder cortical
abnormalities. *Nature communications* **13**, 1-17 (2022).
59   Segal, A. *et al.* Regional, circuit, and network heterogeneity of brain abnormalities in psychiatric
disorders.   *medRxiv*,   2022.2003.2007.22271986   (2022).
https://doi.org/10.1101/2022.03.07.22271986
60   Cai, H. *et al.* Radiosynthesis of 11C-levetiracetam: a potential marker for PET imaging of SV2A
expression. *ACS medicinal chemistry letters* **5**, 1152-1155 (2014).
61   Custer, K. L., Austin, N. S., Sullivan, J. M. & Bajjalieh, S. M. Synaptic vesicle protein 2 enhances
release probability at quiescent synapses. *Journal of Neuroscience* **26**, 1303-1313 (2006).
62   Buckley, K. & Kelly, R. B. Identification of a transmembrane glycoprotein specific for secretory
vesicles of neural and endocrine cells. *The Journal of cell biology* **100**, 1284-1294 (1985).
63   Bajjalieh, S. M., Frantz, G., Weimann, J. M., McConnell, S. K. & Scheller, R. Differential
expression of synaptic vesicle protein 2 (SV2) isoforms. *Journal of Neuroscience* **14**, 5223-5235
(1994).
64   Finnema, S. J. *et al.* Kinetic evaluation and test–retest reproducibility of [11C] UCB-J, a novel
radioligand for positron emission tomography imaging of synaptic vesicle glycoprotein 2A in
humans. *Journal of Cerebral Blood Flow & Metabolism* **38**, 2041-2052 (2018).

Bastin, C. *et al.* In vivo imaging of synaptic loss in Alzheimer's disease with [18F] UCB-H positron emission tomography. *European Journal of Nuclear Medicine and Molecular Imaging* **47**, 390-402 (2020).

Whiteside, D. J. *et al.* Synaptic density affects clinical severity via network dysfunction in syndromes associated with frontotemporal lobar degeneration. *Nature Communications* **14**, 8458 (2023).

Saadabad, F. E. *et al.* in *Neurology.* 3770 (AAN Enterprises).

Chang, Y. Y. *et al.* A study to determine the mechanisms underlying changes in synaptic vesicle glycoprotein 2A density in Alzheimer's disease. *Alzheimer's & Dementia* **19**, e082561 (2023).

Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421-427 (2014). https://doi.org/10.1038/nature13595

Zhou, L. *et al.* Tau association with synaptic vesicles causes presynaptic dysfunction. *Nature Communications* **8**, 15295 (2017). https://doi.org/10.1038/ncomms15295

Parodi, J. *et al.* β-amyloid causes depletion of synaptic vesicles leading to neurotransmission failure. *Journal of Biological Chemistry* **285**, 2506-2514 (2010).

Cabin, D. E. *et al.* Synaptic vesicle depletion correlates with attenuated synaptic responses to prolonged repetitive stimulation in mice lacking α-synuclein. *Journal of Neuroscience* **22**, 8797-8807 (2002).

Soustek, Z. Ultrastructure of cortical synapses in the brain of schizophrenics. *Zentralblatt fur Allgemeine Pathologie u. Pathologische Anatomie* **135**, 25-32 (1989).

Ong, W. & Garey, L. Ultrastructural features of biopsied temporopolar cortex (area 38) in a case of schizophrenia. *Schizophrenia research* **10**, 15-27 (1993).

Rees, S. A quantitative electron microscopic study of the ageing human cerebral cortex. *Acta Neuropathologica* **36**, 347-362 (1976).

Krassner, M. M. *et al.* Postmortem changes in brain cell structure: a review. *Free Neuropathology* **4** (2023).

Onwordi, E. C. *et al.* Synaptic terminal density early in the course of schizophrenia: An in vivo UCB-J positron emission tomographic imaging study of SV2A. *Biological psychiatry* **95**, 639-646 (2024).

Blasco, M. B. *et al.* Synaptic Density in Early Stages of Psychosis and Clinical High Risk. *JAMA psychiatry* **82**, 171-180 (2025).

Schijven, D. *et al.* Large-scale analysis of structural brain asymmetries in schizophrenia via the ENIGMA consortium. *Proceedings of the National Academy of Sciences* **120**, e2213880120 (2023). https://doi.org/doi:10.1073/pnas.2213880120

van Erp, T. G. M. *et al.* Cortical Brain Abnormalities in 4474 Individuals With Schizophrenia and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Consortium. *Biological Psychiatry* **84**, 644-654 (2018). https://doi.org/10.1016/j.biopsych.2018.04.023

van Erp, T. G. M. *et al.* Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Molecular Psychiatry* **21**, 547-553 (2016). https://doi.org/10.1038/mp.2015.63

Olabi, B. *et al.* Are there progressive brain changes in schizophrenia? A meta-analysis of structural magnetic resonance imaging studies. *Biological Psychiatry* **70**, 88-96 (2011). https://doi.org/10.1016/j.biopsych.2011.01.032

Fusar-Poli, P. *et al.* Progressive brain changes in schizophrenia related to antipsychotic treatment? A meta-analysis of longitudinal MRI studies. *Neuroscience and Biobehavioral Reviews* **37**, 1680-1691 (2013). https://doi.org/10.1016/j.neubiorev.2013.06.001

Haijma, S. V. *et al.* Brain Volumes in Schizophrenia: A Meta-Analysis in Over 18 000 Subjects. *Schizophrenia Bulletin* **39**, 1129-1138 (2013). https://doi.org/10.1093/schbul/sbs118

85   Weinberger, D. R., DeLisi, L. E., Perman, G. P., Targum, S. & Jed Wyatt, R. Computed Tomography
in Schizophreniform Disorder and Other Acute Psychiatric Disorders. *Archives of General*
Psychiatry **39**, 778-783 (1982). https://doi.org/10.1001/archpsyc.1982.04290070014004
86   Collins, M. A. *et al.* Accelerated cortical thinning precedes and predicts conversion to psychosis:
The NAPLS3 longitudinal study of youth at clinical high-risk. *Mol Psychiatry* (2022).
https://doi.org/10.1038/s41380-022-01870-7
87   Vita, A., De Peri, L., Deste, G. & Sacchetti, E. Progressive loss of cortical gray matter in
schizophrenia: a meta-analysis and meta-regression of longitudinal MRI studies. *Translational*
Psychiatry **2**, e190-e190 (2012). https://doi.org/10.1038/tp.2012.116
88   Howes, O. D., Cummings, C., Chapman, G. E. & Shatalina, E. Neuroimaging in schizophrenia: an
overview of findings and their implications for synaptic changes. *Neuropsychopharmacology*, 1-
17 (2022).
89   Weinberger, D. R. & Radulescu, E. Structural Magnetic Resonance Imaging All Over Again. *JAMA*
Psychiatry (2020). https://doi.org/10.1001/jamapsychiatry.2020.1941
90   Zipursky, R. B., Reilly, T. J. & Murray, R. M. The myth of schizophrenia as a progressive brain
disease. *Schizophr Bull* **39**, 1363-1372 (2013). https://doi.org/10.1093/schbul/sbs135
91   Weinberger, D. R. & Radulescu, E. Finding the Elusive Psychiatric "Lesion" With 21st-Century
Neuroanatomy: A Note of Caution. *American Journal of Psychiatry* **173**, 27-33 (2016).
https://doi.org/10.1176/appi.ajp.2015.15060753
92   Toyonaga, T. *et al.* The regional pattern of age-related synaptic loss in the human brain differs from
gray matter volume loss: in vivo PET measurement with [11C] UCB-J. *European Journal of*
Nuclear Medicine and Molecular Imaging **51**, 1012-1022 (2024).
93   Hirsch, E., Graybiel, A. M. & Agid, Y. A. Melanized dopaminergic neurons are differentially
susceptible to degeneration in Parkinson's disease. *Nature* **334**, 345-348 (1988).
94   Lotharius, J. & Brundin, P. Pathogenesis of parkinson's disease: dopamine, vesicles and α-synuclein.
Nature Reviews Neuroscience **3**, 932-942 (2002). https://doi.org/10.1038/nrn983
95   Filippi, M. *et al.* Multiple sclerosis. *Nature Reviews Disease Primers* **4**, 43 (2018).
https://doi.org/10.1038/s41572-018-0041-4
96   Mahad, D. H., Trapp, B. D. & Lassmann, H. Pathological mechanisms in progressive multiple
sclerosis. *The Lancet Neurology* **14**, 183-193 (2015).
97   Grothe, M. J., Teipel, S. J. & Initiative, A. s. D. N. Spatial patterns of atrophy, hypometabolism,
and amyloid deposition in Alzheimer's disease correspond to dissociable functional brain networks.
Human brain mapping **37**, 35-53 (2016).
98   Baker, J. T. *et al.* Disruption of Cortical Association Networks in Schizophrenia and Psychotic
Bipolar   Disorder.   *JAMA   psychiatry*   **71**,   109-118   (2014).
https://doi.org/10.1001/jamapsychiatry.2013.3469
99   Chopra, S. *et al.* Generalizable and replicable brain-based predictions of cognitive functioning
across   common   psychiatric   illness.   *Sci   Adv*   **10**,   eadn1862   (2024).
https://doi.org/10.1126/sciadv.adn1862
100   Busatto, G. F. *et al.* Correlation between reduced in vivo benzodiazepine receptor binding and
severity of psychotic symptoms in schizophrenia. *American Journal of Psychiatry* **154**, 56-63
(1997).
101   Selvaraj, S., Arnone, D., Cappai, A. & Howes, O. Alterations in the serotonin system in
schizophrenia: a systematic review and meta-analysis of postmortem and molecular imaging
studies. *Neuroscience & Biobehavioral Reviews* **45**, 233-245 (2014).
102   Frankle, W. G. *et al.* In vivo measurement of GABA transmission in healthy subjects and
schizophrenia patients. *American Journal of Psychiatry* **172**, 1148-1159 (2015).
103   Consortium, T. S. W. G. o. t. P. G., Ripke, S., Walters, J. T. & O'Donovan, M. C. Mapping genomic
loci prioritises genes and implicates synaptic biology in schizophrenia. *medRxiv*,
2020.2009.2012.20192922 (2020). https://doi.org/10.1101/2020.09.12.20192922

104 Owen, M. J., Legge, S. E., Rees, E., Walters, J. T. R. & O'Donovan, M. C. Genomic findings in
schizophrenia and their implications. *Molecular Psychiatry* **28**, 3638-3647 (2023).
https://doi.org/10.1038/s41380-023-02293-8
105 Batiuk, M. Y. *et al.* Upper cortical layer–driven network impairment in schizophrenia. *Science*
Advances **8**, eabn8367 (2022). https://doi.org/doi:10.1126/sciadv.abn8367
106 Ruzicka, W. B. *et al.* Single-cell multi-cohort dissection of the schizophrenia transcriptome. *Science*
384, eadg5136 (2024). https://doi.org/doi:10.1126/science.adg5136
107 Berlekom, A. *et al.* Synapse Pathology in Schizophrenia: A Meta-analysis of Postsynaptic Elements
in Postmortem Brain Studies. *Schizophr Bull* **46**, 374-386 (2020).
https://doi.org/10.1093/schbul/sbz060
108 Glausier, J. R. & Lewis, D. A. Dendritic spine pathology in schizophrenia. *Neuroscience* **251**, 90-
107 (2013). https://doi.org/10.1016/j.neuroscience.2012.04.044
109 Hoftman, G. D., Datta, D. & Lewis, D. A. Layer 3 Excitatory and Inhibitory Circuitry in the
Prefrontal Cortex: Developmental Trajectories and Alterations in Schizophrenia. *Biological*
Psychiatry **81**, 862-873 (2017). https://doi.org/https://doi.org/10.1016/j.biopsych.2016.05.022
110 Pantelis, C. *et al.* Neuroanatomical abnormalities before and after onset of psychosis: a cross-
sectional and longitudinal MRI comparison. *The Lancet* **361**, 281-288 (2003).
https://doi.org/10.1016/S0140-6736(03)12323-9
111 Fusar-Poli, P. *et al.* Neuroanatomy of vulnerability to psychosis: A voxel-based meta-analysis.
Neuroscience & Biobehavioral Reviews **35**, 1175-1185 (2011).
https://doi.org/https://doi.org/10.1016/j.neubiorev.2010.12.005
112 Borgwardt, S. J. *et al.* Structural brain abnormalities in individuals with an at-risk mental state who
later develop psychosis. *The British Journal of Psychiatry* **191**, s69-s75 (2007).
113 Jiang, Y. *et al.* Neurostructural subgroup in 4291 individuals with schizophrenia identified using
the subtype and stage inference algorithm. *Nature communications* **15**, 5996 (2024).
114 Pils, M. *et al.* Disrupted-in-schizophrenia 1 protein aggregates in cerebrospinal fluid are elevated
in patients with first-episode psychosis. *Psychiatry and clinical neurosciences* **77**, 665-671 (2023).
115 Samardžija, B., Renner, É., Palkovits, M. & Bradshaw, N. J. Protein aggregation of DISC1, as
assayed by insolubility, varies across the brain of an individual with schizophrenia and Alzheimer's
disease. *medRxiv*, 2023.2008. 2001.23293413 (2023).
116 Nucifora, L. G. *et al.* Increased Protein Insolubility in Brains From a Subset of Patients With
Schizophrenia. *Am J Psychiatry* **176**, 730-743 (2019).
https://doi.org/10.1176/appi.ajp.2019.18070864
117 Bradshaw, N. J. & Korth, C. Protein misassembly and aggregation as potential convergence points
for non-genetic causes of chronic mental illness. *Molecular psychiatry* **24**, 936-951 (2019).
118 Trossbach, S. *et al.* Misassembly of full-length Disrupted-in-Schizophrenia 1 protein is linked to
altered dopamine homeostasis and behavioral deficits. *Molecular psychiatry* **21**, 1561-1572 (2016).
119 Dahoun, T., Trossbach, S., Brandon, N., Korth, C. & Howes, O. The impact of Disrupted-in-
Schizophrenia 1 (DISC1) on the dopaminergic system: a systematic review. *Translational*
psychiatry **7**, e1015-e1015 (2017).
120 Zhu, S., Abounit, S., Korth, C. & Zurzolo, C. Transfer of disrupted-in-schizophrenia 1 aggregates
between neuronal-like cells occurs in tunnelling nanotubes and is promoted by dopamine. *Open*
Biology **7**, 160328 (2017).
121 Guo, J. L. & Lee, V. M. Cell-to-cell transmission of pathogenic proteins in neurodegenerative
diseases. *Nature medicine* **20**, 130-138 (2014).
122 Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders. (2015).
https://doi.org/10.1038/nrn3901
123 Plitman, E. *et al.* Glutamate-mediated excitotoxicity in schizophrenia: a review. *European*
Neuropsychopharmacology **24**, 1591-1605 (2014).

124   Wolfers, T. *et al.* Mapping the Heterogeneous Phenotype of Schizophrenia and Bipolar Disorder Using Normative Models. *JAMA Psychiatry* (2018). https://doi.org/10.1001/jamapsychiatry.2018.2467

125   Segal, A. *et al.* Regional, circuit and network heterogeneity of brain abnormalities in psychiatric disorders. *Nature Neuroscience* **26**, 1613-1629 (2023). https://doi.org/10.1038/s41593-023-01404-6

126   Maier-Hein, K. H. *et al.* The challenge of mapping the human connectome based on diffusion tractography. *Nature Communications* **8**, 1349 (2017). https://doi.org/10.1038/s41467-017-01285-x

127   Schiavi, S. *et al.* A new method for accurate in vivo mapping of human brain connections using microstructural and anatomical information. *Science advances* **6**, eaba8245-eaba8245 (2020). https://doi.org/10.1126/sciadv.aba8245