Heterogeneous human exposure to arbovirus vectors in an African urban context using immuno-epidemiological biomarker of *Aedes aegypti* bites

Bi Zamblé Hubert ZAMBLE[1,2], Akré Maurice ADJA[3], André Barembaye SAGNA[1,2], François Dipomin TRAORE[2], Mintokapieu Didier Stéphane KPAN[2,3], Négnorogo GUINDO-COULIBALY[3], Affoué Mireille Nadia KOUADIO[2,3], Konan Rodolphe Mardoché AZONGNIBO[2,4], Danielle Dounin ZOH[2,3], Anne POINSIGNON[1], Florence FOURNET[1], Françoise MATHIEU-DAUDE[1,5], Franck REMOUE[1*]

[1] MIVEGEC, University of Montpellier, IRD, CNRS, Montpellier, France
[2] Institut Pierre Richet - Institut National de Santé Publique (IPR-INSP), Bouaké, Côte d'Ivoire
[3] Unité de Formation et de Recherches Biosciences, University Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire
[4] Institut de Géographie Tropicale (IGT), Université Félix Houphouët-Boigny (UFHB), Abidjan, Côte d'Ivoire
[5] UMR SECOPOL, IRD, ILM, Ifremer, University of French Polynesia, Tahiti, French Polynesia

*Corresponding author
Correspondence: franck.remoue@ird.fr

ABSTRACT

Uncontrolled urbanization led to a specific environment, promoting the spread of *Aedes* mosquitoes in many African cities. The objective of the study was to assess human exposure to *Aedes* mosquito bites in the urban area of Abidjan, Côte d'Ivoire, by using antibody-based biomarkers. A cross-sectional study was undertaken during the short rainy season in 3 different neighborhoods (Bromakote, Anoumabo, and Petit-Bassam) presenting various socio-environmental and entomological characteristics. The specific IgG responses to *Aedes* Nterm-34kDa salivary peptide, previously validated as a relevant individual biomarker of exposure to mosquito bites, were assessed in children and analyzed according to neighborhoods and age classes. The specific IgG level was significantly different according to the 3 neighborhoods and higher in Bromakote compared to Anoumabo and Petit-Bassam. No significant difference in specific IgG level was observed according to age. We also noticed an association between the level of specific IgG responses and *Ae. aegypti* densities or various socio-environmental factors. This study indicated that Human exposure to *Aedes* vector appeared to be dependent on neighborhoods within the same city, which could be

41 related to some socio-environmental factors. Antibody-based biomarkers of human
42 exposure to *Aedes* bites could be a helpful tool for assessing the heterogeneity of urban
43 exposure to arbovirus vectors in the framework of monitoring the risk of arbovirus
44 infections, and for evaluating the effectiveness of vector control strategies in national
45 control programs.
46
47 **Keywords:** Urbanization, biomarkers, exposure, Aedes, Africa

                                          Introduction

Emerging and re-emerging diseases such as arboviral infections (dengue, yellow fever, chikungunya,
and Zika) remain major health problems worldwide (Weetman *et al.*, 2018). It is estimated that 2.5
billion people are at risk of dengue infection in the world. Around 200 000 people are affected each
year by yellow fever infection around the world (Weetman *et al.*, 2018). *Aedes aegypti* and *Aedes*
albopictus mosquitoes, the major vectors of arboviruses, are worldwide spread, especially in urban
settings (Vega-Rua *et al.*, 2014).
In Africa, 43% of the population live in urban settings leading to the rapid and uncontrolled
urbanization of many African cities (UN, 2018). Uncontrolled urbanization is often accompanied by
many issues on the spread of vector-borne diseases (Uttara *et al.*, 2012). This urbanization, driven by
the expansion of African cities, creates favorable breeding sites for mosquito proliferation and
potential emergence (Li *et al.*, 2014). Among them, *Aedes aegypti* is the major *Aedes* species in
urban areas in African cities (Li *et al.*, 2014; Fofana *et al.*, 2019).
During the last decades, many epidemics of arboviral diseases occurred in the main cities of some
African countries, as observed in West Africa: in Senegal in 2009 and on major in Ouagadougou,
Burkina Faso in 2023 (Faye *et al.*, 2014; WHO, 2016; Manigart *et al.*, 2024). In Côte d'Ivoire, Abidjan
is the largest and most densely populated city and a prime example of an expanding African
megalopolis with uncontrolled urbanization. In this city, 192 cases of dengue fever were recorded
during the 2017 outbreak. Although poorly documented, yellow fever transmission occurred in Côte
d'Ivoire and some confirmed (n=5) cases were also recorded in the northern part of the country in
2011 (WHO, 2017 and 2011).
No vaccine or medicines are available to control arboviral diseases, except for two efficient vaccines
against yellow fever and Japanese encephalitis. The most effective measures against these diseases
are therefore prevention and vector population control. That involves conventional methods such as
personal protection, community mobilization, insecticide treatments (Andersson *et al.*, 2015), and
innovative approaches such as sterile insect techniques, release of *Wolbachia*-infected *Aedes* (Tur *et*
al., 2021) or biological control of *Aedes* populations (Huang *et al.*, 2017; Utarini *et al.* 2021).
The evaluation of arbovirus transmission risk and vector control effectiveness is crucial. Regarding
Aedes vectors, *Stegomyia* indices such as House Index (HI), Container Index (CI), and Breteau Index
(BI) are calculated and used as indicators for the risk of arbovirus transmission (Bowman *et al.*,
2014). These parameters present several limitations because they represent only an indirect proxy
of the *Aedes* adult density in a study site, and do not directly measure the human-vector contact
(Boyer *et al.*, 2014). In addition, these entomological methods assess human exposure to vector
bites at the population level and do not consider the heterogeneity of inter-individual exposure.
Therefore, WHO underlines the need for the development of alternative indicators to accurately
evaluate the real and direct human exposure to *Ae. aegypti* mosquito bites and to estimate the
potential risk of *Aedes aegypti* borne pathogen transmission in an exposed population (WHO, 2014).
During blood-feeding, concomitantly to blood intake, female mosquitoes inject saliva containing a
cocktail of biologically active proteins that counteract host homeostasis and modulate the
vertebrate immune response (Ribeiro *et al.*, 2003). The injection of vector saliva can also elicit a
human antibody (Ab) response against the salivary proteins (Doucoure *et al.*, 2015; Drame *et al.*,
2010). An interesting approach exploits the immunological properties of mosquito saliva to identify
immuno-epidemiological biomarkers of human exposure to mosquito bites (Fontaine *et al.*, 2011,
Doucoure *et al.*, 2012). To avoid immune cross-reactivity between arthropods and the limitations of
whole saliva use, the identification of antigenic salivary proteins or peptides specific to vector
species/genus was achieved and demonstrated. Specific immunoglobulin G (IgG) antibody responses
have been validated as a serological biomarker of exposure to the main mosquito vectors such as
An. gambiae (IgG response to gSG6 or CE5 salivary proteins, or to the gSG6-P1 salivary peptide), the
vector of human malaria *Plasmodium* (Poinsignon *et al.*, 2008; Rizzo *et al.*, 2014). In the case of
arboviral diseases, several studies have validated the IgG Ab response to only one *Aedes* salivary

peptide (the *Ae. aegypti* Nterm-34 kDa peptide) as a biomarker of human exposure to *Aedes* genus
bites, and thus an indicator for evaluating the risk of arbovirus transmission, and as a relevant tool to
assess vector control strategies (Elanga Ndille *et al.*, 2012; Sagna *et al.*, 2019; Sagna *et al.*, 2018). In a
study in Vientiane city, Laos, the human IgG response against this salivary peptide showed an inter-
individual heterogeneity of exposure to *Ae. aegypti* bites in semi-urban and urban settings (Elanga
Ndille *et al.*, 2014). These results suggested also that the IgG responses can identify areas at risk of
dengue transmission (Elanga Ndille *et al.*, 2014). In another study, the human IgG response to this
salivary peptide has been validated as a relative quantitative indicator to evaluate the efficacy of
vector control strategies (Elanga Ndille *et al.*, 2016). This study highlighted that IgG response to
salivary peptide presented a short-life property with no cumulative Ab response, and a rapid
decreased (2 to 4 weeks) of specific IgG level was observed after the stop of human-*Aedes* contact.
This present study aimed to use the IgG responses to *Ae. aegypti* Nterm-34 kDa salivary peptide to
evaluate the urban heterogeneity of human exposure to the major arbovirus vectors, *Ae. aegypti* in
Abidjan city, Côte d'Ivoire.
114

Material and methods

Study sites
The study was conducted in three urban settings of Abidjan city (Fig.1), the largest and the most
populated city (about 5 million inhabitants) in Côte d'Ivoire. Located in the southern region and
crossed by Ebrie lagoon, Abidjan city is characterized by a sub-equatorial climate, hot and humid,
which includes a main rainy season (May to July), a short rainy season (September to November),
separated by two dry seasons. Rainfall in Abidjan is abundant with more than 1,500 mm per year.
The temperature is around 27°C all year round and the average annual humidity is over 80 %. During
the short rainy season, covered by the present study, the average temperature was 26°C and the
average rainfall 123 mm over 3 months (Kpan *et al.*, 2021).
125

126
127 **Figure 1. Map of the study area in the city of Abidjan, Côte d'Ivoire.**
128 The different study sites (Bromakote, Anoumabo, Petit-Bassam) are indicated by yellow dots.
129
130

131 Three municipalities (Adjame, Marcory and Port-Bouët) of Abidjan were selected and one
132 neighborhood in each municipality was included in the study. In Adjame, the Bromakote
133 neighborhood was selected because of its extensive commercial activities and the numerous
134 exchanges of people (transport) between the different neighborhoods across Abidjan and between

African countries due to the location of the main bus station of the city. This neighborhood also includes the railway station that connects Côte d'Ivoire to Burkina Faso for people and goods. The choice of Anoumabo in Marcory municipality was based on its precarious nature and therefore its environment potentially favorable to the development of mosquitoes. The choice of Petit-Bassam, in Port-Bouet municipality, was driven by the presence of the autonomous port in this neighborhood, resulting in significant traffic of goods and the potential introduction of vector species and pathogens. Besides, the presence of *Ae. albopictus*, a major vector of arbovirus, and cases of dengue infections have already been reported in this neighborhood (Konan *et al.*, 2013).

In the objective to describe the socio-environmental characteristics of the neighborhoods in relation to the risk of proliferation of *Aedes* vector, one hundred households were randomly selected in each neighborhood. The Geographic Information System software (version QGIS 2.14.4) was used to randomly select households. Information on house type, building materials and social practices (water storage, domestic waste and wastewater management, type of toilets) were collected in the three neighborhoods.

Study population

The entire study was carried out on a population of 880 children aged from 6 months to 14 years, enrolled at the school site in each neighborhood, in parallel to a malaria survey on this target age population. This cross-sectional study was performed during the short rainy season (SRS) from September to November 2015. The present immunological study was conducted on a sub-sample of the initial cohort and included 436 children (Table 1). For the immunological assays, blood samples were collected in November 2015 during the SRS, at the fingertips in microtainer tubes (Sarsted, Marnay, France) and serum was obtained, fractionated into aliquots and then frozen at -20 ° C until used.

The Kruskal-Wallis test on children's age indicated a significant difference in the mean of children's age between the studied neighborhoods (p=0.0002). Comparison two-by-two (Dunn's test and adjusted with Bonferroni method) indicated firstly a significant difference in the age mean between children from Bromakote and those living in Petit Bassam, and then between those living in Anoumabo and Bromakote (p-adjusted=0.001 and p-adjusted= 0.0007 respectively).

Aedes aegypti **Nterm-34 kDa salivary peptide**

The Nterm-34 kDa salivary peptide from *Ae. aegypti* was synthesized and purified (> 95 %) by Genepep SA (Saint Jean de Vedas, France). The peptide was shipped in lyophilized form and then suspended in milliQ water and frozen at -20°C until use.

Evaluation of human IgG level to Nterm-34 kDa salivary peptide

IgG level to Nterm-34 kDa peptide was measured by Enzyme-Linked Immuno-Sorbent Assay (ELISA) as previously described (Elanga Ndille *et al.*, 2012). Briefly, MaxiSorp flat-bottom 96-well (Nunc, Roskilde, Denmark) were coated with Nterm-34 kDa peptide at a final concentration of 20 µg/ml in coating buffer (PBS) and incubated for 150 min at 37°C. Plates were blocked with 200 µL of protein-free Blocking Buffer (Thermo Scientific, Rockford, USA) and incubated for 60 min at 37°C. Individual sera were diluted in buffer (PBS-Tween 1%) and incubated at a final dilution of 1/80. Monoclonal mouse biotinylated anti-human IgG (BD Pharmingen, San Diego, CA) was added in at a 1/500 dilution (PBS-Tween 1%) and incubated for 90 min at 37°C. Peroxidase-conjugated Streptavidin (GE Healthcare, Orsay, France) was then added (1/1000 in PBS-Tween 1%) and incubated for 60 min at 37°C. Colorimetric development was carried out with 2,2'-azino-bis ethylbenzothiazoline 6-sulfonic acid diammonium salt (ABTS, Sigma, Saint-Louis, MO, USA) in 50 mM citrate buffer (pH = 4 containing 0.003% H_2O_2) and absorbance (optical density [OD]) was measured at 405 nm (Multisakan GO Thermo Scientific). Each sample was tested in duplicate wells containing Nterm-34 kDa and in a well without antigen (Nterm-30 kDa) to measure non-specific reactions. Individual results were expressed as the ΔOD value: ΔOD = ODx − ODn, where ODx represents the mean of the

186 individual OD value in both wells with salivary antigen and ODn the individual OD value in a blank
187 well containing no antigen.
188
Entomological data
In the present study, three entomological methods were used according to the mosquito stage as
previously described (Kpan *et al.*, 2021). The identification of breeding sites was conducted during
larval survey while ovitraps (eggs were reared until the adult stage at insectarium) and sticky traps
were used for adult mosquitoes evaluation.
Using the random sampling tool of the QGIS 2.14.4, 20 geographical points were generated within
each of the three study sites. One ovitrap and one sticky trap were placed at each geographical point
during the study period (1 month). As a result, 20 ovitraps and 20 sticky traps were used to collect
Aedes eggs and *Ae. aegypti* adults during the studied season in each 3 studied sites. Data in this
present study were collected during the SRS in November 2015.
Ae. aegypti larvae population was estimated through visiting 300 households (100 in each
neighborhood) and by identifying potential breeding containers in each household. The collected
larvae were reared in an insectarium and morphologically identified after emergence. The
Stegomyia indices have been calculated for the risk of arboviruses transmission by *Ae. aegypti*
according to WHO criteria: i) the House Index (HI) represents the percentage of houses infested by
Ae. aegypti larvae; ii) the Container Index (CI) is the percentage of containers infested by *Ae. aegypti*
larvae, and iii) the Bretau Index (BI) is the number of positive breeding sites in 100 visited houses.
According to these *Stegomyia indices*, the WHO criteria indicate that: i) an area is considered to be
at high risk of transmission of dengue and chikungunya viruses by *Aedes aegypti* if HI or BI or CI
exceed 5, 20, and 3%, respectively, and ii) an area is considered to be at high risk of favoring the
transmission of Zika and yellow fever viruses by *Aedes aegypti* if HI or BI or CI exceed 5, 4 and 3%,
respectively (Kpan *et al.*, 2021).
211
Data analysis
Statistical analysis was done using Graph Pad Prism 5® software (San Diego, CA) and R (Version 3.5.3,
R Core Team, Vienna, Austria). After checking that the data did not follow a Gaussian distribution
(normal distribution), the non-parametric Mann-Whitney test was used to compare the Ab levels
between two independent groups and the non-parametric Kruskal-Wallis test was used to compare
the Ab levels between more than two groups. The Dunn's post-hoc test was also performed for two
by two comparisons in more than three groups, then the difference was adjusted with the
Bonferroni method. The prop.test was used to compare proportions between two or more two
groups. The chi-square test was used to compare the HI and CI between study sites. All differences
were considered significant for $p<0.05$.
222

223   **Results**

Socio-environmental characteristics of neighborhoods
Environmental factors and sociological behaviors/practices (housing type, construction materials,
water storage, domestic waste and waste water management) that could potentially be associated
with human exposure to *Aedes* mosquitoes were analyzed and compared according to the three
neighborhoods. A total of 55, 87, and 98 houses in Anoumabo, Bromakote, and Petit-Bassam,
respectively, were randomly selected (among the 100 households by neighborhood of the
entomological surve and visited allowing to identify common and divergent factors between the 3
neighborhoods (Table 1).
232
233
234

235
236
237 **Table 1. Socio-environmental characteristics of the study sites**
238

	Neighborhoods			
	Anoumabo	**Bromakote**	**Petit Bassam**	*p*-value
Neighborhood characteristics n (%)*				
House type	**n.a= 55**	**n.a= 87**	**n.a= 98**	
Common house	48 (87.28)	74 (83.15)	89 (90.82)	NS
Apartment	03 (05.45)	11 (12.36)	02 (02.04)	0.01
Individual house	04 (07.27)	02 (02.25)	06 (06.12)	NS
Construction materials	**n.a= 45**	**n.a=89**	**n.a= 80**	
Cement	38 (84.44)	83 (93.26)	72 (73.47)	NS
Wood	04 (08.89)	05(05.62)	08 (08.16)	NS
Improved banco	02 (04.44)	01 (01.12)	0	NS
Banco (simple and improved)	01 (02.22)	0	0	NS
Water storage containers	**n.a = 48**	**n.a=74**	**n.a= 75**	
Uncovered containers	40 (83.33)	58 (78.38)	57 (76.00)	0.001
Covered containers	01 (02.08)	01(01.35)	10 (13.33)	0.02
Both covered and uncovered containers	07 (14.59)	15 (20.27)	08 (10.67)	NS
Waste management	**n.a = 50**	**n.a=87**	**n.a=97**	
Collected by companies	17 (34.00)	48 (55.17)	24 (24.74)	0.005
Dropped off near the house	09 (18.00)	04 (04.60)	60 (61.86)	0.001
Burned near the house	01 (20.00)	0	01 (01.03)	NS
Collected or dropped off near the house	23 (73.00)	35 (40.23)	12 (12.37)	0.001

239
240

	Neighborhoods			
	Anoumabo	**Bromakote**	**Petit Bassam**	*p*-value
Wastewater management	**n.a = 55**	**n.a = 87**	**n.a= 98**	
Septic pits	21 (38.18)	55 (63.22)	16 (16.33)	<0.001
Sewage channels	05 (09.00)	17 (19.54)	04 (04.08)	0.0001
Septic pits or abandoned wells	09 (16.36)	07 (08.05)	02 (02.04)	0.006
Pour in the street	19 (14.55)	05 (05.75)	71 (72.45)	<0.001
Septic pits or pour in the street	01 (01.82)	02 (02.30)	05 (05.10)	NS
Sewage channels or abandonned wells	0	01(01.15)	0	NS
			0	
Toilet type	**n.a = 55**	**n.a = 89**	**n.a= 94**	
Modern toilet	11 (20.00)	15 (16.85)	28 (28.57)	0.097
Latrine	43 (78.18)	72 (80.9)	66 (67.35)	NS
Modern toilet and latrine	01 (01.82)	02 (02.25)	0	NS

n.a: number of answers

* All percentage calculated for each characteristic in corresponding column

NS : Not Significant between the 3 neighborhoods

241
242
243
There was no clear difference in the housing type between the three neighborhoods, with the
exception of the number of apartments, which was higher in Bromakote than in the two other sites
(12.36 %; *p*=0.01). Most houses were constructed with cement and no significant difference in the
construction materials was observed between the three sites.
Most residents in the three neighborhoods used the stored water for cooking or drinking, and the
comparison of this practice (use of containers for water storage) was not statistically significant

between neighborhoods (*p*=0.262, data not shown). The large majority of water storage containers
found in all neighborhoods were uncovered (>75 %). However, a significant difference was observed
in the proportion of uncovered water storage containers between neighborhoods (*p*=0.001) with
higher proportion recorded in Anoumabo (83.33 %).
Regarding domestic waste management, a higher proportion of inhabitants in Bromakote declared
that their waste was collected by private companies (55.17 %; *p*=0.005). In contrast, domestic waste
was mostly dropped off near the house in Petit-Bassam (61.22 %; *p*<0.001). Fifty-five (55)
inhabitants in Anoumabo, 80 in Bromakote, and 98 in Petit-Bassam provided information about
wastewater management. Among them, a higher proportion in Bromakote (61.8 %) discharged
wastewater into a septic pit compared to other neighborhoods (*p*<0.001). Channels were also used
for discharging wastewater by 19.54 %, (*p*=0.006). Some inhabitants used both septic pit or
abandoned wells with a higher proportion in Anoumabo (16.36 %; *p*<0.001). In addition, most
people used latrine as a type of toilet, and a very low number of modern toilets was recorded in the
three neighborhoods.
Altogether, the results indicated that some sociological practices such as wastewater and waste
management, water storage mode and housing type appeared differently represented between the
three neighborhoods.
267
Specific IgG antibody level

269   IgG levels to Nterm-34 kDa salivary peptide were compared between the 3 studied neighborhoods in
270   the sub-sample of 436 of children (Table 2, Fig. 2).

272   **Table 2. Population characteristics and median of IgG level to Nterm-34 kDa salivary peptide in the study sites**

	Neighborhoods				
	Anoumabo	**Bromakote**	**Petit Bassam**	**Total**	***p*-value**
Demographic data					
n (%)	135 (30.96)	122 (27.98)	179 (41.06)	436	
Age mean (95% CI)	6.50 (5.95-7.06)	4.98 (4.33-5.64)	5.90 (6.43-6.96)	6.05 (5.72-6.40)	0.001
Age groups [years] (%)					
[1-5]	54	73	78		
[6-10]	66	34	70		NS
[11-14]	15	15	30		
Gender ratio (Male/Female)	0.82 (61/74)	0.92 (56/61)	0.69 (73/105)	0.80 (190/240)	0.016
Anti-Nterm-34 kDa IgG median level (95% CI)	0.174 [0.118-0.271]	0.321 [0.261-0.378]	0.138 [0.106-0.173]		0.001
Anti-Nterm-34 kDa IgG median level by age group (95% CI)					
[1-5]	0.15 [0.16-0.34]	0.35 [0.33-0.46]	0.17 [0.19-0.28]	0.21[0.27-0.33]	0.001
[6-10]	0.20 [0.22-0.34]	0.26 [0.25-0.54]	0.12 [0.14-0.26]	0.16[0.22-0.31]	0.001
[11-14]	0.17 [0.14-0.35]	0.20 [0.14-0.37]	0.15 [0.12-0.26]	0.16[0.17-0.27]	NS

NS : Not Significant between the 3 neighborhoods

274   First, high inter-individual heterogeneity of specific IgG responses (ΔOD ranging from 0 to 1.2) was
275   observed within each neighborhood. Secondly, the level of specific IgG responses varied significantly
276   between neighborhoods (*p*<0.001). These IgG levels were significantly higher in children from
277   Bromakote than those from Anoumabo and Petit-Bassam (*p*<0.001). No significant difference in
278   specific IgG levels was observed between Anoumabo and Petit-Bassam (*p*=0.102).

282
283 **Figure 2. IgG responses to *Aedes aegypti* Nterm- 34 kDa salivary peptide in the study sites.**
284 Each dot represents an individual IgG response (ΔOD) against the *Ae. aegypti* Nterm-34 kDa salivary peptide, and
285 the horizontal bar indicates the median value of specific IgG levels for the site under study. The number of
286 processed sera is indicated below (n).
287
The median of IgG level was also compared between the 3 pre-defined age groups ([1-5], [6-10], and
[11-14] years old) within each neighborhood (Fig 3A, Fig 3B and Fig 3C). There was no significant
difference in specific IgG responses according to the age group whatever the neighborhood (*p*>0.05).
However, highly significant variation in the specific IgG level was observed between the three study
sites in children aged [1-5] and [6-10] years-old (*p*=0.001, Table 2). Indeed, as observed for the
whole studied population, the specific IgG responses were higher in Bromakote compared to the
two other neighborhoods in these youngest age groups. In contrast, no significant differences of
specific IgG level were observed between the three neighborhoods in the [11-14] age group.
296

297
298 **Figure 3. IgG responses to *Aedes aegypti* the Nterm- 34 kDa salivary peptide according to age.**
299 Each dot represents the individual IgG response (ΔOD) against the *Ae. aegypti* Nterm-34 kDa salivary peptide and
300 the horizontal bar indicates the median value of specific IgG levels for each age group. Results are presented

according to the 3 studied sites A: Anoumabo, B: Bromakote and C: Petit-Bassam. The number of processed sera is indicated below (*n*).

Entomological indices

Only three mosquito species were collected during the short rainy season across the three neighborhoods, with *Aedes aegypti* being the only species identified within the *Aedes* genus (Table 3). Three entomological methods were employed, each targeting different mosquito life stages. The dipping method was used for larval surveys, while ovitraps (followed by rearing) and sticky traps were used to assess adult mosquito populations.

Table 3. Entomological data in the study sites

Entomological methods	Species	Neighborhoods			p-value
		Anoumabo	Bromakote	Petit-Bassam	
		n	n	n	
	Ae.aegypti	436	48	192	
Larval survey	*Cx quinquefasciatus*	26	0	1	
	Cx tigripes	0	0	1	
Adults survey n (%)					
Ovitraps	*Ae.aegypti*	68 (25.28)	61 (22.68)	140 (52.04)	0.001
Sticky traps	*Ae.aegypti*	21 (07.72)	186 (68.38)	65 (23.90)	0.001
Breeding site types n (%)					
Visited houses		100	100	100	
Potential breeding site		435 (40.17)	361 (33.33)	287 (26.50)	0.001
Positive breeding sites		78 (17.93)	57 (15.80)	62 (21.60)	NS
Positive breeding site types :					
Water containers[a]		27 (34.61)	14 (24.56)	10 (16.13)	NS
Abandoned containers[b]		12 (15.38)	14 (24.56)	8 (12.90)	0.044
Tires		23 (29.50)	20 (35.09)	30 (48.39)	NS
Other containers[c]		16 (20.51)	9 (15.79)	14 (22.58)	0.066
Ae. aegypti Stegomyia Indices					
House Index		45%	39 %	28 %	NS
Container Index		17.93 %	15.80 %	21.60 %	0.042
Breteau Index		78	57	62	
WHO risk scale		[5-7]	[5-6]	[4-6]	

[a] (barrel, basin, pan, cooler, jar, bucket) [b] (all abandoned water storage containers, others abandoned containers for domestic use, wastes of civilization), [c] (flower pots, wells, water troughs, gutters, tarpaulins, septic tank).
* All percentage calculated between neighborhoods (corresponding row) for each entomological parameter
NS : Not Significant between the 3 neighborhoods

The comparison of results using the ovitraps showed a significant difference in the *Ae. aegypti* proportion between the 3 neighborhoods (*p*=0.001), with the highest proportion observed in Petit-Bassam. In contrast, with sticky traps, a significantly higher proportion of adult mosquitoes was observed in Bromakote (68.38 % compared to the two other sites, *p*=0.001). In visited houses, 1083 potential breeding sites were counted (435 in Anoumabo, 361 in Bromakote, and 287 in Petit-Bassam). The proportions of potential breeding sites were significantly higher in Anoumabo compared to the two other neighborhoods (*p*=0.001). The type of positive breeding sites appeared also different according to the neighborhoods with a higher proportion of water storage containers in Anoumabo (NS), abandoned containers in Bromakote (*p*<0.05), and tires in Petit-Bassam (NS).
Finally, the major *Stegomyia* indices were calculated. The HI and BI parameters appeared higher in Anoumabo but the difference was not significant between the three sites. In contrast, CI was different according to the studied sites and higher in Petit-Bassam (*p*<0.05). The WHO risk scale was thus calculated and 5-7, 5-6 and 4-6 scores were obtained in Anoumabo, Bromakote and Petit-Bassam, respectively. These scores suggested a similar risk between neighborhoods even if a slightly higher risk was noticed in Anoumabo as observed for BI.

                                                      Discussion

In the present study, the quantitative and specific IgG responses to the *Aedes aegypti* Nterm-34 kDa
salivary peptide were measured in order to evaluate and compare the level of exposure to *Aedes*
aegypti bites in children from three neighborhoods of Abidjan city.
Firstly, the immunological data showed individual heterogeneity of IgG response to the *Aedes*
salivary peptide within each studied neighborhood. This suggests that children were differently
exposed to *Ae. aegypti* bites in each studied neighborhood. This heterogeneous exposure *to Aedes*
appears to be closely associated with the urban context, where breeding sites are likely to be highly
diverse (Elanga Ndille *et al.*, 2012; Sagna *et al.*, 2019). In addition, various individual factors (such as
personal protection measures or attractiveness to mosquitoes) and household characteristics could
contribute to these results.
Secondly, the major result of this study is that IgG levels to *Aedes aegypti* Nterm-34 kDa salivary
peptide varied among populations living in different neighborhoods within the same city. This
suggests varying levels of human exposure to *Aedes* vectors across the neighborhoods of Abidjan
city. The influence of some social practices or environmental factors (i.e., the mode of water storage
containers) may explain these differences in *Aedes* exposure, as previously reported (Sagna *et al.*,
2018). In addition, some entomological parameters assessed in the present study may be linked to
the observed variations between neighborhoods, including adult *Aedes* caught by the sticky traps,
the types of breeding sites, and particularly abandoned water storage containers. As observed with
the "adult" entomological methods (capture of adult), the specific IgG level was significantly higher
in Bromakote compared to other neighborhoods. The immunological results suggest that children in
Bromakote were more exposed to *Ae. aegypti* bites than those living in Anoumabo and Petit-
Bassam. Interestingly, specific IgG levels appeared to be associated with only some entomological
indicators. This discrepancy may be attributed to the entomological methods used in this study, as
previously reported (Sagna *et al.*, 2018). One hypothesis is that some entomological parameters may
not be well-suited to the African urban context (WHO, 2016; Nascimento *et al.*, 2020).
Thirdly, some environmental factors (waste management, water storage) and social practices
(household type and construction materials) were compared between the studied neighborhoods.
The objective was to identify the factors that could explain the site-dependent levels of *Aedes*
exposure. Indeed*, Ae. aegypti* mosquitoes are known to exploit any abandoned containers that can
hold water, creating ideal conditions for egg laying (Chareonviriyaphap *et al.*, 2003; Hiscox *et al.*,
2013). Unfortunately, no data on a potential association between socio-environmental level and
specific IgG response was available at the individual level. These gaps could not allow statistical
analyzes to be carried out. Our study represented an initial approach at the population and
descriptive levels.
The active and permanent circulation of the dengue virus in West Africa is maintained by *Aedes*
vectors, which find adequate breeding conditions for their development, as reported in
Ouagadougou, Burkina Faso (Fournet *et al.*, 2016). The present study highlighted that water
containers, abandoned containers or tires could be the major breeding sites according to the studied
neighborhoods. In addition, the proportion of children living in an apartment was higher in
Bromakote compared to other studied neighborhoods. Although the number of apartments was low
in the present study, one hypothesis is that this type of housing may also contribute to increased
Aedes exposure, as previously suggested (Poinsignon *et al.*, 2019). Indeed, the presence of potential
breeding sites in apartments and/or the higher concentration of human population could play a
significant role in this type of housing, compared to individual households. Nevertheless, further
studies are needed to support this hypothesis, and particularly at the individual and/or household
levels.
The IgG level to *Aedes aegypti* Nterm-34 kDa salivary peptide was also analyzed according to 3 age
groups ([1-5], [6-10], and [11-14] years old) and the results showed heterogeneity in specific IgG
levels within each age group. Children appeared therefore to be differently exposed to *Ae. aegypti*

bites in each group. The youngest children were differently exposed to *Aedes* between the studied
neighborhoods. At this young age (1 to 5 years old), children typically stay at home, and could be
therefore exposed to the *Aedes* mosquito which breeds in the abandoned containers in the
compound and have daily biting activities (Sagna *et al.*, 2019). Those attending school and aged
between 6 to 14 years could be also bitten by *Ae. aegypti* in the schoolyard or the classroom, as
previously observed (Garcia-Rejon *et al.*, 2011). Nevertheless, specific IgG responses remained
higher in Bromakote compared to the other sites, but only in children of [1-5] and [6-10] year's age
groups, as observed in the entire studied population. It suggests that the younger children (<10
years old) could represent an appropriate target population for identifying the heterogeneity of
human exposure to *Aedes* bites in urban environments. Altogether, the results indicated that
children of all ages were at risk of arboviral virus transmission in urban settings, which should be
considered for implementation of control strategies.
Overall, the present study highlights that the IgG level to *Ae. aegypti* Nterm-34 kDa salivary peptide
could represent a relevant tool in the African urban context. It can be used to identify the
neighborhoods at risk of exposure to *Ae. aegypti* bites and, consequently, at risk of arbovirus
transmission (Elanga Ndille *et al.*, 2014; Sagna *et al.*, 2018 and 2019). Nevertheless, the present
study presents some limitations. Indeed, socio-environmental data were not directly collected from
the household of sampled children which would have been relevant for integrating these variables
into a statistical model to assess their actual impact on specific IgG level.
In conclusion, this study highlights that IgG level against *Ae. aegypti* Nterm-34 kDa salivary peptide
represents a reliable indicator for evaluating the heterogeneity of children exposure to *Ae. aegypti*
bites both at individual and population levels in African urban context. It could also provide an
alternative approach to establish new criteria for measuring the efficacy of specific vector control
strategies in urban settings, enabling more relevant monitoring of the programmes developed by
public health authorities in African regions.
410

Appendices

Ethics statement

Acknowledgements

419 We gratefully acknowledge the populations of Anoumabo, Bromakote, and Petit-Bassam, especially
420 householders and guardians of children, for their kind support and collaboration.

Funding

422 This research was integrated into the ARBORISK multidisciplinary study funded through the C2D
423 program (Contrat de Désendettement et de Développement; 2013). BZH Zamble was supported by a
424 PhD fellowship provided by the Méditerannée Infection Foundation. DD Zoh was supported by a
425 fellowship from the IRD.

Conflict of interest disclosure

427 The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
428 relation to the content of the article.
429

Data, script, code, and supplementary information availability

Data and scripts are not directly accessible online because saved on personnal computer of authors.
However, the data are available in supplementary materials section.

References

Andersson N, Nava-Aguilera E, Arosteguí J, Morales-Perez A, Suazo-Laguna H, Legorreta-Soberanis J,
Hernandez-Alvarez C, Fernandez-Salas I, Paredes-Solís S, Balmaseda A, Cortés-Guzmán AJ, Serrano de
Los Santos R, Coloma J, Ledogar RJ, Harris E (2015) Evidence-based community mobilization for
dengue prevention in Nicaragua and Mexico (Camino Verde, the Green Way): cluster randomized
controlled trial. *British Medical Journal* **351**, 3267. doi: 10.1136/bmj.h3267.
Bowman LR, Runge-Ranzinger S, McCall PJ (2014) Assessing the Relationship between Vector Indices and
Dengue Transmission: A Systematic Review of the Evidence. *PLoS Negl Trop Dis.* **8**(5):e2848. doi:
10.1371/journal.pntd.0002848
Boyer S, Foray C, Dehecq JS (2014) Spatial and Temporal Heterogeneities of *Aedes albopictus* Density in
La Reunion Island: Rise and Weakness of Entomological Indices. *PLoS ONE* **9**(3):e91170. doi:
10.1371/journal.pone.0091170
Chareonviriyaphap T, Akratanakul P, Nettanomsak S, Huntamai S (2003) Larval habitats and distribution
patterns of *Aedes aegypti* (Linnaeus) and *Aedes albopictus* (Skuse), in Thailand. *Southeast Asian J Trop*
Med Public Health **34**, 529–535.
Doucoure S, Mouchet F, Cornelie S, DeHecq JS, Rutee AH, Roca Y, Walter A, Hervé JP, Misse D, Favier F,
Gasque P, Remoue F (2012) Evaluation of the Human IgG Antibody Response to *Aedes albopictus*
Saliva as a New Specific Biomarker of Exposure to Vector Bites. *PLoS Negl Trop Dis.* **6** (2):e1487. doi:
10.1371/journal.pntd.0001487.
Doucoure S, Cornelie S, Drame PM, Marie A, Elanga Ndille EE, Mathieu-Daude F, Mouchet F, Poinsignon
A, Remoue F (2015) Biomarkers of Vector Bites: Arthropod Immunogenic Salivary Proteins in Vector-
Borne Diseases Control. In: Preedy VR, Patel VB, editors. General Methods in Biomarker Research and
their Applications. p. 1177-1205.
Drame PM, Poinsignon A, Besnard P, Le Mire J, Dos-Santos MA, Sow CS, Cornelie S, Foumane V, Toto JC,
Sembene M, Boulanger D, Simondon F, Fortes F, Carnevale P, Remoue F (2010) Human Antibody
Response to *Anopheles gambiae* Saliva: An Immuno-Epidemiological Biomarker to Evaluate the
Efficacy of Insecticide-Treated Nets in Malaria Vector Control. *Am J Trop Med Hyg.* **83**, 115-121. doi:
10.4269/ajtmh.2010.09-0684
Elanga Ndille E, Doucoure S, Damien G, Mouchet F, Drame PM, Cornelie S, Noukpo H, Yamadjako S,
Djenontin A, Moiroux N, Misse D, Akogbeto M, Corbel V, Henry MC, Chandre F, Baldet T, Remoue F
(2012) First Attempt To Validate Human IgG Antibody Response to Nterm-34kDa Salivary Peptide as
Biomarker for Evaluating Exposure to *Aedes aegypti* Bites. *PLoS Negl Trop Dis.* **6**(11):e1905. doi:
10.1371/journal.pntd.0001905
Elanga Ndille EE, Dubot-Pérès A, Doucoure S, Mouchet F, Cornelie S, Sidavong B, Fournet F, Remoue F
(2014) Human IgG antibody response to *Aedes aegypti* Nterm-34 kDa salivary peptide as an indicator
to identify areas at high risk for dengue transmission: a retrospective study in urban settings of
Vientiane city, Lao PDR. *Trop Med Int Health.* **19**, 576–580. doi: 10.1111/tmi.12280
Elanga Ndille E, Doucoure S, Poinsignon A, Mouchet F, Cornelie S, D'Ortenzio E, DeHecq JS, Remoue F
(2016) Human IgG Antibody Response to *Aedes* Nterm-34kDa Salivary Peptide, an Epidemiological
Tool to Assess Vector Control in Chikungunya and Dengue Transmission Area. *PLoS Negl Trop Dis.*
10(12) :e0005109. doi: 10.1371/journal.pntd.0005109
Faye O, Ba Y, Faye O, Talla C, Diallo D, Chen R, Mondo M, Ba R, Macondo E, Siby T, Weaver SC, Diallo M,
Sall AA (2014) Urban Epidemic of Dengue Virus Serotype 3 Infection, Senegal, 2009. *Emerging*
Infectious Diseases **20**, 456–459. doi: 10.3201/eid2003.121885
Fofana D, Beugré JMV, Yao-Acapovi GL, Lendzele SS (2019) Risk of Dengue Transmission in Cocody
(Abidjan, Côte d'Ivoire). *Journal of Parasitological Research.* **14**;2019:4914137. doi:
10.1155/2019/4914137
Fontaine A, Diouf I, Bakkali N, Missé D, Pagès F, Fusai T, Rogier C, Almeras L (2011) Implication of
haematophagous arthropod salivary proteins in host-vector interactions. *Parasite and Vectors* **4**:187.
doi: 10.1186/1756-3305-4-187

Fournet F, Rican S, Vaillant Z, Roudot A, Meunier-Nikiema A, Kassié D, Dabiré RK, Salem G (2016) The
Influence of Urbanization Modes on the Spatial Circulation of Flaviviruses within Ouagadougou
(Burkina Faso). *Int J Environ Res Public Health*. **13**(12):1226. doi: 10.3390/ijerph13121226
García-Rejón JE, Loroño-Pino MA, Farfán-Ale JA, Flores-Flores LF, López-Uribe MP, Najera-Vazquez Mdel
R, Nuñez-Ayala G, Beaty BJ, Eisen L (2011) Mosquito Infestation and Dengue Virus Infection in *Aedes*
aegypti Females in Schools in Mérida, México. *Am J Trop Med Hyg*. **84**, 489–496. doi:
10.4269/ajtmh.2011.10-0654
Huang YS, Higgs S, Vanlandingham DL (2017). Biological Control Strategies for Mosquito Vectors of
Arboviruses. *Insects* **8**(1):21. doi: 10.3390/insects8010021.
Hiscox A, Kaye A, Vongphayloth K, Banks I, Piffer M, Khammanithong P, Sananikhom P, Kaul S, Hill N,
Lindsay SW, Brey PT (2013) Risk Factors for the Presence of *Aedes aegypti* and *Aedes albopictus* in
Domestic Water-Holding Containers in Areas Impacted by the Nam Theun 2 Hydroelectric Project,
Laos. *Am J Trop Med Hyg*. **88**, 1070–1078. doi: 10.4269/ajtmh.12-0623
Konan YL, Coulibaly ZI, Koné AB, Ekra KD, Doannio JM, Dosso M, Odéhouri-Koudou P. (2013) Species
composition and population dynamics of *Aedes* mosquitoes, potential vectors of arboviruses, at the
container terminal of the autonomous port of Abidjan, Côte d'Ivoire. *Parasite* **20**:13. doi:
10.1051/parasite/2013013
Kpan MDS *et al.* (2021) Spatial Heterogeneity and Seasonal Distribution of *Aedes* (Stegomyia) *aegypti* (L)
in Abidjan, Côte d'Ivoire. *Vector Borne and Zoonotic Dis* **21**, 769-776. doi: 10.1089/vbz.2021.0002
Li Y, Kamara F, Zhou G, Puthiyakunnon S, Li C, Liu Y, Zhou Y, Yao L, Yan G, Chen XG (2014) Urbanization
Increases *Aedes albopictus* Larval Habitats and Accelerates Mosquito Development and Survivorship.
PLoS Negl Trop Dis. **8**(11):e3301. doi: 10.1371/journal.pntd.0003301.
Manigart O, Ouedraogo I, Ouedraogo HS, Sow A, Lokossou VK (2024) Dengue epidemici in Burkina
Faso: how can the response improve?. Lancet 23:02803-02809. doi: 10.1016/S0140-
6736(23)02803-9
Poinsignon A, Cornelie S, Mestres-Simon M, Lanfrancotti A, Rossignol M, Boulanger D, Cisse B, Sokhna C,
Arcà B, Simondon F, Remoue F. (2008) Novel Peptide Marker Corresponding to Salivary Protein gSG6
Potentially Identifies Exposure to *Anopheles* Bites. *PLoS ONE* **3**(6):e2472. doi:
10.1371/journal.pone.0002472
Poinsignon A, Boulanger D, Binetruy F, Elguero E, Darriet F, Gallian P, De Lamballerie X, Charrel RN,
Remoue F (2019) Risk factors of exposure to *Aedes albopictus* bites in mainland France using an
immunological biomarker. *Epidemiol Infect*. **147**:e238. doi: 10.1017/S0950268819001286.
Ribeiro JMC, Francischetti IM (2003) Role of Arthropod Saliva in Blood Feeding: Sialome and Post-Sialome
Perspectives. *Annu Rev Entomol*. **48**, 73–88. doi: 10.1146/annurev.ento.48.060402.102812
Rizzo C, Lombardo F, Ronca R, Mangano V, Sirima SB, Nèbiè I, Fiorentino G, Modiano D, Arcà B. (2014)
Differential antibody response to the *Anopheles gambiae* gSG6 and cE5 salivary proteins in individuals
naturally exposed to bites of malaria vectors. *Parasit Vectors*. **7**:549. doi: 10.1186/s13071-014-0549-8.
Sagna AB, Kassié D, Couvray A, Adja AM, Hermann E, Riveau G, Salem G, Fournet F, Remoue F (2019)
Spatial Assessment of Contact Between Humans and Anopheles and Aedes
Mosquitoes in a Medium-Sized African Urban Setting, Using Salivary Antibody–Based Biomarkers. *The*
J Infect Dis **220**:1199–1208. doi: 10.1093/infdis/jiz289
Sagna AB, Yobo MC, Elanga Ndille E, Remoue F (2018) New Immuno-Epidemiological Biomarker of Human
Exposure to *Aedes* Vector Bites: From Concept to Applications. *Trop Med Infect Dis*. **3**(3)80. doi:
10.3390/tropicalmed3030080.
Tur C, Almenar D, Benlloch-Navarro S, Argilés-Herrero R, Zacarés M, Dalmau V, Pla I (2021) Sterile Insect
Technique in an Integrated Vector Management Program against Tiger Mosquito *Aedes albopictus* in
the Valencia Region (Spain): Operating Procedures and Quality Control Parameters. *Insects* **12**(3):272.
doi: 10.3390/insects12030272.
United Nations (2018) Department of Economic and Social Affairs, Population Division. The speed of
urbanization around the world. 2018/1:1–2.
Uttara S (2012) Impacts of urbanization on environment. *Applied Science* **2**,10-16.
Vega-Rúa A, Zouache K, Girod R, Failloux AB, Lourenço-de-Oliveira R (2014) High Level of Vector
Competence of *Aedes aegypti* and *Aedes albopictus* from Ten American Countries as a Crucial Factor
in the Spread of Chikungunya Virus. *J Virol*. **88**, 6294–6306. doi: 10.1128/JVI.00370-14

Weetman D, Kamgang B, Badolo A, Moyes CL, Shearer FM, Coulibaly M, Pinto J, Lambrechts L, McCall PJ
(2018). Aedes mosquitoes and Aedes-borne arboviruses in Africa: Current and future threats. *Int J*
Environ Res Public Health. **15**(2):220. doi: 10.3390/ijerph15020220
WHO (2011) Yellow fever in Côte d'Ivoire. WHO. World Health Organization Available from:
https://www.who.int/csr/don/2011_01_25/en/
WHO (2016) Entomological surveillance for *Aedes* spp. in the context of Zika virus: interim guidance for
entomologists. (Who/Zikv/Vc/16.2): 10 p.
World Health Organizations. (2014)   A global brief on vector-borne diseases. Available from:
www.who.int/world-health-day
WHO (2016) Dengue Fever – Burkina Faso. World Health Organization. Available from:
http://www.who.int/csr/don/18-november--dengue-burkina-faso/en/
WHO (2017) Dengue fever – Côte d'Ivoire. World Health Organization. Available from:
http://www.who.int/csr/don/04-august-2017-dengue-cote-d-ivoire/en/
550
551
552
553