TITLE

A Systematic Review of Measles Virus Transmisability in the Air to Guide Exposure Periods
for Contact Tracing in Public Spaces

AUTHORS

Haley F. Farrie[1], MPH, Sarah Gillani[1], MPH, Fatima Nkempu Ameaka[1], PharmD, MPH, Eric
S. Toner[1,2], MD, Sanjana Ravi[1], PhD, MPH, Sutyajeet Soneja[1,2], PhD, Caitlin M. Rivers[1,2],
PhD, MPH

Corresponding Author:

Haley Farrie, MPH

Johns Hopkins Center for Outbreak Response Innovation

700 East Pratt Street, Suite 900

Baltimore, MD 21202

Phone: 667-306-9796

Email: Hfarrie1@jh.edu

AFFILIATIONS

[1] Johns Hopkins Bloomberg School of Public Health Center for Outbreak Response
Innovation
[2] Johns Hopkins Bloomberg School of Public Health Center for Health Security
19

20 **FUNDING**
21 This analysis was supported by the U.S. Centers for Disease Control & Prevention
22 (Cooperative Agreement #NU38FT000004).
23

24 **ETHICAL STATEMENT**
25 Interviews cited within this systematic review informed its basis and were approved by the
26 Institutional Review Board at the Johns Hopkins Bloomberg School of Public Health (IRB
27 #00028357) and not considered human subjects research.
28

29 **WORD COUNT, TABLES, AND FIGURES**
30 2,996 words; 1 table; 1 figure
31
32

ABSTRACT (250 words)

Objective: Epidemiological evidence was systematically reviewed to determine how long
the measles virus remains transmissible after an infectious case leaves a public space to
address inconsistencies in measles contact tracing exposure window guidelines.

Methods: A systematic literature review following PRISMA guidelines was conducted using
PubMed, EMBASE, Web of Science, and SCOPUS databases for publications from January
1988 to July 2024. Additional sources were identified through reference list reviews and
Google Scholar searches. Studies examining how long the measles virus survives in the air
or remains transmissible after an infectious case leaves a public space were included,
while non-evidence-based recommendations and mathematical models were excluded.

Results: Initial database searches identified 1,054 studies, with none meeting initial
inclusion criteria after screening. Supplemental searches identified five relevant articles
(1964-1987). Two experimental studies demonstrated measles virus survival between 30
and 120 minutes, with increasing survival time for lower humidity levels. Experimental
studies showed survival at two hours in low humidity (12-15%), up to 60 minutes in
moderate humidity (36-37%), and 30 minutes in high humidity (60-70%) at 20°C. Three
publications reviewed how long measles virus was transmissible in real-world settings,
ranging from 60 to 120 minutes.

Conclusions: Limited evidence exists to guide the precise determination of the duration of
measles transmissibility. Current health department guidelines rely on limited research
from 1964 to 1987. Additional studies are needed to understand how long the virus is
transmissible in real-world settings, particularly given the implications for contact tracing
efficiency and resource allocation during outbreak responses.

KEYWORDS

Public health practice, public health preparedness, communicable diseases, disease
outbreaks, epidemiology, public health

SUMMARY BOX

1) What is the current understanding of this subject?
61
The current understanding of how long the measles virus remains transmissible in
the air is based on limited research from 1964-1987, with CDC guidelines

recommending a two-hour exposure window while some health departments use a
one-hour window.

2) What does this report add to the literature?

This report adds a comprehensive systematic review that confirms the scarcity of
evidence on measles virus transmissibility in air, highlighting that no new research
has been published since 1987 to inform contact tracing guidelines.

3) What are the implications for public health practice?

The implications for public health practice include the need for new research to
standardize evidence-based guidelines for measles contact tracing, which would
optimize resource allocation during outbreak responses, potentially reduce
unnecessary interventions, and help foster trust in public health activities.

Introduction

The United States eliminated measles in 2000[1] due to high vaccination rates and effective
public health measures; however, the U.S. has experienced several outbreaks since then,
typically after introduction from international travelers.[2] These situations are managed by
health department teams, which act quickly to identify people exposed to the case,
provide post-exposure prophylaxis (PEP) to individuals without immunity where possible,
and quarantine contacts at risk of developing the infection. These responses are resource-
intensive, require significant staff time and funding, and can detract from routine health
department operations.[3]

Contact tracing for measles presents unique challenges. Like influenza[4] and SARS-CoV-2[5],
measles spreads through airborne and droplet transmission.[6] However, measles also
remains infectious in the air for an extended time, even after the infected person is no
longer present.[7] According to the current U.S. Centers for Disease Control and Prevention
(CDC) measles exposure guidelines, if a person experiences either of the following without
personal protective equipment for any amount of time, they are considered exposed: 1)
occupying the same air space at the same time as someone with measles or 2) entering a
room within 2 hours after a person with measles left.[7] These exposure guidelines are based
on one experimental study of how long the measles virus remains in the air[8], three
publications reviewing the evidence of real-world outbreaks (1985-1987)[9–11], and a
literature review.[12]

The Johns Hopkins Center for Outbreak Response Innovation conducted eleven key informant interviews with U.S. state and local health departments to understand the operational challenges related to contact tracing and outbreak response. Interviews took place between March and October of 2024. These interviews revealed variations in the measles exposure guidelines for contact tracing.[13] While most health departments utilize the two-hour exposure window recommended by the CDC, two relied on a more lenient one-hour exposure window.[14] The exposure window directly impacts contact tracing procedures. Reducing the exposure window from two hours to one hour can substantially reduce the number of people considered exposed and thus reduce the number of contacts requiring notification/quarantine. A shorter exposure window would likely considerably reduce staff hours and costs required to conduct these activities.[15] Moreover, accurately identifying the exposure window facilitates efficient contact tracing within the 72-hour window when PEP remains effective.[16] Accurately identifying this timeframe is also crucial for ensuring that staff time and funding are appropriately managed and that routine health department work is not disrupted unnecessarily. To address this inconsistency and ensure that exposure guidelines are based on the strongest data available, we conducted a systematic review of the evidence for how long the measles virus remains airborne and capable of infection after an infectious case has left an area.

Methods

This systematic literature review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.[17] This study aimed to examine the duration of measles virus infectivity in airborne environments to guide epidemiologists in determining exposure periods for contact tracing in public spaces after the presence of an infectious case.

Literature Search Strategy

The literature search was conducted in Pubmed, EMBASE, Web of Science, and SCOPUS for articles published between January 1988 and July 2024. This starting year was chosen as it marks the commercialization of the taq polymerase enzyme critical for the widespread utilization of polymerase chain reaction (PCR) technology for accurate measles diagnosis.[18] The following search query was used: ("measles" OR "rubeola") AND ("air microbiology" OR "respiratory aerosols and droplets" OR "airborne" OR "exposure period" OR "survival time" OR "environmental stability" OR "aerosol*" OR "droplet*" OR "air") AND ("disease transmission, infectious" OR "transmission" OR "airborne transmission" OR "survival" OR "mortality"). All results were managed through Covidence,

133 a systematic review processing software, for de-duplication, title and abstract screening,
134 full-text review, and extraction.

Study Screening and Eligibility

After de-duplication, two independent reviewers screened titles and abstracts to assess
eligibility for full-text review. Any disagreements were resolved by a third independent
reviewer, who made a final determination.

Studies were included if they examined the duration of measles infectivity in the air after an
infected person has left a space, specifically in public places (e.g., hospitals, airplanes,
grocery stores). Studies that occurred indoors and outdoors were both eligible for
inclusion. Eligible studies included peer-reviewed articles and grey literature from public
health authorities that address contact tracing in relation to exposure periods and review
the relevant scientific evidence. Both human and animal studies, as well as laboratory-
based investigations on measles infectivity in the air, were considered.

Exclusion criteria included recommendations and studies that were not evidence-based
(including mathematical models), unrelated to airborne transmission of measles in public
environments, and publications not in English. Reviews, editorials, opinion pieces, and
duplicate studies or studies with overlapping data sets were also excluded. If an outbreak
or airborne transmission event was discussed in both peer-reviewed articles and grey
literature, for example, journal publications, only the peer-reviewed articles were included
to reduce duplicative reporting of results.

The manuscripts that met inclusion criteria during the title and abstract screening were
advanced to a full text review. During this phase, two reviewers again independently
assessed each manuscript for inclusion. Items that received discordant votes were each
discussed during a live meeting attended by both primary reviewers and a third reviewer. In
this meeting, each reviewer presented their rationale, and the third reviewer made the final
determination after considering all viewpoints. Any manuscripts that passed the full-text
review were subsequently included in the data extraction and analysis phase.

Supplemental Search Strategies

Due to the dearth of results from the systematic literature search, additional search
strategies were conducted to ensure comprehensive coverage of the evidence. These
three additional methods included 1) review of references from literature reviews identified
during the Covidence search, 2) a Google Scholar search utilizing the same search query
as was implemented in the study databases, and 3) a review of all literature used by the
U.S. CDC to inform the established measles exposure guidelines. These supplemental

search strategies followed the same inclusion and exclusion criteria used during the
systematic search, except that there was no limitation applied to the date of publication.

Literature reviews, and therefore their content, identified during the systematic search
were originally excluded because they did not meet the original search criteria for primary
data collection. However, any literature review references that specified primary data
collection of measles exposure data with timeframes were included as part of the
supplemental search; this is referred to later in this write-up as the literature review
snowball sampling. Although the literature reviews themselves are still not included in the
results, the primary research articles themselves are included. The second supplemental
search strategy was the Google Scholar search, which was conducted using key phrases
searched in combination and independently: "Measles Exposure," "Measles Outbreak,"
"Measles Case Study," "Measles Case", "Measles Evidence," "Measles Airborne," and
"Measles Transmission." The third strategy involved reviewing the evidence published on
the U.S. CDC website.[7] To aid the supplemental search, the findings of this systematic
review were also shared with the state and local health department partners identified
through previous interviews, including those that followed both one- and two-hour
exposure windows, and no additional articles were added.[13]

184   Results

Database searches initially identified 1,054 studies from four databases: Scopus (392),
Web of Science (262), PubMed (200), and Embase (200), as shown in Figure 1 below. After
removing 495 duplicates, 559 studies remained for the title and abstract screening. After
the first round of screening, 463 studies were excluded based on criteria outlined in the
methodology, leaving 96 studies for full-text review. However, none of these studies
ultimately met the previously mentioned criteria for data extraction and analysis.

The supplemental search strategies employed identified only five articles containing
evidence that could be used to determine how long measles virus remains infectious in the
air after an infected person leaves a space. All five articles were published between 1965
and 1987, and two were published with the same lead author. None of the five studies
contained primary data that could be used to inform measles contact tracing guidelines for
outdoor spaces.

The first two publications assessing measles survival in the air were experimental studies
published in 1964[8] and 1965[19]. Measles virus survival in the air was used by the authors as
a proxy for the window of time a person may be infected after an infected person leaves an
indoor space. These studies used aerosolized measles virus in a room where air samples

were taken at regular intervals and the surviving virus was estimated by titration. Both
articles were published by the primary author J.G. de Jong one year apart. The first study,
published in 1964, showed that measles virus survived two hours when disseminated in a
room with low humidity (relative humidity 12-15%) but less than 30 minutes in a room with
high humidity (relative humidity 68-70%).[8] This study did not account for factors such as
light or airflow, and temperature was held constant at 20°C (68°F). The second article,
which was an extension of the first, considered additional factors that may affect the
duration of measles infectivity in air, such as additional variations in humidity, slight
variations in temperature, and the presence or absence of artificial light.[19] These findings
reaffirmed conclusions in the previous article and showed that measles virus survives for
less time in the air with increasing levels of humidity when temperature is held constant at
20°C. In the 1965 study, measles was tested at 36-37% relative humidity (RH) and 60-64%
RH, with the virus surviving for at least 60 minutes at the former RH and only 30 minutes at
the latter RH. The presence or absence of artificial light was not found to have any effect on
measles virus survival in the air when temperature and humidity were constant. Measles
virus survival was slightly longer at 15°C than at 20°C, however, the influence of this on the
survival time in the air was not reported.[19]

Twenty years later, three publications reported observational data that included evidence
on the duration of measles virus transmissibility. The 1985 study by Remington et al.
discusses an outbreak of measles in a pediatrician's office. The article identified three
children who developed measles after being exposed in a pediatric office 60 to 75 minutes
after the index case had departed.[11] According to the timeline provided by the authors, one
of the secondary cases arrived at 1:30 PM (60 minutes after the case departed) and
became infected; the next secondary case arrived a little after 1:30 PM (about 65 minutes
after the case departed) and became infected; and the last secondary case arrived around
1:45 PM (about 75 minutes after the case departed) and became infected. Of the three
children that became infected in the 60-75 minutes after the index case departed, two
were under the age of 15 months and one was age 15 months to 18 years. All three were
unvaccinated. A total of 32 children were present within 90 minutes after the case
departed, and only the three children discussed above became infected. An additional 28
children were present in the pediatric office 90 minutes or more after the infected case
departed and none of these individuals developed a measles infection. Interestingly, the
observed attack rates for unvaccinated children present with the index case and
unvaccinated children that arrived less than 90 minutes after the index case are similar,
around 25%, whereas the attack rate for unvaccinated children arriving 90 minutes or more
after the index case is zero. Remington et al. emphasized that factors such as coughing
and inadequate fresh-air ventilation can prolong the duration of airborne infectivity,

increasing the risk of transmission to susceptible individuals who enter the space after an

infected person has left.[11] The second observational publication by Bloch et al. describes a

measles outbreak in a pediatric practice where the index patient, who was in the second

day of rash, was able to transmit measles to other patients through airborne transmission,

even after leaving the office.[10] The article suggests that measles virus can remain

infectious in the air for at least one hour after the index patient has left the area, providing

evidence that measles can be transmitted through airborne spread even in the days after

symptom onset, rather than just in the pre-rash period as previously believed. The specific

duration and time windows are not reported as they had been in the Remington article;

therefore, it is not clear how much greater than one hour the exposure window was. The

final publication summarized how long measles virus remains transmissible after an

infectious case leaves a space in a hospital setting. During this outbreak, individuals who

arrived 29 to 120 minutes after a case was present contracted measles. The minimum

exposure window in this outbreak was observed when an 8-month-old was infected with

measles after entering an emergency room 29 minutes after the index case was no longer

present. The maximum exposure window observed was when a 24-year-old contracted

measles while purchasing medications at the university hospital two hours after an

infected internal medicine resident passed by the pharmacy seeking evaluation for her

illness.[20]

A summary of the results is available in Table 1. Both the smallest and largest exposure

windows were identified from laboratory studies. The smallest infectious exposure window

reported was about 30 minutes, which was dependent on high humidity (68-70%) and

room temperature (20°C) while the largest exposure window was about 120 minutes,

which was dependent on low humidity (12-15%) at room temperature (20°C).[8]

Observational studies reported an exposure window within the range of that reported by

the experimental studies, both within the general range of 60 to 90 minutes. No additional

observational or experimental studies have met the criteria for assessing the exposure

window since 1985.

Discussion

All four hallmark articles that were identified through our search strategy highlight

important features of measles airborne infectivity, though limited in scope. Based on the

combination of formal and informal search strategies, this search is comprehensive and

demonstrates that there is minimal published evidence to guide precise determination of

how long measles remains infectious in the air after an infected individual is no longer

present. It is clear there is a gap in existing evidence on the duration of measles infectivity
in air, and whatever existing research exists is long outdated.

Study Limitations

Search criteria in the systematic review was limited to manuscripts published in English,
potentially excluding useful research in other languages. The search was limited to January
1988 to July 2024 after PCR testing became widespread to reduce potential bias from
improper diagnosis, but may have inadvertently excluded earlier studies that could have
provided valuable insights into airborne transmission. However, our supplemental search
mitigated this risk by including earlier studies. Also, changes in Medical Subject Heading
(MeSH) terms over time may have resulted in the omission of relevant studies that were
indexed under our search terminologies. Additionally, certain older papers lack abstracts
or are available only as scanned PDFs without searchable text, which would not have been
detected by the search engine utilized and would have led these papers to be inadvertently
excluded. Furthermore, while we developed a comprehensive set of search terms, it is
possible that some relevant studies were missed due to variability in terminology used by
authors. These factors may have limited the ability to capture the full breadth of all
available relevant evidence.

Research Needs

Health departments currently rely on CDC measles exposure guidelines based on research
from De Jong,[8,19] Remington,[11] and Bloch[10] between 1964 and 1985. While beneficial,
additional evidence is needed to better understand how long measles remains infectious
in the air after a case is no longer present. This evidence can be gathered through
additional laboratory and observational studies.

Laboratory studies are needed to better understand how the measles virus survives in the
air and on surfaces under different environmental conditions and in various settings.
Current literature only evaluates measles survival at 20°C (68°F) and fails to consider the
diverse factors present in public settings. Additionally, existing studies have only examined
measles survival in controlled environments. Future research should investigate how
measles survives in spaces that mimic real-world environments, such as classrooms,
open-concept offices, grocery stores, restaurants, airplanes, and hospitals. This will help
inform how spatial obstacles and modern HVAC and air filtration systems can influence
the movement and survival of the measles virus in places where people are most likely to
be exposed. Some modeling studies have addressed this to a limited degree. Riley et al. in
1978 developed a mathematical model to analyze a measles outbreak in a suburban
elementary school, highlighting the airborne transmission of the virus.[21]

Additional observational studies gathered from real-world outbreaks are also needed to understand the actual exposure window for measles contacts after cases are no longer present. This information could be gathered through coordinated efforts by state and local health departments conducting measles case investigations in their jurisdictions. Health departments should consider collecting exposure time windows for each contact identified in a case and organizing databases so that epidemiologists can easily determine which contacts became cases. Coordinating entities could then pool de-identified data on the exposure windows for each contact and calculate the attack rates for exposure times.

This systematic review also identified a gap in evidence necessary for contact tracing in outdoor public spaces. Although this systematic review was not limited to indoor settings, the researchers also did not identify any experimental or observational studies that assessed measles virus infectivity in outdoor air spaces that met the other inclusion criteria. Future studies in these settings are needed to inform evidence-based measles contact tracing.

Public Health Implications

Research is needed to clarify how long the measles virus remains transmissible after an infectious case leaves a public space. Robust evidence would enable health departments to refine and standardize evidence-based guidelines rooted in science and tailored to actual risks to optimize contact tracing while minimizing unnecessary interventions. Whether the research confirms the two-hour window or suggests adjustments, such research would improve public health responses' efficiency and effectiveness, fostering trust in evidence-based recommendations.

Acknowledgments

The authors acknowledge our state and local health department partners for inspiring this research. We would like to especially thank those health departments that shared the contact tracing guidance they would follow during a measles outbreak, including the Alameda County Health Department, Chester County Health Department, Los Angeles County Department of Public Health, Marathon County Health Department, Public Health-Dayton & Montgomery, Public Health- Seattle & King County, South Carolina Department of Health, Tennessee Department of Health, and additional state and local health departments that remain anonymous. The research team would like to acknowledge the CDC Insight Net Program for supporting this work. The funders had no input or oversight over the conduct of this work.

Bibliography

1. US CDC. History of Measles. Measles (Rubeola). May 20, 2024. Accessed December 19, 2024. https://www.cdc.gov/measles/about/history.html

2. Mathis AD, Raines K, Masters N, et al. Measles — United States, January 1, 2020–March 28, 2024. *MMWR Morb Mortal Wkly Rep*. 2024;73. doi:10.15585/mmwr.mm7314a1

3. Outbreak response operations during the U.S. measles epidemic, 2017–19. *ResearchGate*. Published online December 9, 2024. doi:10.1186/s12889-021-10652-9

4. US CDC. Infection Prevention and Control Strategies for Seasonal Influenza in Healthcare Settings. Influenza (Flu). November 27, 2024. Accessed January 30, 2025. https://www.cdc.gov/flu/hcp/infection-control/healthcare-settings.html

5. US EPA. Indoor Air and Coronavirus (COVID-19). June 17, 2020. Accessed January 30, 2025. https://www.epa.gov/indoor-air-quality-iaq/indoor-air-and-coronavirus-covid-19

6. US CDC. Clinical Overview of Measles. Measles (Rubeola). January 17, 2025. Accessed January 30, 2025. https://www.cdc.gov/measles/hcp/clinical-overview/index.html

7. US CDC. Measles. Infection Control. November 1, 2024. Accessed March 17, 2025. https://www.cdc.gov/infection-control/hcp/healthcare-personnel-epidemiology-control/measles.html

8. De Jong JG, Winkler KC. Survival of Measles Virus in Air. *Nature*. 1964;201(4923):1054-1055. doi:10.1038/2011054a0

9. Sienko DG, Friedman C, McGee HB, et al. A measles outbreak at university medical settings involving health care providers. *Am J Public Health*. 1987;77(9):1222-1224.

10. Bloch AB, Orenstein WA, Ewing WM, et al. Measles Outbreak in a Pediatric Practice: Airborne Transmission in an Office Setting. *Pediatrics*. 1985;75(4):676-683. doi:10.1542/peds.75.4.676

11. Remington PL, Hall WN, Davis IH, Herald A, Gunn RA. Airborne Transmission of Measles in a Physician's Office. *JAMA*. 1985;253(11):1574-1577. doi:10.1001/jama.1985.03350350068022

12. Biellik RJ, Clements CJ. Strategies for minimizing nosocomial measles transmission. *Bull World Health Organ*. 1997;75(4):367-375.

13. Farrie H. Measles State and Local Response: Challenges, Successes, Innovations, and Standardized Tools for Resiliency. Presented at: Measles Partner Meeting; October 29, 2024; Zoom. N.A

14. Farrie H. *Survey of Health Department Partners*. Johns Hopkins Center for Outbreak
Response Innovation; 2025:1.

15. Farrie H. Measles Processes Key Informant Interview. Published internally March 15,
2024.

16. CDC. Measles Vaccine Recommendations. Measles (Rubeola). October 2, 2024.
Accessed January 30, 2025. https://www.cdc.gov/measles/hcp/vaccine-
considerations/index.html

17. PRISMA statement. PRISMA statement. Accessed December 19, 2024.
https://www.prisma-statement.org

18. The History of PCR - U.S. Accessed December 20, 2024.
https://www.thermofisher.com/us/en/home/brands/thermo-scientific/molecular-
biology/molecular-biology-learning-center/molecular-biology-resource-
library/spotlight-articles/history-pcr.html

19. De Jong JG. The survival of measles virus in air, in relation to the epidemiology of
measles. *Arch Für Gesamte Virusforsch*. 1965;16(1):97-102. doi:10.1007/BF01253797

20. A measles outbreak at University Medical Settings Involving Healthcare Providers.

21. RILEY EC, MURPHY G, RILEY RL. AIRBORNE SPREAD OF MEASLES IN A SUBURBAN
ELEMENTARY SCHOOL. *Am J Epidemiol*. 1978;107(5):421-432.
doi:10.1093/oxfordjournals.aje.a112560

Figure 1. PRISMA Framework with the number of studies identified across each phase.

Table 1. Summary of all publications utilizing primary data that could be used to inform exposure windows for measles contact tracing after an infectious case is no longer present in indoor or outdoor space.

Setting	Environmental Conditions	Infectious Exposure Window	Citation
Laboratory Experiment	Humidity: 12-15% Temperature: 20°C	120 min	De Jong et al. 1964 [a]
	Humidity: 68-70% Temperature: 20°C	<30 min	
	Humidity: 60-64% Temperature: 20°C	30 min	De Jong et al. 1965 [b]
	Humidity: 36-37% Temperature: 20°C	>60 min	
Pediatric Outpatient Office	Humidity: 22% Temperature: 20°C	60-75 min	Remington et al. 1985 [a]
	Humidity: not reported Temperature: not reported	1hr	Bloch et al. 1985 [a]
Hospital Pharmacy	Humidity: not reported Temperature: not reported	29-120 min	Sienko et al. 1987 [a]

[a] Utilized to inform CDC measles contact tracing guidelines.

[b] Not utilized to inform CDC measles contact tracing guidelines.