Title Page

From dysphoria to anhedonia: Age-related shift in the link between cognitive and
affective symptoms

Daniel Harlev[1,2], Aya Vituri[3], Moni Shahar[3], Cam-CAN[4], Noham Wolpe[1,5]

[1]Department of Physical Therapy, The Stanley Steer School of Health Professions, Faculty of
Medical & Health Sciences, Tel Aviv University, Tel Aviv, Israel

[2]Department of Psychiatry, Rambam Health Care Campus, Haifa, Israel

[3]Tel Aviv Center for Artificial Intelligence & Data Science (TAD), Tel Aviv University,
6997801, Israel

[4]Cambridge Centre for Ageing and Neuroscience, University of Cambridge and MRC
Cognition and Brain Sciences Unit, Cambridge, UK.

[5]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv 6997801, Israel.

Send correspondence to Dr Harlev (harlev1@tauex.tau.ac.il), Department of Physical
Therapy, The Stanley Steer School of Health Professions, Faculty of Medicine, Tel Aviv
University, Tel Aviv 6997801, Israel

Keywords: Aging, Depression, Bridging, Cognition, Gray matter volume, Symptoms
Network Analysis

Abstract

Aging affects cognition, emotion, and brain structure, yet how these domains interact remains

unclear. We examined cognitive-affective symptom networks in 756 young (≤45 years) and

1,230 older (≥65 years) adults. Cognition was assessed with the Addenbrooke's Cognitive

Examination Revised, and affective symptoms were assessed with the Hospital Anxiety and

Depression Scale. Graphical LASSO estimated network structure, computing strength

centrality for overall network structure and two independent bridging centrality measures for

measuring cognitive-affective links. While overall network structure was similar across age

groups, depressive (not anxiety) symptoms were more strongly linked to cognition in older

adults. The key bridging symptom shifted from dysphoria in young adults to anhedonia in

older adults. Anhedonia was also associated with the relationship between gray matter

volume and cognition, particularly in older adults. The results suggest an age-related shift in

how cognitive and affective symptoms interact across the lifespan, highlighting the need for

age-specific interventions.

Introduction

Aging leads to significant changes in cognition, emotion, and brain structure. As the general population ages, there is an urgent need to understand how these domains interact over the lifespan (Harada et al., 2013; Liverman et al., 2015; Ram et al., 2011). For example, mental health disorders, such as depression and anxiety, which are highly prevalent in old age (Teachman, 2006), are frequently intertwined with cognitive impairments (Bergmann et al., 2024; Wang & Blazer, 2015). Moreover, the relationships between depression, anxiety and cognition differ across the lifespan (Bergmann et al., 2024).

Cognitive decline with age often co-occurs with affective changes. Common affective changes in ageing, which are also central to the diagnosis of Major Depressive Disorder (MDD) are apathy and anhedonia (Fahed & Steffens, 2021; Husain & Roiser, 2018; Marin, 1991). Apathy reflects a reduction in goal-directed behavior while anhedonia is the reduced ability to enjoy things. These emotional changes are particularly pronounced in Late-Life Depression (LLD), which manifests differently compared to early life depression, with more pronounced cognitive impairments (Unützer, 2007). Neurobiological changes in LLD further illustrate the shared mechanisms linking emotional dysregulation and cognitive dysfunction with age (Alexopoulos, 2019; Bergmann et al., 2024). However, previous studies have primarily examined broad depressive severity rather than specific symptoms, making it difficult to determine which aspects of depression are most relevant for cognitive decline (Bergmann et al., 2024; Du et al., 2022). Given that depression is highly heterogeneous, a symptom-level approach may provide deeper insights into how affective changes shape cognitive function across aging.

Understanding these specific symptom interactions is essential, as cognitive and affective symptoms do not merely co-occur but also influence each other. Mood disorders can exacerbate cognitive decline, while cognitive impairments can heighten the vulnerability to depression (James et al., 2021; Steffens & Potter, 2008; Wang & Blazer, 2015). This complex bidirectional relationship makes it difficult to distinguish between healthy aging processes and neurodegenerative processes, highlighting the need for alternative frameworks to capture and quantify these relationships across the lifespan (Butters et al., 2008; Diniz et al., 2013; Soleimani et al., 2023).

Symptom network analyses investigate how specific symptoms interact with one another (Borsboom & Cramer, 2013; Boschloo et al., 2015). Previous studies investigating these

interactions have focused on younger populations and examined either cognitive or affective

domains separately (Beard et al., 2016; Harlev, Vituri, et al., 2025; Park & Kim, 2020). In

older adults, previous studies have focused on affective symptoms, while excluding cognitive

domains (An et al., 2019; Belvederi Murri et al., 2020; Harlev, Vituri, et al., 2025), and the

few studies combining affective and cognitive domains had methodological limitations. For

example, one study (Bai et al., 2023) used less sensitive cognitive measures like the Mini-

Mental State Examination, while another (Zhou et al., 2023) focused on clinical populations

which limits broader applicability to the general population. Importantly, previous research

lacks consideration of neurobiological measures, such as brain structural integrity measures,

such as gray matter volume (GMV).

Here, we address these gaps by directly comparing cognitive-affective networks in the

general population across the lifespan. We analyzed data from the Cambridge Centre for

Ageing and Neuroscience, which is a population-based cohort of overall healthy individuals

across the lifespan. Cognitive performance was assessed with the Addenbrooke's Cognitive

Examination Revised (ACE-R) to assess performance in key cognitive domains. Affective

symptoms were assessed with the Hospital Anxiety and Depression Scale (HADS). We

compared strength centrality to assess overall network structure and bridge centrality

measures to assess the cognitive-affective links in both young and older adults. We further

tested whether depression and anxiety symptoms differentially contribute to linking cognitive

and affective symptoms across age groups. Lastly, we explored whether the age-related

difference in bridging symptoms moderates the relationship between GMV and cognitive

performance across the lifespan.

Methods

Participants

We analyzed data from the Cambridge Centre for Ageing and Neuroscience (Cam-CAN)—a large-scale project focusing on aging (Shafto et al., 2014). Cam-CAN recruited a population-based sample of approximately 2600 adults aged 18 years and over from the general population via Primary Care Trust's lists within the Cambridge City area in the UK. Ethical approval for the study was granted by the Cambridgeshire 2 Research Ethics Committee (reference: 10/H0308/50), and written informed consent was obtained from all participants prior to their participation. The dataset includes detailed demographic, cognitive, and neuroimaging data.

Participants were categorized into two age groups: young ($\leq$45 years) and older ($\geq$65 years), to test for age-related differences in cognitive-affective networks. Cognitive performance was assessed using the Addenbrooke's Cognitive Examination-Revised (ACE-R) (Mioshi et al., 2006), which evaluates five domains: memory, fluency, language, visuospatial abilities, and orientation. Affective symptoms were measured using the Hospital Anxiety and Depression Scale (HADS) (Bjelland et al., 2002), which includes subscales for depressive (HD1–HD7) and anxiety (HA1–HA7) symptoms. After excluding participants with missing demographic or clinical data, the groups included 756 young adults and 1230 older adults.

Network Analyses

Sparsity-based networks were constructed separately for each age group using the Graphical LASSO method, as done previously (Friedman et al., 2008). We included participants across the full range of HADS scores, including those with low or zero scores, to capture the entire spectrum of affective symptomatology and its relationship with cognitive performance, as done previously (Bai et al., 2023). The use of Graphical LASSO allowed us to include all participants by penalizing weak or noisy connections to ensure network construction was robust across the full range of affective and cognitive scores. Non-significant connections were removed based on the Extended Bayesian Information Criterion (EBIC) (Foygel & Drton, 2010) for model selection. Nodes represented cognitive and affective domains, with edges reflecting regression coefficients. Covariates, including sex and education (number of years in education), were included to account for potentially confounding effects. All non-categorical data were Z-scored before entry into the analysis.

In line with our study goals and with previous research, we computed two key metrics in the
networks (Bringmann et al., 2019; Jones et al., 2021). First, we sought to identify the
importance of each node within the whole network by calculating strength centrality. Strength
centrality quantifies the total connectivity of a node by summing the absolute weights of all
edges connected to it. Higher strength centrality indicates that a node is more integrated
within the network, i.e., it has stronger connections to multiple other nodes. This allowed us
to test whether overall network structure remained similar with age, with similarly important
nodes.

Second, we sought to identify the nodes linking cognitive and affective symptoms by
computing two independent and complementary centrality measures. Betweenness centrality
quantifies how often a node appears on the shortest paths between pairs of other nodes in a
network (where shortest path is the minimum number of steps/lowest-weight sum needed to
travel between two nodes in a network). In other words, betweenness centrality reflects a
node's role as an intermediary. When a node mediates a large proportion of the shortest paths,
it plays a critical role in connecting otherwise separate regions of the network, and is thus
interpreted as a bridging measure (Zhang & Luo, 2017). To complement this approach, we
also computed Bridge Expected Influence (BEI), which measures a node's connections
spanning different pre-defined communities, weighted by their influence on neighboring
nodes (Jones et al., 2021). Unlike betweenness centrality, BEI is domain-sensitive and
requires *a priori* labelling of the different symptom domains (cognitive and affective).

We first assessed the consistency of centrality ranks across age groups using Spearman (rank)
correlation to determine whether the overall structure of centrality values remained similar
between young and older adults. Next, we reported the nodes with the highest centrality value
in each domain (depression, anxiety and cognition) for each age groups to identify possible
age-related differences in the importance of network nodes. Finally, we tested for differences
in centrality measures between depression and anxiety nodes in each group. This tests
whether within each group, there was any significant differences in centrality measures
between depression and anxiety symptoms. To test this, permutation testing with 5,000
iterations was used within each age group to compare centrality measures between
depression-related (HD1–HD7) and anxiety-related (HA1–HA7) nodes. The test was
performed by randomly shuffling the depression and anxiety labels while maintaining the
original network topology, generating a 'null' distribution of expected differences by chance
alone. Statistical significance was determined as the proportion of permutations in which the

shuffled difference exceeded the observed value (with $p < 0.05$ considered statistically
significant).

Post hoc GLM Analyses

Building on the network analyses, we conducted exploratory post hoc Generalized Linear
Models (GLMs) to test whether the age-related difference in bridging symptoms modulated
the relationship between GMV and cognition. While previous research has shown that
depressive symptoms moderate GMV-cognition associations (Du et al., 2022), these studies
did not examine how specific depressive symptoms contribute to this relationship.

The analysis included 733 participants in total, who took part in Cam-CAN Stage 2 and Cam-
CAN Frail, which followed similar neuroimaging protocols in individuals with lower scores
in the ACE-R (Kocagoncu et al., 2022; Shafto et al., 2014). Structural MRI data were
acquired using a 3T Siemens TIM Trio scanner with T1-weighted MPRAGE sequences (TR =
2250 ms, TE = 2.99 ms, TI = 900 ms, flip angle = 9°, isotropic voxel size = 1 mm³) (Taylor et
al., 2017). Preprocessing followed standard protocols in SPM12, including segmentation into
gray matter. To account for individual differences in head size, total intracranial volume
(TIV) was computed from the segmentation, and GMV was included in the model as the ratio
GMV/TIV.

We examined whether anhedonia versus dysphoria moderated the relationship between GMV
and cognitive performance. To quantify this, we computed the anhedonia-to-dysphoria
difference(HD7–HD3), where higher values indicate a greater relative severity of anhedonia
compared to dysphoria (Lee et al., 2019).

The GLM included cognitive performance (total ACE-R score) as the dependent variable.
Age, GMV and anhedonia-to-dysphoria difference were the independent variables. Two-way
interactions were tested to determine whether anhedonia-to-dysphoria difference moderated
the relationship between GMV and cognition across age groups. For consistency with the
network analysis, education and sex were included as covariates, and LASSO regularization
was implemented. To complement the model results. All statistical analyses were conducted
in Python (v3.11.5) (Van Rossum & Drake, 2009).

Results

Participants and item severity

Participant demographics are summarized in Table 1 for both young and older adult groups. Sex distribution was similar across groups, but education level was significantly higher for younger compared to older adults. Depressive symptoms varied between the two groups. Older adults had significantly higher scores on items related to psychomotor slowing and reduced anticipatory anhedonia/enjoyment, while young adults scored higher on measures of dysphoria/cheerfulness and consummatory anhedonia/enjoyment. There were no significant differences between groups for apathy/interest in appearance or laughter. In contrast, anxiety symptoms were consistently higher among young adults across all items. Lastly, young adults had better cognitive performance compared to older adults across all cognitive domains.

Network Analyses

Cognitive-affective networks for young and older adults were constructed for young and older adults. In both age groups, nodes clustered into cognitive and affective domains, with predominantly positive connections within each domain, and largely negative connections between them, although some inter-domain connections were positive (Fig. 1).

To examine the network structure and whether it differs between age groups, we calculated two key centrality measures, namely strength centrality and betweenness centrality. First, strength centrality, which quantifies the overall importance of a node by summing up the overall connection weight of a node, was computed. The pattern of strength centrality values was overall consistent between groups (Fig. 2A). The rank of strength centrality values was indeed highly correlated between young and older adults (Spearman's $\rho = 0.795$, $p = 4.80\text{e-}05$). In terms of their values, the strongest cognitive nodes for both young and older adults were memory (young: 0.848, old: 0.731), fluency (young: 0.694, old: 0.883) and language (young: 1.063, old: 0.991). Among the affective nodes, the nodes with highest strength centrality for both groups were anticipatory anhedonia/enjoyment (HD6) (young: 1.045, old: 1.155) for depression, and worrying thoughts (HA3) (young: 1.031, old: 1.070) for anxiety. A permutation test (n = 5000) comparing mean strength centrality between depression-related nodes (HD1–HD7) and anxiety-related nodes (HA1–HA7) showed no significant differences in either age group (young: mean observed difference = -0.217, $p = 0.618$; old: mean observed difference = -0.207, $p = 0.531$). Together, these results suggest that overall network structure was similar between young and older adults.

Next, we computed two independent and complementary bridging measures. First, we
computed each node's betweenness centrality, which quantifies how often a node appears on
the shortest paths between other nodes. In contrast to strength centrality, betweenness
centrality pattern was different between young and older adults (Fig. 2B). The rank of
betweenness centrality values was not correlated between young and older adults
(Spearman's $\rho = 0.330$, $p = 0.168$). In terms of their value, the symptoms with the highest
betweenness value were dysphoria (HD3) in young adults (0.4575) but anhedonia (HD7) in
older adults (0.2843). A permutation test (n = 5000) comparing mean betweenness centrality
between depression- and anxiety-related nodes showed group-specific differences.
Specifically, in young adults, depression and anxiety nodes did not differ significantly
(observed difference = 0.053, $p = 0.702$). However, in older adults, betweenness centrality
was significantly higher for depression-related nodes compared to anxiety-related nodes
(observed difference = 0.326, $p = 0.033$).

The second bridging measure we computed was BEI, which quantifies bridging based on
predefined node (symptom) labels. The analysis showed similar results to those of
betweenness centrality above (Fig. S1), namely no rank correlation between BEI in young
and older adults (Spearman's $\rho = 0.326$, $p = 0.173$); dysphoria (BEI = -0.0741) and anhedonia
(BEI = -0.2012) had the highest negative BEI values in young and older adults, respectively;
and no differences in young adults (mean observed difference = -0.071, $p = 0.548$) but
significantly higher depression-related BEI compared to anxiety-related BEI in older adults
(mean observed difference = -0.443, $p = 0.0014$). Interestingly, rumination (HA3) showed a
surprisingly positive BEI value in both young (0.0314) and older adults (0.0516)

Bridging symptoms, GMV, and cognitive performance
To examine whether the age-related difference in bridging symptoms moderates the
relationship between GMV and cognitive performance, we conducted a *post hoc* GLM
analysis (Table 2). For the main contrasts of interest, significant effects were observed for the
anhedonia-to-dysphoria difference $\times$ GMV and anhedonia-to-dysphoria difference $\times$ age,
indicating that more anhedonia relative to dysphoria was associated with a more positive
relationship between GMV and cognition. However, the three-way interaction (anhedonia-to-
dysphoria difference $\times$ GMV $\times$ age) was not significant.

Although the three-way interaction was not statistically significant, suggesting that the
relationship between GMV and cognition did not differ systematically across all levels of age

(age included in the GLM as continuous variable), exploratory subgroup analyses revealed

additional meaningful differences. Specifically, in young adults ($\leq$ 45 years old), the

correlation between GMV and cognitive performance was not significant for both people

with higher anhedonia relative to dysphoria ($r = 0.030$, $p = 0.738$) and people with higher

dysphoria relative to anhedonia ($r = -0.256$, $p = 0.151$) (Fig. 3A). In contrast, in older adults

($\geq$ 65 years old) with higher anhedonia relative to dysphoria, GMV was significantly

positively correlated with cognitive performance. ($r = 0.333$, $p = 8.606e-05$), but not for older

adults with higher dysphoria relative to anhedonia ($r = 0.150$, $p = 0.515$) (Fig. 3B). The

correlation between GMV and cognition was indeed significantly stronger in older compared

to young adults for individuals with high anhedonia relative to dysphoria ($z = -2.502$, $p =$

0.012), but not for individuals with higher dysphoria relative to anhedonia ($z = -1.386$, $p =$

0.166). These findings indicate that while the full three-way interaction may lack consistency

to reach statistical significance, subgroup-specific associations—particularly in older adults

with more prominent anhedonia—may still be present.

Discussion

Our study investigated the structure of cognitive-affective symptom network across the lifespan. Overall network structure was similar across groups, Importantly, however, distinct patterns of symptom connectivity emerged for the bridging of affective and cognitive symptoms between age groups. Across two independent bridging measures, dysphoria was the strongest cognitive-affective bridging symptom in young adults, while anhedonia was the strongest bridging symptom for older adults. Moreover, depressive symptoms had significantly higher cognitive-affective bridging values compared to anxiety symptoms, specifically for older adults. Moreover, higher anhedonia-to-dysphoria difference was associated with stronger and more positive relationship between GMV and cognition, with this effect being most pronounced in older adults. These findings suggest an age-related shift in the symptoms that bridge cognitive and affective symptoms.

Model free symptom severity

Cognitive and anxiety items showed a distinct age-related pattern, with anxiety symptoms being more severe in younger adults, while cognitive function was better preserved in this group compared to older adults.

This pattern reflects the generally higher prevalence of anxiety in younger populations within the general population, consistent with prior epidemiological findings (Kessler et al., 2010; Remes et al., 2016). In contrast, depressive symptoms exhibited greater variability, consistent with the large heterogeneity of depression manifestation (Fried, 2017; Goldberg, 2011). Older adults reported more severe psychomotor slowing, while younger adults reported more severe mood-related symptoms. These findings are consistent with known age-related differences in depressive symptom profiles (Schaakxs et al., 2017; Sutin et al., 2013), and our principal aim was to examine cognitive-affective interactions by age using symptom network analyses.

Maintenance of overall cognitive-affective network structure

Strength centrality, which reflects the overall connectivity of nodes, showed remarkable stability across age groups. Within the cognitive nodes, memory and language consistently retained high centrality values in both young and older adults. This likely reflects their role as core cognitive functions that remain relevant regardless of age. Similarly, within the affective nodes, anticipatory anhedonia (HD6) and worrying thoughts (HA3) retained high centrality

values across age groups. This pattern aligns with previous findings indicating that the core
organization of depressive and anxiety symptom networks, including the centrality of key
symptoms, remains largely stable across the lifespan (Harlev, Vituri, et al., 2025).

This cross-domain consistency across the lifespan supports the view that affect and cognition
form an integrated clinical construct that is preserved with age. Indeed, meta-analytic
evidence confirms that depressive symptoms are reliably associated with cognitive control
deficits from childhood through old age (Dotson et al., 2020). This stability suggests that
depression, anxiety, and cognitive symptoms belong to a common clinical syndrome that
remains consistent across age. But while the structure appears similar, the mechanisms
linking these domains may change with age—pointing to potentially distinct pathways.

Age-related shift in cognitive-affective bridging symptoms

Using two independent and complementary bridging centrality measures (betweenness and
BEI), we identified a difference in how affective symptoms integrate with cognition across
age groups. In younger adults, mood-related symptoms, and particularly cheerful feelings,
were strongly connected to cognitive nodes, reflecting the importance of emotional positivity
in cognitive performance during earlier life stages (Knight et al., 2016). In contrast, in older
adults, anhedonia became the principal bridging symptom, reflecting its stronger association
with cognitive deficits in late life (Regier et al., 2013).

A shift from mood-related link to cognitive performance in younger adults to anhedonia-
related link to cognitive performance in older adults may reflect age-related neurobiological
changes in affective regulation. Whereas younger adults show strong reciprocal links between
mood and cognitive function (Elliott et al., 2011), aging is associated with reduced emotional
responsivity but increased motivational deficits (Husain & Roiser, 2018). Consistent with
this, in Late-Life Depression (LLD), apathy—but not dysphoria—exhibits the strongest
association with cognitive dysfunction (Harlev, Singer, et al., 2025).

Building on these results, we conducted a GLM linking cognition with anhedonia-to-
dysphoria, GMV and age. While the three-way interaction between anhedonia-to-dysphoria,
GMV, and age was not significant, post hoc analyses revealed specific patterns that merit
attention. Specifically, a positive association between GMV and cognition was observed in
older adults with greater anhedonia relative to dysphoria. This effect was not evident in

young adults or in individuals with relatively greater dysphoria. The absence of a significant
three-way interaction suggests that these effects do not generalize across all subgroups or
may not follow a consistent interaction pattern across the full range of the sample.
Nonetheless, the subgroup-specific findings suggest that the relationship between brain
structure and cognitive performance may become more tightly coupled in certain clinical or
demographic contexts—such as older individuals experiencing prominent anhedonia.

These results are consistent with observations that in older adults, anhedonia is more closely
linked to cognitive decline and structural brain changes (Turner & Husain, 2022), while in
young adults, emotional reactivity changes are more closely linked to cognitive (Scher et al.,
2005). One possible explanation is cognitive reserve, which reflects individual differences in
the brain's ability to maintain function despite structural decline (Stern, 2009). Notably, the
stronger association between cognition and GMV was seen particularly in older adults.   The
stronger association between lower GMV and poorer cognitive performance in older
individuals with high anhedonia can thus reflect reduced cognitive reserve. As anhedonia
increases, cognition may become more dependent on brain integrity (Capogna et al., 2019;
Zhu et al., 2021).

Beyond structural differences, network-based measures further highlight this shift in
cognitive-affective interactions with aging. While overall network strength remained
comparable between depressive and anxiety symptoms, depressive symptoms in older adults
exhibited stronger connectivity to cognitive nodes compared to anxiety symptoms. This
finding underscores the increasingly tight link between depressive symptoms and cognitive
functioning in older populations (Jellinger, 2023; Koenig et al., 2014).

Together, these findings suggest that in aging, depression—particularly its anhedonic
aspects—may be increasingly intertwined with cognitive decline and underlying brain
vulnerability. Indeed, apathy and anhedonia have been increasingly recognized as key early
features of neurodegenerative conditions, including Alzheimer's disease, Parkinson's disease,
and frontotemporal dementia (Steffens et al., 2022). Their presence in these disorders reflects
disruptions in reward-processing circuits, which are crucial for both cognitive and affective
regulation (Husain & Roiser, 2018). In aging, anhedonia may represent a distinct mechanism
of depression—one more closely tied to neurodegenerative pathways than to mood
dysregulation.

Clinical implications: rethinking late-life depression treatments

If anhedonia reflects a different mechanism of depression in aging, which is linked to
neurodegeneration rather than traditional mood disturbances, then current treatment
approaches may need to be reconsidered. Currently, depression in older adults is treated
similarly to depression in younger adults, despite lower efficacy (Unützer, 2007). While
serotonin-based antidepressants remain standard, anhedonia and apathy may be more strongly
linked to dysfunction in dopamine-mediated reward circuits (Alexopoulos, 2019).
Dopaminergic interventions, cognitive-behavioral strategies emphasizing goal-directed
activity, and neurostimulation approaches have shown promise in targeting motivational
impairments rather than mood dysregulation alone (Yuen et al., 2015). Future research should
investigate whether such approaches can prove efficacious in older adults suffering from
anhedonic depression.

Strengths and limitations

To our knowledge, our study is the first to directly compare differences in cognitive and
affective symptoms in a large population-based study. The results from multiple centrality
measures converged to show both stable and dynamic patterns in cognitive-affective network.
Moreover, by integrating neuroimaging data, the analysis provided supplementary evidence
for a possible biological link underlying cognitive-affective interactions with age.

However, several limitations must be considered. First, the cross-sectional design precludes
causal inferences about the observed shifts in network patterns. Second, our neuroimaging
analyses focused on global GMV rather than region-specific effects, providing a standardized
clinical measure, but limiting insights into the role of specific brain structures in cognitive-
affective interactions. Third, the study did not include a clinical population, which restricts
the generalizability of findings to clinical settings. However, we argue this may better reflect
patterns in the general population. Fourth, reliance on self-reported symptom measures could
introduce biases, particularly in older adults, which would affect the accuracy of symptom
network measures. This limitation is further highlighted by the interpretation of single-item
measures, which may not fully capture the complexity of syndromes like anhedonia and
apathy. This oversimplification, while common to many other symptom network studies, calls
for caution when interpreting the results of these studies.

Conclusion

Depressive symptoms, and particularly anhedonia, become more central in linking cognitive
and affective symptoms with age. Higher anhedonia relative dysphoria was associated with

stronger and more positive relationship between GMV and cognition, particularly in older
adults. These findings highlight how aging alters cognitive-affective interactions and
underscore the need for therapeutic strategies that account for both symptom profiles and
neurobiological factors across the lifespan.

Acknowledgements

Cam-CAN research was supported by the Biotechnology and Biological Sciences Research Council (BB/H008217/1). DH was supported by an Israel Science Foundation Mavri fellowship. NW was supported by an Israel Science Foundation Personal Research Grant (1603/22). We are grateful to the Cam-CAN respondents and their primary care teams in Cambridge for their participation in this study.

Conflict of interest

DH, AV, MS and NW declare no conflict of interest in relation to the subject of this study.

Data and code availability

All data used for this work are publicly available upon signing data sharing agreement on https://cam-can.mrc-cbu.cam.ac.uk/dataset/. Code used to analyse the data and generate the figures is available on https://github.com/dharlev/From-dysphoria-to-anhedonia-Age-related-shift-in-the-link-between-cognitive-and-affective-symptoms.

References

Alexopoulos, G. S. (2019). Mechanisms and treatment of late-life depression. *Translational*

Psychiatry, *9*(1), 188.

An, M. H., Park, S. S., You, S. C., Park, R. W., Park, B., Woo, H. K., Kim, H. K., & Son, S. J.

(2019). Depressive symptom network associated with comorbid anxiety in late-life

depression. *Frontiers in Psychiatry*, *10*, 856.

Bai, W., Zhang, J., Smith, R. D., Cheung, T., Su, Z., Ng, C. H., Zhang, Q., & Xiang, Y.-T.

(2023). Inter-relationship between cognitive performance and depressive symptoms

and their association with quality of life in older adults: A network analysis based on

the 2017–2018 wave of Chinese Longitudinal Healthy Longevity Survey (CLHLS).

Journal of Affective Disorders, *320*, 621–627.

https://doi.org/10.1016/j.jad.2022.09.159

Beard, C., Millner, A. J., Forgeard, M. J. C., Fried, E. I., Hsu, K. J., Treadway, M. T.,

Leonard, C. V., Kertz, S. J., & Björgvinsson, T. (2016). Network analysis of

depression and anxiety symptom relationships in a psychiatric sample. *Psychological*

Medicine, *46*(16), 3359–3369. https://doi.org/10.1017/S0033291716002300

Belvederi Murri, M., Amore, M., Respino, M., & Alexopoulos, G. S. (2020). The symptom

network structure of depressive symptoms in late-life: Results from a European

population study. *Molecular Psychiatry*, *25*(7), 1447–1456.

https://doi.org/10.1038/s41380-018-0232-0

Bergmann, E., Harlev, D., & Wolpe, N. (2024). Depressive symptoms are linked to age-

specific neuroanatomical and cognitive variations. *Journal of Affective Disorders*,

S0165-0327(24)01772-5. https://doi.org/10.1016/j.jad.2024.10.077

Bjelland, I., Dahl, A. A., Haug, T. T., & Neckelmann, D. (2002). The validity of the Hospital

Anxiety and Depression Scale: An updated literature review. *Journal of*

Psychosomatic Research, *52*(2), 69–77. https://doi.org/10.1016/S0022-3999(01)00296-3

Borsboom, D., & Cramer, A. O. J. (2013). Network analysis: An integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology*, *9*, 91–121. https://doi.org/10.1146/annurev-clinpsy-050212-185608

Boschloo, L., Borkulo, C. D. van, Rhemtulla, M., Keyes, K. M., Borsboom, D., & Schoevers, R. A. (2015). The Network Structure of Symptoms of the Diagnostic and Statistical Manual of Mental Disorders. *PLOS ONE*, *10*(9), e0137621. https://doi.org/10.1371/journal.pone.0137621

Bringmann, L. F., Elmer, T., Epskamp, S., Krause, R. W., Schoch, D., Wichers, M., Wigman, J. T. W., & Snippe, E. (2019). What do centrality measures measure in psychological networks? *Journal of Abnormal Psychology*, *128*(8), 892–903. https://doi.org/10.1037/abn0000446

Butters, M. A., Young, J. B., Lopez, O., Aizenstein, H. J., Mulsant, B. H., Reynolds Iii, C. F., DeKosky, S. T., & Becker, J. T. (2008). Pathways linking late-life depression to persistent cognitive impairment and dementia. *Dialogues in Clinical Neuroscience*, *10*(3), 345–357. https://doi.org/10.31887/DCNS.2008.10.3/mabutters

Capogna, E., Manca, R., De Marco, M., Hall, A., Soininen, H., & Venneri, A. (2019). Understanding the effect of cognitive/brain reserve and depression on regional atrophy in early Alzheimer's disease. *Postgraduate Medicine*, *131*(7), 533–538. https://doi.org/10.1080/00325481.2019.1663127

Diniz, B. S., Butters, M. A., Albert, S. M., Dew, M. A., & Reynolds, C. F. (2013). Late-life depression and risk of vascular dementia and Alzheimer's disease: Systematic review and meta-analysis of community-based cohort studies. *The British Journal of Psychiatry*, *202*(5), 329–335.

Dotson, V. M., McClintock, S. M., Verhaeghen, P., Kim, J. U., Draheim, A. A., Syzmkowicz, S. M., Gradone, A. M., Bogoian, H. R., & Wit, L. D. (2020). Depression and Cognitive Control across the Lifespan: A Systematic Review and Meta-Analysis. *Neuropsychology Review*, *30*(4), 461–476. https://doi.org/10.1007/s11065-020-09436-6

Du, Y., Yan, F., Zhao, L., Fang, Y., Qiu, Q., Wei, W., Wang, J., Tang, Y., Lin, X., & Li, X. (2022). Depression symptoms moderate the relationship between gray matter volumes and cognitive function in patients with mild cognitive impairment. *Journal of Psychiatric Research*, *151*, 516–522. https://doi.org/10.1016/j.jpsychires.2022.05.017

Elliott, R., Zahn, R., Deakin, J. F. W., & Anderson, I. M. (2011). Affective Cognition and its Disruption in Mood Disorders. *Neuropsychopharmacology*, *36*(1), 153–182. https://doi.org/10.1038/npp.2010.77

Fahed, M., & Steffens, D. C. (2021). Apathy: Neurobiology, Assessment and Treatment. *Clinical Psychopharmacology and Neuroscience*, *19*(2), 181–189. https://doi.org/10.9758/cpn.2021.19.2.181

Fossati, P., Coyette, F., Ergis, A.-M., & Allilaire, J.-F. (2002). Influence of age and executive functioning on verbal memory of inpatients with depression. *Journal of Affective Disorders*, *68*(2), 261–271. https://doi.org/10.1016/S0165-0327(00)00362-1

Foygel, R., & Drton, M. (2010). Extended Bayesian information criteria for Gaussian graphical models. *Advances in Neural Information Processing Systems*, *23*. https://proceedings.neurips.cc/paper/2010/hash/072b030ba126b2f4b2374f342be9ed44-Abstract.html

Fried, E. (2017). Moving forward: How depression heterogeneity hinders progress in treatment and research. *Expert Review of Neurotherapeutics*, *17*(5), 423–425. https://doi.org/10.1080/14737175.2017.1307737

Friedman, J., Hastie, T., & Tibshirani, R. (2008). Sparse inverse covariance estimation with

the graphical lasso. *Biostatistics*, *9*(3), 432–441.

https://doi.org/10.1093/biostatistics/kxm045

Goldberg, D. (2011). The heterogeneity of "major depression". *World Psychiatry*, *10*(3), 226–

228. https://doi.org/10.1002/j.2051-5545.2011.tb00061.x

Harada, C. N., Natelson Love, M. C., & Triebel, K. L. (2013). Normal cognitive aging.

Clinics in Geriatric Medicine, *29*(4), 737–752.

https://doi.org/10.1016/j.cger.2013.07.002

Harlev, D., Singer, S., Goldshalger, M., Wolpe, N., & Bergmann, E. (2025). Acoustic speech

features are associated with late-life depression and apathy symptoms: Preliminary

findings. *Alzheimer's & Dementia (Amsterdam, Netherlands)*, *17*(1), e70055.

https://doi.org/10.1002/dad2.70055

Harlev, D., Vituri, A., Shahar, M., & Wolpe, N. (2025). *Depression and Anxiety Symptom*

Networks Across the Lifespan (p. 2025.02.17.25322400). medRxiv.

https://doi.org/10.1101/2025.02.17.25322400

Husain, M., & Roiser, J. P. (2018). Neuroscience of apathy and anhedonia: A transdiagnostic

approach. *Nature Reviews Neuroscience*, *19*(8), 470–484.

James, T. A., Weiss-Cowie, S., Hopton, Z., Verhaeghen, P., Dotson, V. M., & Duarte, A.

(2021). Depression and episodic memory across the adult lifespan: A meta-analytic

review. *Psychological Bulletin*, *147*(11), 1184–1214.

https://doi.org/10.1037/bul0000344

Jellinger, K. A. (2023). The heterogeneity of late-life depression and its pathobiology: A brain

network dysfunction disorder. *Journal of Neural Transmission*, *130*(8), 1057–1076.

https://doi.org/10.1007/s00702-023-02648-z

Jones, P. J., Ma, R., & McNally, R. J. (2021). Bridge Centrality: A Network Approach to

Understanding Comorbidity. *Multivariate Behavioral Research*, *56*(2), 353–367.

https://doi.org/10.1080/00273171.2019.1614898

Kessler, R. C., Ruscio, A. M., Shear, K., & Wittchen, H.-U. (2010). Epidemiology of Anxiety

Disorders. In M. B. Stein & T. Steckler (Eds.), *Behavioral Neurobiology of Anxiety*

and Its Treatment (pp. 21–35). Springer. https://doi.org/10.1007/7854_2009_9

Knight, B. G., Rastegar, S., & Kim, S. (2016). Chapter 15 - Age Differences in the

Connection of Mood and Cognition: Evidence from Studies of Mood Congruent

Effects. In K. W. Schaie & S. L. Willis (Eds.), *Handbook of the Psychology of Aging*

(Eighth Edition) (pp. 279–302). Academic Press. https://doi.org/10.1016/B978-0-12-

411469-2.00015-7

Kocagoncu, E., Nesbitt, D., Emery, T., Hughes, L. E., Henson, R. N., & Rowe, J. B. (2022).

Neurophysiological and brain structural markers of cognitive frailty differ from

Alzheimer's disease. *Journal of Neuroscience*, *42*(7), 1362–1373.

Koenig, A. M., Bhalla, R. K., & Butters, M. A. (2014). Cognitive Functioning and Late-Life

Depression. *Journal of the International Neuropsychological Society*, *20*(5), 461–467.

https://doi.org/10.1017/S1355617714000198

Lee, J. R., Suh, S. W., Han, J. W., Byun, S., Kwon, S. J., Lee, K. H., Kwak, K. P., Kim, B. J.,

Kim, S. G., Kim, J. L., Kim, T. H., Ryu, S.-H., Moon, S. W., Park, J. H., Lee, D.-W.,

Youn, J. C., Lee, D. Y., Lee, S. B., Lee, J. J., … Kim, K. W. (2019). Anhedonia and

Dysphoria Are Differentially Associated with the Risk of Dementia in the Cognitively

Normal Elderly Individuals: A Prospective Cohort Study. *Psychiatry Investigation*,

16(8), 575–580. https://doi.org/10.30773/pi.2019.06.07

Liverman, C. T., Yaffe, K., & Blazer, D. G. (2015). *Cognitive aging: Progress in*

understanding and opportunities for action.

https://books.google.com/books?hl=en&lr=&id=wEZmCgAAQBAJ&oi=fnd&pg=PT

15&dq=Aging+is+a+complex+biological+process+accompanied+by+gradual+change

s+in+various+physiological+and+psychological+domains.+These+changes+influence

+cognitive+abilities,+emotional+regulation,+and+mental+health,+often+leading+to+

variability+in+individual+aging+trajectories&ots=qj-

nhGOYrN&sig=Di1hAx4tdzB982CDs6hO-4Nub9k

Marin, R. S. (1991). Apathy: A neuropsychiatric syndrome. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *3*(3), 243–254. https://doi.org/10.1176/jnp.3.3.243

Mioshi, E., Dawson, K., Mitchell, J., Arnold, R., & Hodges, J. R. (2006). The Addenbrooke's Cognitive Examination Revised (ACE-R): A brief cognitive test battery for dementia screening. *International Journal of Geriatric Psychiatry*, *21*(11), 1078–1085. https://doi.org/10.1002/gps.1610

Park, S.-C., & Kim, D. (2020). The Centrality of Depression and Anxiety Symptoms in Major Depressive Disorder Determined Using a Network Analysis. *Journal of Affective Disorders*, *271*, 19–26. https://doi.org/10.1016/j.jad.2020.03.078

Ram, N., Gerstorf, D., Lindenberger, U., & Smith, J. (2011). Developmental change and intraindividual variability: Relating cognitive aging to cognitive plasticity, cardiovascular lability, and emotional diversity. *Psychology and Aging*, *26*(2), 363–371. https://doi.org/10.1037/a0021500

Regier, D. A., Kuhl, E. A., & Kupfer, D. J. (2013). The DSM 5: Classification and criteria changes. *World Psychiatry*, *12*(2), 92–98.

Remes, O., Brayne, C., van der Linde, R., & Lafortune, L. (2016). A systematic review of reviews on the prevalence of anxiety disorders in adult populations. *Brain and Behavior*, *6*(7), e00497. https://doi.org/10.1002/brb3.497

Schaakxs, R., Comijs, H. C., Lamers, F., Beekman, A. T. F., & Penninx, B. W. J. H. (2017).

Age-related variability in the presentation of symptoms of major depressive disorder.

Psychological Medicine, *47*(3), 543–552.

https://doi.org/10.1017/S0033291716002579

Scher, C. D., Ingram, R. E., & Segal, Z. V. (2005). Cognitive reactivity and vulnerability:

Empirical evaluation of construct activation and cognitive diatheses in unipolar

depression. *Clinical Psychology Review*, *25*(4), 487–510.

https://doi.org/10.1016/j.cpr.2005.01.005

Shafto, M. A., Tyler, L. K., Dixon, M., Taylor, J. R., Rowe, J. B., Cusack, R., Calder, A. J.,

Marslen-Wilson, W. D., Duncan, J., & Dalgleish, T. (2014). The Cambridge Centre

for Ageing and Neuroscience (Cam-CAN) study protocol: A cross-sectional, lifespan,

multidisciplinary examination of healthy cognitive ageing. *BMC Neurology*, *14*, 1–25.

Soleimani, L., Hirst, A. K., Gilsanz, P., Whitmer, R. A., Corrada, M. M. M., & Beeri, M. S.

(2023). Association of depression dimensions with cognitive functioning in

community-dwelling oldest old adults of LifeAfter90 study. *Alzheimer's & Dementia*,

19(S18), e079455. https://doi.org/10.1002/alz.079455

Steffens, D. C., & Potter, G. G. (2008). Geriatric depression and cognitive impairment.

Psychological Medicine, *38*(2), 163–175.

Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, *47*(10), 2015–2028.

https://doi.org/10.1016/j.neuropsychologia.2009.03.004

Sutin, A. R., Terracciano, A., Milaneschi, Y., An, Y., Ferrucci, L., & Zonderman, A. B. (2013).

The Trajectory of Depressive Symptoms Across the Adult Life Span. *JAMA*

Psychiatry, *70*(8), 803–811. https://doi.org/10.1001/jamapsychiatry.2013.193

Taylor, J. R., Williams, N., Cusack, R., Auer, T., Shafto, M. A., Dixon, M., Tyler, L. K., Cam-

CAN, & Henson, R. N. (2017). The Cambridge Centre for Ageing and Neuroscience

572 (Cam-CAN) data repository: Structural and functional MRI, MEG, and cognitive data

573 from a cross-sectional adult lifespan sample. *NeuroImage*, *144*, 262–269.

574 https://doi.org/10.1016/j.neuroimage.2015.09.018

575 Teachman, B. A. (2006). Aging and negative affect: The rise and fall and rise of anxiety and

576 depression symptoms. *Psychology and Aging*, *21*(1), 201–207.

577 https://doi.org/10.1037/0882-7974.21.1.201

578 Turner, V., & Husain, M. (2022). Anhedonia in Neurodegenerative Diseases. In D. A.

579 Pizzagalli (Ed.), *Anhedonia: Preclinical, Translational, and Clinical Integration* (Vol.

580 58, pp. 255–277). Springer International Publishing.

581 https://doi.org/10.1007/7854_2022_352

582 Unützer, J. (2007). Late-Life Depression. *New England Journal of Medicine*, *357*(22), 2269–

583 2276. https://doi.org/10.1056/NEJMcp073754

584 Van Rossum, G., & Drake, F. L. (2009). *Python/C Api manual-python 3*. CreateSpace.

585 https://dl.acm.org/doi/abs/10.5555/1593487

586 Wang, S., & Blazer, D. G. (2015). Depression and Cognition in the Elderly. *Annual Review of*

587 *Clinical Psychology*, *11*(1), 331–360. https://doi.org/10.1146/annurev-clinpsy-

588 032814-112828

589 Yuen, G. S., Bhutani, S., Lucas, B. J., Gunning, F. M., AbdelMalak, B., Seirup, J. K.,

590 Klimstra, S. A., & Alexopoulos, G. S. (2015). Apathy in late-life depression:

591 Common, persistent, and disabling. *The American Journal of Geriatric Psychiatry:*

592 *Official Journal of the American Association for Geriatric Psychiatry*, *23*(5), 488–

593 494. https://doi.org/10.1016/j.jagp.2014.06.005

594 Zhang, J., & Luo, Y. (2017). *Degree Centrality, Betweenness Centrality, and Closeness*

595 *Centrality in Social Network*. 300–303. https://doi.org/10.2991/msam-17.2017.68

Zhou, J., Zhou, J., Feng, L., Feng, Y., Xiao, L., Chen, X., Yang, J., & Wang, G. (2023). The

associations between depressive symptoms, functional impairment, and quality of life,

in patients with major depression: Undirected and Bayesian network analyses.

Psychological Medicine, *53*(14), 6446–6458.

https://doi.org/10.1017/S0033291722003385

Zhu, X., Ward, J., Cullen, B., Lyall, D. M., Strawbridge, R. J., Lyall, L. M., & Smith, D. J.

(2021). Phenotypic and genetic associations between anhedonia and brain structure in

UK Biobank. *Translational Psychiatry*, *11*(1), 1–8. https://doi.org/10.1038/s41398-

021-01522-4

Tables and Figures

Variable	Young Adults (mean ± SD or %)	Older Adults (mean ± SD or %)	Test Statistic	p-value (FDR corrected)
Age	33.49 ± 6.87	79.29 ± 7.41		
Sex	42.9%	43.5%	1.26	0.557
Education	None: 1.7%; GCSE: 4.1%; A Level: 8.1%; University: 86.1%	None: 29.5%; GCSE: 7.7%; A Level 3.1%; University: 59.7%	348.68	<0.001
Anhedonia/Enjoyment (HD1)	0.46 ± 0.67	0.61 ± 0.78	-4.65	<0.001
Laughter (HD2)	0.26 ± 0.50	0.27 ± 0.54	-0.25	0.802
Dysphoria/Cheerful Feeling (HD3)	0.35 ± 0.54	0.27 ± 0.52	3.11	0.002
Psychomotor retardation/Slowed Down (HD4)	0.70 ± 0.72	1.36 ± 0.93	-16.47	<0.001
Apathy/Loss of interest in Appearance (HD5)	0.50 ± 0.74	0.47 ± 0.71	0.85	0.434
Anticipatory anhedonia/Look Forward to Things (HD6)	0.37 ± 0.65	0.53 ± 0.72	-4.91	<0.001
Anhedonia/Enjoyment of Book/Media (HD7)	0.28 ± 0.63	0.19 ± 0.52	3.52	<0.001
Feeling of Tension (HA1)	1.06 ± 0.65	0.78 ± 0.62	9.52	<0.001
Frightened Feeling	0.69 ± 0.81	0.53 ± 0.74	4.38	<0.001

(HA2)				
Worrying Thoughts (HA3)	0.98 ± 0.87	0.76 ± 0.80	5.86	<0.001
Relaxed Feeling (HA4)	0.84 ± 0.68	0.60 ± 0.63	7.93	<0.001
Butterflies in Stomach (HA5)	0.61 ± 0.64	0.43 ± 0.59	6.49	<0.001
Restless Feeling (HA6)	1.21 ± 0.90	0.92 ± 0.82	7.36	<0.001
Feeling of Panic (HA7)	0.54 ± 0.67	0.49 ± 0.65	1.68	0.107
Memory	0.93 ± 0.10	0.82 ± 0.17	15.81	<0.001
Fluency	0.89 ± 0.13	0.78 ± 0.19	15.04	<0.001
Language	0.94 ± 0.10	0.92 ± 0.10	3.65	<0.001
Visuospatial	0.98 ± 0.05	0.91 ± 0.14	13.52	<0.001
Orientation	0.97 ± 0.06	0.93 ± 0.10	10.15	<0.001

Table 1. Summary of demographics and item-wise comparisons between young and
older adults. Demographic, cognitive, and affective items for young and older adults are
presented. Education was categorized according to the English system: "None" (no education
beyond age 16), "GCSE" (General Certificate of Secondary Education), "A Levels" (General
Certificate of Education Advanced Level), and "University" (undergraduate or graduate
degree). Continuous variables, including age, and cognitive and affective measures, are
reported as mean ± standard deviation and compared using independent t-tests. Categorical
variables, such as sex distribution and education, are expressed as percentages and compared
with chi-square tests. All *p-values* were corrected for multiple comparisons using FDR
correction.

Predictor	β estimate	SE	t-statistic	p-value	95% CI (Lower)	95% CI (Upper)
Intercept	-0.6187	0.101	-6.149	1.29e-09	-0.816	-0.421
Anhedonia to dysphoria difference)	-0.0453	0.041	-1.106	0.269	-0.126	0.035
Age	-0.2496	0.055	-4.530	6.90e-06 ***	-0.358	-0.141
GMV	0.1274	0.056	2.283	0.023 *	0.018	0.237
Anhedonia-to-dysphoria difference× GMV	0.1329	0.056	2.377	0.018 *	0.023	0.243
Anhedonia-to-dysphoria difference × Age	0.1198	0.057	2.108	0.035 *	0.008	0.231
GMV × Age	0.1858	0.033	5.678	1.98e-08 ***	0.122	0.250
anhedonia-to-dysphoria difference× GMV × Age	-0.0069	0.028	-0.248	0.804	-0.061	0.048
Sex	-0.0300	0.065	-0.461	0.645	-0.158	0.098
Education	0.3062	0.033	9.171	4.80e-19 ***	0.241	0.372

Table 2. Generalized Linear Model (GLM) results for predictors of ACE-R. Results from
a Generalized Linear Model (GLM) predicting cognitive performance (ACE-R) based on
anhedonia-to-dysphoria difference (calculated as (HD7-HD3), age, GMV (calculated as
GMV-to-TIV ratio), and their interactions. All continuous variables were z-scored prior to

model entry. The model includes main effects for anhedonia-to-dysphoria difference, age,

GMV, sex, and education, as well as two-way and three-way interaction terms. Significance

levels: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Model fit: $R^2 = 0.319$, Adjusted $R^2 = 0.310$.

Figure 1. Sparsity-based networks for young and older adults. A) Cognitive-affective symptom network for young adults. Each Node represents a cognitive (blue) or affective (red) item, and edges representing conditional relationships estimated via the Graphical LASSO method. Green edges indicate positive relationships, and orange edges indicate negative relationships. Edge thickness corresponds to the strength of the relationships. B) Same as (A) but for older adults.

Figure 2. Centrality measures across age groups. A) Strength centrality for young (purple) and older adults (green). Strength centrality represents the overall connectivity of each node, calculated as the sum of absolute edge weights connected to the node. Higher values indicate nodes with greater influence across the network. The centrality measures were derived from sparsity-based networks, calculated using the Graphical LASSO method with extended Bayesian Information Criteria thresholding. The x-axis represents centrality values, and the y-axis lists the nodes. B) Same as (A), but for betweenness centrality for young (purple) and older adults (green). Betweenness centrality represents the importance of a node in a network based on how often it appears on the shortest paths between other nodes. A node with high betweenness centrality acts as a key connector within the network.

646
647

Figure 3. Associations between cognitive performance and gray matter volume as a
function of anhedonia/dysphoria by age. A) Scatter plot showing the relationship between
cognitive performance, measured with the Addenbrooke's Cognitive Examination Score –
Revised (ACE-R) and gray matter volume (GMV) for younger adults (≤ 45 years old). Gray
matter volume is shown as a proportion of total intracranial volume. For illustration, data
points are plotted separately for individuals with high anhedonia relative dysphoria (blue) and
high dysphoria relative to anhedonia (red), based on a median split. In the statistical model,
both age and anhedonia-to-dysphoria difference (HD7 - HD3) were analyzed as continuous
variables. Regression lines represent fitted trends for each group, with shaded areas indicating
95% confidence intervals. B) Same as (A) but for older adults (≥ 65 years).

Supplementary Material

Supplementary Figures

Figure S1. Bridge Expected Influence for all nodes across age groups. Bridge Expected
Influence (BEI) for young (purple) and older adults (green). BEI quantifies a node's role in
explicitly bridging the (labelled) cognitive and affective domains by calculating the sum of
edge weights connecting a node to nodes in another domain. Higher values indicate stronger
cross-domain connectivity.