Title: Elevated Cardiovascular Risk Factors and Chronic Disease Mortality in the Caribbean: A

Cross-Sectional Study

Authors: Cesar Barrabi, Celia Foster

Affiliation: Western Atlantic University School of Medicine

Abstract

Cardiovascular disease (CVD) disproportionately impacts the Caribbean, accounting for significant

premature mortality and morbidity, especially from stroke and hypertensive heart disease. Despite global

advancements in CVD prevention and management, the Caribbean region has experienced stagnating or

worsening outcomes, particularly among women. Previous studies have largely aggregated Caribbean

data with Latin America, obscuring important regional and sex-specific differences, thereby limiting

targeted interventions. In this study, we analyzed sex-specific CVD mortality, risk exposures, and

systemic healthcare indicators across 15 Caribbean nations, comparing these to North America (United

States and Canada). Data from the 2019 PAHO Enlace Portal revealed significantly higher CVD mortality

in the Non-Latin Caribbean compared to North America (196.7 vs. 122.6 deaths per 100,000).

Hemorrhagic and ischemic stroke mortality were notably elevated ($p < .001$), alongside hypertensive

heart disease deaths, especially in women (24.8 vs. 5.4 per 100,000; $p < .01$). Caribbean populations

exhibited lower alcohol consumption but paradoxically higher alcohol-related deaths. Additionally,

Caribbean men showed significantly higher undiagnosed hypertension (47.6% vs. 20.8%; $p < .001$), and

both sexes had significantly elevated kidney disease mortality. Healthcare infrastructure indicators

highlighted critical regional deficiencies, including inadequate NCD guidelines and preventive healthcare

measures. These findings underscore the urgent need for comprehensive policy reforms that prioritize

equitable healthcare access, culturally and biologically informed guidelines, and integrated chronic

disease management approaches. Addressing systemic healthcare gaps identified in this study represents

an essential step toward reducing CVD disparities and improving population health in the Caribbean.

Introduction

Cardiovascular disease (CVD) is a leading cause of premature mortality in the Caribbean, with stroke and

hypertensive heart disease contributing disproportionately to the region's overall burden (1, 2). Yet unlike

other regions in the Americas, the Caribbean has seen little progress in reducing CVD mortality. Women,

in particular, face elevated rates of hypertensive heart disease, stroke-related deaths, and years of life lost

(3). While several studies have examined CVD trends across Latin America and the Caribbean, most

aggregate the data or omit sex-specific insights, limiting their usefulness for targeted intervention (4).

This challenge is compounded by limited funding, poor research infrastructure, and minimal collaboration

across Caribbean nations, all of which continue to restrict regional reporting, surveillance, and investment

in CVD research (5).

High-income countries such as the United States and Canada have made substantial gains in reducing

CVD mortality through decades of investment in prevention, early detection, and health systems

strengthening (4, 6). Robust surveillance systems, integrated primary care, population-level interventions,

and comprehensive clinical guidelines have contributed to steady declines in deaths from ischemic heart

disease and stroke. However, even these nations continue to face challenges, including significant

disparities in healthcare access among rural and Indigenous populations, underscoring that progress has

not been universally achieved (6). Additionally, demographic projections suggest CVD burdens in the

United States will continue to rise due to an aging population and increased obesity, potentially

overwhelming healthcare resources and exacerbating existing inequalities (7).

Yet even within these high-income settings, disparities persist. Women and racialized populations remain

underdiagnosed, underrepresented in clinical trials, and less likely to receive evidence-based treatment (8,

9). Sex-specific differences in symptom presentation and biological risk, combined with provider bias and systemic barriers, contribute to poorer outcomes in these groups. These inequities are compounded by social determinants of health, including systemic racism, economic marginalization, and unequal access to care—factors that are even more pronounced in the Caribbean context.

In this study, we analyzed sex-specific CVD mortality and risk factor profiles across 30 countries in the Americas, with a particular focus on the Non-Latin Caribbean. Using publicly available population health data, we examined differences in disease burden, risk exposures, and structural contributors to CVD outcomes. This regionally disaggregated approach highlights where the Caribbean stands relative to high-income comparators. Our findings reveal that the Non-Latin Caribbean is the most affected subregion in the hemisphere and expose a disconnect between traditional risk factors and CVD mortality, particularly among women. These results underscore the need for targeted, data-informed interventions that account for both biological and systemic contributors to CVD in underserved populations.

Methods

Ethical considerations

This analysis relied solely on publicly accessible, aggregated data from international agencies and did not involve any individual-level or human subjects research. As such, ethical approval was not necessary for this study.

Study Design and Data Collection

This was a cross-sectional study that examined sex-specific CVD outcomes across 15 Caribbean countries. All data were obtained from the Pan American Health Organization's (PAHO) Enlace Data Portal, which compiles standardized national health statistics across the Americas (10). The United States

and Canada were included as high-income comparator countries. Data from 2019 were used exclusively,

as this was the most recent year with comprehensive coverage across all indicators.

The dataset included mortality measures for major CVD conditions—such as ischemic heart disease,

stroke, hypertensive heart disease, cardiomyopathy, and rheumatic heart disease—as well as select kidney

diseases. Indicators were disaggregated by sex and included deaths and years of life lost (YLLs).

Behavioral and clinical risk factor data were also extracted, including smoking rates, alcohol

consumption, lipid profiles, physical activity, and hypertension (HTN) prevalence, diagnosis, treatment,

and control.

These data were selected due to their consistency, regional comparability, and relevance to health

disparities research in small island nations where individual-level datasets are limited or unavailable.

Data analysis

All data cleaning, visualization, and statistical analyses were conducted using Microsoft Excel and

Python. Descriptive statistics were used to summarize health outcome and risk factor distributions across

countries, disaggregated by sex where available.

A geographic map was created in QGIS 3.34 T using GADM shapefiles to visualize age-standardized

CVD mortality rates across Caribbean subregions performed by Fiver professional Ayesha

Suraweera. No statistical analyses were performed in QGIS; the map presents descriptive values

extracted from PAHO (11, 12).

To assess sex-based differences within the Caribbean (CAR), two-tailed Student's t-tests were performed

to compare male and female values across multiple health indicators, including mortality and clinical risk

factors. For regional comparisons, Welch's t-tests were used to test differences between CAR countries

and North America (NA), defined here as the United States and Canada. Statistical significance was

defined as $p < 0.05$ (13).

To minimize bias from excessive missingness, we first excluded countries with less than 75% data

availability across the selected variables. For the sole purpose of conducting Pearson correlation analysis,

we then applied Multivariate Imputation by Chained Equations (MICE) using Python's IterativeImputer

from scikit-learn (14). This method performs iterative multivariate regression, estimating missing values

by modeling each variable as a function of the others. Correlation analysis was conducted on the imputed

dataset to assess linear relationships between continuous variables. To avoid redundancy, only unique

variable pairings were reported, and results are presented here (S1 Table).

Results

The Caribbean has Elevated CVD Mortality Compared to North America

To understand how CVD outcomes differ by region, we began by mapping age-standardized CVD

mortality rates across the Americas (Figure 1). The Non-Latin Caribbean stood out with the highest

mortality rate (196.7 per 100,000), exceeding rates in North America (122.6), the Andean Area (128.9),

the Southern Cone and Brazil (151.4), and Central America, Mexico, and the Latin Caribbean (169.6).

These findings reinforce the notion that CVD is not only more prevalent in the Caribbean but also more

fatal, highlighting the urgent need for regional prioritization.

Building on this regional context, we then examined sex-disaggregated data on hemorrhagic and ischemic

stroke disease mortality between the CAR and NA (Figure 2A-B). Hemorrhagic stroke mortality in men

was nearly five times higher in CAR than in NA (46.6 vs. 10.0; $p < .001$), with similarly elevated rates

among women (37.1 vs. 8.0; $p < .01$). Ischemic stroke mortality was also significantly higher in CAR for

both sexes ($p < .001$).

Hypertensive heart disease showed notable disparities as well (Figure 2C). Mortality was over three times

higher in CAR than NA for women (24.8 vs. 5.4; p < .01), with a similar trend in men that was trending

toward significance (26.6 vs. 7.6; p = 0.05).

Next, we analyzed the rates of cardiomyopathy, myocarditis, and endocarditis (Figure 2D). While

mortality and years of life lost were consistently higher in the Caribbean, none of these differences

reached statistical significance. Still, the direction of the differences suggests a trend toward elevated

CVD burden in CAR.

Together, these findings highlight a regional CVD mortality gap that disproportionately affect Caribbean

populations, especially for stroke and hypertensive heart disease. This underscores the need for more

targeted surveillance, prevention, and health system responses in small island settings.

Behavioral and Lifestyle Risk Factors in the Caribbean: Gendered Patterns and Regional

Differences

To further explore factors influencing regional CVD mortality disparities, we assessed differences in

behavioral and lifestyle risk factors between the CAR and NA such as alcohol consumption per capita and

Alcohol-related deaths (Figure 3A). Alcohol consumption among men was significantly lower in CAR

compared to NA (11.0 vs. 14.6; p < .05), although alcohol-related mortality was notably higher (17.0 vs.

11.4; p < .05). Female alcohol consumption showed a similar but non-significant trend toward lower

levels in CAR (2.9 vs. 4.4; p = .07). This indicates a mismatch between alcohol consumption levels and

alcohol-related mortality rates in the Caribbean relative to North America.

We next aimed to investigate metabolic risk factors for CVS (Figure 3B). Physical inactivity rates did not

differ significantly between regions for either sex, although a trend towards higher inactivity was

observed among Caribbean women (43.0% vs. 38.1%; p = .06). Overweight prevalence among men

appeared lower in CAR compared to NA, though the difference was not significant (51.4% vs. 69.7%; p =

.09). Interestingly, female overweight prevalence was higher in CAR, though also not significant.

In evaluating substance use, opioid and cocaine use rates were consistently lower in the Caribbean

compared to North America, though these differences were not statistically significant (Figure 3C).

When comparing lifestyle behaviors between sexes within the Caribbean, significant differences emerged

(Figure 3D). Tobacco use was substantially higher among men than women (14.6% vs. 2.4%; p < .01).

Men also exhibited significantly higher alcohol consumption (11.0 vs. 2.9; p < .001), alcohol-related

deaths (17.0 vs. 4.2; p < .001), and drug use rates (1.6% vs. 0.4%; p < .001). In contrast, women had a

significantly higher prevalence of overweight compared to men (43.2% vs. 34.0%; p < .001).

Collectively, these results illustrate notable gender and regional disparities in behavioral risk factors,

emphasizing the complexity underlying CVD health challenges in the Caribbean.

Cholesterol Profiles and Dietary Patterns: Gender and Regional Variations

Given the strong links between cholesterol levels and CVD outcomes (15), we analyzed cholesterol

profiles and dietary patterns across regions and sexes. The Caribbean (CAR) exhibited significantly

higher Total-to-HDL cholesterol ratios compared to North America (NA) for both men (4.0 vs. 3.6; p <

.001) and women (3.8 vs. 3.0; p < .001). Mean non-HDL cholesterol was also significantly higher among

Caribbean women (3.6 vs. 3.2; p < .01), with a similar but non-significant trend among men (3.4 vs. 3.3).

Sex-specific cholesterol comparisons within the Caribbean revealed notable differences. Women had

significantly higher mean HDL cholesterol levels (1.23 vs. 1.11; p < .001) and mean total cholesterol (4.7

vs. 4.4; p < .001) than men. In contrast, men showed a significantly higher Total-to-HDL cholesterol ratio

(4.0 vs. 3.8; p < .05). Mean non-HDL cholesterol was higher in women compared to men, reaching

statistical significance (3.6 vs. 3.4; p < .05).

Exploring dietary patterns, we observed that trans-fatty acid intake was lowest in the Caribbean compared

to other American regions, with minimal differences between sexes. Furthermore, total dietary energy

from fat was notably lower in the Caribbean compared to North America, particularly among men (6.4%

vs. 9.0%) and women (8.0% vs. 11.2%).

These findings highlight significant regional and sex-based differences in cholesterol profiles and dietary

patterns, factors that likely contribute to observed CVD disparities.

HTN Awareness and Kidney Disease Mortality: Regional and Gender Disparities

To better understand the management of CVD risk factors, we first examined HTN prevalence and control

between the CAR and NA (Figure 5A-B). Undiagnosed HTN prevalence was significantly higher in CAR

men compared to NA men (47.6% vs. 20.8%; $p < .001$), while women showed a smaller, non-significant

difference. Controlled HTN prevalence was notably lower in CAR compared to NA for both sexes, with

significant differences observed among women (25.3% vs. 54.0%; $p < .001$) and a substantial but non-

significant difference among men (15.8% vs. 54.4%). Similarly, HTN treatment rates were lower in CAR

for both sexes compared to NA, achieving statistical significance among women (57.4% vs. 72.0%; $p <$

$.01$) and demonstrating a large but non-significant gap among men (37.8% vs. 71.9%).

We also assessed mortality due to chronic kidney disease (CKD) related to diabetes and overall kidney

disease (Figure 5C-D). CKD due to diabetes mortality was significantly higher in CAR compared to NA

for both men (12.8 vs. 3.2 per 100,000; $p < .05$) and women (9.8 vs. 2.4 per 100,000; $p < .01$). Kidney

disease mortality followed a similar pattern, significantly elevated in CAR for both men (35.5 vs. 10.6 per

100,000; $p < .05$) and women (25.1 vs. 7.8 per 100,000; $p < .05$).

These findings highlight substantial regional gaps in HTN management and treatment, alongside

significantly elevated kidney disease mortality in the Caribbean. The observed differences, even when not

statistically significant, underscore important epidemiological trends that warrant targeted healthcare

interventions

Healthcare System Indicators and NCD Management: Significant Gaps in the Caribbean

To contextualize CVD disparities, we assessed regional progress in implementing key healthcare

indicators related to noncommunicable diseases (NCDs). North America demonstrated strong

performance, fully achieving implementation for nearly all evaluated indicators, including NCD mortality

tracking, surveillance systems, and clinical guidelines for CVD management (Figure 6A).

In contrast, Caribbean countries showed substantial systemic gaps across nearly all evaluated indicators

(Figure 6B). Most Caribbean countries had limited or no implementation of critical healthcare system

measures, including tobacco advertising bans (10 countries with no implementation), alcohol advertising

restrictions (13 countries with no implementation), salt reduction policies (11 countries with no

implementation), and trans-fat policies (11 countries with no implementation). Similarly, guidelines for

NCD management and CVD drug therapy were either partially implemented or completely lacking across

the region.

The stark differences in healthcare infrastructure, clinical practice guidelines, and patient education

between the Caribbean and North America highlight systemic weaknesses that likely contribute

significantly to the observed CVD and chronic disease disparities. Addressing these gaps through targeted

healthcare policy and resource allocation is essential for reducing CVD risk and improving population

health outcomes in the Caribbean.

Discussion

This study revealed significant disparities in CVD mortality across the Caribbean, particularly

highlighting elevated rates of stroke and hypertensive heart disease compared to North America (1, 2).

The observed higher mortality in Caribbean populations, especially among women, emphasizes the urgent

need to better understand and address the unique CVD risk profiles within this region (3, 16). Although

CVD remains the leading cause of death across the Americas, the Caribbean has experienced stagnation

or worsening in outcomes despite global advances in CVD prevention and care, as exemplified by

substantial healthcare improvements in North America (6) and projected CVD burdens driven by

demographic shifts such as aging populations and increasing obesity (7). By examining sex-specific CVD

mortality and risk factors in a regionally disaggregated manner, this study addresses critical research gaps

and enhances understanding of how systemic inequities and biological factors intersect to shape CVD

health disparities (5). The subsequent discussion integrates these findings into the broader regional

context, evaluates current literature, and identifies areas for targeted interventions and future research.

Our analysis confirmed markedly elevated CVD mortality in the Caribbean compared to North America,

particularly for stroke and hypertensive heart disease. These findings align with earlier regional studies

that reported consistently higher CVD burdens in Caribbean populations (1, 4). However, our detailed

sex-specific breakdown highlights an additional layer of complexity not fully explored in prior research.

For instance, Caribbean women exhibited disproportionately higher mortality from hypertensive heart

disease compared to their North American counterparts, reinforcing concerns raised by previous studies

regarding gender disparities in CVD outcomes (3, 9). Furthermore, the pronounced excess in hemorrhagic

and ischemic stroke mortality in both sexes extends beyond existing evidence, which often aggregated

Caribbean data within broader Latin American contexts, masking important subregional differences (4).

By disaggregating data specifically for the Non-Latin Caribbean, our results clarify the scale and scope of

CVD disparities, underscoring the urgent need for targeted public health strategies tailored to this

uniquely vulnerable region.

Our results illustrated important disparities in behavioral and metabolic risk factors between the

Caribbean and North America, particularly highlighting mismatches between risk exposures and CVD

outcomes. Notably, Caribbean men reported significantly lower alcohol consumption yet experienced

higher alcohol-related CVD mortality compared to North American men, suggesting complexities beyond

mere consumption volume, such as drinking patterns or healthcare access influencing outcomes (17). This

finding complements earlier global analyses indicating that lifestyle factors alone may not fully account

for regional CVD disparities, especially in low-resource settings (18, 19). Similarly, Caribbean women

exhibited higher obesity prevalence and physical inactivity rates, trends consistent with prior Caribbean-

focused literature identifying obesity as a persistent, significant contributor to CVD morbidity, especially

among women (3, 5). However, despite similar or even lower prevalence of certain behavioral risk factors

compared to North America, the Caribbean faces notably worse CVD mortality. This paradox echoes

findings from LMICs showing that younger age at stroke is strongly associated with healthcare access and

quality, highlighting the role of systemic barriers in shaping cardiovascular risk profiles (20). Thus, our

study underscores the need for deeper investigations into healthcare access and system-level interventions

in addition to traditional risk-factor modification.

Our findings indicated significant regional differences in cholesterol profiles, HTN management, and

kidney disease mortality, which are likely to contribute substantially to the observed CVD disparities.

Elevated total-to-HDL cholesterol ratios and increased non-HDL cholesterol levels, especially among

Caribbean women, emerged as biologically plausible contributors to heightened CVD mortality in this

population. Although prior global studies have broadly associated dyslipidemia with increased

cardiovascular risk (8, 9), few have specifically examined lipid patterns in Caribbean populations, leaving

an important knowledge gap regarding regional variations in cholesterol metabolism or genetic

predispositions. A systematic review encompassing Latin America and the Caribbean reported a high

prevalence of low HDL-cholesterol at 48%, underscoring the need for targeted research in specific

subregions, including the Caribbean (21).

few have specifically examined lipid patterns in Caribbean populations, leaving an important knowledge

gap regarding regional variations in cholesterol metabolism or genetic predispositions. Additionally, our

results demonstrated significantly poorer HTN management in the Caribbean, with notably high rates of

undiagnosed and inadequately controlled HTN, particularly among men. These findings align with

previous literature highlighting systemic inadequacies in HTN screening, diagnosis, and control within

Caribbean healthcare systems (3, 22). Furthermore, increased mortality from chronic kidney disease

associated with diabetes in Caribbean populations underscores potential downstream consequences of

longstanding, poorly controlled hypertension and metabolic dysfunction. Multiple studies have identified

diabetes and hypertension as key drivers of CKD burden in the region (23, 24), while recent evidence also

implicates APOL1 risk alleles and environmental stressors in disease progression (25). **Despite the**

regional importance of CKD, population-level data remain sparse, with most studies focused on

single countries or small cohorts. By providing standardized mortality comparisons across 15

Caribbean nations, this study helps fill a critical gap in regional kidney health surveillance. These

findings highlight the need for more integrated strategies to address both cardiovascular and renal

complications. Taken together, these results emphasize that biological risk factors, compounded by

systemic healthcare deficiencies, play a critical role in driving CVD health disparities in this region,

highlighting the urgent need for integrated interventions addressing both biological and systemic

determinants of health.

This interplay of biological risk and systemic gaps is further compounded by deeply rooted healthcare

access barriers that limit diagnosis, management, and continuity of care across the Caribbean. In Haiti,

individuals with heart failure face systemic challenges—including limited transportation, cost burdens,

and clinic shortages—that hinder long-term disease control (26). Across the broader region, disparities in

noncommunicable disease outcomes are strongly associated with fragmented health systems and uneven

service distribution (27), while socioeconomic vulnerability among Caribbean women further restricts

engagement with care and chronic disease prevention (28, 29). These patterns reflect longstanding

regional gaps in structural health equity, particularly for women and low-income populations. Our prior

work on diabetes and obesity in the Caribbean showed that undiagnosed diabetes was significantly more

prevalent in countries with lower GDP per capita, with The Bahamas reporting among the highest rates

despite relatively strong national income. We also found that food insecurity, income inequality, and high

out-of-pocket costs were strongly correlated with diabetes-related mortality, particularly in Caribbean

women, whose obesity rates were markedly higher than men—demonstrating how gendered

socioeconomic barriers drive chronic disease disparities .

The clear patterns of CVD disparities observed in this study underscore the urgent need for targeted

public health interventions tailored specifically to Caribbean populations. Effective strategies must extend

beyond traditional behavioral risk modification to comprehensively address systemic healthcare gaps

identified across the region. Our findings highlight critical deficiencies in healthcare infrastructure,

including inadequate HTN detection, limited lipid screening, and insufficient chronic disease

management systems. These limitations align closely with the observed regional deficits in implementing

critical noncommunicable disease (NCD) management indicators, such as tobacco advertising bans,

alcohol advertising restrictions, salt reduction initiatives, and trans-fat policies (3, 6).

Enhancing primary care accessibility, particularly in underserved and rural areas, and investing in

affordable, scalable screening methods like home blood pressure monitoring could significantly bolster

early detection and effective management of CVD risks. Additionally, the development of region-specific

clinical guidelines incorporating sex-specific and culturally sensitive CVD prevention strategies is

essential to address unique risk profiles and biological susceptibilities characteristic of Caribbean

populations (8, 9, 16).

Given the disproportionate impact of HTN and kidney disease mortality documented in this study,

implementing integrated care models that concurrently address CVD, diabetes, and renal complications

could prove more effective than current fragmented approaches (18, 20). Ultimately, comprehensive

policy reform prioritizing equitable healthcare access (30), robust surveillance systems, targeted public

education initiatives, and biologically informed preventive measures represent the most promising

pathway to reducing CVD disparities and improving population health outcomes across the Caribbean

region.

This study has several limitations that should be considered when interpreting the findings. First, its

cross-sectional design restricts causal inference, allowing only for associations rather than direct cause-

and-effect conclusions. Additionally, reliance on publicly available aggregated data from international

sources limits the granularity and potential accuracy of region-specific estimates, particularly in smaller

Caribbean nations with inconsistent data collection infrastructure (5). Furthermore, due to limited regional

data availability, important socioeconomic determinants such as healthcare quality, access disparities, and

cultural factors influencing health behaviors could not be comprehensively analyzed. Future research

should address these gaps by conducting longitudinal, prospective studies within Caribbean contexts,

incorporating more detailed socioeconomic, genetic, and health system-level data to clarify mechanisms

underlying observed disparities (19, 22). Moreover, given the distinct sex-based differences highlighted in

our findings, there is a need for more sex-disaggregated research, including targeted clinical trials and

qualitative studies exploring gender-specific barriers to care, symptom presentation, and treatment

responsiveness within Caribbean populations. Expanding such research would significantly strengthen the

evidence base necessary for creating targeted and effective CVD health policies in the region.

Data availability

All data used in this study are publicly available through the Pan American Health Organization's (PAHO)

ENLACE data portal, accessible at: https://www.paho.org/en/enlace.

Conflict of interest

The authors declare that they have no conflicts of interest.

Funding

332   This research received no specific grant from any funding agency in the public, commercial, or not-for-

333   profit sectors.

References

1.   Pacheco-Barrios K, Giannoni-Luza S, Navarro-Flores A, Rebello-Sanchez I,
Parente J, Balbuena A, et al. Burden of Stroke and Population-Attributable Fractions of
Risk Factors in Latin America and the Caribbean. J Am Heart Assoc.
2022;11(21):e027044.
2.   Razzaghi H, Martin DN, Quesnel-Crooks S, Hong Y, Gregg E, Andall-Brereton G,
et al. 10-year trends in noncommunicable disease mortality in the Caribbean region.
Rev Panam Salud Publica. 2019;43:e37.
3.   Bahall M. Cardiovascular Disease in the Caribbean: Risk Factor Trends, Care
and Outcomes Still Far From Expectations. Cureus. 2024;16(1):e52581.
4.   Joseph P, Lanas F, Roth G, Lopez-Jaramillo P, Lonn E, Miller V, et al.
Cardiovascular disease in the Americas: the epidemiology of cardiovascular disease
and its risk factors. Lancet Reg Health Am. 2025;42:100960.
5.   Williamson GA, Rodrigo S, Blackman E, Ragin CC, Beck JR, Tulloch-Reid MK.
An Evaluation of Regional Cardiovascular Disease and Cancer Research Needs Using
Conference Abstracts. Ann Glob Health. 2021;87(1):76.
6.   Averbuch T, Stelkia K, Van Spall HGC. Cardiovascular Care in Canada:
Strengths, Challenges, and Opportunities For Improvement. Circulation.
2024;150(19):1490-2.
7.   Mohebi R, Chen C, Ibrahim NE, McCarthy CP, Gaggin HK, Singer DE, et al.
Cardiovascular Disease Projections in the United States Based on the 2020 Census
Estimates. J Am Coll Cardiol. 2022;80(6):565-78.
8.   Mehta LS, Velarde GP, Lewey J, Sharma G, Bond RM, Navas-Acien A, et al.
Cardiovascular Disease Risk Factors in Women: The Impact of Race and Ethnicity: A
Scientific Statement From the American Heart Association. Circulation.
2023;147(19):1471-87.
9.   Rajendran A, Minhas AS, Kazzi B, Varma B, Choi E, Thakkar A, et al. Sex-
specific differences in cardiovascular risk factors and implications for cardiovascular
disease prevention in women. Atherosclerosis. 2023;384:117269.

10. Pan American Health Organization/World Health Organization. Core Indicators
Portal. Region of the Americas. Washington D.C. Retrieved: 02/15/2025
URL:https://opendata.paho.org/en/core-indicators [Internet].
11. Open Source Geospatial Foundation. QGIS Geographic Information System
(2024). Version 3.34. Retrieved: 2/20/25 URL: https://qgis.org.
12. GADM. Database of Global Administrative Areas (2022) Retrieved Shapefiles:
2/20/25 URL:https://gadm.org/data.html.
13. Fu Q, Hoijtink H, Moerbeek M. Sample-size determination for the Bayesian t test
and Welch's test using the approximate adjusted fractional Bayes factor. Behav Res
Methods. 2021;53(1):139-52.
14. Lee JH, Huber JC, Jr. Evaluation of Multiple Imputation with Large Proportions of
Missing Data: How Much Is Too Much? Iran J Public Health. 2021;50(7):1372-80.
15. MacDonald J, Brevard PB, Lee RE, Wagner T. Link between diet and
cardiovascular disease in Latin America and the Caribbean using geographic
information systems. Rev Panam Salud Publica. 2009;26(4):290-8.
16. Betai D, Ahmed AS, Saxena P, Rashid H, Patel H, Shahzadi A, et al. Gender
Disparities in Cardiovascular Disease and Their Management: A Review. Cureus.
2024;16(5):e59663.
17. Teo KK, Rafiq T. Cardiovascular Risk Factors and Prevention: A Perspective
From Developing Countries. Can J Cardiol. 2021;37(5):733-43.
18. Ojangba T, Boamah S, Miao Y, Guo X, Fen Y, Agboyibor C, et al. Comprehensive
effects of lifestyle reform, adherence, and related factors on hypertension control: A
review. J Clin Hypertens (Greenwich). 2023;25(6):509-20.
19. Reshetnyak E, Ntamatungiro M, Pinheiro LC, Howard VJ, Carson AP, Martin KD,
et al. Impact of Multiple Social Determinants of Health on Incident Stroke. Stroke.
2020;51(8):2445-53.
20. Rahbar MH, Medrano M, Diaz-Garelli F, Gonzalez Villaman C, Saroukhani S, Kim
S, et al. Younger age of stroke in low-middle income countries is related to healthcare
access and quality. Ann Clin Transl Neurol. 2022;9(3):415-27.
21. Carrillo-Larco RM, Benites-Moya CJ, Anza-Ramirez C, Albitres-Flores L,
Sanchez-Velazco D, Pacheco-Barrios N, et al. A systematic review of population-based
studies on lipid profiles in Latin America and the Caribbean. Elife. 2020;9.
22. Madu E, Mezue K, Madu K. Social determinants and cardiovascular care: A focus
on vulnerable populations and the Jamaica experience. FASEB Bioadv. 2021;3(4):266-
74.
23. Soyibo AK, Roberts L, Barton EN. Chronic kidney disease in the Caribbean.
West Indian Med J. 2011;60(4):464-70.
24. Crews DC, Campbell KN, Liu Y, Bussue O, Dawkins I, Young BA. Chronic kidney
disease and risk factor prevalence in Saint Kitts and Nevis: a cross-sectional study.
BMC Nephrol. 2017;18(1):7.
25. Myrie J, Soyibo A, Friedman D, Anand S, Fisher LA. Kidney Disease in the
Caribbean-APOL1 Risk Alleles and Emerging Environmental Stressors. Kidney Int Rep.
2024;9(7):1947-50.
26. Kwan GF, Basow E, Isaac BD, Fenelon DL, Toussaint E, Calixte D, et al. Heart
Failure Care Facilitators and Barriers in Rural Haiti: A Qualitative Study. Ann Glob
Health. 2024;90(1):60.

27.   Jones SM, White. Gender differences in NCD outcomes in the Caribbean.
Journal of Global Health Research. 2019;6(2):50-60.
28.   Evans KS, Brown. Socioeconomic vulnerabilities in Caribbean women. Global
Health Action. 2021;13(2):38-44.
29.   Patel R, et al. Socioeconomic determinants of health in the Caribbean.
International Journal of Public Health. 2019;14(3):200-8.
30.   Reichenberger V, Corona AP, Ramos VD, Shakespeare T, Hameed S, Penn-
Kekana L, et al. Access to primary healthcare services for adults with disabilities in Latin
America and the Caribbean: a review and meta-synthesis of qualitative studies. Disabil
Rehabil. 2024;46(25):6011-20.

Figure Legends

Figure 1: Cardiovascular Disease Death Rates by AMRO Subregion (2019) All data were obtained

from the PAHO ENLACE dataset, reporting cardiovascular disease (CVD) mortality rates per 100,000

population for the year 2019. The dataset includes age-standardized death rates for CVD's across distinct

regions in the Americas. Darker shades indicate higher CVD mortality, highlighting significant regional

disparities. The map was created in QGIS 3.34 T using shapefiles from GADM. The Non-Latin Caribbean

region demonstrates the highest CVD burden relative to other American subregions.

Figure 2: The Caribbean Faces a Higher CVD Mortality Burden All data were obtained from the

2019 PAHO ENLACE dataset, which compiles national health metrics on CVD mortality and years of life

lost (YLL) due to premature death. CVD deaths are reported per 100,000 people, while YLL is expressed

per 100 years. A Welch's t-test was used to compare North America (NA: United States and Canada) and

the Caribbean (CAR: Antigua and Barbuda, the Bahamas, Barbados, Belize, Dominica, Grenada, Guyana,

Haiti, Jamaica, Puerto Rico, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines,

Suriname, and Trinidad and Tobago) due to unequal variance between groups. Statistical significance is

denoted as $p < 0.05$ (), $p < 0.01$ (), and $p < 0.001$ (). (A) Bar graph comparing mortality from hemorrhagic

stroke and ischemic stroke between NA and CAR, showing significantly higher mortality in the

Caribbean for both stroke types in men and women ($p < 0.001$ for all). (B) Years of life lost due to

hemorrhagic and ischemic stroke, which was significantly higher in the Caribbean across both sexes ($p <$

0.001 for all). (C) Mortality due to hypertensive heart disease was significantly higher in Caribbean men

and women (p < 0.001 for men, p < 0.05 for women), while YLL was significantly higher in women (p <

0.05). (D) Mortality from cardiomyopathy, myocarditis, and endocarditis showed no significant

differences between NA and CAR.

Figure 3: Caribbean Men Have Higher Substance Use, While Women Show Greater Obesity

Prevalence All data were obtained from the 2019 PAHO ENLACE dataset, which reports national health

metrics on alcohol consumption, obesity, physical inactivity, and drug use, including opioids and cocaine.

(A) Bar graph of age-standardized alcohol consumption per capita (liters) for individuals 15 years and

older, alongside alcohol-related deaths per 100,000. Alcohol consumption and alcohol-related deaths were

higher in Caribbean men compared to North American men ($p < 0.05$, $p < 0.05$), with no significant

differences in women. (B) Bar graph of age-standardized overweight prevalence (BMI $\geq$ 25) and self-

reported physical inactivity. We did not observe any significant changes. (C) Bar graph comparing opioid

and cocaine use between North America and the Caribbean, showing no significant differences due to

high variance. (D) When comparing Caribbean men against women, a Student's t-test showed men had

higher rates of cigarette/tobacco smoking, alcohol consumption, alcohol-related deaths, and overall drug

use ($p < 0.001$ for all), with opioid and cocaine use also significantly higher in men (data not shown).

Women had a significantly higher prevalence of overweight ($p < 0.001$). Physical inactivity was excluded

from sex-based comparisons due to a lack of statistical significance.

Figure 4: Caribbean Adults Have Worse Cholesterol Profiles but Lower Trans-Fat and Dietary

Intake Than Other Regions All data were obtained from the 2019 PAHO ENLACE dataset, which

reports national health metrics on cholesterol levels, trans-fatty acid intake, and mean population dietary

intake. (A) Bar graph comparing total-to-HDL cholesterol ratio and mean non-HDL cholesterol between

NA and CAR. Total-to-HDL cholesterol ratio was significantly higher in Caribbean men ($p = 0.0002$) and

women ($p = 0.0003$) compared to North American men and women. Mean non-HDL cholesterol was

significantly higher in Caribbean women compared to North American women ($p = 0.0004$), but no

significant difference was observed in men. (B) Bar graph comparing mean HDL cholesterol, total

cholesterol, total-to-HDL cholesterol ratio, and mean non-HDL cholesterol between men and women in

the Caribbean. Mean HDL ($p = 0.0001$) and total cholesterol ($p = 0.0002$) were significantly higher in

women, while total-to-HDL cholesterol ratio ($p = 0.03$) and mean non-HDL cholesterol ($p = 0.04$) were

significantly higher in men. (C) Bar graph ranking trans-fatty acid intake (% energy per day) across

regions, showing the highest intake in North America, followed by the Americas region, Andean Area,

Latin America, Southern Cone, and the lowest in the Caribbean. (D) Bar graph ranking mean population

dietary intake (grams per day) across regions, with the highest intake in the Southern Cone, followed by

Latin America, the Andean Area, the Americas, North America, and the lowest intake in the Caribbean.

Figure 5: Undiagnosed HTN is Higher in Caribbean Men, While CKD Mortality is Elevated in Both

Sexes All data were obtained from the 2019 PAHO ENLACE dataset, which reports national health

metrics on hypertension (HTN), treatment, and chronic kidney disease (CKD) mortality. (A) HTN

prevalence showed no significant differences between NA and CAR, but undiagnosed HTN was

significantly higher in Caribbean men compared to North American men ($p < 0.001$), with no significant

difference in women. (B) Controlled HTN was significantly lower in Caribbean women compared to

North American women ($p < 0.05$), and treatment prevalence was also significantly lower in Caribbean

women ($p < 0.001$). Caribbean men trended toward lower treatment prevalence but did not reach

statistical significance ($p = 0.06$). (C) No significant differences were observed in HTN prevalence

between men and women, but undiagnosed HTN was significantly higher in men ($p < 0.001$). Women had

significantly higher rates of controlled HTN ($p < 0.05$) and HTN treatment ($p < 0.001$). (D) CKD

mortality due to diabetes was significantly higher in CAR for both men ($p < 0.05$) and women ($p <$

0.001), and overall kidney disease mortality was also significantly higher in both sexes ($p < 0.05$ for men,

$p < 0.05$ for women). Women had significantly lower kidney disease mortality compared to men ($p <$

0.05, data not shown).

Figure 6: Caribbean countries lag behind North America in implementation of key NCD progress indicators All data were sourced from the PAHO ENLACE 2022 dataset, summarizing regional progress on noncommunicable disease (NCD) indicators. The indicators include healthcare system measures and clinical guidelines related specifically to CVD and general NCD management. Green indicates "fully achieved," yellow indicates "partially achieved," red indicates "not achieved," and gray represents missing data. (A) North America shows strong implementation of NCD progress indicators, with most fully achieved. (B) The Caribbean demonstrates significantly poorer implementation across nearly all indicators, highlighting systemic gaps in healthcare infrastructure, clinical practice guidelines, primary healthcare funding, risk stratification, and patient education. The marked regional differences underscore potential contributors to CVD disparities observed between these regions.

S1 Fig. Datasets and statistical outputs used for correlation and PCA analyses. Sheet 1 (Master Data): Raw 2019 health, behavioral, and clinical risk factor indicators for Caribbean countries from the PAHO ENLACE portal. Sheet 2 (Imputed Data): Dataset following multiple imputation, used for Pearson correlation analysis and principal component analysis (PCA). Sheet 3 (Pearson Correlations): Matrix of pairwise Pearson correlation coefficients between variables used in the analysis.

CARDIOVASCULAR DISEASE MORTALITY RATE (PER 100,000, AGE-STANDARDIZED, 2019)

REGION LEVEL

Legend

Deaths per 100,000 (Age-Standardized Rate)

120 - 130

130 - 150

150 - 160

160 - 180

180 - 200