Pathways to optimize a pediatric telemedicine and medication delivery service: A multi-level qualitative study in Haiti

Authors
Frantz Emmanuel Garilus(BS)[1], Kerven Cassion(MPH)[1], Youseline Cajusma(BSN)[1], Katelyn E. Flaherty(PhD)[1,2], Jude Ronald Beausejour(MD)[1], Lerby Exantus(MD)[3], Valery M. Beau de Rochars(MD)[4], Chantale Baril(MD)[3], Torben K. Becker(MD, MBA, PhD)[2], Rochelle K. Rosen(PhD)[5], Eric J. Nelson(MD, PhD)[1], Molly B. Klarman(MPH)[1]*

Affiliations
[1]Departments of Pediatrics and Environmental and Global Health, University of Florida, Gainesville, FL, USA
[2] Department of Emergency Medicine, University of Florida, Gainesville, FL, USA
[3] Université d'État d'Haiti- Faculté de Médecine et de Pharmacie, Port-au-Prince, Haiti
[4] Department of Health Services Research, Management and Policy, University of Florida, Gainesville, FL, USA
[5]Center for Behavioral and Preventive Medicine, The Miriam Hospital, Providence, RI, USA

*Corresponding author
Molly B Klarman, MPH
2055 Mowry Road, Emerging Pathogens Institute, University of Florida
Gainesville, FL 32611
Email: mklarman@ufl.edu

Keywords: framework matrix analysis, telemedicine, pediatrics, scaling

Word counts:
Abstract: 186 words
Text: 4,719 words
Figures: 2
Tables: 2
Supplemental text: 2

ABSTRACT

Background. While telemedicine has become an established component of healthcare delivery globally, challenges to scaling emerging initiatives persist across multiple levels. Over the last 5 years, our team has developed a pediatric telemedicine and medication delivery service (TMDS) in Haiti that integrates clinical guidance with rapid access to medications. Building on successful proof-of-concept studies, we are now well positioned to characterize both general challenges to telemedicine scale-up and those unique to the TMDS model.

Methods. In this study, we conducted focus group discussions and administered written questionaries to TMDS staff, including physicians, nurses, and medication delivery drivers. Using framework matrix analysis we identified key challenges and opportunities associated with the TMDS model.

Results. Areas for improvement related to obtaining quality information from virtual exams, the reliability of technology and communication infrastructure, conditions necessary for effective in-person exams, the limited scope of the TMDS workflow and clinical resources, and uncertainty surrounding long-term sustainability. These insights informed the development of targeted action items categorized into three domains: conceptual, physical and mission-oriented.

Conclusion. The findings will guide our internal scale-up strategy and may offer guidance to similar telemedicine initiatives.

INTRODUCTION

In the wake of the COVID-19 pandemic, telemedicine became a central fixture in healthcare infrastructure [1]. A post-pandemic analysis of the utilization of telehealth and telemedicine services found at least 77 countries had adopted the approach [2]. Telemedicine, generally defined as '*the use of technology to provide healthcare services between parties that are geographically separated'* [3, 4], is rapidly diversifying in scope [5]. One categorization of telemedicine uses five application types: tele-consultation, tele-diagnosis, tele-ambulance, tele-monitoring, and digital self-management. The perspective considers six additional descriptive components: personnel, target population, setting, technology, data provision and intended outcome [6]. Permutations of these components demonstrate the broad range of possible telemedicine applications.

The benefits of telemedicine vary depending on the specific application or use case. One use case is a type of synchronous (real-time) teleconsultation between provider and patient known as 'triage and treat'. In this model, patients are remotely screened to determine the appropriate location and method of treatment [7, 8]. In low-resource settings, the triage and treat approach can increase access to specialists, address shortages of healthcare workers, and reduce transportation related barriers, including time and cost [9-11]. With these goals in mind, our team developed a pilot pediatric telemedicine and medication delivery service (TMDS) that has been deployed in Haiti and Ghana [12, 13]. In the TMDS model, patients are first triaged and then assessed through a virtual exam, followed by the delivery of medications directly to their homes. To the best of our knowledge, there are few other (if any) affordable, high-

quality, evidence-based, pediatric TMDS interventions deployed in resource-limited

settings. As with many aspects of telemedicine's evolution, this is likely to change.

Scaling novel healthcare interventions, such as telemedicine, is challenging,

particularly in low-resource settings [14-16]. A broadly used definition of 'scaling up' a

health innovation is, '*deliberate efforts to increase the impact of successfully tested*

health innovations, so as to benefit more people and to foster policy and program

development on a lasting basis' [17]. Scaling is unlikely to occur organically, even with

the most promising innovations [14]. Barriers to scale can be numerous and present

across various implementation domains[18].

One common approach to identify these barriers is through frameworks such as

the Consolidated Framework for Implementation Research (CFIR) [19]. CFIR covers

five domains: innovation characteristics, outer setting, inner setting, characteristics of

individuals, and implementation process. These domains can help assess barriers to

implementing telemedicine interventions. Within the 'innovation domain', issues with

poor interoperability are common [20, 21]. Within the 'outer setting' many low-resource

countries lack nationwide policies and regulatory standards for telemedicine [22-24].

Within the 'inner setting' challenges to obtain and maintain information and

communication technology infrastructure (ICT) are often cited [23, 25]. Within the

'individuals' domain both providers and patients have expressed hesitancy to accept

telemedicine [26, 27]. Within the 'implementation process' domain many telemedicine

programs were rapidly implemented in response to COVID-19 and did not undergo the

necessary planning stages to support sustained success [28]. In addition to identifying

barriers, stakeholders are also committed to recognizing the facilitators that support

successful, and scalable implementation.

Qualitative methods play an integral role in implementation science, allowing

implementors to systematically identify barriers and facilitators to successful initiatives.

The methods allow for systematic and efficient description of what is happening and

why it is happening [29]. Proctor et al. proposed eight outcome measures for

implementation research, of which five recommend qualitative research methods as the

primary measurement tool [30]. Similar to quantitative research, the research question

leads decisions on which data collection and analysis methods are most appropriate

[31]. Among various approaches, framework matrix analysis is especially effective for

producing rapid, streamlined summaries that support deductive learning and can be

easily shared between diverse stakeholders [32].

With the objective of prioritizing early access to pre-emergent healthcare to

prevent the need for more costly and complex emergency care, our team developed a

nighttime pediatric telemedicine and medication delivery service (TMDS) in Haiti. The

service has evolved within the Improving Nighttime Access to Care and Treatment- Haiti

(INACT-H) study series. The TMDS model was designed to overcome the healthcare

access barriers of distance and nighttime [33]. From 2019 to 2021, we validated this

innovative healthcare delivery model, finding that it was clinically safe, feasible to

implement, and supported by reliable clinical resources [12, 34]. From 2021 to 2023, we

focused on positioning the TMDS to scale by modifying the workflow. The resulting

scalable model maintained strong clinical and operational outcomes [35]. A cost-

effectiveness analysis confirmed that a scaled TMDS was a cost-effective alternative to

hospital emergency care for pre-emergency pediatric conditions at night in Haiti [36]. In

this context, we sought to qualitatively identify challenges and opportunities that could

improve both cognitive and physical factors to optimize the TMDS for large-scale

implementation.

METHODS

Study setting. A TMDS called MotoMeds was launched in the semi-urban and rural

region of Gressier, Haiti in September 2019. The workflow of the TMDS is as follows

(Figure 1): a parent calls the call center regarding their sick child. A TMDS nurse

(registered nurse or family nurse practitioner) collects demographic and clinical findings

from the parent to complete a virtual exam by phone. Nurses use TMDS specific clinical

resources to generate an assessment and treatment plan that includes illness severity,

disposition, medication/fluid recommendations, and advice for follow-up care. Illnesses

triaged as severe are referred directly to emergency services. Among children who

reside within a delivery zone, those with higher clinical uncertainty are offered an in-

person household exam with delivery and patients with low clinical uncertainty are

offered a delivery of medications/fluids. Children who reside outside of a delivery zone

are provided with guidance alone. All children receive a follow-up call at 10-days and

patients with severe illness also receive a call at 24-hours. On-call physicians are

available by phone to consult on cases beyond the scope of the TMDS clinical

resources.

In October 2021, a second delivery zone was established in the urban center of

Les Cayes, located 175 Km west of Gressier. The workflow of this delivery zone differs

in that treatment plans generated at the central call center in Gressier are relayed to on-

call nurses in Les Cayes who either dispatch motorcycle delivery drivers to transport

medications/fluids or conduct in-person household exams. Once proofs of concept were

established through three years of evaluation, iteration, and growth, our team pivoted

towards positioning the TMDS for scale. We sought qualitative feedback from TMDS

staff as a critical component of this process. The research team consisted of EJN, the

PI who is a pediatrician and scientist with qualitative research experience [32, 37], MBK,

the study director who is an international public health specialist, and two bilingual

Haitian American facilitators, FG and KC. Each team member was coached in

qualitative methods by RKR, an expert in qualitative methods including matrix analyses

applied in global health settings challenged by technological and situational limitations

[32, 38].

Study design

Focus groups. EJN, MBK, FG, and KG developed a written agenda (Additional File 1) to

guide focus group discussions for each of four types of TMDS staff/participants.

Participant types included: i) call center nurses, who conduct virtual exams at the call

center and relay treatment plans, ii) delivery nurses, who receive and enact virtual

treatment plans by dispatching a motorcycle delivery driver or conducting in-person

household exams, iii) on-call physicians, who provide consultations by phone, and iv)

motorcycle delivery drivers, who transport nurses and deliver medications/fluids to

households. Most questions and discussion prompts were specific to participant type,

but a subset was shared across all participants. Topic areas were identified *a priori* as

either factors that could enable or impede scaling (clinical, technical, and logistical

challenges and solutions) or that qualitatively assess impact (family expectations and

satisfaction, and value to TMDS staff and community). The agenda was translated to

Haitian Creole by FG and KC. MBK reviewed the agenda questions with a TMDS staff

member to ensure the questions were relevant to the staff experience.

In June and July 2022, FG and KC conducted nine focus group discussions via

video conferencing software (Zoom) with facilitators based in the United States and

participants located in Haiti. All active staff members were offered participation and

were placed in groups of 3-4 based on type and location. At the beginning of each

session a facilitator read a waiver of documentation of written consent statement to

each person individually. After expressing verbal agreement, participants were given a

study ID. The video sessions were recorded.

Questionnaires. Two versions of a nurse questionnaire were developed by MBK with 26

short answer questions for call center nurses and 15 for delivery nurses (Additional File

2). The questionnaire focused on identifying ways to improve the clinical resources and

procedures that guide virtual and in-person exams. Given the growing interest in clinical

decision support (CDS) tools, particularly digital versions of World Health Organization

guidelines, the questionnaire also assessed interest in integrating electronic clinical

decision support into the TMDS workflow.

Data analysis

Data preparation. FG and KC translated (Haitian Kreyol to English) and transcribed the

focus group video recordings in a single step. One facilitator completed a focus group in

its entirety and the other reviewed it for accuracy, which included confirming the

translation and correct identification of the speakers. Discrepancies were discussed

between the facilitators and if necessary, a third team member (MBK) was consulted.

Questionnaire responses were translated into English using the same procedure

applied to the video recordings.

Matrix analysis. Four matrices were deductively constructed in Microsoft Excel. The

column headers were drawn from the focus group agenda and nurse questionnaire and

the rows corresponded to each participant. To populate the matrices, one facilitator read

a transcript or questionnaire question by question and summarized the response by

each participant in the corresponding cell, separating unique ideas with bullet points.

The opposite facilitator reviewed the transcript or questionnaire and the matrix for

completeness and a third team member reviewed the matrices for clarity. At the bottom

of the matrix 4 additional 'summary rows' were created; in the 1st row, MBK summarized

the responses from all participants for each question, in the 2nd, 3rd and 4th rows FG,

KC, and MBK suggested possible action items to address ideas presented by

participants. Through Zoom the four-member team reviewed the 'summary row' cells

question by question and through discussion established a 'consensus of summary' and

a 'consensus of action items' which were entered into additional rows of the matrix

spreadsheets. Common domains relating to barriers and facilitators for scaling, and

potential action items to improve the service were identified. FG and KC reviewed focus

group transcripts and identified quotes that supported the common domains.

RESULTS

Study overview. All active staff (14 nurses, 13 drivers, 3 on-call physicians) agreed to

participate in one of nine focus group discussions (Table 1). In addition, 14 nurses

completed a written questionnaire. Participant input was captured within five domains:

1) Virtual exams at the call center, 2) In-person exams at the household, 3) Clinical

resources, 4) Communication, and 5) Expectations and impact.

Virtual exams. When discussing how nurses perceive the information parents relay

over the phone, nurses stated they felt confident in the information families shared when

responses were provided without hesitation, consistent over time, and aligned with other

responses. They doubted the accuracy of the information when they had to repeat a

question multiple times, and when they received inconsistent or implausible responses.

One nurse explained, *"…sometimes we ask questions, and they* [callers] *are bouncing*

between 'yes' or 'no'. This makes it difficult for us to have confidence in the information

we are receiving. What also happens, when we do not have confidence, we mark 'not

confident' [on the case report form]*. For example, if you have asked your first question*

and the answer is 'yes', and for things to correlate, the answer to the second question

must also be 'yes', but the parents say 'no'. Thus, there is a lack of correlation, and this

makes us lack confidence in the information we are receiving." In these situations,

nurses stated that they attempted to improve the data quality by rewording questions

and explaining the importance of providing accurate responses. If they remained unsure

about a particular response, they marked 'not confident' on the case report form and

that response did not factor into the treatment plan. Nurses reported that specific

components of the virtual exam were particularly challenging to complete. They felt that

families were often resistant to identifying a chief complaint among multiple complaints

and struggled to measure the respiratory and heart rates. Multiple nurses noted that

questions related to respiratory problems (e.g., stridor, retractions) were difficult for

parents to understand and often required repeat explanations and/or exploratory follow-

up questions that were not part of the structured virtual exam.

Call center characteristics. When reflecting on the work environment, nurses

attributed the effective operation of the call center to the consistent availability of

necessary materials and strong teamwork among the nursing staff. In contrast,

unreliable internet and cellular signal were obstacles to performing call center

responsibilities. Although technology is integral to the TMDS workflow, frequent

changes to user-interfaces, such as laptop password expirations and software updates,

along with unfamiliar information technology (IT) issues were cited as impeding call

center operations. One nurse commented, *"…problems we have are that in the*

[software] applications, you connect to your account in a specific way and in a moment

it changes. This means the nurse must have knowledge in informatics. If all the nurses

did not have knowledge in informatics, this would be a big problem for us. I can come

today to sign into my account to replace the nurse that was here last night, and the way

I am used to signing in, I can't sign in that way anymore." IT management strategies

involved relying on contingencies (secondary internet carrier) and contacting the Study

Director.

In-person exams. During in-person exams nurses faced both technical and clinical

challenges. Technical challenges included inadequate lighting and space, a lack of flat

surfaces to place a scale to weigh patients, and missing materials due to failure to verify

supply levels before departure. Clinical challenges included difficulty obtaining vital

signs from young children, parental hesitancy to obtain temperature rectally, and a

feeling of incompetency when the TMDS clinical resources or formulary did not cover a

patient's situation or when contacting the on-call physician in the presence of the

parents. One nurse said, *"A difficulty we face performing in-person exams is when I*

encounter a complex case and I need to call the on-call physician in front of the family,

they might think I am incompetent". Despite these challenges, nurses valued several

aspects of the in-person exam, including obtaining vital signs, exploring the patient's

medical history, and administering the first dose of medication.

Household logistics. Nurses commonly reported challenges reaching participant

households, which they attributed to the difficulties of nighttime travel by motorcycle,

poor road quality, remote household locations, and inability to connect with the parent

by phone upon arrival. Nurses reported spending substantial time explaining the

consent process, educating parents on how to administer medications, and discussing

the rationale for the TMDS guidelines (e.g., why it only serves children ≤10 years or only

covers specific illnesses). A mismatch between the parameters of the TMDS and

parental expectations also required dialogue. Nurses described parents wanting certain

medications that were not indicated (e.g., an antibiotic for a cough), requesting in-

person exams for additional unenrolled children at the household, and encountering a

different clinical situation than what was relayed during the virtual exam. One nurse

commented, "*The problem I face is that when we reach their house, they tell us that*

they have other children that are sick and if we could see them also. We tell the parent

to call the call center first, but they don't agree. They want us to examine all their

children, although they did not originally call for that. Another issue that we face is that

the problem the parent calls for is not fully accurate. When we get there, we see that the

child has more problems than they [originally] told us." The TMDS clinical guidelines

recommended that patients with severe illness seek emergency level care. However,

parents of children with severe conditions frequently reported being unable to do so due

to financial constraints. In these situations, families still expected to receive

medications, despite delivery nurses not being equipped to manage severe conditions.

One nurse shared, "*Another problem is that MotoMeds does not treat every problem,*

some problems that a child may have, they are in the red [severe] category, you must

send them to the hospital. Sometimes when we get to the house, with the state of the

child, we can't give them anything and we must send them to the hospital. But the

parents don't have the means and they are not happy that we didn't give their child any

medication. The parents feel as though they didn't get any service because of this."

Clinical resources. The nurses generally expressed satisfaction with the clinical

resources, stating that they rarely encountered clinical situations beyond their capacity

to manage. When faced with situations outside the scope of the clinical resources, they

contacted an on-call physician, as the workflow intended. However, nurses did note

occasional challenges related to medication dosing. They expressed concern about the

inability to divide pills evenly, or at all, if the medication was only available in capsule

form. Further, nurses were concerned that the estimated weights they used to perform

dosing calculations could misrepresent patients' actual weights in the context of

undernutrition.

The role of the on-call physician was evaluated from both the nurses' and

physicians' perspectives. Nurses suggested that physicians may not be fully aware of

the constraints they face, such as being limited to the medications listed in the

formulary. In some instances, physicians recommended medications not included in the

formulary, even when equally effective alternatives were available within it. Despite the

known limitations of telemedicine, physicians reported receiving sufficient case details

to offer consultation. However, they noted that having a mechanism to receive photos or

videos would be beneficial. It was broadly acknowledged that the on-call physician role

is crucial but could be improved by developing a formal tool for gathering and relaying

case details, as well as providing physicians with feedback on the outcome of each

case they consult. Additionally, despite feeling connected to project leadership through

WhatsApp groups and weekly email updates, on-call physicians felt disconnected to the

broader TMDS team.

In response to the possibility of incorporating an electronic clinical decision

support (eCDS) tool into the TMDS workflow, all nurses supported the adoption of this

approach. They believed it would increase efficiency, conserve resources, and improve

the quality of clinical records/data collection. However, it was noted that the tool should

not depend on a stable internet connection.

Communication. Given the critical importance of effective communication across the

multiple channels within the TMDS workflow, all participants were asked about their

experience communicating with staff and families. Call center nurses consistently cited

an unstable cellphone network as their primary barrier to communication with delivery

nurses and drivers. Similarly, delivery drivers also identified the unstable cellphone

network as their main obstacle to communicating with nurses, other drivers, and

families. The inability to reliably reach families by phone posed challenges to locating

homes, particularly in a setting without formal addresses. In such situations, drivers

relied on regional landmarks obtained during the virtual exams, along with their own

local knowledge to find households. Drivers suggested the use of radios or higher

quality cellphones to improve communication. They also recommended that nurses

prioritize collecting formal names of landmarks and individuals instead of nicknames.

Call center nurses, in turn, suggested several improvements, including ensuring that

delivery nurses and drivers are prepared at the beginning of each shift, having checked

that phones are charged and have sufficient cell minutes and data. The delivery nurses

proposed working from a dedicated office rather than from home, and additional training

with technology.

Expectations. Through both virtual and in-person interactions with families, nurses and

drivers identified several expectations families had for the TDMS. Nurses reported that

families generally expected and received reliable, timely care, and quality medication.

However, nurses felt that families also held expectations for services beyond the

program's scope, including care for patients older than 10 years, specific medications

(e.g., cough syrup), service during daytime hours, and financial assistance to access

emergency care when referred. As one nurse explained, *"Sometimes when a parent*

calls with a problem that surpasses our limits, we ask them to go to the hospital. The

parent usually expects that we would do something for them before they go to the

hospital. But if we send them to the hospital, there is no visit or medication that is sent

to that house." Drivers reported observing expectations similar to those noted by nurses

but also described that families expected to receive medications with every call. They

were generally unaware of the specific criteria (e.g., illness, severity, and location) that

determined eligibility for medication delivery.

Impact. All staff reported that their involvement with the TMDS had a positive impact on

their personal and professional lives. On a personal level, staff expressed increased

satisfaction from helping their communities and benefitting from higher income. Nurses

gained specific skills, including experience with the principles of telemedicine, pediatric

care, dosing medication, technology applications, and awareness of self-education

methods (e.g., referencing guidelines). Physicians shared that they found personal

satisfaction in providing more affordable and accessible healthcare to their community.

They also noted that exposure to telemedicine brought an innovative dimension to their

practice. Delivery drivers reported feeling greater respect and a sense of security from

belonging to an institution. Professionally, drivers increased their familiarity with

technology and gained a deeper geographic knowledge of their communities.

Staff reported that the TMDS had positively impacted the community as well.

They emphasized that the TMDS reduces barriers (e.g., financial, security,

transportation) to accessing healthcare, prevents illnesses from progressing into

emergencies, reduces overspending on hospital care, creates jobs in the community,

increases respect for disadvantaged people, and improves health literacy among the

community. Despite the overall positive impact of the TMDS on staff and the

community, there existed several challenges to the durability and scalability of the

intervention. Staff safety, potential for burnout, limited scope of the TMDS model,

funding, and the humanitarian and political crises in Haiti were presented by nurses and

drivers as risks to the long-term sustainability of the TMDS.

DISCUSSION

In this study, call center and delivery nurses, delivery drivers, and on-call physicians

shared their perspectives on the barriers and facilitators to scaling the TMDS healthcare

model through focus group sessions and written questionnaires. The areas of

concentration spanned the entire workflow from virtual exams at the call center to in-

person exams at the household, as well as communication systems connecting staff to

each other and families. Expectations and impact were assessed to determine the

extent of alignment with the broader goal of improving accesses to pre-emergent care

for children. The results have enabled staff and study leadership to develop solutions

aimed at refining the service in a way that anticipates scaling (Table 2). The findings

can also provide direction to other teams faced with similar challenges.

The process of evaluation through focus groups, written questionnaires and a

framework matrix analysis inspired an unanticipated higher-ordered synthesis and

visualization of the strengths and weaknesses of the TMDS model (Figure 2). This

visualization divides the findings into two domains: procedural workflow and governing

mission. Once the workflow is set to a virtual or in-person encounter, nurses must have

immediate situational awareness and adapt accordingly. This involves a cognitive

process of decision making and a series of physical actions. In parallel, the situational

awareness and adaptations require communication with a spectrum of different types of

people with variable access to connectivity (e.g., cellular networks). These activities are

hard, stressful and carry risk. Motivation to succeed with these difficult tasks is founded

in a commitment to mission. This mission extends from self to community, motivates

personal and team-based creative problem-solving, and requires recognition that the

service has limited scope and capacity.

With this visualization in mind, several solutions were identified, relating to both

the cognitive and physical challenges. Cognitive challenges included struggling to

effectively phrase questions to conduct productive virtual exams, as well as difficulty

managing situations outside the scope of the established workflow and clinical

resources. While nurses mitigated these challenges through experience, the time

required to gain sufficient experience poses a barrier to scaling. To address this,

consolidating acquired knowledge into a 'tip sheet' and protocolizing how to handle

edge-cases might expedite the training of new providers. Additionally, implementing an

eCDS tool that digitizes the clinical guidelines may streamline onboarding and provide

structured guidance for conducting exams. Finally, refining components of the virtual

and in-person exam processes could further enhance the quality of treatment plans.

Solutions to address the physical challenges focused on strengthening the

technology infrastructure and innovating resources. In settings like Haiti, where

telecommunication services are unreliable yet critical for telemedicine, improving the

telecommunication infrastructure at its source may not be feasible. However,

establishing multiple contingencies at each touchpoint ('hardening') can help minimize

communication barriers. Additionally, households where in-person exams are

conducted lack the resources typically found in clinics. To improve the working

conditions, innovating materials to provide adequate lighting, a designated workspace,

and improved organizational capacity would strengthen the quality and efficiency of in-

person encounters.

Finally, unmet expectations from both families and staff, along with risks to the

long-term sustainability of the TMDS, present obstacles to the mission. During this initial

pilot phase, the scope of the TMDS was intentionally limited in terms of age, clinical

offerings, hours of operation, and workflow to establish proofs of concept. As the TMDS

is positioned for scale, the scope may expand. Until then, aligning the scope with staff

and community expectations can be achieved through clear and consistent messaging.

Furthermore, an uncertain path to sustainability poses a barrier to maintaining high

levels of impact and trust among both staff and community members. To mitigate these

risks, increasing and diversifying partnerships, while ensuring high standards for

working conditions and professional development, may improve the likelihood of long-

term scalability and sustainability.

Our results align with, as well as differ from, the experiences of other

telemedicine services. In the telemedicine literature, staff and end-user acceptability are

cited as common barriers [39]. While we did not encounter opposition to telemedicine,

misaligned expectations suggest underlying concerns related to acceptability. Another

frequently cited barrier involves factors related to the implementation environment, such

as nationwide policies and legal liability [28, 40]. These concerns did not apply to our

controlled pilot implementation but may become relevant as the service scales. The

limited technical abilities of staff were identified as a barrier in our study and is

consistent with findings from other research [39, 41]. Additionally, most telemedicine

services, particularly those operating in low resource settings, face persistent

challenges related to unreliable telecommunication infrastructure [40, 41].

This study was not without limitations. First, data on family expectations and

community impact were based solely on staff impressions, which may have introduced

bias. Future research could address this limitation by directly enrolling families as

participants to obtain firsthand insights. Second, the translation process could have

introduced information bias, potentially altering the nuances and exact meanings of

responses from the original language. To mitigate this risk we worked with bilingual

facilitators and had three team members involved in translating and confirming

transcriptions before analysis began. Third, conducting focus groups via video

conference may have introduced a second form of information bias, as some sections of

the recordings were inaudible due to unstable connectivity. Despite these limitations,

the study yielded important findings with both internal value for the TMDS and broader

relevance to the telemedicine research community.

CONCLUSION

Telemedicine has secured a permanent role in healthcare delivery globally. However,

scaling novel telemedicine applications presents challenges. To better understand these

challenges, we used a framework matrix analysis to synthesize insights from focus

group discussions and written questionnaires completed by TMDS staff. The findings

highlighted key areas for improvement and led to the development of essential action

items, organized into three domains; conceptual, physical and mission-oriented. These

action items will guide internal efforts to position the MotoMeds TMDS for scale and will

hopefully catalyze external improvements among like-minded telemedicine initiatives.

 TABLES

 Table 1. TMDS staff participant characteristics.

Characteristic	Call center nurses[a] n=8	Delivery nurses[b] n=6	On-call physicians[c] n=3	Delivery drivers[d] n= 13
Age; median yrs (min, max) or values	31 (27-39)	31 (27-42)	36, 51, 66	36 (27-40)
Relevant professional experience; median yrs (min, max) or values	9 (3-11)	7 (1-14)	10, 15, 34	12 (5-22)
Experience working for TMDS; median yrs (min, max) or values	3 (1-3)	1	3	3 (1-3)

[a] Call center nurses conduct virtual exams at the call center and relay treatment plans to delivery nurses.
[b] Delivery nurses receive and enact virtual treatment plans by conducting in-person household exams or dispatching a motorcycle delivery driver.
[c] On-call physicians provide consultations by phone.
[d] Delivery drivers transport nurses and deliver medications/fluids to households by motorcycle.

430 **Table 2. TMDS staff perspectives on barriers to scale and potential solutions.**

Domain	Challenges	Solutions
Virtual exam	• Lack of confidence in the accuracy of patient details provided by parents over the phone	• Request 'read back' from parents to ensure two-way understanding • Provide a 'tip sheet' with common follow-up and exploratory questions • Establish a standardized pathway for handling situations when the nurse is uncertain about the quality of information • Eliminate or modify unproductive components of the virtual exam
Call center environment	• Technological barriers (e.g., unstable cellular signal, changing user interfaces)	• Enhance the robustness of technology infrastructure and establish contingencies • Strengthen and diversify technology training
In-person exam	• Restrictive scope of workflow and clinical resources	• Develop strategies to address barriers that prevent families from seeking emergency care • Facilitate interactions with hospital access points • Refine the workflow to accommodate situations in which additional patients are encountered during a household visit
Household environment	• Maneuvers to arrive at the household and function effectively once there	• Assure dispatch of nurses to conduct in-person exams is indicated • Innovate nurse equipment to increase lighting, maximize space, and improve durability
Clinical resources	• Limitations on medication dosing and edge cases • Informal workflow for consulting on-call physicians	• Incorporate clinical decision support tools for medication dosing • Protocolize edge cases within the clinical resources • Create structured systems for presenting cases and outcomes to on-call physicians
Communication	• Unstable cellular connectivity disrupts communication chains • Unreliable patient addresses	• Create an office in the delivery zone, rather than utilize mobile workstations • Focus on obtaining proper names for individuals and locations
Expectations	• Family expectations exceed the scope of the TMDS	• Improve messaging on the scope of a TMDS with staff and within the community
Impact	• Risks to long-term sustainability could limit impact	• Foster in-depth partnerships and distribute risk • Ensure standards of professionalism are supported and continue offering professional development opportunities

 FIGURES

 FIGURE 1

Figure 1. General workflow of a telemedicine and medication delivery service (TMDS). The parent of a sick child contacts the call center. A TMDS nurse receives the call and conducts a virtual exam by asking the parent targeted questions about the child and their symptoms. The nurse references clinical resources to create a treatment plan that includes disposition, medications/fluids, and follow-up recommendations. Families are contacted by phone at 24 hours and/or 10 days. A version of this figure was published previously[35].

447 **FIGURE 2**
448
449

450
451 **Figure 2.** Positioning a TMDS for scale; barriers and solutions between operations and
452 mission.
453
454

ADDITIONAL FILES

Additional File 1. PDF. Agenda to guide focus group discussions.

Additional File 2. PDF. Questionnaire for call center and delivery nurses (INACT3)

ADDITIONAL INFORMATION

Abbreviations. telemedicine and medication delivery service (TMDS), Consolidated Framework for Implementation Research (CFIR), communication technology infrastructure (ICT), Improving Nighttime Access to Care and Treatment- Haiti (INACT-H), information technology (IT), electronic clinical decision support (eCDS)

Ethics statement. This study was reviewed and approved by the University of Florida Institutional Review Board (202002693) as well as the Comité National de Bioéthique (National Bioethics Committee of Haiti; Ref2021-11). Consent process described in manuscript.

Consent for publication. Not applicable.

Availability of data and materials. The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: Authors declare that they have no competing interests.

Code availability. Not applicable.

Funding. This work was supported by grants to EJN by the Children's Miracle Network at University of Florida and National Institutes of Health [5R21TW012332-02]. The funders had no direct role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Private donations were made to the University of Florida Foundation to support the TMDS service in Haiti.

Author contributions.

Conceptualization: EJN, MBK, TKB, RKR,

Implementation of the study: FEG, KC, MBK

Statistical analyses and visualizations: FEG, KC, KF, EJN, MBK

Funding acquisition: MBK, EJN

Supervision: MBK, EJN, VBR, JRB, LE, CB

Writing – original draft: MBK, EJN

Writing – review & editing: RKR, FEG, KC, YC, KEF, JRB, TKB, LE, VBR, CB, MBK,

EJN

All authors read and approved the final manuscript.

Acknowledgements: We would like to thank the TMDS staff for their enrollment and

participation in this study. We would also like to acknowledge the University of Florida

Department of Pediatrics administrators Randy Autrey, Krista Berquist and Brittney

Johnson. Additionally, we express our appreciation for the academic support provided

by Glenn Morris at the Emerging Pathogens Institute, and Rashmin Savani Chair of the

Department of Pediatrics.

REFERENCES

1. Bhaskar S, Bradley S, Chattu VK, Adisesh A, Nurtazina A, Kyrykbayeva S, Sakhamuri S, Yaya S, Sunil T, Thomas P *et al*: **Telemedicine Across the Globe-Position Paper From the COVID-19 Pandemic Health System Resilience PROGRAM (REPROGRAM) International Consortium (Part 1)**. *Frontiers in Public Health* 2020, **8**.
2. Ndwabe H, Basu A, Mohammed J: **Post pandemic analysis on comprehensive utilization of telehealth and telemedicine**. *Clinical eHealth* 2024, **7**:5-14.
3. Sood S, Mbarika V, Jugoo S, Dookhy R, Doarn CR, Prakash N, Merrell RC: **What is telemedicine? A collection of 104 peer-reviewed perspectives and theoretical underpinnings**. *Telemed J E Health* 2007, **13**(5):573-590.
4. **Telemedicine Opportunities and Developments in Member States** [https://apps.who.int/iris/bitstream/handle/10665/44497/9789241564144_eng.pdf]
5. Liu P, Wang F, Xu W, Li Y, Li B: **Trends and frontiers of research on telemedicine from 1971 to 2022: A scientometric and visualisation analysis**. *Journal of Telemedicine and Telecare* 2023, **29**(9):731-746.
6. Harst L, Otto L, Timpel P, Richter P, Lantzsch H, Wollschlaeger B, Winkler K, Schlieter H: **An empirically sound telemedicine taxonomy – applying the CAFE methodology**. *Journal of Public Health* 2022, **30**(11):2729-2740.
7. **Telehealth for emergency departments. Tele-triage** [https://telehealth.hhs.gov/providers/best-practice-guides/telehealth-for-emergency-departments/tele-triage#:~:text=Tele%2Dtriage%20can%20be%20used,come%20to%20the%20emergency%20department]
8. McConnochie KM, Conners GP, Brayer AF, Goepp J, Herendeen NE, Wood NE, Thomas A, Ahn DS, Roghmann KJ: **Effectiveness of Telemedicine in Replacing In-Person Evaluation for Acute Childhood Illness in Office Settings**. *Telemedicine and e-Health* 2006, **12**(3):308-316.
9. Mahmoud K, Jaramillo C, Barteit S: **Telemedicine in Low- and Middle-Income Countries During the COVID-19 Pandemic: A Scoping Review**. *Front Public Health* 2022, **10**:914423.
10. Alnasser Y, Proaño A, Loock C, Chuo J, Gilman RH: **Telemedicine and Pediatric Care in Rural and Remote Areas of Middle-and-Low-Income Countries: Narrative Review**. *J Epidemiol Glob Health* 2024, **14**(3):779-786.
11. Moyo J, Madziyire G: **Use of telemedicine in obstetrics and gynaecology in Zimbabwe during a lockdown period**. *Pan Afr Med J* 2020, **35**(Suppl 2):89.
12. Klarman MB, Flaherty KE, Chi X, Cajusma Y, Capois AC, Vladimir Dofiné MD, Exantus L, Friesen J, Beau de Rochars VM, Baril C *et al*: **Implementation of a pediatric telemedicine and medication delivery service in a resource-limited setting: A pilot study for clinical safety and feasibility**. *The Journal of Pediatrics* 2022.
13. Flaherty KE, Klarman MB, Zakariah AN, Mahama MN, Osei-Ampofo M, Nelson EJ, Becker TK: **Evaluating the prerequisites for adapting a paediatric nighttime telemedicine and medication delivery service to a setting with high malarial burden: A cross-sectional pre-implementation study**. *Trop Med Int Health* 2023, **28**(9):763-770.
14. McCannon CJ, Berwick DM, Massoud MR: **The Science of Large-Scale Change in Global Health**. *JAMA* 2007, **298**(16):1937-1939.
15. Hanson K, Ranson MK, Oliveira-Cruz V, Mills A: **Expanding access to priority health interventions: a framework for understanding the constraints to scaling-up**. *Journal of International Development* 2003, **15**(1):1.

16.   Zomahoun HTV, Ben Charif A, Freitas A, Garvelink MM, Menear M, Dugas M,
Adekpedjou R, Légaré F: **The pitfalls of scaling up evidence-based interventions in**
health. *Glob Health Action* 2019, **12**(1):1670449.
17.   **ExpandNet** [https://expandnet.net/]
18.   Bulthuis SE, Kok MC, Raven J, Dieleman MA: **Factors influencing the scale-up of**
public health interventions in low- and middle-income countries: a qualitative
systematic literature review. *Health Policy Plan* 2020, **35**(2):219-234.
19.   Damschroder LJ, Reardon CM, Widerquist MAO, Lowery J: **The updated Consolidated**
Framework for Implementation Research based on user feedback. *Implement Sci*
2022, **17**(1):75.
20.   Huang F, Cheng S, Du X, Chen W, Scano F, Falzon D, Wang L: **Electronic recording**
and reporting system for tuberculosis in China: experience and opportunities. *J*
Am Med Inform Assoc 2014, **21**(5):938-941.
21.   Ndlovu K, Scott RE, Mars M: **Interoperability opportunities and challenges in linking**
mhealth applications and eRecord systems: Botswana as an exemplar. *BMC*
Medical Informatics and Decision Making 2021, **21**(1):246.
22.   Spicer N, Berhanu D, Bhattacharya D, Tilley-Gyado RD, Gautham M, Schellenberg J,
Tamire-Woldemariam A, Umar N, Wickremasinghe D: **'The stars seem aligned': a**
qualitative study to understand the effects of context on scale-up of maternal and
newborn health innovations in Ethiopia, India and Nigeria. *Globalization and Health*
2016, **12**(1):75.
23.   Adebayo PB, Oluwole OJ, Taiwo FT: **COVID-19 and Teleneurology in Sub-Saharan**
Africa: Leveraging the Current Exigency. *Frontiers in Public Health* 2021, **8**.
24.   Mengiste SA, Antypas K, Johannessen MR, Klein J, Kazemi G: **eHealth policy**
framework in Low and Lower Middle-Income Countries; a PRISMA systematic
review and analysis. *BMC Health Services Research* 2023, **23**(1):328.
25.   Pérez-Noboa B, Soledispa-Carrasco A, Padilla VS, Velasquez W: **Teleconsultation**
Apps in the COVID-19 Pandemic: The Case of Guayaquil City, Ecuador. *IEEE*
Engineering Management Review 2021, **49**(1):27-37.
26.   Jiang Y, Sun P, Chen Z, Guo J, Wang S, Liu F, Li J: **Patients' and healthcare**
providers' perceptions and experiences of telehealth use and online health
information use in chronic disease management for older patients with chronic
obstructive pulmonary disease: a qualitative study. *BMC Geriatrics* 2022, **22**(1):9.
27.   Aydemir S, Ocak S, Saygılı S, Hopurcuoğlu D, Haşlak F, Kıykım E, Aktuğlu Zeybek Ç,
Celkan T, Demirgan EB, Kasapçopur Ö *et al*: **Telemedicine Applications in a Tertiary**
Pediatric Hospital in Turkey During COVID-19 Pandemic. *Telemedicine and e-Health*
2020, **27**(10):1180-1187.
28.   Dodoo JE, Al-Samarraie H, Alzahrani AI: **Telemedicine use in Sub-Saharan Africa:**
Barriers and policy recommendations for Covid-19 and beyond. *International*
Journal of Medical Informatics 2021, **151**:104467.
29.   Hamilton AB, Finley EP: **Qualitative methods in implementation research: An**
introduction. *Psychiatry Res* 2019, **280**:112516.
30.   Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R, Hensley
M: **Outcomes for Implementation Research: Conceptual Distinctions,**
Measurement Challenges, and Research Agenda. *Administration and Policy in Mental*
Health and Mental Health Services Research 2011, **38**(2):65-76.
31.   Busetto L, Wick W, Gumbinger C: **How to use and assess qualitative research**
methods. *Neurological Research and Practice* 2020, **2**(1):14.
32.   Rosen RK, Gainey M, Nasrin S, Garbern SC, Lantini R, Elshabassi N, Sultana S, Hasnin
T, Alam NH, Nelson EJ *et al*: **Use of Framework Matrix and Thematic Coding**

Methods in Qualitative Analysis for mHealth: The FluidCalc app. *International*
Journal of Qualitative Methods 2023, **22**:16094069231184123.
33.   Klarman M, Schon J, Cajusma Y, Maples S, Beau de Rochars VEM, Baril C, Nelson EJ:
Opportunities to catalyse improved healthcare access in pluralistic systems: a
cross-sectional study in Haiti. *BMJ Open* 2021, **11**(11):e047367.
34.   Klarman MB, Chi X, Cajusma Y, Flaherty KE, Capois AC, Dofiné MDV, Exantus L,
Friesen J, Beau de Rochars VM, Becker T *et al*: **Development and evaluation of a**
clinical guideline for a paediatric telemedicine service in a low-resource setting.
BMJ Paediatr Open 2024, **8**(1).
35.   Klarman MB, Chi X, Cajusma Y, Flaherty KE, Beausejour JR, Exantus L, Beau de
Rochars VM, Baril C, Becker TK, Gurka MJ *et al*: **Evaluation of a Scalable Design for**
a Pediatric Telemedicine and Medication Delivery Service: A Prospective Cohort
Study in Haiti. *medRxiv* 2024:2024.2008.2016.24312128.
36.   Flaherty KE, Klarman MB, Cajusma Y, Schon J, Exantus L, Beau de Rochars VM, Baril
C, Becker TK, Nelson EJ: **A Nighttime Telemedicine and Medication Delivery**
Service to Avert Pediatric Emergencies in Haiti: An Exploratory Cost-
Effectiveness Analysis. *The American Journal of Tropical Medicine and Hygiene* 2022,
106(4):1063-1071.
37.   Biswas D, Hossin R, Rahman M, Bardosh KL, Watt MH, Zion MI, Sujon H, Rashid MM,
Salimuzzaman M, Flora MS *et al*: **An ethnographic exploration of diarrheal disease**
management in public hospitals in Bangladesh: From problems to solutions.
Social Science & Medicine 2020, **260**:113185.
38.   Rosen RK, Garbern SC, Gainey M, Lantini R, Nasrin S, Nelson EJ, Elshabassi N, Alam
NH, Sultana S, Hasnin T *et al*: **Designing a Novel Clinician Decision Support Tool**
for the Management of Acute Diarrhea in Bangladesh: Formative Qualitative
Study. *JMIR Hum Factors* 2022, **9**(1):e33325.
39.   Scott Kruse C, Karem P, Shifflett K, Vegi L, Ravi K, Brooks M: **Evaluating barriers to**
adopting telemedicine worldwide: A systematic review. *J Telemed Telecare* 2018,
24(1):4-12.
40.   Mamuye A, Nigatu AM, Chanyalew MA, Amor LB, Loukil S, Moyo C, Quarshie S,
Antypas K, Tilahun B: **Facilitators and Barriers to the Sustainability of eHealth**
Solutions in Low- and Middle-Income Countries: Descriptive Exploratory Study.
JMIR Form Res 2023, **7**:e41487.
41.   Sagaro GG, Battineni G, Amenta F: **Barriers to Sustainable Telemedicine**
Implementation in Ethiopia: A Systematic Review. *Telemedicine Reports* 2020,
1(1):8-15.
629