Effects of emotional arousal on amygdala neural response and cigarette craving reduction

during repeated exposure to graphic warning labels

Astrid P. Ramos-Rolón, Daniel D. Langleben, Kevin G. Lynch, Corinde E. Wiers, Zhenhao Shi

Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

Corresponding author:

Zhenhao Shi, PhD

3535 Market Street, Suite 500

Philadelphia, PA 19104

Email: zhshi@upenn.edu

ABSTRACT

Graphic warning labels (GWLs) have been implemented on cigarette packaging worldwide. In the U.S., GWLs

have encountered legal obstacles based on the tobacco industry arguments that their aversive imagery

unnecessarily triggers strong emotional arousal. This longitudinal study evaluated the effect of the high-arousal

GWLs on cigarette craving and the neural substrates of emotional processing. 158 adult smokers were exposed

to either high-arousal (n=79) or low-arousal (n=79) GWLs that were attached to their own cigarette packs for 4

weeks. Craving and brain activity in response to GWLs and control stimuli were measured using functional

magnetic resonance imaging before and after the 4-week exposure. The amygdala, which plays a key role in

emotional processing, served as the a priori region of interest. Results indicate that, at baseline, high-arousal

GWLs elicited a larger reduction in craving and stronger amygdala neural response compared to low-arousal

GWLs; however, at week 4, GWL-induced craving reduction and amygdala response became comparable

between the high-arousal and low-arousal groups. Amygdala response mediated GWLs' effects on craving

reduction, which was moderated by arousal and time in such a way that the amygdala's mediating role was more

pronounced for high-arousal than low-arousal GWLs at baseline but did not differ between groups at week 4.

Together, the results suggest that the impact of emotional arousal on cigarette cravings decreases over time,

potentially due to the amygdala's diminishing responsivity to repeated presentations of high-arousal imagery.

Low-arousal GWLs may represent a more feasible approach for tobacco control efforts in the U.S.

1   INTRODUCTION

Tobacco use is a major public health issue, with over 1.2 billion users worldwide in 2022 (World Health

Organization, 2024). Combustible cigarette smoking remains the most prevalent form of tobacco consumption

(World Health Organization, 2024). In the past two decades, many countries have implemented graphic warning

labels (GWLs) on cigarette packages (Hiilamo et al., 2014). The U.S. first adopted text-only warnings in 1966,

and in 2009, Congress tasked the Food and Drug Administration (FDA) with GWL implementation (Hiilamo et

al., 2014). Using data from an online survey (Nonnemaker et al., 2010), the FDA selected the GWLs that evoked

the highest emotional response (ER) and mandated them in a Final Rule (U.S. Food and Drug Administration,

2011). However, this Final Rule was overturned in court after tobacco companies successfully argued that the

GWLs were overly emotionally provocative and infringed on their First Amendment rights to commercial free

speech ("R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al.," 2012). In 2020,

the FDA introduced a new set of 11 GWLs, which have not been tested for ER (U.S. Food and Drug

Administration, 2022). The new GWLs faced similar legal challenges ("R.J. Reynolds Tobacco Company, et al.

v. U.S. Food & Drug Administration, et al.," 2022; "R.J. Reynolds Tobacco Company, et al. v. U.S. Food &

Drug Administration, et al.," 2024b) and were blocked again in January 2025 ("R.J. Reynolds Tobacco

Company, et al. v. U.S. Food & Drug Administration, et al.," 2025).

This debate has spurred research into the risk vs. benefit of using highly emotional imagery in GWLs. Studies

show that GWLs evoking a higher ER lead to a greater immediate reduction in cigarette craving and better GWL

recall compared to lower-ER GWLs (Wang, Lowen, et al., 2015; Wang, Romer, et al., 2015). However, research

to date has largely focused on the short-term effects of GWLs, although in real life, smokers would be exposed

to GWLs in a repeated and chronic manner. To bridge this gap, our group examined the longer-term effects of

exposure to high-ER vs. low-ER GWLs (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023). We found that

compared to high-ER GWLs, a month-long exposure to low-ER GWLs was associated with a greater reduction

in cigarette consumption. These findings appear to contradict the cross-sectional evidence that postulated greater

efficacy of high-ER GWLs largely based on intermediate outcome variables such as recall, attitudes and

intentions. Reconciling the cross-sectional and longitudinal findings is crucial for a full mechanistic

understanding of the dynamic impact of ER on GWL efficacy.

Craving is a central characteristic of substance use disorders (American Psychiatric Association, 2013) and a

key motivational driver of drug seeking (Robinson & Berridge, 1993). Craving for cigarettes robustly

precipitates subsequent smoking (Carter et al., 2008; Motschman et al., 2018; Shiffman et al., 2002) and predicts

cessation failure (Fidler et al., 2011). While smoking temporarily alleviates craving (Carter et al., 2008), chronic

smoking exacerbates craving in the long term which can persist after extended abstinence (Robinson &

Berridge, 1993). Reducing craving is an important goal in the clinical and public health interventions of tobacco

use (Do & Galván, 2014; Shiffman et al., 2004; Wang, Lowen, et al., 2015; Wang, Romer, et al., 2015). We

previously showed that, cross-sectionally, high-ER GWLs produced greater reductions in craving and cue-

reactivity than low-ER GWLs (Wang, Lowen, et al., 2015; Wang, Romer, et al., 2015). However, the salience of

affective stimuli (e.g., GWLs) decreases after repeated exposure due to habituation of the brain's emotion

circuits (Fischer et al., 2000; Fridriksson et al., 2018; Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023).

Specifically, using functional magnetic resonance imaging (fMRI), we showed that GWL-induced neural

response in the amygdala, a key region for emotional processing (LeDoux, 2007), was initially greater for high-

ER GWLs than low-ER GWLs, but the difference was markedly reduced after one month of chronic GWL

exposure (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023). Such habituation may attenuate high-ER

GWLs' greater efficacy in reducing craving compared to low-ER ones.

The aim of the present study was twofold: to supplement cross-sectional data (Wang, Lowen, et al., 2015) with

longitudinal evidence on the relative efficacy of high-ER vs. low-ER GWLs; and to extend our prior research on

cigarette consumption (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023) by examining the motivational

underpinning (i.e., craving) and neural correlates (Wang, Lowen, et al., 2015). Adult smokers who were

randomly assigned to the high-ER and low-ER groups underwent fMRI while viewing GWLs and control

stimuli and reported momentary craving. The fMRI assessment was repeated after four weeks of naturalistic

GWL exposure, as described elsewhere (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023). Briefly,

exposure was modeled by providing participants with cigarette packs carrying the high-ER or low-ER GWLs.

We hypothesized that, while high-ER GWLs would initially induce greater amygdala response and greater

reduction in cigarette craving compared to low-ER GWLs, the impact of ER would diminish after repeated

GWL exposure due to emotional habituation (Fischer et al., 2000; Fridriksson et al., 2018; Shi, Wang, Fairchild,

Aronowitz, Padley, et al., 2023). We further hypothesized that amygdala neural activity would be associated

with craving reduction and mediate the diminishing effect of ER on GWL efficacy.

MATERIALS AND METHODS

Participants

168 adult tobacco smokers were recruited from the Philadelphia area and randomly assigned to either the high-

ER group or the low-ER group (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023). Eight participants were

excluded from the current analysis due to missing fMRI data, and two were excluded due to excessive head

motion during fMRI, resulting in a total of 158 participants included in the final analysis (n=79 for each group).

Inclusion criteria were: 1) 18 to 65 years old; 2) currently smoking ≥2 cigarettes per day; 3) ≥100 cigarettes

smoked during lifetime; 4) not currently enrolled in any smoking cessation programs; 5) not currently using

other nicotine products (chewing tobacco, cigars, electronic cigarettes, etc.); 6) not a member of the household

or friend of an already enrolled study participant; 7) no medical or psychiatric disorders or treatments that may

affect the cerebrovascular system; 8) no contraindications for fMRI; 9) (if female) not pregnant or breast-

feeding. Participants were instructed to avoid using any other nicotine products throughout the study. All

participants provided written informed consent to participate in the protocol approved by the University of

Pennsylvania Institutional Review Board.

Study design

Participants were enrolled seven days before GWL exposure (day −7) and received a week's supply of cigarettes

of their preferred brands in their original packaging without GWLs. Upon enrollment, demographic information

and age of smoking initiation were collected, and nicotine dependence was measured by the Fagerström Test for

Nicotine Dependence (FTND) (Heatherton et al., 1991). On days 0, 7, 14, and 21, participants received weekly

cigarette supply in packages that carried either high- or low-ER GWLs based on their assigned experimental

group (see **Figure 1**). To discourage participants from stockpiling the cigarettes, all packages had been unsealed

prior to being dispensed. Brain activity and cigarette craving in response to GWLs were measured using fMRI

immediately before (day 0) and after (day 28) the four weeks' GWL exposure. Other measures included the

number of cigarettes smoked per day during the past seven days (CPD), urine concentration of nicotine

metabolite cotinine, desire to quit smoking, self-efficacy, and perceived GWL effectiveness. Details about these

measures and their results were reported in Shi, Wang, Fairchild, Aronowitz, Padley, et al. (2023).

[Figure 1]

Study stimuli

The FDA evaluated the ER to 36 GWLs composed of colored images paired with one of nine textual warnings

using an online survey with over 18,000 respondents (Nonnemaker et al., 2010). Out of these 36 GWLs, we

selected twelve GWLs with higher ER scores and twelve with lower ER scores (mean±SD=18.21±1.25 versus

16.06±1.09, Mann-Whitney U=207, p=0.001) that carried the same textual warnings and were comparable in

terms of image brightness, contrast, color saturation, depiction of human faces, depiction of smoking cues, and

usage of photographs as opposed to infographics (ps>0.12) (Shi, Wang, Fairchild, Aronowitz, Padley, et al.,

2023). The GWLs covered 50% of the front and back of the packages and were rotated using a Latin square

design so that each participant received approximately the same proportion of each specific GWL during the

four weeks of exposure.

fMRI data acquisition

fMRI was performed on a Siemens (Siemens, Erlangen, Germany) TIM Trio 3T system in 50 participants (26

high ER, 24 low ER) and a Siemens Prisma 3T system in 108 participants (53 high ER, 55 low ER; $\chi^2(1)=0.03$,

p=0.86). We previously showed that inter-scanner variability and its impact on neural activity measures were

minimal, allowing us to combine data in subsequent analysis (Shi, Wang, Fairchild, Aronowitz, Padley, et al.,

2023). The scanners used identical parameters outlined below. Blood oxygenation level-dependent (BOLD)

fMRI used a whole-brain, single-shot gradient-echo echo-planar sequence with the following parameters:

repetition time (TR)/echo time (TE)=2000/30 ms, field of view (FOV)=220 mm², matrix=64×64, slice

thickness/gap=3.4/0 mm, 32 slices, effective voxel resolution of 3.4×3.4×3.4 mm³. After BOLD fMRI, 5-minute

T1-weighted 3D magnetization-prepared gradient-echo structural images were acquired with the following

parameters: TR/TE=1810/3.51 ms, FOV=240×180 mm², matrix=256×192, effective voxel resolution of

0.94×0.94×0.94 mm³. An oblique acquisition oriented along the anterior commissure–posterior commissure line

allowed coverage of the entire brain except the lower cerebellum.

Participants completed the fMRI task at day 0 and day 28. To ensure that participants were in a non-deprived

state during the scans, they were escorted outside approximately 30 minutes before each fMRI session to smoke

one of their own cigarettes. The fMRI task followed a block design, with stimuli presented in six blocks of

GWLs and six blocks of control images. Each GWL block comprised six of the twelve high-ER or low-ER

GWLs, depending on the participant's group assignment. Each GWL was presented three times throughout the

task. Similarly, each control block contained six of twelve smoking-irrelevant control images (e.g., buildings)

paired with smoking-irrelevant text (e.g., "WARNING: Buildings are hazardous"), matching the anti-smoking

text warnings on the GWLs in terms of word count, font style, and font size. Each GWL and image was

presented for 2 seconds against a black background, resulting in a block duration of 12 seconds. Immediately

before and after each block, participants were asked to indicate their current level of cigarette craving by

answering the question "How much do you want to smoke a cigarette right now?" using a visual analog scale

from 0 ("not at all") to 10 ("extremely"). They were given 4 seconds to provide their response. Inter-block

intervals, indicated by a white crosshair in the center of the screen, lasted for 9.5 seconds. All stimuli were

delivered using Presentation (Neurobehavioral System Inc, Albany, CA) and presented using a rear projector

system that was viewed through a mirror mounted on the head coil in the scanner.

fMRI data processing

fMRI data were analyzed using SPM 12 (Wellcome Trust Centre for Neuroimaging, London, UK). Images were

adjusted for slice timing, realigned to the first scan to correct for head motion, normalized into stereotactic

Montreal Neurological Institute (MNI) space with 3-mm isotropic cubic voxels, and spatially smoothed by a

Gaussian filter with full-width at half-maximum parameter set to 8 mm. Severe head motion was defined as

motion greater than one voxel, a commonly used cutoff in the task-fMRI literature. Such a cutoff ensured the

robustness of SPM's default motion correction algorithm (spm_realign.m) (Friston et al., 1995; Friston et al.,

1996). The root-mean-squared framewise displacement of the remaining participants was minimal (range=0.02–

0.35 mm) (Jenkinson et al., 2002).

After preprocessing, we conducted individual-level general linear model (GLM) analysis using the "beta series"

approach (Zanto et al., 2011) to model each block with the canonical hemodynamic response function as a

single regressor. High-pass filter of 128 sec was applied. Six rigid body motion parameters (x, y, z, pitch, roll,

yaw) and their first-order derivatives were included as covariates, and the rest periods were treated as the

implicit baseline. The GLMs resulted in a total of twelve parameter-estimate maps for each participant and each

fMRI session (six for the GWL blocks and six for the control image blocks). The bilateral amygdala was

anatomically defined as an a priori regions of interest using the Automated Anatomical Labeling atlas (Tzourio-

Mazoyer et al., 2002). For each participant and each fMRI session, the percent signal change in the amygdala in

response to each GWL and control block was extracted for subsequent statistical analysis.

Statistical analysis

Statistical analysis was performed using R. We compared baseline participant characteristics between the high-

ER and low-ER groups using χ^2 tests for categorical variables and Mann-Whitney U-tests for numerical

variables. We then evaluated the external validity of the craving measure. To do so, we examined the association

of mean craving with nicotine dependence measured by FTND and with smoking severity indexed by CPD.

Mean craving was calculated by averaging all 24 craving ratings (i.e., before and after the six GWL and six

control image blocks) for each subject and each fMRI session. CPD was log-10 transformed due to high right-

skewness. A linear model examined the effect of nicotine dependence on mean craving at day 0 while

controlling for group (i.e., mean_craving ~ 1 + group + FTND). A linear mixed-effects (LME) model examined

the effect of CPD on mean craving across both the day-0 and day-28 time points with random subject-specific

intercepts while controlling for time and group [i.e., mean_craving ~ 1 + time + group + CPD + (1|subject)].

LME analysis was performed using the lme4 and lmerTest packages (Bates et al., 2015; Luke, 2017). Statistical

significance was evaluated with the Satterthwaite approximation for degrees of freedom (Luke, 2017).

The effect of GWLs and control images on cigarette craving during the fMRI task was calculated as the change

in craving ratings from before to after each GWL block and each control image block, respectively. Therefore,

there was a total of twelve scores of craving reduction (i.e., Δcraving) for each participant and each fMRI

session (six for the GWL blocks and six for the control image blocks). Similarly, twelve values of amygdala

percent signal change were extracted per participant per session. A series of LME models were conducted to test

for the study hypotheses. For all models, random intercepts were included to account for repeated measures

across GWL and control image blocks within subjects and subject-time combinations. We first tested the

diminishing role of ER in GWLs' ability to reduce cigarette craving after repeated exposure. Specifically, the

model examined the effects of time (day 0 vs. day 28), group (high-ER vs. low-ER), and stimulus (GWL vs.

control), on smoking reduction [i.e., the $c_{(\bullet)}$ paths in **Figure 5B**: Δcraving ~ 1 + time*group*stimulus +

(1|subject) + (1|subject:time)]. Post-hoc analysis was conducted to decompose three-way interactions using the

emmeans package, and multiple comparisons were adjusted using the Šidák method. For completeness and as a

replication of our previously reported analysis (Shi, Wang, Fairchild, Aronowitz, Padley, et al., 2023), a similar

LME model was used to examine the effects of time, group, and stimulus on amygdala neural activity [i.e., the

$a_{(\bullet)}$ paths in **Figure 5B**: amygdala ~ 1 + time*group*stimulus + (1|subject) + (1|subject:time)]. Next, we tested

the association between amygdala response and craving reduction [i.e., the b paths in **Figure 5B**: Δcraving ~ 1 +

amygdala + (1|subject) + (1|subject:time)].

Lastly, we tested whether amygdala response mediated high-ER GWLs' decreasing ability to reduce craving

after repeated exposure. We first tested an unmoderated mediation model, where stimulus was the input

variable, amygdala response was the mediator, and craving reduction was the outcome. We then tested a

moderated mediation model adapted from Hayes (2015) with time and group being the moderators (see **Figure**

5). Both models involved an LME analysis for the mediator (i.e., the $a_{(\bullet)}$ paths) and another LME analysis for

the outcome (i.e., the b and $c_{(\bullet)}$ paths combined). The index of moderated mediation was calculated as $a_7 \times b$. The

conditional indirect effect of stimulus on craving through amygdala activity, which varied as a function of time

and group, was calculated as $(a_3 + a_5 \times time + a_6 \times group + a_7 \times time \times group) \times b$. Bootstrap using case resampling

(10000 iterations) (Reiser et al., 2017) was performed to determine the bias-corrected and accelerated

confidence interval and significance of effects.

RESULTS

Participant characteristics

Participants of the high-ER and low-ER groups were comparable with regard to demographics and baseline

characteristics (see **Table 1**). All participants completed the first fMRI session, and 140 participants (89%)

completed the second fMRI session. The completion rate for the second fMRI session was comparable between

the high-ER and low-ER groups (n=71 vs. 69, $\chi^2(1)$=0.06, p=0.80).

12   [Table 1]

External validity of craving

Mean cigarette craving across all ratings during the MRI session at day 0 was positively associated with baseline

nicotine dependence indexed by FTND (β=0.22, 95% confidence interval [CI]=[0.06,0.39]; F(1,153)=7.18,

p=0.008). Mean cigarette craving was also positively associated with smoking severity indexed by CPD across

day 0 and day 28 (β=2.21, 95% CI=[0.88,3.53]; F(1,205.34)=10.61, p=0.001). There was a trend for a main

effect of time on mean craving, with marginally higher mean craving at day 0 than day 28 (4.12 vs. 3.71,

difference=0.41, 95% CI=[–0.003,0.83]). The main effect of group and the time×group interaction were not

significant (p=0.68 & 0.78).

Effect of GWLs on craving

There was a significant time×group×stimulus interaction on craving reduction (F(1,2919.96)=9.944, p=0.002)

(see **Figure 2**). Post-hoc analysis showed significant group×stimulus interaction at day 0 (p<0.001), such that

the reduction of craving induced by the GWLs (vs. control images) was greater in the high-ER group (–0.81 vs.

0.37, difference=–1.18, 95% CI=[–1.44,–0.92]) than the low-ER group (–0.50 vs. 0.09, difference =–0.59, 95%

CI=[–0.85,–0.33]). The group×stimulus interaction was not significant at day 28 (p=0.86), with comparable

effect of GWLs vs. control images on the reduction of craving between the high-ER group (–0.38 vs. 0.00,

difference=–0.38, 95% CI=[–0.65,–0.12]) and the low-ER group (–0.49 vs. –0.04, difference=–0.45, 95% CI=[–

0.72,–0.19]).

8   [Figure 2]

Effect of GWLs on amygdala neural activity

There was a significant time×group×stimulus interaction on amygdala neural activity ($F_{(1,3274)}$=5.36, p=0.021)

(see **Figure 3**). Post-hoc analysis showed significant group×stimulus interaction at day 0 (p<0.001), such that

amygdala response to GWLs (vs. control images) was greater in the high-ER group (0.18 vs. –0.03,

difference=0.21, 95% CI=[0.13,0.29]) than the low-ER group (0.04 vs. 0.02, difference=0.03, 95% CI=[–

0.05,0.10]). The group×stimulus interaction was not significant at day 28 (p=0.69), with comparable amygdala

response to GWLs vs. control images between the high-ER group (0.26 vs. 0.13, difference=0.13, 95%

CI=[0.04,0.21]) and the low-ER group (0.08 vs. –0.01, difference=0.09, 95% CI=[0.01,0.17]). These results

were consistent with the original analysis reported in Shi, Wang, Fairchild, Aronowitz, Padley, et al. (2023).

18   [Figure 3]

19   *Association between craving reduction and amygdala neural activity*

20   Greater amygdala response was associated with greater reduction in cigarette craving (β=–0.15, 95% CI=[–

21   0.22,–0.08]; $F_{(1,703.36)}$=16.79, p<0.001) (see **Figure 4**). The association remained significant when controlling

22   for the effects of time, group, stimulus, and their interactions (β=–0.11,95% CI=[–0.19,–0.04];

23    $F_{(1,721.39)}$=9.56, p=0.002).

1   [Figure 4]

Mediation analysis

We found a significant unmoderated indirect effect of stimulus on craving reduction through amygdala neural

activity (effect=−0.006, 95% CI=[−0.01,−0.001]; p=0.022), suggesting a key role of amygdala in mediating

GWLs' impact on craving. We then examined the moderated mediation model, as shown in **Figure 5**. We found

a significant moderated mediation, such that the mediator effect was moderated by the group×time interaction

(effect=−0.002, 95% CI=[−0.005,−0.0001]; p=0.035) (see **Figure 6**). Post-hoc analysis showed that, at day 0, the

indirect effect of stimulus on craving reduction through amygdala activity was stronger for the high-ER group

(β=−0.011, 95% CI=[−0.021,−0.001]) than the low-ER group (β=−0.0004, 95% CI=[−0.005,0.003]; p=0.020).

At day 28, the indirect effect was comparable between the high-ER group (β=−0.007, 95% CI=[−0.018,−0.001])

and the low-ER group (β=−0.005, 95% CI=[−0.013,−0.0005]; p=0.41).

12   [Figure 5]

13   [Figure 6]

DISCUSSION

We found that high-ER GWLs initially elicited a stronger amygdala response and greater craving reduction

compared to low-ER GWLs, similar to what we observed in our prior cross-sectional study (Wang, Lowen, et

al., 2015; Wang, Romer, et al., 2015). However, these differences diminished with repeated GWL exposure,

such that by day 28, amygdala response and craving reduction became comparable between the high-ER and

low-ER groups. Mediation analyses further revealed that amygdala response mediated the effects of GWLs on

craving reduction. The mediating effect was stronger for high-ER than low-ER GWLs at baseline but not at day

28.

The diminishing effect of ER on amygdala response is consistent with emotional habituation, a process in which

an individual's reaction to repeatedly presented affective stimuli gradually decreases over time. While our data demonstrated amygdala habituation over four weeks of chronic GWL exposure, previous studies have reported more rapid amygdala adaptation occurring within seconds when identical GWLs were presented consecutively (Fridriksson et al., 2018). Habituation is a fundamental form of learning that optimizes cognitive resource allocation and facilitates the differentiation between familiar and novel stimuli. fMRI studies have specifically shown amygdala habituation to emotional stimuli across various modalities, including visual (Breiter et al., 1996; Plichta et al., 2014; Wright et al., 2001), auditory (Mutschler et al., 2010), and olfactory (Poellinger et al., 2001). Despite the decreasing amygdala activity, we previously showed that high-ER GWLs were better remembered over time (Wang et al., 2019). This indicates that while amygdala-mediated emotional processing may facilitate encoding, it may not be necessary for retention. Taken together, our findings highlight the importance of considering emotional habituation in the design of anti-smoking warnings, such as GWLs, that are intended for prolonged exposure among smokers. Strategies to mitigate habituation, such as reducing exposure frequency and increasing stimulus diversity, have shown promise; however, their effectiveness also wanes over time (Rankin et al., 2009). Further research is needed to evaluate the utility and feasibility of alternative approaches for enhancing GWLs' ability to emotionally engage smokers in the long term.

In addition to habituation, prolonged exposure may trigger reactance, a psychological resistance to perceived persuasion attempts, leading smokers to consider the warnings as threatening and thus dismiss or counterargue them (Brehm & Brehm, 2013). GWLs have been shown to lead to reactance more so than textual anti-smoking warnings (Erceg-Hurn & Steed, 2011; LaVoie et al., 2015; Maynard et al., 2014). Reactance to GWLs has been associated with reduced intention to quit smoking and more cigarettes smoked per day (Hall et al., 2018; Hall et al., 2017) [but see Cho et al. (2016)]. Moreover, chronic GWL exposure may also lead to message fatigue, where individuals become tired of seeing the warnings and perceive them as annoying and uninformative (So et al., 2017). Over time, message fatigue can result in reduced attention, lower emotional engagement, and decreased motivation to change behaviors (Keating & Skurka, in press). Our finding that high-ER GWLs' ability to reduce craving diminished over time suggest that ER may contribute to reactance and message fatigue elicited by GWLs. Although we did not explicitly measure reactance and message fatigue, our recent

exploratory analysis of the same dataset showed that, after completing the four weeks' GWL exposure, smokers

expressed a moderate sense of relief from having to see the GWLs (Shi et al., 2024). While such sense of relief

did not differ between the high-ER and low-ER groups, it was only within the high-ER group that greater relief

predicted less smoking (Shi et al., 2024). Therefore, addressing reactance and message fatigue may potentially

help enhance the behavioral efficacy of high-ER GWLs.

Our previous work examined GWL effects on smoking behavior (Shi, Wang, Fairchild, Aronowitz, Padley, et

al., 2023), while the current study focused on a key motivational underpinning of smoking, i.e., craving. Both

studies indicate that low-ER GWLs are non-inferior to high-ER GWLs during chronic naturalistic exposure.

However, while we previously showed that low-ER GWLs was more effective in reducing smoking (indexed by

the number cigarettes per day), here we found comparable efficacy between high-ER and low-ER GWLs in

reducing craving after the exposure, despite high-ER ones being more effective initially. Several factors may

explain the differences in the two studies' findings. Smoking and craving function on different time scales—

while smoking behavior is a discrete, observable action that unfolds over a longer period (e.g., minutes or

hours), craving is a fluctuating internal state that can change moment to moment. The temporal resolution of

fMRI (approximately one second) allowed us to integrate momentary assessments of craving in the fMRI task

and measure its association with neural activity, providing insight into the immediate psychological and

neurobiological effects of GWLs. Beyond craving, smoking behavior is also shaped by a range of additional

factors, such as cigarette availability, environmental constraints (Pentz et al., 1989), nicotine metabolism

(Pianezza et al., 1998), stress (McKee et al., 2010), and social influence (East et al., 2021). These factors may

potentially add to or interact with the effects of craving on actual smoking during GWL exposure (Childs & de

Wit, 2010; Dimoff et al., 2019; Gass et al., 2014). Understanding these complex, interrelated factors will be

crucial for interpreting the findings of our previous (ref) and current studies and will refine future interventions

aimed at reducing smoking through GWLs.

Craving is a multi-dimensional phenomenon involving cognitive, affective, sensory, and physiological

components (Kober & DeLeone, 2011; Miele et al., 2023; Rosenberg, 2009). It is one of the eleven diagnostic

criteria for tobacco use disorder in DSM-5 (American Psychiatric Association, 2013). Despite some criticisms

over the accuracy of self-report measures of craving (Sayette et al., 2000), we found that average craving was

positively correlated with nicotine dependence and smoking severity, supporting the external validity of the

measure. Factors that contribute to cigarette craving include stress (Childs & de Wit, 2010), nicotine withdrawal

(Hughes & Hatsukami, 1986), exposure to smoking-related conditioned stimuli (i.e., cues) (Betts et al., 2021;

Shi, Wang, Fairchild, Aronowitz, Lynch, et al., 2023), and spontaneous smoking-related thoughts (e.g., intrusive

thoughts and imaginations of cigarette smell) (Aslan et al., 2023). Our previous cross-sectional study showed

greater reduction in smoking cue-induced craving after viewing high-ER GWLs compared to low-ER ones

(Wang, Romer, et al., 2015). In the current study, craving was measured without the presentation of smoking

cues and without the induction of stress or withdrawal. Therefore, our current data reflect diminishing utility of

high-ER GWLs in reducing basal craving, which, compared to cue-induced craving, was shown to be more

closely related to nicotine dependence and smoking severity (Dunbar et al., 2014). Nevertheless, it would be

interesting for future studies to examine whether our longitudinal findings can be generalized to GWL effects on

cue-, withdrawal-, and stress-induced craving.

Our study comes with a number or limitations. Firstly, the four-week exposure period, though substantial for

neuroimaging, is brief compared to potentially years of exposure to GWLs if implemented. Future studies

should extend this timeframe and incorporate real-world measures, such as ecological momentary assessment, to

better understand the dynamic and long-term impact of GWLs on craving and smoking outside fMRI. Secondly,

given our focus on emotional processing and the amygdala's well-established role in this domain (LeDoux,

2007), we selected the amygdala as our sole region of interest. Future studies should examine other brain

regions and their interregional connectivity, which may further account for GWLs' efficacy. Thirdly, our study

participants were exclusive combustible cigarette smokers who were not currently using other nicotine products.

However, the number of electronic cigarette users has been rapidly increasing (Vahratian et al., 2025), and the

dual use of electronic and combustible cigarettes has become highly prevalent (Coleman et al., 2022).

Examining high-ER and low-ER GWLs' efficacy in the dual user population would be important for a

comprehensive understanding of their public health impact.

The major argument against GWLs used by the tobacco companies in court has been that the highly aversive

images included in the GWLs "do not convey any warning information at all … [and] are unabashed attempts to

evoke emotion" ("R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al.," 2012),

that the GWLs "are designed to shock rather than inform", and that the "graphics go far beyond what would be

necessary to communicate a simple factual message" ("R.J. Reynolds Tobacco Company, et al. v. U.S. Food &

Drug Administration, et al.," 2024a). Our past studies provide both behavioral and neurobiological evidence

supporting the effectiveness of GWLs. Notably, we found that under prolonged, real-world exposure conditions,

low-ER GWLs are non-inferior to high-ER GWLs in reducing smoking motivation and behavior. Thus, our

findings do not support maximizing the ER in GWLs for the greatest public health benefit and suggest that any

future GWLs should be pre-tested for ER and their neurobehavioral effects prior to implementation. These

findings offer a potential pathway to balancing public health objectives with the constitutional concerns

surrounding GWL enforcement.

DECLARATION OF INTERESTS

The authors report no conflicts of interest in this work.

DATA AVAILABILITY

The data underlying this article can be shared on reasonable request to the corresponding author.

FUNDING

This work was supported by the NARSAD Young Investigator Grant from the Brain & Behavior Research

Foundation (#30780) (ZS) and the following National Institutes of Health grants: T32AG076411 (APRR),

R01DA036028 (DDL), R00AA026892 (CEW), and K01DA051709 (ZS). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the NIH or the FDA.

AUTHOR CONTRIBUTIONS

Astrid P. Ramos-Rolón: formal analysis (equal), investigation (equal), methodology (equal), software (equal), validation (equal), visualization (equal), original draft preparation (equal), review & editing (equal). **Daniel D. Langleben**: conceptualization (lead), data curation (lead), funding acquisition (lead), investigation (equal), methodology (supportive), project administration (lead), resources (lead), supervision (lead), review & editing (equal). **Kevin G. Lynch**: formal analysis (supportive), investigation (supportive), methodology (supportive), software (supportive), review & editing (equal). **Corinde E. Wiers**: conceptualization (supportive), investigation (supportive), methodology (supportive), resources (supportive), original draft preparation (supportive), review & editing (equal). **Zhenhao Shi**: conceptualization (supportive), data curation (supportive), formal analysis (equal), investigation (equal), methodology (equal), project administration (supportive), resources (supportive), software (equal), supervision (supportive), validation (equal), visualization (equal), original draft preparation (equal), review & editing (equal).

REFERENCES

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®).* American Psychiatric Association.

Aslan, M., Sala, M., Gueorguieva, R., & Garrison, K. A. (2023). A network analysis of cigarette craving. *Nicotine & Tobacco Research*, *25*(6), 1155-1163. https://doi.org/10.1093/ntr/ntad021

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1-48. https://doi.org/10.18637/jss.v067.i01

Betts, J. M., Dowd, A. N., Forney, M., Hetelekides, E., & Tiffany, S. T. (2021). A meta-analysis of cue

reactivity in tobacco cigarette smokers. *Nicotine & Tobacco Research*, *23*(2), 249-258.

https://doi.org/10.1093/ntr/ntaa147

Brehm, S. S., & Brehm, J. W. (2013). Psychological Reactance: A Theory of Freedom and Control.

Academic Press.

Breiter, H. C., Etcoff, N. L., Whalen, P. J., Kennedy, W. A., Rauch, S. L., Buckner, R. L., Strauss, M.

M., Hyman, S. E., & Rosen, B. R. (1996). Response and habituation of the human amygdala

during visual processing of facial expression. *Neuron*, *17*(5), 875-887.

https://doi.org/10.1016/S0896-6273(00)80219-6

Carter, B., Lam, C., Robinson, J., Paris, M., Waters, A., Wetter, D., & Cinciripini, P. (2008). Real-time

craving and mood assessments before and after smoking. *Nicotine & Tobacco Research*, *10*(7),

1165-1169. https://doi.org/10.1080/14622200802163084

Childs, E., & de Wit, H. (2010). Effects of acute psychosocial stress on cigarette craving and smoking.

Nicotine & Tobacco Research, *12*(4), 449-453. https://doi.org/10.1093/ntr/ntp214

Cho, Y. J., Thrasher, J. F., Swayampakala, K., Yong, H.-H., McKeever, R., Hammond, D., Anshari,

D., Cummings, K. M., & Borland, R. (2016). Does reactance against cigarette warning labels

matter? Warning label responses and downstream smoking cessation amongst adult smokers in

Australia, Canada, Mexico and the United States. *PLOS One*, *11*(7), e0159245.

https://doi.org/10.1371/journal.pone.0159245

Coleman, S. R. M., Piper, M. E., Byron, M. J., & Bold, K. W. (2022). Dual use of combustible

cigarettes and e-cigarettes: a narrative review of current evidence. *Current Addiction Reports*,

9(4), 353-362. https://doi.org/10.1007/s40429-022-00448-1

Dimoff, J. D., Sayette, M. A., & Levine, J. M. (2019). Experiencing cigarette craving with a friend: a

shared reality analysis. *Psychology of Addictive Behaviors*, *33*(8), 721-729.

https://doi.org/10.1037/adb0000519

Do, K. T., & Galván, A. (2014). FDA cigarette warning labels lower craving and elicit frontoinsular

activation in adolescent smokers. *Social Cognitive and Affective Neuroscience*, *10*(11), 1484-

1496. https://doi.org/10.1093/scan/nsv038

Dunbar, M. S., Shiffman, S., Kirchner, T. R., Tindle, H. A., & Scholl, S. M. (2014). Nicotine

dependence, "background" and cue-induced craving and smoking in the laboratory. *Drug and*

Alcohol Dependence, *142*, 197-203.

East, K., McNeill, A., Thrasher, J. F., & Hitchman, S. C. (2021). Social norms as a predictor of

smoking uptake among youth: a systematic review, meta-analysis and meta-regression of

prospective cohort studies. *Addiction*, *116*(11), 2953-2967.

Erceg-Hurn, D. M., & Steed, L. G. (2011). Does exposure to cigarette health warnings elicit

psychological reactance in smokers? *Journal of Applied Social Psychology*, *41*(1), 219-237.

Fidler, J. A., Shahab, L., & West, R. (2011). Strength of urges to smoke as a measure of severity of

cigarette dependence: comparison with the Fagerström Test for Nicotine Dependence and its

components. *Addiction*, *106*(3), 631-638. https://doi.org/10.1111/j.1360-0443.2010.03226.x

Fischer, H. Ê., Furmark, T., Wik, G., & Fredrikson, M. (2000). Brain representation of habituation to

repeated complex visual stimulation studied with PET. *NeuroReport*, *11*(1), 123-126.

Fridriksson, J. F., Rorden, C., Newman-Norlund, R. D., Froeliger, B., & Thrasher, J. F. (2018).

Smokers' neurological responses to novel and repeated health warning labels (HWLs) from

cigarette packages. *Frontiers in Psychiatry*, *9*, 319.

Friston, K. J., Ashburner, J., Frith, C. D., Poline, J. B., Heather, J. D., & Frackowiak, R. S. J. (1995).

Spatial registration and normalization of images. *Human Brain Mapping*, *3*(3), 165-189.

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S. J., & Turner, R. (1996). Movement-related

effects in fMRI time-series. *Magnetic Resonance in Medicine*, *35*(3), 346-355.

Gass, J. C., Motschman, C. A., & Tiffany, S. T. (2014). The relationship between craving and tobacco

use behavior in laboratory studies: a meta-analysis. *Psychology of Addictive Behaviors*, *28*(4),

1162-1176. https://doi.org/10.1037/a0036879

Hall, M. G., Sheeran, P., Noar, S. M., Boynton, M. H., Ribisl, K. M., Parada Jr, H., Johnson, T. O., &

Brewer, N. T. (2018). Negative affect, message reactance and perceived risk: how do pictorial

cigarette pack warnings change quit intentions? *Tobacco Control*, *27*(e2), e136-e142.

https://doi.org/10.1136/tobaccocontrol-2017-053972

Hall, M. G., Sheeran, P., Noar, S. M., Ribisl, K. M., Boynton, M. H., & Brewer, N. T. (2017). A brief

measure of reactance to health warnings. *Journal of Behavioral Medicine*, *40*(3), 520-529.

https://doi.org/10.1007/s10865-016-9821-z

Hayes, A. F. (2015). An index and test of linear moderated mediation. *Multivariate Behavioral

Research*, *50*(1), 1-22. https://doi.org/10.1080/00273171.2014.962683

Heatherton, T. F., Kozlowski, L. T., Frecker, R. C., & Fagerstrom, K. O. (1991). The Fagerstrom Test

for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *British Journal

of Addiction*, *86*(9), 1119-1127.

Hiilamo, H., Crosbie, E., & Glantz, S. A. (2014). The evolution of health warning labels on cigarette

packs: the role of precedents, and tobacco industry strategies to block diffusion. *Tobacco Control*, *23*(1), e2. https://doi.org/10.1136/tobaccocontrol-2012-050541

Hughes, J. R., & Hatsukami, D. (1986). Signs and symptoms of tobacco withdrawal. *Archives of General Psychiatry*, *43*(3), 289-294. https://doi.org/10.1001/archpsyc.1986.01800030107013

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage*, *17*(2), 825-841.

Keating, D. M., & Skurka, C. (in press). Meta-analytic evidence that message fatigue is associated with unintended persuasive outcomes. *Communication Research*. https://doi.org/10.1177/00936502241287875

Kober, H., & DeLeone, C. M. (2011). Smoking and neuroimaging: a review. *Current Cardiovascular Risk Reports*, *5*(6), 484-491. https://doi.org/10.1007/s12170-011-0201-5

LaVoie, N. R., Quick, B. L., Riles, J. M., & Lambert, N. J. (2015). Are graphic cigarette warning labels an effective message strategy? A test of psychological reactance theory and source appraisal. *Communication Research*, *44*(3), 416-436. https://doi.org/10.1177/0093650215609669

LeDoux, J. (2007). The amygdala. *Current Biology*, *17*(20), R868-R874.

Luke, S. G. (2017). Evaluating significance in linear mixed-effects models in R. *Behavior Research Methods*, *49*(4), 1494-1502. https://doi.org/10.3758/s13428-016-0809-y

Maynard, O. M., Attwood, A., O'Brien, L., Brooks, S., Hedge, C., Leonards, U., & Munafò, M. R. (2014). Avoidance of cigarette pack health warnings among regular cigarette smokers. *Drug and Alcohol Dependence*, *136*, 170-174.

McKee, S. A., Sinha, R., Weinberger, A. H., Sofuoglu, M., Harrison, E. L. R., Lavery, M., & Wanzer,

J. (2010). Stress decreases the ability to resist smoking and potentiates smoking intensity and reward. *Journal of Psychopharmacology*, *25*(4), 490-502. https://doi.org/10.1177/0269881110376694

Miele, C., Cabé, J., Cabé, N., Bertsch, I., Brousse, G., Pereira, B., Moulin, V., & Barrault, S. (2023). Measuring craving: a systematic review and mapping of assessment instruments. What about sexual craving? *Addiction*, *118*(12), 2277-2314. https://doi.org/10.1111/ADD.16287

Motschman, C. A., Germeroth, L. J., & Tiffany, S. T. (2018). Momentary changes in craving predict smoking lapse behavior: a laboratory study. *Psychopharmacology*, *235*(7), 2001-2012. https://doi.org/10.1007/s00213-018-4898-4

Mutschler, I., Wieckhorst, B., Speck, O., Schulze-Bonhage, A., Hennig, J., Seifritz, E., & Ball, T. (2010). Time scales of auditory habituation in the amygdala and cerebral cortex. *Cerebral Cortex*, *20*(11), 2531-2539. https://doi.org/10.1093/cercor/bhq001

Nonnemaker, J., Farrelly, M., Kamyab, K., Busey, A., & Mann, N. (2010). Experimental study of graphic cigarette warning labels (Contract No. HHSF-223-2009-10135G, RTI Project No. 0212305.007.003).

Pentz, M. A., Brannon, B. R., Charlin, V. L., Barrett, E. J., MacKinnon, D. P., & Flay, B. R. (1989). The power of policy: the relationship of smoking policy to adolescent smoking. *American Journal of Public Health*, *79*(7), 857-862. https://doi.org/10.2105/AJPH.79.7.857

Pianezza, M. L., Sellers, E. M., & Tyndale, R. F. (1998). Nicotine metabolism defect reduces smoking. *Nature*, *393*(6687), 750-750. https://doi.org/10.1038/31623

Plichta, M. M., Grimm, O., Morgen, K., Mier, D., Sauer, C., Haddad, L., Tost, H., Esslinger, C.,

Kirsch, P., & Schwarz, A. J. (2014). Amygdala habituation: a reliable fMRI phenotype.

NeuroImage, *103*, 383-390.

Poellinger, A., Thomas, R., Lio, P., Lee, A., Makris, N., Rosen, B. R., & Kwong, K. K. (2001).

Activation and habituation in olfaction—an fMRI study. *NeuroImage*, *13*(4), 547-560.

R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al., 696 F.3d 1205,

Nos. 11-5332, 12-5063 (U.S. Court of Appeals for the DC Circuit 2012).

R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al., No. 6:20-cv-00176

(U.S. District Court for the Eastern District of Texas 2022).

R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al., No. 24-189

(Supreme Court of the United States 2024a).

R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al., 94 F.4th 863, No.

23-40076 (U.S. Court of Appeals for the Fifth Circuit 2024b).

R.J. Reynolds Tobacco Company, et al. v. U.S. Food & Drug Administration, et al., No. 6:20-cv-00176

(U.S. District Court for the Eastern District of Texas 2025).

Rankin, C. H., Abrams, T., Barry, R. J., Bhatnagar, S., Clayton, D. F., Colombo, J., Coppola, G.,

Geyer, M. A., Glanzman, D. L., Marsland, S., McSweeney, F. K., Wilson, D. A., Wu, C. F., &

Thompson, R. F. (2009). Habituation revisited: an updated and revised description of the

behavioral characteristics of habituation. *Neurobiology of Learning and Memory*, *92*(2), 135-138.

Reiser, M., Yao, L., Wang, X., Wilcox, J., & Gray, S. (2017). A Comparison of bootstrap confidence

intervals for multi-level longitudinal data using Monte-Carlo simulation. In D.-G. Chen & J. D.

Chen (Eds.), *Monte-Carlo simulation-based statistical modeling* (pp. 367-403). Springer Nature.

Robinson, T. E., & Berridge, K. C. (1993). The neural basis of drug craving: an incentive-sensitization

theory of addiction. *Brain Research Reviews*, *18*(3), 247-291. https://doi.org/10.1016/0165-

0173(93)90013-P

Rosenberg, H. (2009). Clinical and laboratory assessment of the subjective experience of drug craving.

Clinical Psychology Review, *29*(6), 519-534.

Sayette, M. A., Shiffman, S., Tiffany, S. T., Niaura, R. S., Martin, C. S., & Shadel, W. G. (2000). The

measurement of drug craving. *Addiction*, *95*(Suppl 2), S189-S210.

Shi, Z., Wang, A.-L., Fairchild, V. P., Aronowitz, C. A., Lynch, K. G., Loughead, J., & Langleben, D.

D. (2023). Addicted to green: priming effect of menthol cigarette packaging on brain response to

smoking cues. *Tobacco Control*, *32*(e1), e45-e52. https://doi.org/10.1136/tobaccocontrol-2021-

056639

Shi, Z., Wang, A.-L., Fairchild, V. P., Aronowitz, C. A., Padley, J. H., Lynch, K. G., Loughead, J., &

Langleben, D. D. (2023). Effects of emotional arousal on the neural impact and behavioral

efficacy of cigarette graphic warning labels. *Addiction*, *118*(5), 914-924.

Shi, Z., Wang, A.-L., Liu, J., Audrain-McGovern, J., Lynch, K. G., Loughead, J., & Langleben, D. D.

(2024). Delayed effects of cigarette graphic warning labels on smoking behavior. *medRxiv*,

2024.2001.2006.24300835. https://doi.org/10.1101/2024.01.06.24300835

Shiffman, S., Gwaltney, C. J., Balabanis, M. H., Liu, K. S., Paty, J. A., Kassel, J. D., Hickcox, M., &

Gnys, M. (2002). Immediate antecedents of cigarette smoking: an analysis from ecological

momentary assessment. *Journal of Abnormal Psychology*, *111*(4), 531-545.

https://doi.org/10.1037/0021-843X.111.4.531

Shiffman, S., West, R. J., & Gilbert, D. G. (2004). Recommendation for the assessment of tobacco

craving and withdrawal in smoking cessation trials. *Nicotine & Tobacco Research*, *6*(4), 599-614.

https://doi.org/10.1080/14622200410001734067

So, J., Kim, S., & Cohen, H. (2017). Message fatigue: conceptual definition, operationalization, and

correlates. *Communication Monographs*, *84*(1), 5-29.

https://doi.org/10.1080/03637751.2016.1250429

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer,

B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a

macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, *15*(1),

273-289.

U.S. Food and Drug Administration. (2011). Required warnings for cigarette packages and

advertisements. Final rule. *Federal Register*, *76*(120), 36628-36777.

U.S. Food and Drug Administration. (2022). Tobacco products; required warnings for cigarette

packages and advertisements; delayed effective date. *Federal Register*, *87*(40), 11295.

Vahratian, A., Briones, E. M., Jamal, A., & Marynak, K. L. (2025). Electronic cigarette use among

adults in the United States, 2019–2023 (NCHS Data Brief, volume 524).

Wang, A.-L., Lowen, S. B., Romer, D., Giorno, M., & Langleben, D. D. (2015). Emotional reaction

facilitates the brain and behavioural impact of graphic cigarette warning labels in smokers.

Tobacco Control, *24*(3), 225-232. https://doi.org/10.1136/tobaccocontrol-2014-051993

Wang, A.-L., Romer, D., Elman, I., Turetsky, B. I., Gur, R. C., & Langleben, D. D. (2015). Emotional

graphic cigarette warning labels reduce the electrophysiological brain response to smoking cues.

Addiction Biology, *20*(2), 368-376. https://doi.org/10.1111/adb.12117

Wang, A.-L., Shi, Z., Fairchild, V. P., Aronowitz, C. A., & Langleben, D. D. (2019). Emotional salience of the image component facilitates recall of the text of cigarette warning labels. *European Journal of Public Health*, *29*(1), 153-158.

World Health Organization. (2024). WHO global report on trends in prevalence of tobacco use 2000–2030.

Wright, C. I., Fischer, H., Whalen, P. J., McInerney, S. C., Shin, L. M., & Rauch, S. L. (2001). Differential prefrontal cortex and amygdala habituation to repeatedly presented emotional stimuli. *NeuroReport*, *12*(2), 379-383.

Zanto, T. P., Rubens, M. T., Thangavel, A., & Gazzaley, A. (2011). Causal role of the prefrontal cortex in top-down modulation of visual processing and working memory. *Nature Neuroscience*, *14*(5), 656-661. https://doi.org/10.1038/nn.2773

Table 1. Participant characteristics at baseline (count or mean±SD).

Variable	High-ER (n=79)	Low-ER (n=79)	p-value[a]
Age	29.4±9.6	27.8±8.0	0.42
Sex	37 female, 42 male	38 female, 41 male	>0.99
Race	44 White, 20 AA, 6 Asian, 9 other	44 White, 18 AA, 6 Asian, 11 other	0.96
Ethnicity	11 Hispanic	5 Hispanic	0.19
Age of smoking initiation (years)	16.5±3.2	17.6±4.4	0.099
Nicotine dependence (FTND)[b]	4.6±2.6	4.3±2.4	0.39
Number of cigarettes smoked per day[b]	13.5±8.6	12.0±6.6	0.44

Abbreviations: ER, emotional response; FTND, Fagerström Test for Nicotine Dependence; AA, African American.

[a] P-values were obtained from group comparisons using χ^2 tests for categorical variables and Mann-Whitney U-tests for numerical variables.

[b] Baseline FTND and number of cigarettes was missing in two and one participant, respectively.

Cigarettes only	Cigarettes + High ER GWLs	Cigarettes + High ER GWLs	Cigarettes + High ER GWLs	Cigarettes + High ER GWLs
Cigarettes only	Cigarettes + Low ER GWLs	Cigarettes + Low ER GWLs	Cigarettes + Low ER GWLs	Cigarettes + Low ER GWLs

day -7   day 0 (MRI)   day 7   day 14   day 21   day 28 (MRI)

2 **Figure 1.** Study timeline schematic

5 **Figure 2.** Difference in cigarette craving reduction between GWL blocks and control image blocks across time

6 (day 0 vs. day 28) and group (high-ER vs. low-ER). Abbreviations: GWL, graphic warning label; ER, emotional

7 response.

2 **Figure 3.** Amygdala activity in response to GWLs vs. control images across time (day 0 vs. day 28) and group

3 (high-ER vs. low-ER). Abbreviations: GWL, graphic warning label; ER, emotional response.

6 **Figure 4.** Significant association between change in cigarette craving and amygdala neural activity.

7 Abbreviation: ER, emotional response.

2   **Figure 5.** Moderated mediation model presented in the conceptual form (A) and the statistical form (B) (adapted

3   from Hayes (2015)).

6   **Figure 6.** Conditional indirect effect of stimulus (GWLs vs. control images) on Δcraving through amygdala

7   neural activity across time (day 0 vs. day 28) and group (high-ER vs. low-ER). Abbreviations: GWL, graphic

8   warning label; ER, emotional response.