How should we study the indirect effects of antimicrobial treatment strategies? A causal perspective.

Juan Gago[a,1], Christopher Boyer[a,b,c,1], Marc Lipsitch[a]

[a]*Department of Epidemiology, Harvard T.H. Chan School of Public Health, 677 Huntington Avenue, Boston, 02115, MA, USA*
[b]*Department of Medicine, Case Western Reserve School of Medicine, 9501 Euclid Avenue, Cleveland, 44106, OH, USA*
[c]*Department of Quantitative Health Sciences, Cleveland Clinic, 9211 Euclid Avenue, Cleveland, 44195, OH, USA*

Abstract

Effective antimicrobial stewardship requires unbiased assessment of the benefits and harms of different treatment strategies, considering both immediate patient outcomes and patterns of antimicrobial resistance. In principle, these benefits and harms can be expressed as causal contrasts between treatment strategies and therefore should be ideally suited for study under the potential outcomes framework. However, causal inference in this setting is complicated by interference between individuals (or units) due to the indirect effects of antibiotic treatment, including the spread of resistant bacteria to others. These indirect effects complicate the assessment of trade-offs as benefits are mostly due to the direct effects among those treated, while harms are more diffuse and therefore harder to measure. While causal frameworks and study designs that accommodate interference have previously been proposed, they have been applied predominately to the study of vaccines, which differ from antimicrobial interventions in fundamental ways. In this article, we review these existing approaches and propose alternative adaptations tailored to the study of antimicrobial treatment strategies.

Keywords: causal inference, interference, drug resistance, antibiotics

---

[1]These authors contributed equally to this work.

March 28, 2025

1. Introduction

Antimicrobial resistance (AMR) is a complex and growing threat to global health. It increases the risk of serious illness and mortality for patients and complicates clinicians' efforts to choose effective antimicrobial treatments. At the population level, AMR threatens the global efficacy of antimicrobials, the therapeutic cornerstone of modern infectious disease management, and jeopardizes procedures that rely on antimicrobial prophylaxis. Furthermore, AMR disproportionately affects vulnerable populations, such as immunosuppressed patients [1], magnifying the clinical and public health burden. While efforts are underway to better understand the biological basis for the emergence and spread of AMR [2], effective interventions are also necessary to limit resistance while continuing to maximize the benefits of antimicrobial therapy [3].

To evaluate the harms and benefits of interventions to curb AMR, we need a framework for formalizing the causal effects of antimicrobial treatments on microbiological and clinical outcomes and then study designs that can be used to empirically estimate these effects. Modern causal inference methods, based on the potential outcomes framework, have become essential tools for formalizing treatment effects. Study designs for eliciting them have become widespread in the health sciences. However, their application to AMR, where treatment and broader transmission dynamics are closely intertwined, has been limited.

A major methodological challenge to developing a causal framework for AMR, common across many infectious diseases, is the presence of interference, where the treatment of one individual can affect the outcomes of others, violating standard assumptions [4]. Indirect effects are particularly important in AMR. Ignoring them can lead to an incomplete assessment of an intervention's benefits and harms by capturing only individual-level outcomes and overlooking how treatment policies shape infection incidence and resistance patterns throughout the population—whether or not individuals are in the study. Indeed, many measures that affect resistance are partly or wholly justified by their impact on others, such as isolating carriers of resistant organisms [5], treatment as prevention [6], and antimicrobial cycling [7].

While these interventions have been tested and studied using transmission-dynamic models, a formal causal framework for them does not yet exist.

Over the last three decades, methods for causal inference on infectious disease interventions that allow for interference have been proposed [8–11]. However, they have primarily focused on vaccination, which differs in important ways from AMR interventions. Most notably, in a vaccine trial the intervention is typically assigned prior to an individual's exposure to infection as a form of inoculation against future infection or adverse sequelae. By contrast, many antimicrobial interventions are applied after exposure or infection has already occurred and only among those seeking treatment. In addition, the course of treatment may be complex as individuals may first be treated "empirically", i.e. without knowledge of the pathogenic source, and may be dynamically updated based on the results of microbial culture or other clinical parameters. As a result, there is no widely adopted causal framework for evaluating indirect and overall effects of antimicrobial treatments or other interventions to limit AMR.

In this paper, we aim to address this gap by extending causal inference frameworks that account for interference to the context of antimicrobial treatments and AMR. Specifically, we aim to establish a causal framework to study antimicrobial treatment effects, both direct and indirect, considering that the effects of interest can be different measures of clinical success (e.g., microbiological clearance) as well as adverse outcomes, such as treatment failure and development of AMR infections. We propose modifications to study designs —such as the two-stage randomization used in vaccine studies— to capture the indirect effects of antibiotics. Additionally, we highlight the benefits of estimating well-defined per-exposure effects to obtain accurate estimates of direct effects. Using directed acyclic graphs (DAGs) and single-world intervention graphs (SWIGs), we analyze both direct and indirect effects of antimicrobial treatment strategies and discuss specific challenges that researchers might encounter. The insights from this framework are valuable for designing and evaluating interventions and policies aimed at controlling the spread of AMR and optimizing antibiotic

use.

2. Motivating example: empiric treatment of bloodstream infections

Vancomycin is commonly used as an empiric treatment (i.e. given before identification of the pathogen) for bloodstream infections to cover *Staphylococcus aureus* species, including methicillin-resistant *S. aureus* (MRSA). However, its use can inadvertently promote the emergence of vancomycin-resistant Enterococcus (VRE) by suppressing susceptible enterococci and altering gut microbiota [12]. In contrast, daptomycin, a last-resort antibiotic effective against both MRSA and VRE, is typically reserved for resistant infections to preserve its utility [13]. There would be trade-offs in using daptomycin versus vancomycin as empiric therapy, particularly in settings with a high prevalence of MRSA and VRE.

Let's consider a hypothetical study to compare these two antimicrobial treatment strategies in an ICU, vancomycin versus daptomycin, for the empiric treatment of suspected bloodstream infection (BSI). Traditionally, this would be evaluated in a randomized trial, often set up as a non-inferiority or superiority trial, in which patients meeting the criteria for suspected BSI would be randomly assigned to receive one of the two empiric treatments. Primary and secondary outcomes for assessing the relative merits of the two treatments might include microbiological clearance of infection, clinical resolution of the event, patient mortality and the development of antimicrobial resistance, specifically an infection that yields a subsequent positive culture for vancomycin-resistant *Enterococci* (VRE). Another possible outcome would be incident colonization by VRE, which could be measured if there are samples for baseline values.

Typically, these trials only measure results in patients who were directly enrolled and treated. However, the indirect effects of empiric treatment on the contacts of the index treated patient are important mechanisms for the spread of resistance. For instance, administering vancomycin to one patient may increase the growth of VRE in the patient's bowel by killing off some of its competitors, including vancomycin-susceptible *Enterococci*. The patient

may also start to shed VRE facilitating its transmission to others, increasing the circulation of VRE, as well as the likelihood that the contacts get exposed and develop a VRE infection [14]. This may be particularly salient in settings where nosocomial infection is common such as hospitals or nursing homes. This scenario is an example of interference: the treatment of an index patient affects the outcomes of others. Specifically, the use of vancomycin in the index patient might influence the response of other patients to the same treatment because treating the index patient makes those other patients more likely to be infected with resistant (rather than susceptible) *Enterococci*. Note that if the trial entry criterion is suspected bloodstream infection, these other indivduals might become study participants or might not (for example if the VRE transmitted to them by a treated participant in the study causes a urinary tract infection rather than a bloodstream infection). Whether counted as a participant or not, the fact that they were infected by VRE, resulting from treatment of another participant, would not be accounted for in trial outcomes.

Quantifying the indirect effects of empiric vancomycin versus daptomycin on subsequent vancomycin-resistance is important to capture the full range of risks and benefits of the two antimicrobial treatment strategies. However, these effects cannot be easily studied in a conventional, individually-randomized trial. They also pose similar challenges for real-world or observational studies that seek to emulate them.

3. Defining direct and indirect causal effects under interference

To begin fixing concepts, consider briefly the simple scenario in which there are just two possible individuals interacting in a hospital setting (sequential patients admitted to an isolated ICU room, for example) indexed by i and j. At admission, we define variables representing each individual's true infection status (I_{i0}, I_{j0}) as well as an indicator of whether infection is suspected (S_{i0}, S_{j0}) by their care team –e.g., because they are symptomatic. A suspected infection can trigger an initial blood culture (Y_{i0}, Y_{j0}) which, for simplicity, we assume is measured without error. The possible values of both infection status and

culture result are no infection/negative culture, susceptible infection/culture, and resistant infection/culture. Thus, for example, Y_{it} for individual i at time t is defined as

$$Y_{it} = \begin{cases} 2 & \text{Resistant culture} \\ 1 & \text{Susceptible culture} \\ 0 & \text{Negative culture} \end{cases}$$

Let (A_{i0}, A_{j0}) denote the empiric antimicrobial treatment of person i and j prior to pathogen identification and (A_{i1}, A_{j1}) denote the concordant treatment decision at time 1 after the results of the culture are obtained. Following our example, assume that there are two possible antibiotic treatments, one class for which resistant species of bacteria exist (vancomycin) and another where resistance is uncommon (daptomycin). Therefore, define for indiviudal i at time t

$$A_{it} = \begin{cases} 2 & \text{Antimicrobial covering resistant pathogens (daptomycin),} \\ 1 & \text{Antimicrobial covering susceptible but not resistant pathogens (vancomycin),} \\ 0 & \text{No treatment.} \end{cases}$$

At a future follow up —say 14 days after admission— we define final infection status (I_{i14}, I_{j14}) and culture result (Y_{i14}, Y_{j14}). In general, we assume that treatment status and culture results are fully observed while true exposure and infection status must be inferred. We also define an intermediate microbiome result (M_{i1}, M_{j1}) to reflect the possibility of carriage (but not infection). The assumed causal relationship between these variables is captured in the causal directed acyclic graph (DAG) of Figure 1.

As in our motivating example, consider the causal effect of empiric treatment. The potential outcomes framework defines causal effects in terms of contrasts of hypothetical interventions, which may be contrary to fact. Let $Y_{i14}^{a_i, a_j}$ denote the potential culture status of person i if the empiric treatment status of person i and person j were set to (a_i, a_j). Thus

for example $Y_{i14}^{1,2}$ is the potential outcome of person i at the 14-day follow up if person i receives empiric vancomycin while person j receives empiric daptomycin; and $Y_{j14}^{0,0}$ denotes the potential outcome of person j if both person i and person j are untreated. In general, each individual has 9 potential outcomes: $Y_{j14}^{2,2}$, $Y_{j14}^{2,1}$, $Y_{j14}^{2,0}$, $Y_{j14}^{1,2}$, $Y_{j14}^{1,1}$, $Y_{j14}^{1,0}$, $Y_{j14}^{0,2}$, $Y_{j14}^{0,1}$, and $Y_{j14}^{0,0}$. However, we focus on the case that person i is admitted prior to person j, in which case transmission can only occur from person i to person j. Therefore, person i is unaffected by the treatment of person j implying $Y_{i14}^{2,a_j} = Y_{i14}^{1,a_j} = Y_{i14}^{0,a_j}$ for all a_j and thus we can simplify to only 3 potential outcomes for person i: $Y_{i14}^{a_i=2}$, $Y_{i14}^{a_i=1}$, $Y_{i14}^{a_i=0}$. Similarly, because initial culture result for person j is unaffected by their empiric treatment, assuming initial cultures are always taken prior to initiation of treatment, we have $Y_{j0}^{a_i,0} = Y_{j0}^{a_i,1} = Y_{j0}^{a_i,2}$ for all a_i and we can again simplify to $Y_{j0}^{a_i=2}$, $Y_{j0}^{a_i=1}$, $Y_{j0}^{a_i=0}$.

Notice, under this notation, the potential outcomes of person j depend on the treatment status person i. This is referred to as *interference* and is contrary to most introductory literature in causal inference where it is assumed that a person's potential outcome may be defined independently of the treatment status of others. However, this assumption is implausible for an infectious disease because pathogens are generally transmitted from one individual to another. Returning to our motivating example, comparing effects of vancomycin vs. daptomycin, consider the following plausible scenario: when person i is treated with vancomycin it can lead to an overgrowth of VRE which they then transmit to person j. When person i is instead treated with daptomycin all *enterococci* are eliminated and therefore person j has no infection. This implies that $Y_{j0}^{1,a_j} \neq Y_{j0}^{2,a_j}$ and therefore the no interference assumption is invalid.

We can now formalize the notion of direct and indirect effects of antimicrobial treatments by focusing, in turn, on the outcomes of person i and person j. To introduce the notion of expectation, we expand consideration to a super population of K pairs of admitted patients where interaction is allowed within but not between pairs.

The direct effects of treatment for person i may be defined as contrasts between $Y_{i14}^{a_i=2}$

and $Y_{i14}^{a_i=1}$, such as

$$\Pr[Y_{i14}^{a_i=2} = 2] - \Pr[Y_{i14}^{a_i=1} = 2],$$

representing resistant infection 14 days after admission when treated empirically with dapto-mycin versus vancomycin. When combined with their culture result at admission this creates three strata in which we can compare the direct effects of vancomycin vs. daptomycin on person i with some of the most sensible contrasts being:

$$\Pr[Y_{i14}^{a_i=2} = 0 | Y_{i0} = 0] - \Pr[Y_{i14}^{a_i=1} = 0 | Y_{i0} = 0] \qquad \text{No infection}$$

$$\Pr[Y_{i14}^{a_i=2} = 0 | Y_{i0} = 1] - \Pr[Y_{i14}^{a_i=1} = 0 | Y_{i0} = 1] \qquad \text{S cleared}$$

$$\Pr[Y_{i14}^{a_i=2} = 0 | Y_{i0} = 2] - \Pr[Y_{i14}^{a_i=1} = 0 | Y_{i0} = 2] \qquad \text{R cleared}$$

$$\Pr[Y_{i14}^{a_i=2} = 1 | Y_{i0} = 1] - \Pr[Y_{i14}^{a_i=1} = 1 | Y_{i0} = 1] \qquad \text{S persistence}$$

$$\Pr[Y_{i14}^{a_i=2} = 2 | Y_{i0} = 2] - \Pr[Y_{i14}^{a_i=1} = 2 | Y_{i0} = 2] \qquad \text{R persistence}$$

$$\Pr[Y_{i14}^{a_i=2} = 2 | Y_{i0} = 1] - \Pr[Y_{i14}^{a_i=1} = 2 | Y_{i0} = 1] \qquad \text{R emergence}$$

When the presenting infection type is not controlled and the effects of antibiotic choice are heterogeneous across indiviudals its possible that, for instance, the stratum-specific effects above may vary depending on the types of individuals who present with susceptible or resistant infections (discussed further in the next section).

The indirect effects of treatment of person i on the outcome of person j may be defined as contrast between Y_{jt}^{2,a_j} and Y_{jt}^{1,a_j}, such as

$$\Pr[Y_{j0}^{a_i=2} = 2] - \Pr[Y_{j0}^{a_i=1} = 2]$$

and

$$\Pr[Y_{j14}^{a_i=2,a_j=1} = 2] - \Pr[Y_{j14}^{a_i=1,a_j=1} = 2],$$

representing resistant infection in person j at baseline and follow up when person i is treated

empirically with daptomycin versus vancomycin and acknowledging that, depending on the mechanism, the treatment of person i may affect either the initial (Y_{j0}) or follow up culture result (Y_{j14}) or both.

Both direct and indirect effects are represented graphically via the DAGs in Figure 1. Building on the work of Ogburn and VanderWeele [15], arrows connecting nodes of the same individual (e.g., from I_{i0} to A_{i0}, or from A_{i0} to M_i) represent *direct effects*. Arrows *between* individuals (e.g., from I_{i0} or M_i to E_{j0}) capture *indirect effects*—the essence of interference—where one person's infection or pathogen shredding can expose another person and cause a new infection. Figure 1 represents several possible interference pathways. For example, one route may be that subject i's infection persists or relapses (node I_{i14}), thereby exposing subject j. Another possibility is that A_{i0} or A_{i1} modifies M_i, creating the conditions to allow overgrowth and transmission to j of a pathogen. In reality, multiple pathways can operate simultaneously, as indicated by the cross-subject arrows from i to j. Furthermore, we can extend our graphs to include potential outcomes through single world intervention graphs (SWIGs) as shown in Figure 2. SWIGs represent hypothetical interventions on nodes through a node-splitting transformation [16]. All nodes that are descendants of the split node (intervention node) are then potential outcomes under the specified intervention.

4. How should we study causal effects of antimicrobial treatments under interference?

Broadly speaking, there are two approaches to studying causal effects under interference in the infectious disease literature. One approach is to directly manipulate the mechanisms through which interference operates (i.e. transmission), for instance by isolating participants and exposing them to the pathogen of interest in a controlled environment to experimentally separate direct and indirect effects. The second approach is to identify clusters or groups of interacting individuals with sufficient "buffer" between them such that interactions, and therefore interference, occurs within but not across groups and then use two stages of ran-

Figure 1: Causal DAGs showing possible interference mechanisms for antimicrobial treatment among two individuals indexed by i and j. Each subject i (top) and j (bottom) has cultures for at least two time points: 0 and 14. The node E_{i0} represents an *exposure* to a pathogen at baseline, which can lead to an *infection* I_{i0} suspected by the presence of *symptoms* denoted by S_{i0}. This prompts an *empiric treatment* A_{i0}, and the initiation of microbiological laboratory tests, which provide an *initial culture result* Y_{i0}. Guided by Y_{i0}, the clinician may adjust the treatment, yielding A_{i1}. Both A_{i0} and A_{i1} can alter the *microbiome* M_{i1}, which, in turn, can influence the final culture result Y_{i14} at follow-up. Notably, there is *interference* across subjects, as one subject's infection or microbiome changes can affect another's exposure. Specifically, I_{i0}, I_{i14} or M_i may alter E_{j0} if subject i transmits the pathogen to subject j. These dashed cross-subject arrows capture real-world interference pathways in which clinical interventions for one person may have downstream effects on the exposures and infection risks of others.

Figure 2: Single world intervention graph (SWIG) showing hypothetical interventions on empiric treatment of individuals i and j. The influence of the treatment of i on j is shown explicitly as all j nodes are now adorned by a_i indicating the potential outcome of person j when person i is treated with empiric treatment a_i.

domization to assign (1) groups to different treatment allocation schemes and (2) individuals within groups to treatments based on their group's allocation scheme. Direct effects are then evaluated by comparing individuals within clusters assigned to same group allocation scheme while indirect effects are evaluated by comparing individuals across clusters with different allocation schemes. We adapt both to the setting of antimicrobial resistance below. In both cases, we start by describing the ideal study design (i.e. the target trial) and then discuss its emulation via observational data when, as often is the case, the ideal design is infeasible.

5. Approach 1: Isolating direct and indirect effects in individuals via controlled exposure to pathogen

The first approach to interference attempts to isolate direct and indirect effects through controlled exposure. To study the direct effects of treatment an individual is placed in isolation and experimentally exposed to both the pathogen of interest and the treatment. This experimental setup is commonly referred to as a "challenge trial". Effectively it measures the joint effect of isolation (e.g., individual sterile rooms, PPE etc.), controlled exposure to the pathogen (i.e. a "challenge"), and antimicrobial treatment. It removes the possibility of interference because isolation prevents all indirect effects via transmission from external sources, provided that the isolation is sufficiently secure. While the design has been conducted in the past under certain restricted scenarios [17], it is often infeasible due to ethical and logistical considerations.

We can formalize antimicrobial effects under a challenge trial by defining exposure E_{i0} for individual i as:

$$
E_{i0} = \begin{cases} 2 & \text{Exposure to resistant pathogen} \\ 1 & \text{Exposure to susceptible pathogen} \\ 0 & \text{No Exposure} \end{cases}
$$

where, in a real challenge trial, the method and dose of each inoculum would be specified. For

example, in a human challenge trial of a pneumoccocal vaccine participants were inoculated with 80,000 CFU/100 μl of *Streptococcus pneumoniae* (6B) per naris. Traditionally, subjects are individually isolated in which case the design targets the so-called "per-exposure effect" [18], i.e. the direct effect of exposure and treatment on the same individual. Because all interaction with others has been eliminated, potential outcomes may be simplified to only depend on an individual's intervention values (Note: this can be seen in the SWIG in Figure 2 by imagining an intervention on E_{j0} which splits the node and blocks the indirect effects from person i to person j). We denote by $Y_{it}^{e_i, a_i}$ the potential outcome for individual i at time t of setting $E_{i0} = e_i$ and $A_{i0} = a_i$.

5.1. Direct effects

An advantage of the randomized challenge trial is that we can estimate the direct effects of the various antimicrobial treatments under different exposures and therefore estimate the true clearance rates of susceptible and resistant strains. These could then be used to quantify direct benefit and how it varies based on local exposure environment. By contrast in an uncontrolled setting, we may not know the true exposure history of those treated with different antibiotic classes (e.g. if we have no baseline culture result) or we may be concerned that there may be unmeasured confounding factors that determine both exposure and heterogeneity in responses to a particular treatment that make the effects among subgroups based on baseline culture non-causal.

We can represent hypothetical challenge interventions of interest (here there are six) via a SWIG where we intervene on exposure (Figure 3). Contrasts between potential outcomes under these interventions could then be used to quantify the direct harms or benefits for the treated individual of different treatment strategies. For example, we could contrast the probability of eradication at day 14 ($Y_{i14} = 0$) when using a treatment option including antimicrobials covering resistant strains ($a_i = 2$) compared to using antimicrobials covering

only susceptible strains ($a_i = 1$) given an exposure to resistant pathogen ($e_i = 2$),

$$\Pr[Y_{i14}^{e_i=2,a_i=2} = 0] - \Pr[Y_{i14}^{e_i=2,a_i=1} = 0],$$

or a susceptible pathogen ($e_i = 1$)

$$\Pr[Y_{i14}^{e_i=1,a_i=2} = 0] - \Pr[Y_{i14}^{e_i=1,a_i=1} = 0].$$

Controlled exposure allows us to study the possible heterogeneity in response to antibiotic treatment based on many factors, including a unique set of comorbidities or other patient characteristics (e.g., age, pregnancy, etc.). Few study designs directly confirm that results from *in vitro* susceptibility tests translate effectively *in vivo*. Examining the per-exposure effect in a challenge trial like this would be the closest equivalent to estimating, in humans, the direct *in vivo* effect of an antimicrobial on an infection.

5.2. Indirect effects

Although, historically, the focus of the challenge trial has been to isolate direct effects on the exposed individual, one could imagine building on the design to study the effects of treatment on transmission and therefore possible indirect effects. First, we could measure the amount of susceptible and resistant pathogen an infected individual sheds through contact with their environment while in isolation and how this is affected by antimicrobial treatment. Second, we could imagine experimentally replicating the process of transmission through controlled contact with a second individual [19]. That is, at the appropriate time a second individual may be placed in the same isolation ward as the index individual and then monitored for the development of the outcome over a sufficient follow up period. Through thoughtful selection of experimental conditions, including the controlled exposure and treatment of both individuals, one could isolate different types of indirect effects and estimate their relative frequency.

(a) direct effect

(b) direct effect and indirect effects

Figure 3: SWIG for possible joint intervention on E_{i0} and A_{i0} in a randomized human challenge trial. The top panel shows the case where individual i is isolated, receives challenge exposure $E_{i0} = e_i$, and then is empirically treated with $A_{i0} = a_i$. It focuses on direct effects on person i. The bottom panel considers an alternative challenge trial in which person i and j are put in isolation together and i receives the challenge exposure while j is allowed to interact with them "naturally".

As before, possible interventions of interest could be represented via SWIGs (Figure 3) in which nodes representing the outcome Y_j of person j are re-introduced. Here we also represent the exposure of person j when they are placed in contact with person i with E_j. If, as assumed, this is done in a way that mimics "natural" exposure, i.e. we just place them in the same room and allow them to touch surfaces, interact, breathe same air, etc., then this second exposure is not "intervened" on as such and, therefore, we do not split the node in the SWIG.

5.3. Observational emulation

As mentioned, challenge trials such as those suggested above are rarely conducted in practice. These trials are typically only ethically justifiable when the risk of severe complications of infection are exceedingly low. In the absence of trial data, another option is to use observational data to try to emulate the trial one would like to run but cannot. At one extreme, one could imagine identifying index outpatient cases of suspected bacterial infection requiring treatment and enrolling them and other members of their household. To reduce external transmission, they could be asked to isolate at home together. In an inpatient setting, you might enroll suspected bacterial infections at admission and attempt to isolate them throughout the course of treatment. In both cases, to estimate the per-exposure effect, we would need information about exposure, presumably via screening tests or culture taken at baseline. Furthermore, as neither exposure nor treatment is experimentally assigned we would need to either identify quasi-random variation or measure a sufficient set of covariates to adjust for confounding. In this case, special care must be taken to identify common causes of the exposure and the outcome as well as treatment and the outcome [18].

5.4. Limitations and links to mathematical modeling

Although per-exposure effects, such as those above, allow for clearer identification of direct and indirect effect mechanisms through controlled manipulation, a limitation is that they do not immediately say what the expected outcome might be when applied to a pop-

ulation where exposure is not controlled (i.e. occurs "naturally"). A possible remedy is to simulate population effects through individual-based or mathematical modeling. These models use per-exposure effects as inputs, along with a mechanism for assigning exposure (e.g. contact-network between agents), to simulate infection dynamics under antimicrobial interventions. However, the validity of this approach depends on (1) the types of exposures studied in the real or emulated challenge trial being "representative" of those that occur in the target population in which the intervention is to be applied and (2) the mechanism for assigning exposure being "exchangeable" with the one that's active in the target population. Additional considerations may be necessary when the per-exposure effect is not constant over time [18].

6. Approach 2: Studying population effects under real-world exposures via two-stage randomization

A second approach to studying causal effects under interference is to identify aggregate groupings or clusters of individuals such that individuals within clusters interact, but those across clusters do not. Once identified, direct and indirect effects can be studied through two stages of randomization. In the first stage, clusters are randomized to different treatment strategies characterized by a certain saturation level of individual treatments (e.g. treat 30% with vancomycin and 70% with daptomycin). In the second stage, individuals within groups are randomized to receive treatment according to the assigned saturation levels of their group. This design was first described in canonical work by Halloran & Struchiner [4] as applied to vaccines, but it has also been used to study, for instance, the equilibrium effects of economic interventions. In theory, the design could also accommodate more complex group-level interventions such as cycling of antibiotics or combination therapies.

In the following sections, we describe several possible adaptations of the two-stage design to the study of antimicrobial treatments.

6.1. Design 1: Mass drug administration

Perhaps the most straightforward adaptation of the two-stage design for vaccines would be a design to capture the use of antibiotics irrespective of infection such as in the mass drug administration of azithromycin [20]. Originally used for treatment of trachoma, where it was judged more efficient to treat all than to test and treat only those with trachoma [21], mass azithromycin distribution to under-5 year olds in areas of high child mortality has been shown to reduce mortality risk. In this case the intervention does not condition on being previously exposed, infected, or presenting to a health facility, but rather is typically done in community-based campaign in lower resourced settings where healthcare and drug treatments may not be readily available. Because the intervention closely resembles a vaccination campaign, very few changes are needed from the cannonical two-stage vaccine design.

Following Hudgens & Halloran [9] and VanderWeele & Tchetgen Tchetgen [11], we expand our previous setup as follows. Consider a sample of K clusters, indexed by $k = 1, \ldots, K$, where now each cluster is comprised of n_k individuals (instead of just two). As before, individuals within clusters interact but there is (approximately) no interaction across clusters. The antimicrobial treatment received by individuals in cluster k is represented by the vector (vectors represented in **bold**) $\mathbf{A}_k = (A_{k1}, \ldots, A_{kn_k})$, where A_{kj} is the treatment received by individual j. In the MDA setting treatment only occurs at one time point and, for simplicity, we define each A_{kj} to be binary (0: no treatment, 1: azithromycin). We define $\mathbf{A}_{k,-j}$ as the treatments received by the other individuals in cluster k excluding individual j. We refer to $\mathbf{A}_k$ as the cluster-level treatment or allocation program. The set of all possible treatments for cluster k is $\mathcal{A}(n_k)$, meaning $\mathbf{A}_k$ can take one of 2^{n_k} possible values. Let $Y_{kj}^{\mathbf{a}_k}$ denote the potential outcome of individual j in cluster k at the end of follow up (e.g. 30 days) under allocation program $\mathbf{a}_k$ and let $Y_{kj}^{a, \mathbf{a}_{k,-j}}$ denote the potential outcome when individual j has treatment status a and the rest of the group k has treatment status $\mathbf{a}_{k,-j}$. As before we define Y_{kj} to be tri-level culture result (0: negative culture, 1: susceptible culture, 2: resistant culture). To ease notation, consider a simple scenario where $Y_{kj}^{\mathbf{a}_k}$ and $Y_{kj}^{a, \mathbf{a}_{k,-j}}$ only depends

on $\mathbf{a}_k$ and $\mathbf{a}_{k,-j}$ via the proportion of individuals who are treated which we denote by π, i.e. $g(\mathbf{a}_k) = \frac{1}{n_k} \sum_{j=1}^{n_k} a_{kj}$. In this case, we can simplify potential outcomes to Y_{kj}^π and $Y_{kj}^{a,\pi}$. We also allow for Y_{kj}^π and $Y_{kj}^{a,\pi}$ to be random rather than fixed by considering sampling from a superpopulation of clusters.

Under this design we can define the direct effect on resistance (i.e blood culture at 14 days after administration as positive for resistant strains) as

$$\Pr[Y_{kj}^{a=1,\pi} = 2] - \Pr[Y_{kj}^{a=0,\pi} = 2],$$

which is identified under randomization by the within-cluster comparison of those treated with antimicrobials ($A_{kj} = 1$) versus those untreated ($A_{kj} = 0$) among clusters assigned to treatment saturation π, i.e.

$$\Pr[Y_{kj} = 2 | A_{kj} = 1, \pi(\mathbf{a}_{k,-j}) = \pi] - \Pr[Y_{kj} = 2 | A_{kj} = 0, \pi(\mathbf{a}_{k,-j}) = \pi].$$

For the direct effects on other outcomes like mortality or susceptible infection we simply replace Y_{kj} with the appropriate outcome.

We can also define the indirect effects at different saturation levels π versus π' through

$$\Pr[Y_{kj}^{a,\pi} = 2] - \Pr[Y_{kj}^{a,\pi'} = 2],$$

which is identified under randomization by the between-cluster comparison of clusters assigned to saturation π versus π', i.e.

$$\Pr[Y_{kj} = 2 | A_{kj} = a, g(\mathbf{a}_{k,-j}) = \pi] - \Pr[Y_{kj} = 2 | A_{kj} = a, g(\mathbf{a}_{k,-j}) = \pi'],$$

where we could consider indirect effects on antimicrobial saturation separately on those treated or on those untreated.

Finally, we can define the population-level overall effect of saturation level π versus π' as

$$\Pr[Y_{kj}^{\pi} = 2] - \Pr[Y_{kj}^{\pi'} = 2],$$

which is identified under randomization by the marginal between-cluster comparison of clusters assigned to saturation π versus π', i.e.

$$\Pr[Y_{kj} = 2 | g(\mathbf{a}_k) = \pi] - \Pr[Y_{kj} = 2 | g(\mathbf{a}_k) = \pi'].$$

Such studies have been performed for multiple outcomes [20, 22]

6.2. Design 2: Post-exposure treatment of suspected infection

As in our motivating example, many common antimicrobial treatments are applied to those with suspected infections. This leads to several challenges in adapting the canonical design as only a portion of the population is eligible for treatment at each time point, i.e. those who are possibly infected. This means the allocation programs are *dynamic* interventions rather than the *static* interventions considered in the canonical two-stage design. Furthermore, eligibility evolves over the follow up period and is affected by treatment in previous time periods because eligibility is based on suspected infection (which is acquired from exposure to previously infected and possibly treated individuals). Finally, post-exposure studies involve complex sampling schemes: we typically only enroll and measure outcomes in those who present as a suspected infection at some point over the follow up period, we often do not measure outcomes of individuals in the cluster who do not. The latter has implications for the types of indirect effects that we can study. In the DAG in Figure 1 we represent the minimal case of interference between two individuals. The following design expands this to multiple individuals and different clusters, where we assume partial interference.

We enrich the previous setup by indexing the time each variable is measured using

$t = 0, \ldots, T$. To reduce notational burden, we suppress indices for cluster unless neces-sary. Clusters in this setting could represent distant geographies each served by a fixed health facility or health system as in a multicenter trial. We define an example dynamic post-exposure allocation program which, each day, assigns individuals within clusters who present with suspected infection to empiric antimicrobial treatment $A_t = 1$ (vancomycin) with probability π and empiric treatment $A_t = 2$ (daptomycin) with probability $1 - \pi$. That is, we assign treatment program $\mathbf{A}_t$ in each cluster at time t according to the probability mass function,

$$g(\mathbf{A}_t | \mathbf{S}_t; \pi) = \prod_{j=1}^{n} \mathbb{1}(S_{jt} = 1) \times \left\{ \pi^{\mathbb{1}(A_{jt}=1)}(1 - \pi)^{\mathbb{1}(A_{jt}=2)} \right\}.$$

This implies that, at each time point, on average, $\pi \sum_{j=1}^{n} S_{jt}$ individuals are treated and $(1-\pi) \sum_{j=1}^{n} S_{jt}$ individuals are untreated within each cluster. Taken over time this essentially defines a series of nested trials among those admitted with suspected infection at each time point. Let Y_{jt}^{π} denote the potential outcome of an individual j within a cluster at time t under an intervention that assigns antimicrobial treatment according to $g(\mathbf{A}_t | \mathbf{S}_t; \pi)$ and let $Y_{jt}^{a_t, \pi}$ denote the potential outcome of an individual assigned treatment status $A_t = a_t$ at time t while the rest of their cluster is assigned treatment according to $g(\mathbf{A}_{t,-j} | \mathbf{S}_{t,-j}; \pi)$. As before we assume potential outcomes only depend on the proportion treated within cluster and are invariant to which specific individuals are treated.

Under this design, we can define a direct effect on resistance at time $t + 14$, i.e. 14 days after admission, as

$$\Pr[Y_{j,t+14}^{a_t=2, \pi} = 2 | S_{jt}^{\pi} = 1] - \Pr[Y_{j,t+14}^{a_t=1, \pi} = 2 | S_{jt}^{\pi} = 1].$$

with other direct effects defined based on follow up and initial culture results as in Section

3, e.g. the direct effect on the emergence of resistance is

$$\Pr[Y_{j,t+14}^{a_t=2,\pi} = 2 | S_{jt}^{\pi} = 1, Y_{jt}^{\pi} = 1] - \Pr[Y_{j,t+14}^{a_t=1,\pi} = 2 | S_{jt}^{\pi} = 1, Y_{jt}^{\pi} = 1]$$

and the head-to-head comparison of daptomycin versus vancomycin on clearance of susceptible baseline infection is

$$\Pr[Y_{j,t+14}^{a_t=2,\pi} = 0 | S_{jt}^{\pi} = 1, Y_{jt}^{\pi} = 1] - \Pr[Y_{j,t+14}^{a_t=1,\pi} = 0 | S_{jt}^{\pi} = 1, Y_{jt}^{\pi} = 1].$$

In contrast with the canonical two-stage design, here there are multiple direct effects (one for each time point) each defined for a potentially non-overlapping subgroup composed of those with suspected infection at time t. These direct effect estimands are essentially what is being estimated in a classic single-stage trial such as a non-inferiority or superiority trial of a new antibiotic class where interference is ignored. That is, the single-stage trial can be conceived of as a two-stage trial with one cluster or group. However, typically these trials report a single effect estimate which pools over the direct effects at each time point. When follow up is short relative to the dynamics of transmission this may be inconsequential. However, because exposure affected by prior antimicrobial treatment assignment in the cluster, in general we would expect the effects to vary over time as the composition of resistance among those presenting with suspected bacterial infection changes over time. Under randomization, the direct effects on resistance above are identified by

$$\Pr[Y_{j,t+14} = 2 | S_{jt} = 1, A_{jt} = 2, g(\mathbf{a}_{t,-j}) = \pi] - \Pr[Y_{j,t+14} = 2 | S_{jt} = 1, A_{jt} = 1, g(\mathbf{a}_{t,-j}) = \pi],$$

which is the within-cluster comparison of those treated versus untreated after presenting with suspected bacterial infection. The results for direct effect on both AMR emergence and effectiveness in treating the infection are also identified by the within-cluster comparison among their admission culture subgroups.

Because the treatment is applied after exposure and affects subsequent exposure, the arm-specific cross-cluster comparisons considered in the mass drug administration trial do not make as much sense. Instead, we focus on the overall effect on culture results at the time infection is first suspected and a culture taken, as these have yet to be contaminated by direct treatment, but are indirectly affected by accumulating effects of treatment of previous infections, that is the potential that resistant bacteria selected by prior treatment of other patients have infected a focal patient. However, assessment of the overall effect in a post-exposure trial will depend on the sampling scheme. When all infections present to or are acquired within a health facility and the underlying source population is fully enumerated, the overall effect on resistance at time t, defined as contrasts such as

$$\Pr[Y_{jt}^{\pi} = 2] - \Pr[Y_{jt}^{\pi'} = 2],$$

is identified under randomization by the marginal between-cluster comparison of incidence of resistant infection at admission in clusters assigned to saturation π versus π', i.e.

$$\Pr[Y_{jt} = 2|g(\mathbf{a}_t) = \pi] - \Pr[Y_{jt} = 2|g(\mathbf{a}_t) = \pi'].$$

Alternatively, when we do not observe the outcome for all individuals in a cluster or when the sampling fractions are unknown, the assessment of the overall effect is more challenging. This may be the case for health facility based interventions where the catchment area is not fully enumerated or when both facility-based and community spread of infection or commensal carriage is possible, but only individuals with severe enough infection to seek care are observed. One work around when all suspected infections are cultured upon admission is to use the between cluster ratio of odds that presenting infection is susceptible or resistant to treatment as a proxy for the overall effect of treatment saturation π versus π', i.e.

$$\frac{\Pr[Y_{jt}^{\pi} = 2]\Pr[Y_{jt}^{\pi'} = 1]}{\Pr[Y_{jt}^{\pi} = 1]\Pr[Y_{jt}^{\pi'} = 2]} = \frac{\Pr[Y_{jt} = 2|S_{jt} = 1, g(\mathbf{a}_t) = \pi]\Pr[Y_{jt} = 1|S_{jt} = 1, g(\mathbf{a}_t) = \pi']}{\Pr[Y_{jt} = 1|S_{jt} = 1, g(\mathbf{a}_t) = \pi]\Pr[Y_{jt} = 2|S_{jt} = 1, g(\mathbf{a}_t) = \pi']}.$$

Figure 4: Modified two-stage randomized trial to estimate effects for antimicrobial treatments

6.3. Observational emulation

Much like the challenge trials considered previously, two-stage randomized trials are rarely conducted in practice. As before, in the absence of trial data, another option is to use observational data to attempt to emulate a two-stage trial. Emulations of two-stage trials are rarely discussed in the literature, but they are possible in theory. They would involve collecting observational data across a number of clusters with sufficient separation or buffer that the partial interference assumption holds. In this case, the observational study can be viewed as essentially a two-stage trial where allocation program $\mathbf{A}_t$ is assigned according to $f(\mathbf{A}_t|\mathbf{L}_t, \mathbf{S}_t)$ where $\mathbf{L}_t$ are pre-treatment covariates, which is a group-level generalization of the standard conditional randomization assumption in emulations of individually-randomized trials. A consequence of this is that, unlike emulations of individually-randomized trials, emulations of two-stage trials must account for potential confounding via individual and group-level covariates. Another complication of two-stage emulations is that the group-level allocation plan that each cluster pursue may be ill-defined unless there are agreed upon treatment guidelines or policies within each cluster. Emulations require data from

settings in which there is natural variation in antimicrobial strategies that were pursued across clusters. An example could be when there's a shortage or introduction of a particular antimicrobial treatment. Emulations may also be more informative in instances where there were coordinated policies at the cluster-level as this may permit more nuanced assessment of effects of policy alone versus full compliance with policy (akin to intention-to-treat versus per protocol effects in randomized trials). Regardless, data must also include information on confounders at both the individual and cluster levels, including details about antimicrobials not covered by the emulated trial.

7. Discussion

The use of causal inference methods to answer questions in acute infectious diseases where transmission plays an important role has not seen the same exponential growth as in other areas, such as the use of this methods in the study of chronic diseases [23]. Part of the difficulty in applying causal methods for ID studies stems from the fact that the assumption of no interference is often violated [24]. Additionally, most of the work in causal inference with interference has focused mainly on studies where vaccine is the primary intervention of interest.

In this paper, building on previous work in causal inference with interference, we focused on presenting how this framework could be implemented for the study of antimicrobial treatments strategies and AMR. We aimed to formalize cases of antimicrobial use employing DAGs and SWIGs to help those with interest in ID research present problems in a more unified way using tools form causal inference methods. This work continues the efforts of Ogburn and VanderWeele [15] by applying graphical representations to research problems involving interference. We also discussed how challenge trials and a two-stage randomization design (and potentially their observational emulations) could be adapted to study the indirect effects of antimicrobial treatment strategies. Traditionally, evaluations of new antimicrobial strategies have focused on assessing non-inferiority compared to the standard of care [25].

However, there is increasing interest in understanding how these strategies might affect the spread of AMR, particularly through indirect effects arising from interference between patients. There are settings where the indirect effects might play a crucial role such as the mass drug administration approach used to reduce chilhood mortality in Niger [26].

In settings such as ICUs and nursing homes—where antimicrobials are extensively used among vulnerable populations— the potential for patient-to-patient transmission is also significant [27, 28]. Expanding study designs to account for these indirect effects is key for a more comprehensive assessment of treatment impacts. By explicitly incorporating these effects, we can better evaluate how interventions influence both individual outcomes and the broader dynamics of resistance, making stronger stewardship recommendations.

Using DAGs and SWIGs as we have presented facilitates the clear specification of causal contrasts and assumptions. These graphical tools make the underlying causal structures explicit, allowing researchers to identify and address potential biases in studies assessing antimicrobial treatments and AMR. By clarifying the causal pathways under consideration and formalizing the nodes commonly present in ID studies, our approach enhances the evaluation of causal effect estimates in scenarios where interference is present, aligning with methodologies discussed by previous work [9, 11, 24]

The application of the steps described here are future steps that this research group plan to apply to different current clinical problems. We also hope that the present work will encourage the broader use of causal inference methods in the study of acute infectious diseases.

References

1. Organization world health, w. h. *Antimicrobial resistance: Report by the Director-General* tech. rep. (World Health Organization, 2020).

2. Souque, C., González Ojeda, I. & Baym, M. From Petri Dishes to Patients to Populations: Scales and Evolutionary Mechanisms Driving Antibiotic Resistance. eng. *Annual*

Review of Microbiology **78,** 361–382. doi:10.1146/annurev-micro-041522-102707 (2024).

3. Lipsitch, M. & Samore, M. H. Antimicrobial use and antimicrobial resistance: a population perspective. *Emerg Infect Dis* **8,** 347–354. doi:10.3201/eid0804.010312 (2002).

4. Halloran, M. E. & Struchiner, C. J. Causal Inference in Infectious Diseases. *Epidemiology* **6,** 142–151 (1995).

5. Huskins, W. C. *et al.* Intervention to Reduce Transmission of Resistant Bacteria in Intensive Care. *New England Journal of Medicine* **364,** 1407–1418. doi:10.1056/NEJMoa1000373 (2011).

6. Kunkel, A., Colijn, C., Lipsitch, M. & Cohen, T. How could preventive therapy affect the prevalence of drug resistance? Causes and consequences. *Philosophical Transactions of the Royal Society B: Biological Sciences* **370,** 20140306. doi:10.1098/rstb.2014.0306 (2015).

7. Duijn, P. J. v. *et al.* The effects of antibiotic cycling and mixing on antibiotic resistance in intensive care units: a cluster-randomised crossover trial. English. *The Lancet Infectious Diseases* **18,** 401–409. doi:10.1016/S1473-3099(18)30056-2 (2018).

8. Sobel, M. E. What Do Randomized Studies of Housing Mobility Demonstrate?: Causal Inference in the Face of Interference. *Journal of the American Statistical Association* **101,** 1398–1407 (2006).

9. Hudgens, M. G. & Halloran, M. E. Toward Causal Inference With Interference. *Journal of the American Statistical Association* **103,** 832–842. doi:10.1198/016214508000000292 (2008).

10. Tchetgen Tchetgen, E. J. & VanderWeele, T. J. On causal inference in the presence of interference. eng. *Statistical Methods in Medical Research* **21,** 55–75. doi:10.1177/0962280210386779 (2012).

11. VanderWeele, T. J. & Tchetgen Tchetgen, E. J. Effect partitioning under interference in two-stage randomized vaccine trials. *Statistics & probability letters* **81,** 861–869. doi:`10.1016/j.spl.2011.02.019` (2011).

12. Chanderraj, R. *et al.* Vancomycin-Resistant Enterococcus Acquisition in a Tertiary Care Hospital: Testing the Roles of Antibiotic Use, Proton Pump Inhibitor Use, and Colonization Pressure. *Open Forum Infectious Diseases* **6,** ofz139. doi:`10.1093/ofid/ofz139` (2019).

13. Adamu, Y., Puig-Asensio, M., Dabo, B. & Schweizer, M. L. Comparative effectiveness of daptomycin versus vancomycin among patients with methicillin-resistant Staphylococcus aureus (MRSA) bloodstream infections: A systematic literature review and meta-analysis. *PLOS ONE* **19,** e0293423. doi:`10.1371/journal.pone.0293423` (2024).

14. Heath, M. R. *et al.* Gut colonization with multidrug resistant organisms in the intensive care unit: a systematic review and meta-analysis. *Critical Care* **28,** 211. doi:`10.1186/s13054-024-04999-9` (2024).

15. Ogburn, E. L. & VanderWeele, T. J. Causal Diagrams for Interference. *Statistical Science* **29,** 559–578. doi:`10.1214/14-STS501` (2014).

16. Richardson T, R. J. *Single World Intervention Graphs (SWIGs) : A Unification of the Counterfactual and Graphical Approaches to Causality* in (2013).

17. Adams-Phipps, J. *et al.* A Systematic Review of Human Challenge Trials, Designs, and Safety. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* **76,** 609–619. doi:`10.1093/cid/ciac820` (2022).

18. O'Hagan, J. J., Lipsitch, M. & Hernán, M. A. Estimating the Per-Exposure Effect of Infectious Disease Interventions. en-US. *Epidemiology* **25,** 134. doi:`10.1097/EDE.0000000000000003` (2014).

19. Eyal, N. & Lipsitch, M. Testing SARS-CoV-2 vaccine efficacy through deliberate natural viral exposure. eng. *Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases* **27,** 372–377. doi:`10.1016/j.cmi.2020.12.032` (2021).

20. Peterson, B. *et al.* Assessment of Spillover of Antimicrobial Resistance to Untreated Children 7–12 Years Old After Mass Drug Administration of Azithromycin for Child Survival in Niger: A Secondary Analysis of the MORDOR Cluster-Randomized Trial. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* **79,** 1136–1143. doi:`10.1093/cid/ciae267` (2024).

21. Mariotti, S. P. New Steps toward Eliminating Blinding Trachoma. *New England Journal of Medicine* **351,** 2004–2007. doi:`10.1056/NEJMe048205` (2004).

22. O'Brien, K. S. *et al.* Azithromycin distribution and childhood mortality in compliance-related subgroups in Niger: complier average causal effect and spillovers in a cluster-randomized, placebo-controlled trial. *International Journal of Epidemiology* **51,** 1775–1784. doi:`10.1093/ije/dyab198` (2022).

23. Scola, G. *et al.* Implementation of the trial emulation approach in medical research: a scoping review. en. *BMC Medical Research Methodology* **23,** 186. doi:`10.1186/s12874-023-02000-9` (2023).

24. Halloran, M. E. & Hudgens, M. G. Causal Inference for Vaccine Effects on Infectiousness. en. *The International Journal of Biostatistics* **8,** 1–40. doi:`10.2202/1557-4679.1354` (2012).

25. Lanini, S. *et al.* Non-inferiority versus superiority trial design for new antibiotics in an era of high antimicrobial resistance: the case for post-marketing, adaptive randomised controlled trials. eng. *The Lancet. Infectious Diseases* **19,** e444–e451. doi:`10.1016/S1473-3099(19)30284-1` (2019).

26. Mack, I. *et al.* Antimicrobial Resistance Following Azithromycin Mass Drug Administration: Potential Surveillance Strategies to Assess Public Health Impact. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* **70,** 1501–1508. doi:10.1093/cid/ciz893 (2020).

27. Van den Dool, C., Haenen, A., Leenstra, T. & Wallinga, J. The Role of Nursing Homes in the Spread of Antimicrobial Resistance Over the Healthcare Network. *Infection Control and Hospital Epidemiology* **37,** 761–767. doi:10.1017/ice.2016.59 (2016).

28. Wunderink, R. G. *et al.* Antibiotic Stewardship in the Intensive Care Unit. An Official American Thoracic Society Workshop Report in Collaboration with the AACN, CHEST, CDC, and SCCM. *Annals of the American Thoracic Society* **17,** 531–540. doi:10.1513/AnnalsATS.202003-188ST (2020).