1   **Title**

2   The Oldenburg Hearing Health Record (OHHR)

3   **Authors**

4   Sumbul Jafri[1,2], Daniel Berg[2,3], Mareike Buhl[4], Matthias Vormann[2,5], Samira Saak[2,3],

5   Kirsten C. Wagener[2,5], Christiane M. Thiel[1,2], Andrea Hildebrandt[1,2] & Birger Kollmeier[2,3]

6   **Affiliation**

7   1. Department of Psychology, School of Medicine and Health Sciences, Carl von Ossietzky

8   Universität Oldenburg, Germany

9   2. Cluster of Excellence "Hearing4All", Carl von Ossietzky Universität Oldenburg, Germany

10   3. Department of Medical Physics and Acoustics, School of Medicine and Health Sciences, Carl von

11   Ossietzky Universität Oldenburg, Germany

12   4. Université Paris Cité, Institut Pasteur, AP-HP, Inserm, CNRS, Fondation Pour l'Audition,

13   Institut de l'Audition, IHU reConnect, F-75012 Paris, France

14   5. Hörzentrum Oldenburg gGmbH, Germany

15   **Corresponding authors**

16   Correspondence should be addressed to Andrea Hildebrandt or Birger Kollmeier.

17   Email: andrea.hildebrandt@uni-oldenburg.de, birger.kollmeier@uni-oldenburg.de

Abstract

Hearing health is shaped by both measurable auditory function and the perceived ability to navigate daily life. To fully understand its complexities, objective assessments of hearing and functional performance must be complemented by subjective reports on lived hearing experiences. The Oldenburg Hearing Health Record (OHHR) was developed to unite these measures, offering a comprehensive and open-access resource for hearing health. It contains data from 581 adults aged 18–86 years (*255 females; mean age = 67.31 years; SD = 11.93*) with varying degrees of hearing loss. Data were collected between 2013 and 2015 at the Hörzentrum Oldenburg in collaboration with Hearing4all. OHHR includes audiometric tests (Pure Tone Audiometry, Loudness Scaling, Speech in Noise tests), self-reports on hearing difficulties, lifestyle, technology use, and cognitive assessments (DemTect, Vocabulary size test). These measurements remain relevant in clinical and research settings. The dataset supports cross-disciplinary analyses linking hearing ability with cognition and quality of life, contributing to personalized hearing healthcare and advancing precision medicine.

Background

Hearing loss is a major global health concern[1–3]. More than 1.5 billion people experience some degree of hearing loss during their lifetime, with at least 430 million requiring hearing health care. In Europe alone, approximately 190 million individuals (20% of the population) are affected to varying degrees by hearing loss or deafness[1]. Beyond its sheer prevalence, hearing loss has a profound impact on an individual's health and quality of life. It has been associated with an elevated risk of cognitive decline, increased perception of cognitive load, and is a contributing factor to accelerated brain atrophy, particularly in regions critical for language and memory[4–7]. Neuroimaging studies show that even mild-to-moderate hearing loss increases listening effort and alters functional brain connectivity in areas supporting attention and cognitive control[8]. This increased listening effort reflects the additional cognitive resources required to process degraded auditory input and maintain speech understanding[9]. Hearing loss also has significant psychosocial consequences, including isolation, depression, and reduced social engagement, all of which further harm well- being[10,11]. Indeed, hearing loss disrupts every aspect of daily life, from difficulties understanding speech in noisy environments to safety concerns linked to poor sound localization. The sustained effort required to listen can be exhausting, with studies showing significantly elevated fatigue levels in individuals with hearing loss[12], often leading to social withdrawal. In many cases, the social stigma surrounding hearing aid use, and the added burden of comorbidity with tinnitus can make coping with hearing loss even more challenging. In workplace and educational settings, hearing loss can impede performance and achievement, further diminishing quality of life[13].

All these multifaceted associations between hearing loss, psychosocial well-being, and health underscore the need for a data-driven investigation and simultaneous modelling of these multivariate associations. Such an approach requires large, shared databases that incorporate subjective reports, multiple objective auditory measures (e.g., speech-in-noise perception, loudness tolerance), assessment of cognitive performance, comorbid health conditions and psychosocial factors to improve and promote an individualized understanding of the impact of hearing loss. The promotion of comprehensive hearing health datasets will facilitate the extension of knowledge beyond the confines of traditional and small-scale audiometric assessment and provide insight into complex multivariate associations.

An overview of existing hearing data repositories was recently published by Sethukumar and colleagues (2023)[14]. Their catalogue brings together a wide range of international publications, including studies on hearing loss, ageing, neurodegenerative disorders, healthcare and disease registries, birth cohorts and population-based studies. Each study comprises one or a few limited types of hearing data, which taken together include pure-tone audiometry, speech-in-noise measurements, otoacoustic emissions, auditory brainstem response, speech discrimination, loudness recruitment, and other clinical and self-reported hearing measures. Notably, sixty percent of the studies included in the catalogue refer to objective hearing data, and in some cases, even genetic testing. The catalogue also highlights datasets with extensive audiological testing, such as the Japanese Intractable Inner Ear Databank[15], the Swedish National Databank on Sudden Sensorineural Hearing Loss[16], and the Childhood Development after Cochlear Implant Dataset from Johns Hopkins University, USA[17]. Some cohort studies also incorporate broader biopsychosocial data along with hearing related measures from larger populations, thus providing potential for valuable insights into hearing health.

Despite these substantial contributions to the study of hearing loss, the advancement of
hearing research is hampered by the lack of comprehensive datasets that integrate extensive
audiological test data with cognitive, social, lifestyle, and medical variables in a single
record that is openly available for scientific discovery. Many of the studies catalogued by
Sethukumar and colleagues (2023) do not have access to the participants' lifestyle and
medical history. Although the UK Biobank[18] for example offers an extensive biomedical
dataset, it lacks pure tone audiometry data and uses only the Digit Triplet Test as a screening
tool for speech in noise testing, limiting its utility in evaluating functional hearing abilities.
Other studies, such as the British 1970 Cohort[19] and the AGES Reykjavik Study[20], provide
audiometric and self-reported data but do not include any speech-in-noise measurements,
which are crucial for understanding real-world communication challenges. The Rotterdam
Study[21] is a longitudinal investigation of the incidence and determinants of chronic disease
and disability in older adults. It combines data on hearing, cognition, physical and mental
health, genetic testing, neuroimaging, and epidemiological data. However, the primary focus
of this study in the field of otolaryngology is alone age-related hearing loss or presbycusis.
This focus limits its applicability to other forms of hearing impairment, such as noise
induced or ototoxic hearing loss and congenital conditions. The ongoing Rheinland study[22]
also has a special focus, namely on aging and neurodegenerative disorders. Both studies
employ the Mini-Mental State Examination for cognitive screening, which cannot detect
subtle cognitive challenges faced by individuals with hearing problems[23]. Although
the standard Mini-Mental State Examination remains widely used; research indicates
that the Montreal Cognitive Assessment and the Dementia Detection test are more
sensitive in detecting early cognitive decline in diverse populations[24,25]. Importantly,
access to many of these datasets requires formal and complex authorization, some of
which are financially burdensome, and others have institutional affiliation restrictions,
all of which limit their availability for widespread and accelerated research. It is worth
noting, however, that some of these studies are ongoing and their policies and access options
may update in the future.

To address the limitations of existing hearing health datasets and their limited accessibility,
we describe and share here a comprehensive open access dataset. The Oldenburg
Hearing Health Data Record (OHHR) was collected between 2013 and 2015 at the Hörzentrum
Oldenburg gGmbH in collaboration with the Cluster of Excellence "Hearing4all" in Germany.
OHHR is focused on individuals with well-defined hearing loss and comprises a hearing loss
cohort of 442 out of a total of 581 participants. Having a well-defined cohort of individuals
who are deaf or hard of hearing, and those who are unaffected allows researchers to isolate
the effects of hearing impairment from potential confounding variables. Although other
datasets, such as birth cohorts or disease-specific studies, offer valuable insights, they may
not capture the same level of detail on hearing loss and associated health and well-being. The
current data resource offers an extensive array of self-reported and examiner-driven
measures in conjunction with pure tone audiometry, thereby providing researchers with a
multifaceted perspective on the nature of hearing loss and associated factors. It also includes
a precise speech-in-noise test, i.e. the Göttingen Sentence Test[26] to characterize the
individual´s functional hearing deficit (to be treated with a hearing device) and the Digit
Triplet Test, which is widely used as a hearing screening tool for assessing speech recognition
in noisy environments[27–29]. The Dementia Detection test was used to identify cognitive

impairment, primarily assessing fluid cognitive abilities such as memory and attention. In
addition, a vocabulary size test was administered as a proxy for crystallized intelligence and to
estimate premorbid cognitive ability[30]. This distinction is particularly useful when interpreting
screening outcomes, as it helps differentiate between long-standing individual differences
(reflected by crystallized ability) and clinically relevant early cognitive decline (reflected by fluid
ability)[31,32]. To achieve a holistic assessment, the dataset considers not only clinical
evaluations but also draws from self-reported information using standardized questionnaires.
Thus, OHHR captures a broad spectrum of data, including socio-demographic characteristics,
hearing health history and impact on daily life, noise exposure, hearing aid use and
satisfaction, mental and physical health status, and potential comorbidities of hearing loss. Its
increased granularity facilitates the exploration of the interplay between hearing loss,
functional hearing deficit, cognitive decline, and overall quality of life and health.

Precision medicine in audiology undoubtedly benefits from extensive datasets such as the
OHHR. This dataset has supported several significant studies within the field, demonstrating its
utility in identifying individual predictors of speech-in-noise performance[33], exploring
determinants of hearing aid uptake[34], developing Common Audiological Functional
Parameters[35–37] and advancing auditory profiling for clinical decision support systems[38–40].
However, these publications relied on internal, unstructured versions of the dataset and did not
involve all the variables or any data for public sharing. Their focus was primarily on audiological
outcomes, with limited integration of the broader cognitive, psychosocial, or demographic data
included in the resource. In contrast, the curated subset described in this paper has
undergone extensive refinement in terms of structure, consistency, exhaustiveness and
documentation. It includes comprehensive variable-level metadata, detailed test
descriptions, and quality control protocols that were not publicly available before. While this
release draws from the same broader data source used in prior publications; it now offers a more
complete, transparent, and accessible version of the OHHR that is suitable for reuse by the
wider research community.

The release of OHHR in a standardized and accessible format creates opportunities for
interdisciplinary research across audiology, cognitive science, psychology, and public health. The
dataset's integration of audiological, psychosocial, cognitive, and demographic data enables a
wide range of applications—from developing predictive models for personalized rehabilitation
to informing strategies for hearing loss prevention and public health planning. Researchers can
use OHHR to identify high-risk populations, evaluate factors influencing hearing aid adoption,
and assess the broader impact of hearing loss on quality of life and daily functioning.

The inclusion of patient-reported outcomes, together with comprehensive data on hearing aid
use, provides a strong foundation for advancing patient-centered care models and evaluating
assistive technologies in real-world contexts. OHHR also supports the exploration of diverse
factors that shape the perception and management of hearing difficulties, contributing to a
more nuanced understanding of hearing loss. Its structured format and thorough documentation
promote reproducibility and cross-study comparisons, making it a valuable resource for data-
driven approaches in clinical and public health research. By making this curated dataset openly
available, we aim to support research on hearing loss by providing resources for early detection,
intervention development, and rehabilitation studies.

Methods

Participants

Participants aged 18 years and older were recruited from an existing database of volunteers at Hörzentrum Oldenburg gGmbH (abbreviated as HZO in the following). This non-profit organization is affiliated with the Carl von Ossietzky Universität Oldenburg and partners with the "Hearing4all" Cluster of Excellence funded by the German Research Foundation (https://hearing4all.de/en/). It works with a global network of scientific and industrial collaborators to develop innovative audiological methods and relies on the support of about 2,000 local volunteers. After screening for completeness, the final dataset prepared for public release via Zenodo (https://zenodo.org/records/14177902)[41] includes 581 individuals, aged 18 to 86 years (n = 255 female; median age = 70.0 years; mean age = 67.31 years; standard deviation (SD) = 11.93 years). The data was anonymized in accordance with the General Data Protection Regulation (GDPR; Regulation (EU) 2016/679). Each participant was assigned a unique identifier to maintain anonymity while enabling multivariate analyses. Data protection approval for the preparation and open release of the dataset was granted by the Data protection and Information Security Management Office at the Carl von Ossietzky University of Oldenburg, Germany.

While the OHHR cohort includes a higher proportion of individuals with hearing loss than would be expected in the general population, its purpose was not to reflect population-level prevalence of hearing loss. Rather, the aim was to establish a large cohort oversampling a wide range of hearing impairments, offering a robust foundation for research focused on the deaf and hard-of-hearing individuals. This contrasts with many population-based datasets, which often include a majority of normal-hearing individuals. Additional information on the hearing loss prevalence and how the sample compares to population-level estimates is provided in the *Technical validation* section.

Procedure

The data collection process began with an 11-page questionnaire, referred to as the "home questionnaire", which volunteers completed at home. It was sent via postal mail to potential participants along with study information and consent forms, with instructions to complete everything using pencil and paper and return it via prepaid mail. Participants who returned the completed home questionnaire were then invited to a one-hour lab-based test session. This initial process ensured that participants understood the nature of the data they would be sharing prior to any in-person testing. The test session included a face-to-face interview to discuss the individual's medical history, as well as several assessments. Besides a clinically standard pure tone audiogram, supra-threshold loudness perception was determined with the Adaptive Categorical Loudness Scaling[42]. In addition, two speech-recognition-in-noise tests (the Göttingen Sentence Test - GÖSA[26] and the German Digit Triple Test - DTT[28]), and two cognitive tests (Vocabulary size test or Wortschatztest - WST[43] and the Dementia Detection test - DemTect[25]) were performed. All measures were undertaken by trained professional staff members. Hearing assessments, including Pure Tone Audiometry, Adaptive Categorical Loudness Scaling, the GÖSA, and the German DTT, were performed

205 without hearing aids, even for regular users. However, cognitive assessments and the
206 questionnaires, where optimal hearing was required were conducted with hearing aids for those
207 who typically used them[33].

Ethics and data anonymization

The local ethics committee at the Carl von Ossietzky Universität Oldenburg reviewed and
authorized the original data collection between 2013 and 2015, ensuring compliance with the
ethical standards described in the Declaration of Helsinki[44], except for the requirement for
preregistration of the study. Participants were compensated with 12€ per hour for their
involvement.

To prepare the current publicly available dataset, additional ethical approvals and data
protection measures were implemented. All records were anonymized to protect participant
privacy. To minimize the risk of re-identification, k-anonymity (k=4) was applied, ensuring
that each participant's quasi-identifiers were indistinguishable from at least three others.
Further safeguards were introduced, including grouping categories with fewer than four cases
into broader classifications. The full anonymization process was reviewed and approved by the
university's data protection officer, and documentation outlining the procedures and risk
assessments are available upon request. At the time of the original data collection,
approximately 40% of participants had consented to the storage of their contact
information. These individuals were re-contacted and provided explicit consent for the
publication of their anonymized data. For the remaining 60%, whose data had already been
pseudonymized and who could not be re-contacted, a waiver of consent for data sharing was
granted by the data protection officer following the completion of the anonymization process.

Data protection approval for the preparation and public release of the dataset was
subsequently granted by the Data Protection and Information Security Management Office
at Carl von Ossietzky Universität Oldenburg (Application Number: DSM-H4A Open
Dataset/20241113-0009).

Self-reports included in the home questionnaire (HQ)

Subjective experience of hearing loss related consequences reflects the functional impact of hearing loss on an individual's life, an aspect that cannot be captured by audiometric measures alone. There has been much discussion in the literature about the low association between examiner-driven ("objective") and self-reported ("subjective") measures of hearing loss. However, it is crucial to also characterize individuals with hearing loss in terms of their subjective hearing-related experience, along with their subjective health status (chronic diseases), self-reported demographics (age, gender), personality traits (self-efficacy, self-esteem), mood (depressive symptoms) and social context (social network size)[45]. Research by Wang and colleagues (2020)[46] show that underreporting of hearing problems is associated with age, ear problems, lifestyle, and subjective beliefs about hearing health. Notably, older adults tend to underreport their hearing difficulties[47]. A combined approach of self-reports and audiometric measures is also effective in detecting high-frequency hearing loss. Furthermore, self-reports are particularly sensitive in contributing to the identification of mid-frequency hearing loss[48]. Understanding the factors that influence self-reported hearing problems, including cognitive abilities, education, and lifestyle, enables researchers and practitioners to interpret these reports more precisely and use them as complementary measures to audiometry for prediction and for deriving individualized treatment recommendations[49]. The HQ covered several key assessments, which are described in detail below.

Hearing anamnesis

The HQ collected detailed information on the individual's hearing history, including diagnosis and duration of hearing impairment, subjective ratings of hearing problems in both quiet and noisy environments, and noise exposure. Individuals were asked about the causes of their hearing difficulties (see Figure 1), hearing aid use (past and present), duration of use, and any history of sudden sensorineural hearing loss, for example due to middle ear infections. Age-related hearing loss (ARHL) was among the most frequently reported causes. In the context of this dataset, ARHL also known as presbycusis is a bilateral, symmetrical, and progressive sensorineural hearing loss that occurs with advancing age. It is characterized by reduced hearing sensitivity, especially for high-frequency sounds, and may be accompanied by reduced speech understanding, particularly in noisy environments. Ear noise/tinnitus was also addressed. This included its occurrence, duration and the physical or mental discomfort it causes. Furthermore, to evaluate the comfort and experience of hearing aid use, the questionnaire included three items from the seven-item International Outcome Inventory for Hearing Aids[50].

Figure 1. Self-reported causes of hearing difficulties

Note. Absolute frequencies of self-reported causes of hearing difficulties in the OHHR dataset (N = 581).
A total of 370 individuals reported a perceived cause of hearing loss, while 72 reported not knowing the
reason and 139 reported normal hearing. Age-related hearing loss (ARHL) or presbycusis showed the
highest prevalence among the reported causes (n = 134). Other reported causes included noise induced
(n = 48), sudden deafness (n = 43), explosion or firing practice (n = 40), congenital causes (n = 16), and
various other factors (e.g., ear surgery, hereditary factors, tinnitus, infections, concerts, medication,
injury, otosclerosis; n = 2–15).

General health and lifestyle

Individuals provided information about their overall health, including any limitations in daily
activities due to health problems, how their physical and mental health had affected task
performance in the four weeks prior to data collection, pain impacting daily activities,
emotional well-being, and perceived memory issues. They also reported any chronic
conditions diagnosed in their lifetime and in the past 12 months, such as respiratory disorders,
cardiovascular diseases, diabetes, musculoskeletal, or psychiatric conditions (see Table 1).
Some related conditions were grouped together for brevity in Table 1 (e.g., heart attack, angina
pectoris, and cardiac insufficiency as cardiovascular conditions), although they were asked
separately to reflect how patients typically recognize these conditions and to enable more
detailed information beyond a single general measure. These data are valuable, for example, for
assessing the burden of comorbid conditions and for calculating a comorbidity index according
to the German National Health Examination Survey[51]. This index quantifies the impact of
chronic conditions on the use of health care services, on quality of life, and on other health-
related outcomes.

Table 1. Prevalence of health problems and hearing difficulties, overall and by gender

Health problems	*n*	% in the sample	% of men	% of women
Hearing Difficulties	443	75.0	46.87	28.09
Musculoskeletal conditions	325	55.0	28.43	26.57
Respiratory conditions	79	13.4	6.09	7.28
Cardiovascular conditions	77	13.0	10.49	2.54
High blood pressure/cholesterol	317	53.6	32.49	21.15
Diabetes mellitus	52	8.8	5.41	3.38
Neurological disease, e.g., Parkinson's disease	33	5.6	4.57	1.02
Sight disorder/blindness	83	14.0	8.46	5.58
Risk of falling/dizziness	75	12.7	5.08	7.61
Chronic gastric-intestinal disease	65	11.0	5.41	5.58
Cancer	61	10.3	5.92	4.4
Psychiatric conditions	71	12.0	5.58	6.43
Other chronic diseases (kidney, bladder, skin, etc.)	172	29.1	17.77	11.34
Physical handicap (arms/legs)	108	18.3	9.31	8.97

Note. Cardiovascular conditions refer to clinically diagnosed disorders affecting the heart or vascular
system (e.g., heart attack, cardiac insufficiency or angina pectoris), whereas high blood pressure and
cholesterol represent individual cardiovascular risk factors that may occur independently of a formal
disease diagnosis. Reporting separately allows for a more refined analysis of the relationship between
comorbid health conditions, risk factors, and hearing health.

Short form 12 health survey (SF-12)[52]

The health status section of the HQ also included the SF-12 health survey, which is a concise
version of the 36-item health survey, but comprehensive enough to provide an overview of
an individual's physical and mental health. The SF-12 contains 12 questions, yielding two
summary scores. The physical component score (PCS) quantifies limitations in physical
functioning, role limitations due to physical health problems, bodily pain, general health
perceptions, and social functioning limitations due to physical problems. The mental
component score (MCS) indicates vitality, emotional well-being, and social functioning
limitations due to emotional problems, as well as interference with daily activities caused by
mental health problems. The PCS and MCS raw scores are separately transformed into
standardized scores and included in the OHHR as separate scores (*T*-values, with *M = 50,*
SD= 10).

Media consumption and device usage

A specific section of the HQ aimed to assess the participant's media consumption habits and
preferences. It included questions about headphone use for TV and radio, detailing the types
of hardware and software used (e.g., radio, Bluetooth, infrared) at the time of data collection
and how often they were used. Individuals also reported their use of technology for work and
personal activities, both with and without hearing aids. In addition, they were asked about
their experiences with sound levels, including situations where sounds were perceived as too
loud with or without hearing aids. The survey also addressed the demand and perceived
sound quality of hearing aids, the frequency of increasing TV volume without them, and
individual musical preferences and expectations for sound quality.

Technology readiness

The HQ also included a 12-item questionnaire to measure the use of adaptive technology,
particularly among older adults[53]. The measurement instrument is based on the theoretical
concepts outlined in the technology acceptance model[54], which postulates how perceived
usefulness and ease of use influence technology adoption. Furthermore, it also incorporates
concepts from Ajzen's (1991)[55] work on personal beliefs about competence and control.
Neyer and colleagues (2012)[53] extended these concepts by developing a comprehensive
model of technology readiness that integrates both attitude-oriented and personality-
theoretical perspectives. The main construct measured by the 12 items is technology
readiness, or the willingness to invest time and effort in learning and using modern
technology. Additionally, there are three subscales that can be measured with this
questionnaire. Technology competence reflects the perceived ability and confidence in using
technology effectively[56]. Technology acceptance quantifies an individual's relationship with
modern technologies, primarily in terms of interest in technical innovations[54]. Technology
control reflects the degree to which individuals feel that they have control over their
technology use and can overcome challenges[56].

Demographic and socio-economic status

The demographic information was collected in both the clinical interview (age, gender,
language status, etc.) and the HQ (household composition and living situation). Participants
also reported socio-economic details about themselves, such as their educational degree,
profession, occupation, and household income. The Scheuch-Winkler Index (SWI) was used
to quantify socio-economic status (SES) by summing up education, occupation, and net
income, resulting in a score ranging from 3 to 21[57]. This index is crucial for describing health
disparities by investigating how socio-economic factors influence health outcomes. These
scores were categorized into three SES groups—low, medium, and high—to represent
different socio-economic strata within the German population.

Overall, the HQ collected extensive data that allows for a comprehensive characterization of
the impact of hearing loss on an individual's quality of life and their biopsychosocial
characteristics, making this dataset highly valuable for informing precision audiology. By
including validated instruments along with newly designed questions, the HQ provided a
thorough dataset for analyzing the complex, multivariate interactions between hearing
health, general health, technology use, and socio-economic factors. All of this is
complemented by further objective measures of hearing and cognition, which are described
in the following.

Clinical interview for anamnesis

To gain a detailed overview about the participants' hearing health, beyond the survey data
recorded with the HQ, an expert interviewer conducted a structured face-to-face interview
with the participants. The interview was designed to record further demographic information
and questions pertaining to the onset, progression, and specific characteristics of the
individual's hearing loss, along with details on cochlear implants or any other implant, hearing
aid use (access and frequency), family history of hearing loss, and history of ear infections.

Audiological tests

Pure Tone Audiometry

Pure tone audiometry is a standard tool used in clinical settings to identify hearing
impairments. The resulting audiogram provides a comprehensive assessment of hearing
sensitivity, conductive hearing loss, cochlear function, and discomfort levels. The audiometry
procedure for OHHR was performed in the laboratory, with each ear assessed independently.
Air conduction hearing thresholds were measured using a Siemens Unity II audiometer with
Sennheiser HDA200 headphones, while bone conduction thresholds were evaluated with a
RadioEar B71 bone conduction transducer. This was done with sinusoids for frequencies
ranging from 125 Hz to 8000 Hz in an acoustically shielded audiometry booth. Bone
conduction measurements were limited to 500 to 4000 Hz, following standard clinical
practice. Below 500 Hz, the vibration threshold is too close to the hearing threshold, making
measurements unreliable. Frequencies above 4000 Hz are excluded due to excessive
distortion produced by the transducer. A standard clinical threshold method was used with a
step size of 5 dB, where a threshold was accepted if the probe tone was not heard twice at
lower levels and detected twice at higher levels. To summarize overall hearing sensitivity, the

Pure Tone Average (PTA) was calculated as the mean hearing threshold across 500, 1000,
2000, and 4000 Hz. Additionally, measurements of uncomfortable loudness levels (UCL) were
performed at 500, 1000, 2000, and 4000 Hz. Tone levels were increased in 5 dB increments
and participants were asked to indicate when the sound became uncomfortably loud by
pressing a button. The respective presentation level was recorded, and the presentation
sequence was continued for the next test frequency at a presumably comfortable level. The
complete assessment took approximately 13 minutes.

Adaptive Categorical Loudness Scaling

Adaptive Categorical Loudness Scaling (ACALOS) is a method used in clinical audiology to
measure an individual's subjective perception of loudness according to Brand and Hohmann
(2002)[42]. It is particularly adept at diagnosing loudness recruitment, which is commonly
encountered among individuals with sensorineural hearing loss. This phenomenon is
characterized by an excessive increase in perceived loudness as the sound level increases.

The stimulus signals used were one-third-octave bands of noise (duration 2 seconds)
presented through Sennheiser HDA 200 headphones for 20 trials in a pseudo-random order.
Participants were asked to rate their perceived loudness of each stimulus level using an 11-
category response scale, ranging from "not heard" to "too loud". The categories
corresponded to 0, 5, 15, 25, 35, 45 and 50 categorical units, with four unnamed categories
situated between them (10, 20, 30, 40 cu). The presentation levels were adaptively adjusted
based on the participants' previous responses, thereby ensuring that the test assessed the
whole individual range of loudness perception[42]. Measurements were recorded successively
for 1500 and 4000 Hz narrow-band noise stimuli for the left and right ear separately. A
loudness function can be fitted to the responses of each of the ACALOS measurements using
either the Brand and Kollmeier (2002)[58] staircase method or the BTUX fitting method[59],
depending on the researcher's preference[33].

The ACALOS procedure was designed to provide a reliable and efficient estimation of loudness
functions with a small number of trials. It has good test-retest reliability, with intraindividual
standard deviations for loudness levels ranging from 4-5 dB, comparable to or slightly better
than other procedures that require more trials. The adaptive approach allows for efficient use
of trials and better coverage of the auditory dynamic range, particularly benefiting people
who are hard of hearing. Additionally, ACALOS eliminates the need for prior measurements
of hearing thresholds, simplifying the testing procedure[42].

While Pure tone audiometry and ACALOS are effective in assessing hearing sensitivity,
everyday communication depends on the ability to recognize and understand speech,
particularly in noisy environments. Individuals with similar ages and audiogram thresholds
can still exhibit substantial differences in speech comprehension in noisy settings. This
underscores the importance of including speech-in-noise measures as part of a
comprehensive audiological assessment such as the OHHR intends to provide. These
additional measures are described below.

Digit Triplet Test

The Digit Triplet Test (DTT) employs a closed-set response format that facilitates convenient self-testing via telephone or the internet, primarily for hearing screening purposes. It consists of lists of twenty-seven-digit triplets spoken in background noise used to adaptively determine the speech recognition threshold (SRT). In the adaptive procedure, the signal-to-noise ratio (SNR) for each trial is influenced by the participant's previous performance. The initial SNR was set at 0 dB, with the noise level remaining constant. The speech presentation level, however, varied: it was decreased following correct identifications and increased if the listener had difficulty understanding any digit[28]. For the current database, the test was administered in the presence of an expert in a controlled laboratory environment. The DTT takes approximately three minutes to complete. The background noise was presented at 65 dB sound pressure level (SPL) for most participants. For those with severe hearing loss, where 65 dB SPL was not sufficiently audible, the presentation level was increased to 80 dB SPL to ensure the test stimuli could be perceived[60]. The purpose of this test is to estimate the SRT, which is defined as the speech level at which a person can hear 50% of the numbers correctly. The DTT offers several advantages, including its brevity, familiarity with the stimuli employed, efficiency, online accessibility, and availability in multiple languages. It has also been shown to be an effective tool for both adults and children with cochlear implants, with of reliability and resilience to learning effects, linguistic abilities, and personal factors such as educational background[61].

Göttingen Sentence Test

The Göttingen Sentence Test (GÖSA) is an open-set speech intelligibility test in which listeners hear 20 meaningful sentences, each presented one at a time in background noise and consisting of three to seven words. The 20-sentence list is randomly selected from a pool of 10 predefined sets of everyday sentences. It provides a more realistic representation of communication situations than tests that have single words or sounds presented in noise. Each sentence is meticulously designed to maintain perceptual equivalence and consistent difficulty levels across different sets, ensuring reliable and comparable results[26,62].

During the testing session, the sentences were presented with test-specific speech-shaped noise via a free-field loudspeaker in a sound-attenuated room. Participants were instructed to repeat as many parts of the sentence as they could after each sentence presentation. The GÖSA takes about 5 minutes to complete. The noise level during testing is set at 65 dB SPL and increased to 80 dB SPL for participants who were unable to perceive the stimuli adequately at the lower level due to severe hearing loss. The GÖSA also employs an adaptive procedure to determine the SRT with precision in the range of ±1 dB. The noise level is kept constant, and speech levels are adjusted based on the listener's responses, facilitating accurate assessment of speech perception abilities. In summary, the test offers a comprehensive assessment of speech intelligibility, using ecologically valid materials and carefully calibrated background noise to simulate real-world listening conditions[58].

Cognitive measures

Screening tests for dementia and cognitive assessment in general require the participant to hear and comprehend the test items sufficiently well. Similarly, audiological tests, except visual inspection of the ears, demand cognitive abilities such as attention, working memory, and semantic knowledge. These abilities are essential for attending to, comprehending, remembering, executing instructions, and communicating with the examiner. Therefore, an accurate diagnosis of cognitive and audiological conditions is contingent upon a thorough examination of both[63].

Dementia Detection Test

The Dementia Detection test (DemTect) is a neuropsychological screening tool employed to assess cognitive impairment. It is recognized for its high sensitivity and time efficiency, given that it can be administered in 8 to 10 minutes. DemTect comprises five subtests that cover a wide range of cognitive abilities, including immediate and delayed recall of verbal information, working memory, language and number processing, and executive functioning. In the Word List subtest, participants are presented with a 10-item word list over two trials to assess both immediate and delayed recall, thus targeting memory function, which is an essential domain for detecting mild cognitive impairment (MCI) and dementia[64–66]. The Number Conversion task assesses executive function and language processing by requiring participants to switch between different representations of numbers, such as words and numerals. It captures a range of errors, including those related to language impairment, lexical and syntactic processing, and literacy difficulties. Such impairments have been frequently observed in individuals with dementia[67,68]. In the Semantic Fluency Task ('Supermarket'), participants are asked to generate and name items belonging to a specific category, such as supermarket items, within a limited time. It assesses a range of cognitive abilities including attention, working memory, cognitive flexibility, problem solving, semantic memory, language production and processing speed. Studies show that verbal fluency is often impaired in the early stages of dementia[69–72]. The Digit Span task requires participants to repeat sequences of numbers in reverse order and is a measure of working memory. Deficits in this area are considered one of the earliest signs of dementia[25,73,74]. Finally, in the Delayed Recall task, participants are asked to recall the 10-item word list presented at the beginning. This task assesses long-term memory.

The overall score is calculated by summing the scores of each subtest, with each subtest incorporating age-based scoring adjustments for participants under and over 60 years of age to account for age-related differences in cognitive abilities. DemTect has demonstrated robust construct validity and high reliability in both test-retest and inter-rater assessments. Notably, it excels in detecting mild Alzheimer's disease (AD) with a sensitivity of 100% and mild cognitive impairment with a sensitivity of 80%. In contrast, the widely used Mini-Mental State Examination shows limited efficacy in identifying MCI, with a sensitivity of only 69%[23,25]. This tool enables the identification of potential cognitive decline or impairment of participants in OHHR.

Vocabulary size test

The vocabulary size test (in German Wortschatztest-WST[43]) was used to assess the verbal
intelligence and language comprehension of the participants included in the OHHR. Factor
analytic validation studies have revealed that indicators derived from the vocabulary tests
load highly on a general cognition factor (commonly called the g-factor), meaning that the
scores from WST can be interpreted as effective indicators of g[33,43]. Therefore, this
assessment has the additional purpose of estimating premorbid intelligence levels in
individuals with mild to moderate cognitive impairment and tracking the progression of
dementia.

The WST is a 10-minute test consisting of 40-word recognition tasks. In each task, participants
must identify the real word presented alongside five similar non-words. The tasks are
arranged in a line-by-line format, with increasing difficulty. The raw score is determined by
the number of correctly identified words. This test demonstrates high reliability, as evidenced
by a split-half reliability coefficient (r = .95) and Cronbach's Alpha (α = .94). The score is largely
independent of age, exhibiting a very low correlation with age (r = .08), but it is positively
correlated with educational and vocational qualifications (r = .60). It has been normed for a
wide age range (20-90 years) and has been standardized using a representative sample (N =
572) and Rasch scaling to ensure equivalent measurement of abilities across items.

Data records

The OHHR is accessible via Zenodo[41] and is arranged into three principal folders, including **data**,
metadata, and **tools**. These folders provide all the essential resources needed to access,
interpret, and use the dataset easily and effectively. The tools folder contains conversion
scripts that read and join the audiogram and loudness scaling JSON tables for potential use in
Excel, MATLAB, Python, and SQL, thereby facilitating data management and visualization. The
metadata folder includes detailed description files that define all the variables within
each .json file, while comprehensive schema files ensure data consistency and enable
accurate interpretation across varying platforms. Below is an overview of the principal folders
and their structure (see Figure 2), as well as the file names within Table 2.

All the data tables can be found (.json files) inside the data folder and their corresponding
schema files in the metadata folder. There are also description files (.rtf format) in the
metadata folder that explain the various columns inside each data table. In addition to the
primary dataset, the data folder contains six taxonomical .json files that provide verbal
representations of different categorical data. The most relevant files are listed and explained
in Table 2.

Table 2. Overview of the files providing verbal explanations for levels in categorical data

File name	Representation
tax_education.json	Contains verbal representations of education levels
tax_job.json	Contains verbal representations of job levels
tax_net_income.json	Contains verbal representations of income levels
tax_school.json	Contains verbal representations of school levels
tax_tech.json	Contains verbal representations of responses from the technical survey, extracted from the original home questionnaire
tax_yes_no_unknown.json	Contains verbal representations for yes, no, and unknown responses

526   **Figure 2. Overview of the data folders and tools of OHHR**

527   *Note*. (a) The data folder contains the core JSON dataset files, including audiogram and loudness scaling tables
528   with linked components (indicated by a downward arrow) that must be joined for use. Taxonomy files define
529   response categories. The tools folder provides resources for data visualization and processing using MATLAB,
530   Python, Excel, and SQL.

b.

(b) The metadata folder includes documentation and schema files. RTF files describe each dataset, while JSON schema files define their structure and format.

Technical validation

Standardized testing protocol

A standardized testing protocol was followed to ensure consistency and reliability in data collection. First, the order and timing of the assessments was standardized. Before any laboratory testing, the HQ was duly completed and returned. The subsequent interviews and assessments were primarily conducted on the same day. However, in certain instances, the assessments were done on alternative days when necessary or to obtain any missing information. To ensure adherence to the established protocol, the dates for each test were recorded. Second, standardized questionnaires and measurements were used wherever possible – see descriptions of the instruments above.

Medical device certification

The audiometric testing software used was certified as a medical device according to EU regulations and bears the CE mark. Classified as a legacy device under the EU directive, it is suitable for diagnostics in clinical settings and by hearing care professionals. This certification guarantees compliance with stringent safety and performance standards. More details can be found on the Hörzentrum Oldenburg website.

Calibration procedures

The audiometers were calibrated according to established standards and guidelines (ANSI S3.6-2010; ISO 389-8-2004), ensuring the precision and consistency of our auditory measurements. The speech tests and ACALOS were conducted using the Research and Development version of the Oldenburg Measurement Applications (OMA), designed for scientific research. This software operates on a commercially available PC with a high-quality sound card, supporting multi-channel capability, speech tests in various languages, and customizable noise signals. More details can be found on the Hörzentrum Oldenburg website.

Data completeness

During the validation process, we thoroughly inspected missing data across both auditory and non-auditory variables. All critical measures and questionnaires were complete, except for SF-12, which had 13 missing entries. The SF-12 was part of the HQ filled out by participants without supervision. Some entries are missing, especially in free response sections if the questions were not relevant to the participant.

Data quality assessment

We have implemented a comprehensive technical validation process to ensure the highest standards of data quality and reliability. This section details the applied methods and validation steps, emphasizing compliance with prior established knowledge.

First, where applicable, comparisons with normative data were completed to contextualize the study results within established benchmarks or reference points. For instance, the noise level was calibrated to 80 dB SPL during speech-in-noise testing for participants with severe hearing loss who could not adequately perceive the standard 65 dB SPL. Additionally, age-specific scoring criteria were applied in the DemTect assessment to account for age-related

changes in cognition. Supplementary Table 1 presents the descriptive statistics for the
standardized measurements in the dataset, including the number of items, measurement
intention, mean (±SD), and range of values. Normative reference data are primarily
available from normal-hearing population samples. The measurements are sensitive enough
to differentiate between normal hearing and impairment. However, normative data for
specific classes of hearing impairment in the German population were not consistently
available in the literature; such data, wherever found, were included in the table[75–77].

Second, the univariate and bivariate distributions of the audiological measures were analysed
according to self-reported normal hearing status or a professional diagnosis of hearing loss.
This analysis was conducted to validate that the audiological measures in the OHHR
effectively capture group differences. Figure 3 illustrates these distributions, showing clear
differences in mean values between groups. Additionally, the figure demonstrates
approximately Gaussian distributions within groups and as expected, strong bivariate
associations between different measures of hearing ability, supporting the reliability and
validity of the audiological data. It is important to note that this figure is intended solely to
validate the consistency of hearing-related measures, rather than to assess differences in
other audiological or non-audiological variables between groups.

Third, we explored whether the expected low, but well-established association between
examiner-assessed and self-reported measures of hearing loss, as reported in the literature,
was also present for the variables included in the OHHR. To this end, Figure 4 illustrates the
distribution of Pure Tone Average thresholds, as well as Speech Recognition Thresholds
from the Göttingen Sentence Test and Digit Triplet Test, across varying levels of self-
reported hearing problems. These measures are presented for both quiet and noise
conditions. The figure demonstrates a small but consistent positive association between
self-reported hearing difficulties and objective measures, including bone and air conduction
thresholds, as well as speech-in-noise performance. These findings validate the OHHR
hearing-related variables against the established literature.

Figure 3. Univariate and bivariate distributions of audiological measures stratified by hearing status

Note: PTA (Pure Tone Average): Measures hearing sensitivity across frequencies (0.5, 1, 2, 4 kHz) using air (AC) and bone conduction (BC) thresholds for both left and right ears, expressed in decibels Hearing Level (dB HL). SRT_DTT (Speech Reception Threshold – Digit Triplet Test): Assesses the ability to understand digit sequences in background noise at 50% accuracy, expressed in decibels Signal-to-Noise Ratio (dB SNR). SRT GÖSA (Speech Reception Threshold – Göttingen Sentence Test): Evaluates speech recognition at 50% accuracy with sentences presented in noise, expressed in dB SNR. Off-diagonal plots display the bivariate scatterplots between each pair of measures for normal-hearing (NH; blue) and deaf or hard of hearing (D/HH; red) individuals. Both axes represent the respective measurement values in dB HL for PTA and dB SNR for DTT and GÖSA. Diagonal plots show the distribution of each measure by hearing status, with the x-axis representing measurement values (in dB HL or dB SNR) and the y-axis representing probability density. Correlation coefficients (Pearson's *r*) for each pair are shown in the upper triangle: Overall, NH, and D/HH., illustrating their relationships within and across groups.

Figure 4. Bivariate distributions of objective auditory measures (PTA Air and Bone conduction, Speech in noise measures) by self-reported hearing severity

Note. Comparison of self-reported hearing difficulties and objective auditory measures across three
domains: Air Conduction (a), Bone Conduction (b), and Speech-in-Noise Measures (c). The violin plots
display the distribution of Pure Tone Average (PTA) thresholds at key frequencies (500, 1000, 2000,
and 4000 Hz) for both ears, measured through bone and air conduction, as well as Speech Recognition
Thresholds (SRT in dB SNR) from the Digit Triplet Test (DTT) and Göttingen Sentence Test (GÖSA). Self-
reported hearing problems are categorized by severity in quiet and noise conditions. Black dots
represent individual data points, and solid lines indicate the Mean ± Standard deviation. These plots
illustrate how subjective perceptions of hearing difficulties align with objective auditory thresholds
and speech-in-noise performance across different tests.

Representativeness and prevalence

To contextualize the OHHR dataset within the German population, we compared key
sample characteristics with those reported in the Gutenberg Health Study (GHS)[2]. The GHS is
a large, population-based cohort from Mainz and the Mainz-Bingen district, comprising
5,024 individuals aged 25–86 years (*mean age = 61.2, SD ±13.4 years*; 48.4% women). While
the GHS provides population-level estimates, the OHHR intentionally oversampled
individuals with hearing loss to capture a broader spectrum of severity for clinical research.

In the OHHR, 76.1% of participants reported being diagnosed with hearing loss, substantially
higher than the 34.5% prevalence in the GHS. Applying Röser's four-frequency
classification[78] to the better ear, the OHHR showed a greater proportion of moderate
(22.0%), severe (61.1%), and profound (15.1%) hearing loss, while mild and normal hearing
cases were few (1.0% and 0.7%, respectively). In contrast, the GHS reported higher rates of
mild loss (22.5%) and lower rates for other more severe categories. Bilateral hearing loss
affected 86.6% of OHHR participants, compared to 28.5% in the GHS, and 47% of OHHR
participants reported tinnitus, almost double the 25.5% prevalence in the GHS. Bilateral
hearing aid provision was also higher (41.8% vs. 7.7%). Sex-specific differences were
consistent with the GHS findings, showing hearing loss more common in men (84.7%) than
women (65.1%). Hearing loss prevalence in the OHHR increased with age, rising from 44.3%
in participants under 55 years to over 80% in those aged 75 and older. This pattern is similar to
the age-related trends reported in the GHS, where prevalence ranged from 16.7% in
the 55–59 year group to 71.1% in the 75–79 year group.

In the subgroup of participants who reported being diagnosed with hearing loss (n = 442), most
belonged to the middle (n = 223) and high (n = 158) socioeconomic status (SES) categories as
classified by the Scheuch-Winkler Index, with fewer individuals in the low SES group (n = 61).
Profound and severe hearing loss was more common across all SES levels: profound hearing loss
affected 54.1% (low SES), 50.7% (middle SES), and 51.3% (high SES), while severe hearing loss
ranged from 29.5% (low SES) to 38.6% (middle SES) and 34.8% (high SES). Unlike the GHS, which
reported significantly higher hearing loss prevalence in lower SES groups (p < 0.001), the OHHR
cohort showed no clear SES gradient. This likely reflects the clinical nature of the sample and the
greater likelihood of individuals from more advantaged backgrounds to participate in research.
A similar pattern emerged for education. Among participants with hearing loss, most had
middle-level education (n = 279; school students or those with technical or vocational training)
or high-level or university education (n = 142), while only 21 individuals had low or no formal
vocational qualification. Profound hearing loss was slightly more common in the low education
group (54.1%), but both middle (50.5%) and high (51.4%) education groups also showed
substantial proportions of profound hearing loss. The small size of the low education group
warrants caution in interpreting this difference.

Overall, while our cohort is not representative of the general population in terms of hearing
loss prevalence, it is highly representative of the clinical population most affected
by hearing impairment. The inclusion of individuals with a broad range of severity, a high
prevalence of bilateral hearing loss, and frequent comorbid tinnitus makes this dataset
particularly valuable for research on hearing rehabilitation and patient-centered care. The
comparison with the GHS highlights these distinctions and situates our findings within the

668   broader context of hearing health in Germany.

669 Usage notes

670 We recommend verifying the completeness of any columns of interest before analysis. If
671 missing data is identified, it is advisable to either impute the missing cases[79] or exclude them
672 to ensure accuracy.

673 Documentation and version control

674 Comprehensive documentation of data collection methods, transformations, and version
675 control procedures have been prepared. This ensures transparency and accuracy in handling
676 the dataset throughout its lifecycle. Users can access the latest version of the dataset on
677 Zenodo[41] (see below for access information), where all version histories are recorded.

678 Compliance with the FAIR principles

The dataset adheres to the FAIR principles (Findable, Accessible, Interoperable, and
Reusable), ensuring usability for a wide range of research. Findability is achieved through its
deposition in a public repository such as the "Hearing4all" Zenodo Community, with the
present description as a supporting reference. Accessibility is ensured through its open-access
status (under CC BY 4.0 license), allowing unrestricted access to all data files, which are stored
in machine-readable JSON format and accompanied by comprehensive documentation.
Interoperability is supported by storing all data in JSON format including JSON schema files
and taxonomies, providing standardized data structures and definitions to facilitate
integration with various analytical systems and tools. Reusability is ensured by rich metadata,
clear licensing, and custom tools that support data access across popular analysis platforms
(Python, MATLAB, Excel, SQL), supporting easy adaptation for future research.

690 Temporal consideration and outlook

The data included in the OHHR were collected between 2013 and 2015 and should therefore
be interpreted in the context of the period in which they were acquired. Core clinical methods,
such as pure-tone audiometry, have remained consistent over time and continue to be widely
applied in both clinical practice and hearing research. As a result, these clinical measures do
not become outdated as quickly as some research methodologies, which tend to evolve more
rapidly. Other components of the dataset, including sentence recognition thresholds in noise
and categorical loudness scaling, were collected using methods that were relatively new at the
time but have since become more firmly established in hearing research and, to some extent,
in clinical audiology. A persistent challenge in expanding the clinical application of such
measures are the limited availability of comprehensive reference data from well-characterized
hearing-impaired populations, which is precisely the gap the OHHR aims to address.

Although no substantial changes in unaided hearing patterns are currently known, users of the
dataset should remain mindful of potential demographic shifts, advancements in hearing aid
technology, and evolving healthcare practices that may influence the interpretation and
applicability of findings to contemporary populations. Future research could benefit from
combining the OHHR with more recent datasets to examine such trends over time, potentially
offering valuable insights into changing patterns in hearing health and related outcomes.

Code availability

The dataset is available in a machine-readable JSON format, ensuring easy accessibility across different platforms. Custom example scripts are provided to access individual audiogram and loudness scaling files from Python, MATLAB, and Excel, allowing users to analyze and visualize the data effortlessly. Additionally, an SQL script is available for importing the entire dataset into an SQL database for advanced querying. These tools support smooth integration into a wide range of analysis environments.

Excel scripts

- **Excel/Audiogram.xlsx**: An example for using audiograms in Excel. Reads the JSON tables (live data connection), joins the tables as needed, and shows audiogram plots with the possibility to select an observation by its identification number (ID). The relative path of the .json files can be configured on the 'Setup' sheet.
- **Excel/LoudnessScaling.xlsx**: An example for using loudness scaling measurements in Excel. Reads the JSON tables (live data connection), joins the tables as needed, and shows loudness function plots with raw data and fits with the possibility to select an observation by ID.

MATLAB scripts

- **MATLAB/audiogram_getaudiogram.m:** Returns an audiogram for a user specified observation by joining relevant tables.
- **MATLAB/audiogram_plot.m**: Example script for calling 'audiogram_getaudiogram.m' and plotting the audiogram.
- **MATLAB/loudness_scaling_getls.m**: Returns a loudness scaling measurement for a passed observation by joining relevant tables.
- **MATLAB/loudness_scaling_plot.m**: Example script for calling 'loudness_scaling_getls.m' and plotting up to four loudness functions with up to two fits and the raw data points.

Python scripts

- **Python/plot_audiogram.py**: A script for plotting audiograms for an observation by joining relevant tables.
- **Python/plot_loudness_scaling.py**: A script for plotting loudness scaling for an observation by joining relevant tables.

SQL script

- **SQL/import.sql**: A script including documentation for importing the database into an SQL Database.

The full dataset, metadata and the above-mentioned scripts are available under CC BY 4.0 license at https://zenodo.org/records/14177902.

Acknowledgements

We extend our gratitude to all participants who generously contributed their data for the development of OHHR. We also acknowledge Anja Gieseler and Maike Tahden for their efforts in consolidating the specification document detailing the data codes in the previous version of this dataset, which significantly facilitated the description and refinement of the database. This research has been funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2177/1 - Project ID 390895286.

Author contributions

Conception, design, and preparation of the manuscript: Sumbul Jafri, Daniel Berg, Christiane Thiel, Andrea Hildebrandt

Contribution to study design: Kirsten C. Wagener, Matthias Vormann, Birger Kollmeier

Recruiting participants and data acquisition: Matthias Vormann, Kirsten C. Wagener

Database management: Daniel Berg, Matthias Vormann, Mareike Buhl, Samira Saak

All authors critically reviewed and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Rights and permissions

References

1. World Health Organization. *World Report on Hearing*. (2021).

2. Döge, J. & others. The Prevalence of Hearing Loss and Provision with Hearing Aids in the Gutenberg Health Study. *Deutsches Ärzteblatt International* **120**, 99 https://doi.org/10.3238/arztebl.m2022.0385 (2023).

3. Hackenberg, B. & others. Hearing loss and its burden of disease in a large German cohort—hearing loss in Germany. *The Laryngoscope* **132**, 1843–1849 https://doi.org/10.1002/lary.29980 (2022).

4. Arlinger, S. Negative consequences of uncorrected hearing loss—a review. *International Journal of Audiology* **42**, 17–20 https://doi.org/10.3109/14992020309074639 (2003).

5. Ray, J., Popli, G. & Fell, G. Association of cognition and age-related hearing impairment in the English longitudinal study of ageing. *JAMA Otolaryngology–Head & Neck Surgery* **144**, 876–882 https://doi.org/10.1001/jamaoto.2018.1656 (2018).

6. Wang, H. F. & others. Hearing impairment is associated with cognitive decline, brain atrophy and tau pathology. *EBioMedicine* **86**, https://doi.org/10.1016/j.ebiom.2022.104336 (2022).

7. Pichora-Fuller, M. K. & others. Hearing impairment and cognitive energy: The framework for understanding effortful listening (FUEL). *Ear and Hearing* **37**, 5S-27S https://doi.org/10.1097/AUD.0000000000000312 (2016).

8. Rosemann, S. & Thiel, C. M. The effect of age-related hearing loss and listening effort on resting state connectivity. *Scientific Reports* **9**, 2337 https://doi.org/10.1038/s41598-019-38816-z (2019).

9. Rönnberg, J., Holmer, E. & Rudner, M. Cognitive hearing science and ease of language understanding. *International Journal of Audiology* **58**, 247–261 https://doi.org/10.1080/14992027.2018.1551631 (2019).

10. Shukla, A. & others. Hearing loss, loneliness, and social isolation: a systematic review. *Otolaryngology–Head and Neck Surgery* **162**, 622–633 (2020).

11. Nordvik, Ø. & others. Generic quality of life in persons with hearing loss: a systematic literature review. *BMC Ear, Nose and Throat Disorders* **18**, 1–13 https://doi.org/10.1186/s12901-018-0051-6

(2018).

12. Hornsby, B. W. Y. The effects of hearing aid use on listening effort and mental fatigue associated

with sustained speech processing in adults with hearing loss. *Ear and Hearing* **34**, 523–534

https://doi.org/10.1097/AUD.0b013e31828003d8 (2013).

13. Dixon, P. R. & others. Health-Related Quality of Life Changes Associated With Hearing Loss. *JAMA*

Otolaryngology–Head & Neck Surgery **146**, 630–638 https://doi.org/10.1001/jamaoto.2020.0674 (2020).

14. Sethukumar, P. & others. Cataloging Existing Hearing Loss Cohort Data to Guide the Development of

Precision Medicine for Sensorineural Hearing Loss: A Systematic Review. *International Archives of*

Otorhinolaryngology https://doi.org/10.5152/iao.2023.22690 (2023) doi:10.5152/iao.2023.22690.

15. Kanatani, Y. & others. National registry of designated intractable diseases in Japan: present status and

future prospects. *Neurologia Medico-Chirurgica* **57**, 1–7 https://doi.org/10.2176/nmc.st.2016-0135 (2017).

16. Nosrati-Zarenoe, R., Hansson, M. & Hultcrantz, E. Assessment of diagnostic approaches to idiopathic

sudden sensorineural hearing loss and their influence on treatment and outcome. *Acta Oto-Laryngologica*

130, 384–391 https://doi.org/10.3109/00016480903161541 (2010).

17. Fink, N. E. & others. Childhood development after cochlear implantation (CDaCI) study: Design and

baseline characteristics. *Cochlear Implants International* **8**, 92–116 https://doi.org/10.1002/cii.333 (2007).

18. Sudlow, C. & others. UK biobank: an open access resource for identifying the causes of a wide range

of complex diseases of middle and old age. *PLoS Medicine* **12**, e1001779

https://doi.org/10.1371/journal.pmed.1001779 (2015).

19. Brown, M. 1970 British cohort study. *Open Health Data* **2**, e6 (2014).

20. Fisher, D. E. & others. Sex-specific predictors of hearing-aid use in older persons: The age,

gene/environment susceptibility-Reykjavik study. *International Journal of Audiology* **54**, 634–641

https://doi.org/10.3109/14992027.2015.1024889 (2015).

21. Ikram, M. A. & others. The Rotterdam Study. Design update and major findings between 2020 and

2024. *European Journal of Epidemiology* **39**, 183–206 https://doi.org/10.1007/s10654-023-01094-1

(2024).

22. Xu, P. & others. Hypothalamic substructural integrity is associated with age, sex and cognitive

function across lifespan: A comparative analysis of two large population-based studies.

https://doi.org/10.1002/alz.084672 (2024) doi:10.1002/alz.084672.

23. Pinto, T. C. & others. Is the Montreal Cognitive Assessment (MoCA) screening superior to the Mini-

Mental State Examination (MMSE) in the detection of mild cognitive impairment and Alzheimer's

Disease (AD) in the elderly? *International Psychogeriatrics* **31**, 491–504

https://doi.org/10.1017/s1041610218001370 (2019).

24. Tang-Wai, D. F. & others. CCCDTD5 recommendations on early and timely assessment of

neurocognitive disorders using cognitive, behavioral, and functional scales. *Alzheimer's Dementia (NY)* **6**,

https://doi.org/10.1002/trc2.12057 (2020).

25. Kalbe, E. & others. DemTect: a new, sensitive cognitive screening test to support the diagnosis of

mild cognitive impairment and early dementia. *International Journal of Geriatric Psychiatry* **19**, 136–143

https://doi.org/10.1002/gps.1042 (2004).

26. Kollmeier, B. & Wesselkamp, M. Development and evaluation of a German sentence test for

objective and subjective speech intelligibility assessment. *Journal of the Acoustical Society of America*

102, 2412–2421 https://doi.org/10.1121/1.419624 (1997).

27. Smits, C., Kapteyn, T. S. & Houtgast, T. Development and validation of an automatic speech-in-noise

screening test by telephone. *International Journal of Audiology* **43**, 15–28

https://doi.org/10.1080/14992020400050004 (2004).

28. Zokoll, M. A., Wagener, K. C., Brand, T., Buschermöhle, M. & Kollmeier, B. Internationally

comparable screening tests for listening in noise in several European languages: The German digit triplet

test as an optimization prototype. *International Journal of Audiology* **51**, 697–705

https://doi.org/10.3109/14992027.2012.690078 (2012).

29. Van den Borre, E., Denys, S., van Wieringen, A. & Wouters, J. The digit triplet test: a scoping review.

International Journal of Audiology **60**, 946–963 https://doi.org/10.1080/14992027.2021.1902579 (2021).

30. Lehrl, S., Triebig, G. & Fischer, B. Multiple choice vocabulary test MWT as a valid and short test to

estimate premorbid intelligence. *Acta Neurologica Scandinavica* **91**, 335–345

https://doi.org/10.1111/j.1600-0404.1995.tb07018.x (1995).

31. Overman, M. J., Leeworthy, S. & Welsh, T. J. Estimating premorbid intelligence in people living with

dementia: a systematic review. *International Psychogeriatrics* **33**, 1145–1159

https://doi.org/10.1017/S1041610221000302 (2021).

32. McDonough, I. M. & others. Discrepancies between fluid and crystallized ability in healthy adults: a

behavioral marker of preclinical Alzheimer's disease. *Neurobiology of Aging* **46**, 68–75

https://doi.org/10.1016/j.neurobiolaging.2016.06.011 (2016).

33. Gieseler, A. & others. Auditory and Non-Auditory Contributions for Unaided Speech Recognition in

Noise as a Function of Hearing Aid Use. *Frontiers in Psychology* **8**, 219

https://doi.org/10.3389/fpsyg.2017.00219 (2017).

34. Tahden, M. A., Gieseler, A., Meis, M., Wagener, K. C. & Colonius, H. What keeps older adults with

hearing impairment from adopting hearing aids? *Trends in Hearing* **22**, 2331216518809737

https://doi.org/10.1177/2331216518809737 (2018).

35. Buhl, M., Warzybok, A., Schädler, M. R., Majdani, O. & Kollmeier, B. Common Audiological

Functional Parameters (CAFPAs) for single patient cases: Deriving statistical models from an expert-

labelled data set. *International Journal of Audiology* **59**, 534–547

https://doi.org/10.1080/14992027.2020.1728401 (2020).

36. Buhl, M., Warzybok, A., Schädler, M. R. & Kollmeier, B. Sensitivity and Specificity of Automatic

Audiological Classification Using Expert-Labelled Audiological Data and Common Audiological

Functional Parameters. *International Journal of Audiology* **60**, 16–26

https://doi.org/10.1080/14992027.2020.1817581 (2021).

37. Buhl, M. Interpretable clinical decision support system for audiology based on predicted Common

Audiological Functional Parameters (CAFPAs). *Diagnostics* **12**, 463

https://doi.org/10.3390/diagnostics12020463 (2022).

38. Saak, S. K., Hildebrandt, A., Kollmeier, B. & Buhl, M. Predicting Common Audiological Functional

Parameters (CAFPAs) as Interpretable Intermediate Representation in a Clinical Decision-Support System

for Audiology. *Frontiers in Digital Health* **2**, 596433 https://doi.org/10.3389/fdgth.2020.596433 (2020).

39. Mousavi, H., Buhl, M., Guiraud, E., Drefs, J. & Lücke, J. Inference and learning in a latent variable

model for beta distributed interval data. *Entropy* **23**, https://doi.org/10.3390/e23050552 (2021).

40. Saak, S., Huelsmeier, D., Kollmeier, B. & Buhl, M. A flexible data-driven audiological patient

stratification method for deriving auditory profiles. *Frontiers in Neurology* **13**, 959582

https://doi.org/10.3389/fneur.2022.959582 (2022).

41. Jafri, S. & others. OHHR – The Oldenburg Hearing Health Record [dataset], version 1.2.0.0.

https://doi.org/10.5281/zenodo.15727482 (2025) doi:10.5281/zenodo.15727482.

42. Brand, T. & Hohmann, V. An adaptive procedure for categorical loudness scaling. *Journal of the*

Acoustical Society of America **112**, 1597–1604 https://doi.org/10.1121/1.1502902 (2002).

43. Schmidt, K.-H. & Metzler, P. *Wortschatztest*. (Beltz, 1992).

44. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for

medical research involving human subjects. *JAMA* **310**, 2191–2194

https://doi.org/10.1001/jama.2013.281053 (2013).

45. Pronk, M., Deeg, D. J. & Kramer, S. E. Explaining discrepancies between the digit triplet speech-in-

noise test score and self-reported hearing problems in older adults. *Journal of Speech, Language, and*

Hearing Research **61**, 986–999 https://doi.org/10.1044/2018_jslhr-h-17-0124 (2018).

46. Wang, D. & others. Analysis of influential factors of self-reported hearing loss deviation in young

adults. *Journal of Public Health* **28**, 455–461 (2020).

47. Kamil, R. J., Genther, D. J. & Lin, F. R. Factors associated with the accuracy of subjective

assessments of hearing impairment. *Ear and Hearing* **36**, 164–167

https://doi.org/10.1097/aud.0000000000000075 (2015).

48. Louw, C., Swanepoel, D. W. & Eikelboom, R. H. Self-Reported Hearing Loss and Pure Tone

Audiometry for Screening in Primary Health Care Clinics. *Journal of Primary Care & Community Health*

9, 2150132718803156 https://doi.org/10.1177/2150132718803156 (2018).

49. Gibson, W. K., Cronin, H., Kenny, R. A. & Setti, A. Validation of the self-reported hearing questions

in the Irish Longitudinal Study on Ageing against the Whispered Voice Test. *BMC Research Notes* **7**, 361

https://doi.org/10.1186/1756-0500-7-361 (2014).

50. Cox, R. M. & Alexander, G. C. The International Outcome Inventory for Hearing Aids (IOI-HA):

Psychometric Properties of the English Version. *International Journal of Audiology* **41**, 30–35

https://doi.org/10.3109/14992020209101309 (2002).

51. Bellach, B. M., Knopf, H. & Thefeld, W. Der Bundes-Gesundheitssurvey 1997/98. *Gesundheitswesen*

Sonderheft **60**, S59–S68 (1998).

52. Ware, J. E., Kosinski, M. & Keller, S. D. A 12-Item Short-Form Health Survey: Construction of

Scales and Preliminary Tests of Reliability and Validity. *Medical Care* **34**, 220–233

https://doi.org/10.1097/00005650-199603000-00003 (1996).

53. Neyer, F. J., Felber, J. & Gebhardt, C. Entwicklung und Validierung einer Kurzskala zur Erfassung

von Technikbereitschaft. *Diagnostica* **58**, 87–99 https://doi.org/10.1026/0012-1924/a000067 (2012).

54. Davis, F. D. Perceived Usefulness, Perceived Ease of Use, and User Acceptance of Information

Technology. *MIS Quarterly* **13**, 319–340 https://doi.org/10.2307/249008 (1989).

55. Ajzen, I. The Theory of Planned Behavior. *Organizational Behavior and Human Decision Processes*

(1991).

56. Krampen, G. *Fragebogen Zu Kompetenz-Und Kontrollüberzeugungen (FKK)*. (Hogrefe, Verlag für

Psychologie, 1991).

57. Winkler, J. & Stolzenberg, H. Adjustierung des Sozialen-Schicht-Index für die Anwendung im

Kinder- und Jugendgesundheitssurvey (KiGGS). (2009).

58. Brand, T. & Kollmeier, B. Efficient adaptive procedures for threshold and concurrent slope estimates

for psychophysics and speech intelligibility tests. *Journal of the Acoustical Society of America* **111**, 2801–

2810 https://doi.org/10.1121/1.1479152 (2002).

59. Oetting, D., Brand, T. & Ewert, S. D. Optimized loudness-function estimation for categorical

loudness scaling data. *Hearing Research* **316**, 16–27 https://doi.org/10.1016/j.heares.2014.07.003 (2014).

60. Wagener, K. C. & Brand, T. Sentence intelligibility in noise for listeners with normal hearing and

hearing impairment: Influence of measurement procedure and masking parameters. *International Journal*

of Audiology **44**, 144–156 https://doi.org/10.1080/14992020500057517 (2005).

61. Cullington, H. E. & Aidi, T. Is the Digit Triplet Test an Effective and Acceptable Way to Assess

Speech Recognition in Adults Using Cochlear Implants in a Home Environment. *Cochlear Implants*

International **18**, 97–105 https://doi.org/10.1080/14670100.2016.1273435 (2017).

62. Kuehnel, V., Kollmeier, B. & Wagener, K. Entwicklung und Evaluation eines Satztests für die

deutsche Sprache I: Design des Oldenburger Satztests. *Zeitschrift Für Audiologie* **38**, 4–15 (1999).

63. Lemke, U. Hearing Impairment in Dementia – How to Reconcile Two Intertwined Challenges in

Diagnostic Screening. *Audiology Research* **1**, https://doi.org/10.4081/audiores.2011.e15 (2011).

64. Petersen, R. C. & others. Mild cognitive impairment: clinical characterization and outcome. *Archives*

of Neurology **56**, 303–308 (1999).

65. Petersen, R. C. & others. Current concepts in mild cognitive impairment. *Archives of Neurology* **58**,

1985–1992 (2001).

66. Petersen, R. C. & others. Practice parameter: Early detection of dementia: Mild cognitive impairment

(an evidence-based review) [RETIRED]. *Neurology* **56**, 1133–1142 https://doi.org/10.1212/wnl.56.9.1133

(2001).

67. Deloche, G. & Seron, X. From Three to 3: A Differential Analysis of Skills in Transcoding Quantities

Between Patients with Broca's and Wernicke's Aphasia. *Brain* **105**, 719–733

https://doi.org/10.1093/brain/105.4.719 (1982).

68. Seron, X. & Deloche, G. From 4 to four: a supplement to 'From three to 3'. *Brain* **106**, 735–744

https://doi.org/10.1093/brain/106.3.735 (1983).

69. Tröster, A. I., Salmon, D. P., McCullough, D. & Butters, N. A comparison of the category fluency

deficits associated with Alzheimer's and Huntington's disease. *Brain and Language* **37**, 500–513

https://doi.org/10.1016/0093-934x(89)90032-1 (1989).

70. Chertkow, H. & Bub, D. Semantic Memory Loss in Dementia of Alzheimer's Type: What Do Various

Measures Measure? *Brain* **113**, 397–417 https://doi.org/10.1093/brain/113.2.397 (1990).

71.   Monsch, A. U. & others. Comparisons of verbal fluency tasks in the detection of dementia of the

Alzheimer type. *Archives of Neurology* **49**, 1253–1258

https://doi.org/10.1001/archneur.1992.00530360051017 (1992).

72.   Cerhan, J. H. & others. Diagnostic Utility of Letter Fluency, Category Fluency, and Fluency

Difference Scores in Alzheimer's Disease. *Clinical Neuropsychologist* **16**, 35–42

https://doi.org/10.1076/clin.16.1.35.8326 (2002).

73.   Belleville, S., Peretz, I. & Malenfant, D. Examination of the working memory components in normal

aging and in dementia of the Alzheimer type. *Neuropsychologia* **34**, 195–207

https://doi.org/10.1016/0028-3932(95)00097-6 (1996).

74.   Baddeley, A. D., Baddeley, H. A., Bucks, R. S. & Wilcock, G. K. Attentional control in Alzheimer's

disease. *Brain* **124**, 1492–1508 https://doi.org/10.1093/brain/124.8.1492 (2001).

75.   von Gablenz, P. & Holube, I. Hearing loss and speech recognition in the elderly. *Laryngo-Rhino-*

Otologie **96**, 759–764 https://doi.org/10.1055/s-0043-119388 (2017).

76.   Thiele, C., Sukowski, H., Lenarz, T. & Lesinski-Schiedat, A. Göttinger Satztest im Störgeräusch für

verschiedene Gruppen von Schwerhörigkeit. *Laryngo-Rhino-Otologie* **91**, 782–788

https://doi.org/10.1055/s-0031-1295419 (2012).

77.   Drixler, K., Morfeld, M., Glaesmer, H., Brähler, E. & Wirtz, M. Validierung der messung

gesundheitsbezogener lebensqualität mittels des short-form-health-survey-12 (SF-12 Version 2.0) in einer

deutschen normstichprobe. *Zeitschrift für Psychosomatische Medizin und Psychotherapie* **66**, 272–286

https://doi.org/10.13109/zptm.2020.66.3.272 (2020).

78.   Boenninghaus, H. G. & Röser, D. New tables for the determination of percentile loss of speech

hearing. *Zeitschrift für Laryngologie, Rhinologie, Otologie und ihre Grenzgebiete* **52**, 153–161 (1973).

79.   Jäger, S., Allhorn, A. & Bießmann, F. A benchmark for data imputation methods. *Frontiers in Big*

Data **4**, 693674 https://doi.org/10.3389/fdata.2021.693674 (2021).

Absolute frequencies observed for different self-reported causes of hearing loss in the OHHR

a.

📁 data

- anamnesis.json
- homeqsn.json
- audiogram.json
- ↓ audiogram_line.json
- ↓ audiogram_point.json
- loudness_scaling.son
- ↓ loudness_scaling_data.json
- ↓ loudness_scaling_data_point.json

Fit functions:
- ↓ loudness_scaling_fit_brand.json
- ↓ loudness_scaling_fit_point_brand.json
- loudness_scaling_fit_oetting.json
- ↓ loudness_scaling_fit_point_oetting.json
- digit_triplets_test.json
- goettingen_sentence_test.json
- demtect.json
- verbal_intell.json
- sf12.json
- tech.json
- soc_econ_stat.json
- swi.json

----Taxonomy files----
- tax_education.json
- tax_job.json
- tax_net_income.json
- tax_school.json
- tax_tech.json
- tax_yes_no_unknown.json

📁 tools

- 📁 **Excel**
  - Audiogram.xlsx
  - LoudnessScaling.xlsx

- 📁 **MATLAB**
  - LoadJSON.m
  - audiogram_getaudiogram.m
  - audiogram_plot.m
  - loudness_scaling_getls.m
  - loudness_scaling_plot.m

- 📁 **Python**
  - plot_audiogram.py
  - plot_loudness_scaling.py

- 📁 **SQL**
  - import.sql

Legend

↓	Tables that are linked together
	JSON data files
	JSON taxonomy files and schema files
	RTF data description files
	Excel files for visualizing audiogram and loudness scaling data
	SQL files for data import
	MATLAB files for visualizing audiogram and loudness scaling data
	Python files for visualizing audiogram and loudness scaling data

b.

📁 **metadata**

📄 anamnesis_description.rtf	📄 anamnesis_schema.json
📄 homeqsn_description.rtf	📄 homeqsn_schema.json
📄 audiogram_description.rtf	📄 audiogram_schema.json
	📄 audiogram_line_schema.json
	📄 audiogram_point_schema.json
	📄 loudness_scaling_schema.json
	📄 loudness_scaling_data_schema.json
	📄 loudness_scaling_data_point_schema.json
📄 loudnessscaling_description.rtf	📄 loudness_scaling_fit_brand_schema.json
	📄 loudness_scaling_fit_point_brand_schema.json
	📄 loudness_scaling_fit_oetting_schema.json
	📄 loudness_scaling_fit_point_oetting_schema.json
📄 digit_triplets_test_description.rtf	📄 digit_triplets_test.json
📄 goettingen_sentence_test_description.rtf	📄 goettingen_sentence_test.json
📄 demtect_description.rtf	📄 demtect.json
📄 verbal_intell_description.rtf	📄 verbal_intell.json
📄 sf12_description.rtf	📄 sf12.json
📄 tech_description.rtf	📄 tech.json
📄 soc_econ_stat_description.rtf	📄 soc_econ_stat.json
📄 swi_description.rtf	📄 swi.json

Legend

↓   Tables that are linked together

📄   JSON data files

📄   JSON taxonomy files and schema files

📄   RTF data description files

📊   Excel files for visualizing audiogram and loudness scaling data

📄   SQL files for data import

🔺   MATLAB files for visualizing audiogram and loudness scaling data

🐍   Python files for visualizing audiogram and loudness scaling data

Air conduction

Left ear

Right ear

Self-reported hearing problems in Quiet

Self-reported hearing problems in Noise

PTA (dB HL)

Legend

Very slight
Slight
Medium
Strong
Very strong

PTA (500, 1000, 2000, 4000 Hz) values
Mean ± SD

Bone conduction

Left ear

Right ear

Self-reported hearing problems in Quiet

PTA (dB HL)

Very slight   Slight   Medium   Strong   Very strong

Self-reported hearing problems in Noise

Legend

Very slight

Slight

Medium

Strong

Very strong

PTA (500, 1000, 2000, 4000 Hz) values

Mean ± SD

Speech in noise measures

Digit Triplet Test

Göttingen Sentence Test

Self-reported hearing problems in Quiet

Speech Recognition Threshold (SRT)

Self-reported hearing problems in Noise

Very slight   Slight   Medium   Strong   Very strong

Legend

Very slight
Slight
Medium
Strong
Very strong
Individual SRT values
Mean ± SD