The potential of attractive insecticide-treated nets (ITNs) in reducing malaria transmission: a modeling study

Nicolas Moiroux*[1,2], Cédric Pennetier[1]

[1] MIVEGEC, IRD, CNRS, Univ. Montpellier - Montpellier, France
[2] Pôle de Zoologie Médicale, Institut Pasteur Dakar - Dakar, Sénégal

*Corresponding author
Correspondence: nicolas.moiroux@ird.fr

ABSTRACT

Introduction: Recent studies suggest that insecticide-treated nets (ITNs) may actively attract malaria vectors, increasing their likelihood of coming into contact with the insecticide while potentially reducing personal protection. The impact of such attractive ITN on malaria transmission at the community level remains unclear. In this study, we developed a model to compare the effectiveness of attractive, inert and deterrent ITNs under varying levels of human usage and different degrees of physiological or behavioral resistance to insecticides in malaria vectors.

Methods: We developed a model to simulate the host-seeking, feeding and mortality (HSFM) of mosquito vectors facing ITNs. This model allows mosquitoes to choose between two rooms based on the presence and remote influence (attractive, inert or deterrent) of ITNs. The HSFM model was then integrated into a malaria transmission model to compare the *Plasmodium* transmission potential of mosquitoes exposed to these different type of ITNs under various scenarios of ITN coverage, physiological resistance, quantitative behavioral resistance, and qualitative behavioral resistance.

Results: The model predicts that attractive ITNs consistently reduce malaria transmission potential of vectors more effectively than inert or deterrent ITNs, even in the presence of resistant vector phenotypes. For instance, at intermediate use rate (50%), strongly attractive ITNs are expected to reduce transmission by up to 67% compared to deterrent ITNs. In resistant vector populations, attractive ITNs remained more effective overall, though the reduction in transmission were less pronounced.

Conclusion: Our findings suggest that both inert and attractive ITNs could enhance malaria control efforts, outperforming current deterrent ITNs, even in resistant vector populations. Shifting from deterrent to inert or attractive ITNs could significantly improve vector control strategies, warranting further research and product development to fully explore and optimize this approach.

Keywords: ITN, LLIN, vector control, behavior, plasmodium falciparum, resistance

Introduction

Malaria is a parasitic disease caused by *Plasmodium* protozoa transmitted between humans by *Anopheles* mosquitoes. In 2022 alone, an estimated 249 million cases of malaria and 608 000

deaths were reported worldwide [1]. Thanks to Insecticide-Treated Nets (ITNs), the malaria targets of the 2015 Millennium Development Goals were achieved. Between 2000 and 2015, an estimated 663 million cases were averted, with 68% of this reduction attributed to ITNs [2]. During this period, ITNs were the most widespread intervention, with usage rates increasing globally from less than 2% to 54% [1].

However, despite this progress, malaria remains a serious public health concern. In fact, while the number of deaths remains stable, global malaria cases have actually been rising since 2014, with an increase of 19 million cases in 2022 compared to 2014 [1]. Furthermore, sub-Saharan Africa continues to bear more than 90% of the disease burden, with most of the reported cases occurring in this region [1].

Several factors may be contributing to this resurgence, including insufficient ITN coverage and usage, and the spread of insecticide resistance mechanisms among malaria vector populations [3]. Resistance to pyrethroids - the only insecticide class used in ITNs until recently – has become widespread in sub-Saharan African [4]. The prevalence of both physiological and behavioral resistant phenotypes has increased among malaria vector populations, allowing mosquitoes to survive or avoid contact with treated ITNs, thereby enabling continued transmission to persist. Physiological resistance mechanisms, which enhance mosquito survival after ITN contact, are the most extensively studied. Knockdown resistance (*kdr)* mutations and the overexpression of detoxification enzymes have been well characterized, both of which reduce mosquito susceptibility to pyrethroids [5]. In contrast, behavioral resistance mechanisms are less studied but can be classified as qualitative or quantitative [6]. Qualitative behavioral resistance mechanisms allow mosquitoes to avoid insecticide contact through spatial, temporal, or trophic change. Examples include change in biting time or increased exophagy following implementations of ITN or indoor residual spraying (IRS) [7,8]. Quantitative behavioral resistance, on the other hand, occurs after insecticide contact and involve behavioral responses that reduce insecticide absorption. A well-documented example is the irritant effect of certain pyrethroids [9], which induces escape behaviors and decreases contact duration.

ITNs are mosquito bed nets made of polyester, polyethylene, or polypropylene fibers that are either coated with pyrethroid insecticides or have the insecticide incorporated into the fibers during manufacture. A person sleeping under an ITN acts as a lure for mosquitoes, which are intended to be killed upon contact with the insecticide-treated fibers. Thus, ITNs operate as a "pull-to-kill" vector control strategy. However, paradoxically, most WHO-recommended ITNs exhibit repellent properties, reducing mosquito attraction to human dwellings and hosts. This property, known as deterrence, is measured in experimental huts and defined as the reduction in mosquito entry rates into a hut with a human under an ITN compared to a control hut with a human under an untreated net [10].

While deterrence increases the personal protection provided by ITNs by reducing the number of bites, it also lowers the likelihood of mosquitoes contacting the insecticide, thereby decreasing mosquito mortality rates. The impact of deterrence on community-wide malaria transmission has been evaluated using mathematical models [11], which theoretically predict that although deterrent ITNs offer better personal protection, they are less effective than inert ITNs in reducing malaria transmission.

Conversely, field studies have shown unexpectedly high levels of mosquito attraction to human hosts protected by certain WHO-recommended ITNs, with greater numbers of mosquitoes collected in huts with ITNs than in control huts with untreated nets [12]. These ITNs, which enhance mosquito attraction to hosts, could potentially increase bite exposure, reducing personal protection under sub-optimal usage conditions or when nets are physically deteriorated. Vector control strategies are designed and evaluated primarily for their community-level effects and "attractive" ITNs could substantially influence both vector mortality and disease transmission dynamics. Understanding the impact of the remote (attractive, inert or deterrent) effect of ITNs at the community level is therefore essential for guiding product development and public health policy. Extrapolating findings of Killeen *et al.* [11], whose model did not account for the possibility that ITNs might increase mosquito attraction, we hypothesize that attractive ITNs could contribute to reducing malaria

transmission, offering opportunities for novel ITN designs that may help address recent malaria
resurgences.
In order to explore the theoretical potential of attractive ITNs to reduce malaria transmission in
communities, we developed a modelling framework that combines a host-seeking, feeding and
mortality (HSFM) model for vectors with a malaria transmission model. The HSFM model allows
Anopheles mosquitoes to choose between two human hosts depending on the presence of ITNs
and their remote effects (attractive, inert or deterrent). By integrating the HSFM model into a
transmission model (adapted from Glunt *et al.* [13]), we compared the *Plasmodium* transmission
potential by mosquitoes exposed to attractive, inert, or deterrent ITNs. Our analysis explored a
wide range of realistic conditions that represent major sources of residual malaria transmission,
including different ITN usage rates and varying levels of physiological resistance (ITN-induced
mortality), quantitative behavioral resistance (indoor escaping behaviors), and qualitative
behavioral resistance (spatial-temporal avoidance of ITNs). To our knowledge, this is the first study
to explicitly model implications of attractive ITNs in terms of malaria transmission.

Methods

Host-seeking, feeding and mortality (HSFM) model

114 The HSFM model (Figure 1) is a static and deterministic model of the adult *Anopheles* feeding
115 cycle with three physiological states (host-seeking HS, blood-fed BF, and resting R). From this
116 model, we calculated transition probabilities among physiological states, which were then used in
117 the transmission potential model.
118

Figure 1 - Anopheles Host-Seeking, Feeding, and Mortality (HSFM) model used to
feed a malaria transmission model.
Blue circles represent physiological states of Anopheles mosquitoes (HS: Host-
seeking; BF: Blood-fed; R: Resting). An Anopheles vector looking for a blood-meal
(in state Host-Seeking: HS) will either enter the room of an ITN-protected human
(with probability E_p) or the room of an unprotected one (with probability E_u),
depending on the remote effect of the ITN (attractive, inert or repulsive) and the

human ITN usage rate in the community. After entering a hut, a vector will either be
diverted (impaired and postponed to host-seeking the next night, with probability D)
or attempt to feed, with two possible outcomes: feeding (with probability φ_1) or death
(with probability μ_1). Then, blood-fed (BF) vectors will either die due to post-bite
mortality (μ_2) or survive (S_2), entering the Resting state (R). A vector that enters state
R will complete its gonotrophic cycle (g) with probability S^g and enter state HS to
begin a new cycle.
134

The model considers a community of humans of known size N_h in which a known proportion U_h
is using an ITN with a known 'true' protective efficacy of Π (i.e. the proportion of normal exposure
to mosquito bites upon humans lacking ITNs, which occurs indoors at times when nets would
normally be in use [11]). The average number N_p of people protected by an ITN is therefore:

(1) $N_p = U_h \Pi N_h$

And the average number N_u of unprotected people is:

(2) $N_u = N_h - N_p$

An individual *Anopheles* vector in the host-seeking (HS) state (Figure 1) in this human
community will either enter the room of an ITN-protected human or that of an unprotected one,
depending on the remote effect of the ITN (attractive, inert or repulsive) and the relative availability
of each host type (ITN-protected or unprotected). We expressed the remote effect of ITNs as the
mosquito preference (p_{ITN}) for an ITN-protected host over an unprotected one. This parameter
could be measured in a dual choice olfactometer (i.e. p_{ITN} would be the proportion of *Anopheles*
mosquitoes that choose the ITN-protected human over the total number of *Anopheles* having made
a choice) or in experimental hut trials (i.e. p_{ITN} would be the proportion of *Anopheles* mosquitoes
that entered the hut equipped with an ITN over the sum of *Anopheles* mosquitoes that entered the
hut with an ITN or the control hut with an untreated net). Thus:
-   if p_{ITN} > 0.5, the ITN is attractive, i.e. it increases host attraction,
-   if p_{ITN} = 0.5, the ITN is inert (neither attractive nor repulsive), i.e. it does not interfere with
host attraction,
-   if p_{ITN} < 0.5, the ITN is repulsive, i.e. it reduces host attraction,
156

We assumed that a host-seeking *Anopheles* mosquito is always faced with two potential hosts,
each of whom may be protected or unprotected, and that it must choose between them. It cannot
avoid the choice, continue searching, or switch to a non-human host. Under this assumption, the
mosquito encounters one of the following cases, each with its associated probability:
- (a) two ITN-protected hosts, with probability C_{pp} or
- (b) one ITN-protected and one unprotected host, with probability C_{pu} or
- (c) two unprotected hosts, with probability C_{uu}.
164

Probabilities C_{pp}, C_{pu} and C_{uu} depend on the numbers N_u and N_p of, respectively, unprotected
and ITN-protected hosts in the community. Assuming that users and non-users of ITNs are
randomly distributed in space, probabilities C_{pp}, C_{pu} and C_{uu} follow a hypergeometric distribution
with parameters $N = N_h$, $K = N_p$, and $n = 2$. The hypergeometric distribution describes the probability
of getting *k* successes in *n* draws without replacement from a population of size *N* containing exactly
K successes. We calculated the probabilities that $k = 2$, $k = 1$, and $k = 0$ corresponding to cases
(a), (b), and (c), respectively, using the following formula:

(3) $P(X = k) = \dfrac{\binom{N_p}{k}\binom{N_h - N_p}{2-k}}{\binom{N_h}{2}} = \dfrac{\binom{N_u}{k}\binom{N_u}{2-k}}{\binom{N_h}{2}}$

Therefore, probabilities C_{pp}, C_{pu} and C_{uu} are calculated as follow:

(4) $C_{pp} = P(X = 2) = \dfrac{\binom{N_p}{2}\binom{N_u}{0}}{\binom{N_h}{2}}$

(5) $C_{pu} = P(X = 1) = \dfrac{\binom{N_p}{1}\binom{N_u}{1}}{\binom{N_h}{2}}$

(6) $C_{uu} = P(X = 0) = \dfrac{\binom{N_p}{0}\binom{N_u}{2}}{\binom{N_h}{2}}$

With $C_{pp} + C_{pu} + C_{uu}$ = 1.
180
181   From this, we can compute the probabilities E_u and E_p that a host-seeking vector will enter a
182   house with an unprotected or an ITN-protected human, respectively, as follows:

183   (7) $E_u = C_{uu} + C_{pu}(1 - p_{ITN})$
184
(8) $E_p = C_{pp} + C_{pu}p_{ITN} = 1 - E_u$

After entering a room with an unprotected host, a vector will either be diverted (impaired and
postponed to host-seeking until the next night) with probability D_u, or it will attempt to bite (with
probability 1-D_u). If it attempts to bite, the vector will either be killed (pre-bite mortality) with
probability μ_{1u}, or it will take a blood meal with probability ϕ_{1u}, thereby entering state blood-fed
(BF). Knowing D_u and μ_{1u}, we deduce ϕ_{1u} and calculate ϕ_u, the overall probability of successful
feeding when entering a room with an unprotected host:

(9) $\phi_{1u} = 1 - \mu_{1u}$
193
(10) $\phi_u = (1 - D_u)\phi_{1u}$

Then, vectors that blood-fed on an unprotected host will either die with probability μ_{2u} (post-bite
mortality) or survive and enter the resting state (R) with probability S_{2u}. Knowing μ_{2u}, we have:

(11) $S_{2u} = 1 - \mu_{2u}$

The parameters μ_{1u}, D_u, ϕ_u, ϕ_{1u}, S_{2u} and μ_{2u} have their equivalents μ_{1p}, D_p, ϕ_p, ϕ_{1p}, S_{2p} and
μ_{2p}, respectively, for mosquitoes entering a room with an ITN-protected host. The following
expressions (12)-(14) correspond to expressions (9)-(11), but for a room with an ITN-protected
host:

(12) $\phi_{1p} = 1 - \mu_{1p}$
203
204   (13) $\phi_p = (1 - D_p)\phi_{1p}$
205
206   (14) $S_{2p} = 1 - \mu_{2p}$

207
208   From this HSFM model described above, we then derived the variables needed to feed the
209   "Vector average lifetime infectious bites" model of Glunt *et al.* [13], as described below.
210

First we calculate the probability P_d that a host-seeking (HS) vector will fail to feed, become
diverted, and postpone its feeding attempt until the next night (conditional on surviving). P_d is the
sum of the probabilities that a mosquito will be diverted in rooms with ITN-protected and
unprotected hosts:

(15) $P_d = E_u D_u + E_p D_p$

Diverted mosquitoes will survive until the next night with probability S_d which is equal to the
baseline daily survival rate S of *Anopheles*:

(16) $S_d = S$

Next we calculate the probability $P_{f,u}$ that an HS vector will successfully bite an unprotected
human during the same night and enter the blood-fed state (BF). $P_{f,u}$ is the product of the probability
E_u of entering a room with an unprotected host and the probability ϕ_u of feeding in that room:

(17) $P_{f,u} = E_u \phi_u$

Similarly, the probability $P_{f,p}$ that an HS vector will successfully bite an ITN-protected human
during the same night is obtained by multiplying the probability E_p of entering a room with an ITN-
protected host by the probability ϕ_p of feeding in that room:

(18) $P_{f,p} = E_u \phi_p$

The sum of $P_{f,u}$ and $P_{f,p}$ gives the probability P_f that a host-seeking (HS) vector will successfully
bite during the same night:

(19) $P_f = P_{f,u} + P_{f,p}$

The proportion F_u of mosquitoes that fed on unprotected hosts is:

(20) $F_u = P_{f,u}/P_f$

According to the HSFM model, a proportion of blood-fed (BF) mosquitoes die shortly after
feeding, while the remainder survive to enter the resting (R) state (Figure 1), with probabilities S_{2u}
or S_{2p} depending on whether they fed on an unprotected or ITN-protected host. Until the end of the
gonotrophic cycle (of duration g), mosquitoes survive at a constant daily rate S. Therefore the
probability S_f of surviving from a successful feeding to the next gonotrophic cycle in the host-
seeking state (HS) is:

(21) $S_f = \left(F_u S_{2u} + (1 - F_u) S_{2p} \right) S^g$
239

Transmission potential (TP) model

Variables calculated above from the HSFM model are used in the model described by Glunt *et*
al. [13] to estimate the transmission potential, defined as the average number of infectious bites a
vector is expected to give during its lifetime. The following formulae differ slightly in notation from
those used by Glunt *et al.*.
First we need to determine P_{fA}, the probability that an HS vector survives to take a feed.
Theoretically, a vector may experience zero to an infinite number of diversions before feeding.
Therefore, P_{fA} is given by a geometric series with first term P_f (the probability of successfull biting

during the same night) and ratio $P_d S_d$ (the probability of being diverted and surviving to HS state
the following night):

(22) $\quad P_{fA} = P_f/(1 - P_d S_d)$

Next, we need the average number B_A of subsequent bites that an HS vector survives to give.
B_A is given by a geometric series with first term P_{fA} (the probability that an HS vector survives to
take a feed) and ratio $S_f P_{fA}$ (the probability of taking a feed and surviving to the following
gonotrophic cycle in the HS state):

(23) $\quad B_A = P_{fA}/(1 - S_f P_{fA})$

The probability *c* that a vector will become infected by *Plasmodium falciparum* while taking a
blood-meal is equal to the product of the infectiousness *k* (i.e., the probability of acquiring infection
while biting an infected human) and the prevalence I_h of *Plasmodium falciparum* infection in the
human population:

(24) $\quad c = k I_h$

Therefore, the probability P_I that a vector will acquire *Plasmodium falciparum* during its lifetime
is a geometric series with first term $P_{fA}c$ (the probability that an HS vector survives to take a feed
and acquire *Plasmodium falciparum*) and ratio $S_f P_{fA}(1 - c)$ (the probability of taking a non-
infectious feed and surviving to the next gonotrophic cycle in the HS state):

(25) $\quad P_I = P_{fA}c/\left(1 - S_f P_{fA}(1 - c)\right)$

Considering a newly infected vector, it will experience a combination of diversions (*i*) and
gonotrophic cycles (of duration *g*) during the sporogonic period (of duration *n*) until becoming an
infectious vector. As demonsrated by Glunt *et al.* [13], the probability S_I that a newly infected vector
will survive to the HS state as an infectious individual is given by the following equation:

(26) $\quad S_I = S_f \sum_{i=0}^{\lceil n-g \rceil} (P_f S_f)^{\left\lceil \frac{n-g-i}{g} \right\rceil} (P_d S_d)^i \dfrac{\left(i + \left\lceil \frac{n-2g-i}{g} \right\rceil + if\left(\left\lceil \frac{n-g-i}{g} \right\rceil g + i = n-g, 1, 0\right)\right)!}{\left(\left\lceil \frac{n-2g-i}{g} \right\rceil + if\left(\left\lceil \frac{n-g-i}{g} \right\rceil g + i = n-g, 1, 0\right)\right)! \, i!}$

We finally computed the transmission potential TP (i.e., the average number of infectious bites
during a vector's lifetime, denoted V_{LAIB} in [13]) by multiplying the probability P_I that a vector
acquires *Plasmodium falciparum* during its lifetime, the probability S_I that a newly infected vector
survives to the HS state as an infectious individual, and the average number B_A of subsequent bites
a HS vector is expected to survive to deliver:

(27) $\quad TP = P_I S_I B_A$

We calculated the relative transmission potential (RTP) between two scenarios as the ratio of
the transmission potential TP_{test} computed for the tested scenario to the transmission potential
TP_{ref} computed for the reference scenario:

(28) $\quad RTP = \dfrac{TP_{test}}{TP_{ref}}$

Finally, the percentage change in malaria transmission potential ($RedTP$) between tested and
reference scenarios was computed based on RTP:

(29) $RedTP = (RTP - 1) \times 100$

Parameterization

The models have 15 input parameters, whose default values were defined according to the
literature (Table 1). The default pre-bite and post-bite mortalities in both types of rooms (with or
without ITNs), as well as diversion probability in a room with an ITN, were set to the mean values
calculated from experimental hut trials in [12] (Supplementary figure 2). Because the diversion
probability in a hut without an ITN (D_u) is expected to be overestimated in recent experimental hut
trials (as these used untreated holed nets as reference), we extracted values from older trials
carried out during the 1980s [14,15], where unprotected peoples (i.e., with no nets at all) were used
as the control. From these data, the diversion probability D was calculated as the proportion of
unfed and alive *Anopheles* mosquitoes out of the total number of mosquitoes entering the hut. Pre-
bite mortality μ_1 was calculated as the proportion of unfed and dead *Anopheles* mosquitoes out of
those that were not diverted. Post-bite mortality probability μ_2 was calculated as the proportion of
dead, blood-fed *Anopheles* mosquitoes out of all blood-fed *Anopheles* (alive or dead). Regarding
other entomological parameters, we set the baseline daily survival rate S (during resting and after
diversion) to 0.9, and the duration of the gonotrophic cycle (g) to 3 days, based on field data for
Anopheles gambiae [16–18]. The default value for the duration of sporogony (n) was set to 11 days,
corresponding to the value given by Detinova's formula [19] at a temperature of 26°C for
Plasmodium falciparum. The infectiousness (k) of mosquitoes when biting an infected human was
set to 0.1 according to modelling work by Churcher *et al.* [20]. The default ITN use rate in the human
population (U_h) was set to 50%, corresponding to the current ITN use rate in sub-Saharan Africa
[1]. The default proportion of human exposure to bites that occurs while ITNs are in use (Π) was
set to 0.75, an intermediate value among those recorded in field studies across Africa [21–24]. We
set the default value for the human community size (N_h) to 1000 inhabitants and the prevalence of
infection (I_h) in this population to 50%, a common infection level in sub-Saharan Africa [25].

Table 1: Baseline parameters used in Host-Seeking, Feeding and Mortality (HSFM) and transmission models.

Symbol	Definition	Default value	Range [min, max] or values used in simulations	Range [min, max] and distribution used in sensitivity analysis	Refs.*
S	Daily survival rate (*An. gambiae s.l.*)	0.9	-	[0.61, 0.98]; uniform	[16]
g	Average duration of gonotrophiq cycle (days)	3	-	[2, 5]; discrete uniform	[17,18]
p_{ITN}	Preference of vector for ITN-protected human (against unprotected human) as it might be recorded in a dual choice olfactometer or in experimental hut trials. If p_{ITN} > 0.5, ITN is attractive; if p_{ITN} = 0.5, ITN is inert (neither attractive nor repulsive); if p_{ITN} < 0.5, ITN is repulsive.	0.5, 0.36	0.5, 0.6, 0.7	-	[12,26]
μ_{1u}	Pre-bite mortality probability when attempting to bite an unprotected human	0.03	-	[0, 0.05]; uniform	[12]
μ_{1p}	Pre-bite mortality probability when attempting to bite an ITN-protected human	0.72	[0, 1]	[0.22, 1]; uniform	[12]
D_u	Diversion probability when entering a room of an unprotected human	0.12	-	[0, 0.3]; uniform	[14,15]
D_p	Diversion probability when entering a room of an ITN-protected human	0.31	[0, 1]	[0, 0.51]; uniform	[12]
μ_{2u}	Post-bite mortality probability after a successful feeding on an unprotected human	0.005	-	[0, 0.028]; uniform	[12]
μ_{2p}	Post-bite mortality probability after a successful feeding on an ITN-protected human	0.29	-	[0, 1]; uniform	[12]
N_h	Size of the human community (N inhabitants)	1000	-	[300, 5000]; uniform	-
I_h	Probability of *Plasmodium falciparum* infection in the human population	0.5	-	[0.1, 0.9]; uniform	[25]
k	Infectiousness: probability that a vector become infected (exposed) while taking a blood meal on an infectious host	0.1	-	[0.02, 0.2]; uniform	[20]
n	Duration of extrinsic incubation period of *Plasmodium falciparum* (days)	11	-	[8, 15]; discrete uniform	[19,27,28]
U_h	ITN usage probability in the human population	0.5	[0, 1]	[0.1, 0.9]; uniform	[25]
Π	Proportion of normal exposure to mosquito bites upon humans lacking ITNs, which occurs indoors at times when nets would normally be in use (= ITN protective efficacy)	0.75	[0, 1]	[0.45, 0.9]; uniform	[21–24]

310 *for default values and/or ranges used in sensitivity analysis.

Simulations

Relative transmission potential between attractive and inert ITNs, or between the latter and deterrent ITNs, was simulated while varying (i) ITN use rate (U_h) in the human population, (ii) mosquito pre-bite mortality (μ_{1p}) in rooms equipped with ITN (representing different levels of physiological resistance), (iii) mosquito diversion (D_p) when entering a room equipped with an ITN (representing varying levels of quantitative behavioral resistance), and (iv) the proportion (Π) of human exposure to bites that occurs while ITNs are in use (representing varying levels of mosquito spatiotemporal avoidance of ITNs, i.e., qualitative behavioral resistance).

Attractive ITNs induced vector preferences of $p_{ITN} = 0.6$ or $p_{ITN} = 0.7$, meaning that 60% or 70% of malaria vectors, respectively, selected a human under an ITN when given a choice against an unprotected human. These levels of attraction have been observed multiple times in field studies (Supplementary Figure 1) [12,26]. The inert ITN induced a vector preference of $p_{ITN} = 0.5$, indicating no preference (i.e., vectors were equally likely to select an ITN-protected or unprotected human). In contrast, the deterrent ITN induced a vector preference of $p_{ITN} = 0.36$, meaning that only 36% of malaria vectors choose a human under an ITN when presented against an unprotected human. This value represent the mean deterrence effect observed among significantly deterrent ITNs reported in the literature (Supplementary Figure 1) [12].

Relative transmission potential between deterrent, inert or attractive ITNs and no nets (i.e. 0% use) was also simulated in the same scenarios as above.

Uncertainty and sensitivity analysis

We conducted an uncertainty and sensitivity analysis to evaluate the robustness of the model results and the impact of input parameters beyond the specific scenarios tested in prior simulations. Based on the literature, we defined plausible ranges of values for the parameters (Table 1 and Supplementary Figure 2). We then employed Latin Hypercube Sampling [29] to generate 500 simulations comparing moderately attractive ITNs to deterrent ITNs. We assumed that parameter values were sampled from uniform probability distributions, either discrete or continuous. We calculated partial rank correlation coefficients (PRCC) between outcomes and parameters, as well as partial slope coefficients (PSC; i.e., linear regression coefficients) with their 95% bootstrapped confidence intervals.

Software:

Models were coded in R [29]. The uncertainty and sensitivity analysis was performed using the 'LHS' function from the 'pse' package version 0.4.7 [30].

Results

We simulated the impact of providing human communities with attractive ITNs on malaria transmission potential, expressed as the average number of infectious bites a vector is expected to give during its lifetime, and compared the results to communities receiving either inert or deterrent ITNs.

Effect of ITN usage rates

According to our model, attractive ITNs are expected to reduce malaria transmission more effectively than inert or deterrent ITNs, regardless of usage levels. With all other parameters set to default values based on literature and field data (Table 1 and Supplementary Figure 2), the greatest transmission reduction was observed at use rates of 92–94%. At these levels, a highly attractive ITN ($p_{ITN} = 0.7$) is expected to reduce *Plasmodium* transmission potential, expressed as the average number of infectious bites a vector delivers during its lifetime, by 56% compared to an inert ITN (Figure 2A) and by 74% compared to a deterrent ITN (Figure 2B). Similarly, at 92-94% use rate, moderately attractive ITNs ($p_{ITN} = 0.6$) are predicted to reduce malaria transmission by 33% and 60% compared to inert and deterrent ITNs, respectively (Figure 2A and 2B). At 50% ITN

use, inert and attractive ITNs are expected to reduce transmission by 33-63% compared to deterrent ITNs (Figure 2B). Compared to no nets (0% use), attractive LLINs used by 50% of the population are predicted to reduce transmission by 93%, which is 12 percentage points (p.p.) more than deterrent ITNs (81% reduction; supplementary Figure 3A).

Figure 2 - Reduction in *Plasmodium* transmission potential ($RedTP$) induced (A) by attractive Insecticide-Treated Nets (ITNs) compared to inert ITNs and (B) by attractive and inert ITNs compared to deterrent ITN for varied levels ITN usage rates in the human population (U_h). *Vector preference p_{ITN} of 0.6 indicates that 60% of malaria vectors choose an ITN-protected human if presented against an unprotected human. The reference inert ITN (panel A) induces a vector preference of 0.5 (vectors are equally likely to select an ITN-protected or unprotected human) and the reference deterrent ITN (panel B) induces a vector preference of 0.36 (only 36% of the vectors choose an ITN-protected human against an unprotected human).*

Effect of physiological resistance

Physiological resistance was defined as the vector's ability to survive a feeding attempt in a dwelling equipped with an ITN. In our model, the probability of survival during a feeding attempt corresponds to the likelihood of successfully taking a blood meal, conditioned on not being diverted (Figure 1). Our simulations indicate that, regardless of the level of physiological resistance in the vector population, attractive ITNs consistently outperform inert and deterrent ITNs in reducing malaria transmission (Figure 3), although the reduction in transmission declines as resistance levels increase. In the absence of physiological resistance (i.e., 0% survival), and with all other parameters held constant (including ITN use rate at 50%; Table 1), moderately attractive ITNs (p_{ITN} = 0.6) and highly attractive ITNs (p_{ITN} = 0.7) are predicted to reduce malaria transmission by 32% and 55%, respectively, relative to an inert (p_{ITN} = 0.5) ITN (Figure 3A). When compared to a deterrent ITN (p_{ITN} = 0.36) under the same conditions (no resistance), an inert ITN is expected to reduce malaria transmission by 41%, while moderately attractive (p_{ITN} = 0.6) and highly attractive ITNs (p_{ITN} = 0.7) are expected to reduce transmission by 60% and 73%, respectively (Figure 3B). Under high resistance conditions (82% survival i.e. the highest level observed in field data [12]; supplementary Figure 2B), both moderately and highly attractive ITNs still provide better community protection but yield moderate reductions in transmission potential: of 13% and 25%, respectively, relative to inert ITNs (Figure 3A) and of 29% and 38%, respectively, relative to deterrent ITNs (Figure 3B). Compared to no nets (0% use), attractive LLINs (50% use) are

predicted to achieve a 74% reduction in transmission potential under high resistance conditions
(82% survival), which is 17 percentage points (p.p.) more than deterrent ITNs (supplementary
Figure 3B).

Figure 3 - Reduction in *Plasmodium* transmission potential ($RedTP$) induced (A) by
attractive Insecticide-treated Nets (ITNs) compared to inert ones and (B) by
attractive and inert ITNs compared to deterrent ITNs for varied levels of
physiological resistance in the vector population (feedbing attempt survival, $1 -$
μ_{p1}). *Vector preference* p_{ITN} *of 0.6 indicates that 60% of malaria vectors choose an*
ITN-protected human if presented against an unprotected human. The reference
inert ITN (panel A) induces a vector preference of 0.5 (vectors are equally likely to
select an ITN-protected or unprotected human) and the reference deterrent ITN
(panel B) induces a vector preference of 0.36 (only 36% of the vectors choose an
ITN-protected human against an unprotected human). Lines become grey in the
range [82; 100%], corresponding to survival levels that were never observed in [12]
(supplementary Figure 2).

Effect of quantitative behavioral resistance (escape behavior)

We then analyzed the impact of quantitative behavioral resistance on the efficacy of attractive
ITNs. Specifically, we simulated increasing levels of indoor escape behavior by varying the
diversion probability (D_p) in our model (Figure 1). A diverted mosquito postpones host-seeking until
the following night. The simulations demonstrated that, regardless of the level of indoor escape,
attractive ITNs consistently outperform inert and deterrent ITNs in reducing malaria transmission.
However, as indoor escaping increases, the relative efficacy of attractive ITNs decreases (Figure
4). When compared to an inert ITN (Figure 4A), a moderately attractive ITN is expected to reduce
malaria transmission by 21% (under high resistance, i.e. 51% diversion that is the highest level of
diversion observed in field data [12]; supplementary Figure 2B) to 30% (under no resistance, 0%
diversion), while a highly attractive ITN is expected to reduce transmission by 39 (high resistance,
51% diversion) to 51% (no resistance, 0% diversion). When compared to a deterrent ITNs (Figure
4B), malaria transmission reduction ranged from 28 to 39% for inert ITN, from 43 to 57% for
moderately attractive ITNs ($p_{ITN} = 0.6$) and from 56 to 70% for highly attractive ITNs ($p_{ITN} = 0.7$),
depending on the level of resistance (i.e. between 0 and 51% diversion). Compared to no nets (0%
use), attractive LLINs (50% use) are predicted to achieve a 88% reduction in transmission potential

under high resistance conditions (51% diversion), which is 15 percentage points (p.p.) more than deterrent ITNs (supplementary Figure 3C).

Figure 4 - Reduction in *Plasmodium* transmission potential ($RedTP$) induced (A) by attractive Insecticide-treated Nets (ITNs) when compared to inert ones and (B) by attractive and inert ITNs when compared to deterrent ITNs for varied levels of quantitative behavioral resistance (escape behavior expressed as percent diversion, D_p) in the vector population. *Vector preference p_{ITN} of 0.6 indicates that 60% of malaria vectors choose an ITN-protected human if presented against an unprotected human. The reference inert ITN (panel A) induces a vector preference of 0.5 (vectors are equally likely to select an ITN-protected or unprotected human) and the reference deterrent ITN (panel B) induces a vector preference of 0.36 (only 36% of the vectors choose an ITN-protected human against an unprotected human). Lines become grey in the range [51; 100%], corresponding to diversion levels that were never observed in* [12] *(supplementary Figure 2).*

Effect of qualitative behavioral resistance (spatiotemporal avoidance)

Mosquito vectors of malaria can alter their behavior to seek blood meals at different times or locations (i.e. outdoors), reducing their exposure to ITNs. We simulated this spatiotemporal avoidance of ITNs by varying the parameter Π, which represents the proportion of normal exposure to mosquito bites upon humans lacking ITNs, which occurs indoors at times when nets would normally be in use. In other words, Π quantifies the fraction of total bites that ITNs effectively prevent. As spatiotemporal avoidance increases, Π decreases. Regardless of the level of spatiotemporal avoidance, attractive ITNs consistently reduce malaria transmission compared to both inert and deterrent ITNs (Figure 5). However, as spatial-temporal avoidance increases, the relative efficacy of attractive ITNs decreases. At 62% spatial-temporal avoidance (i.e. the highest value reported in literature [31]), attractive ITNs are expected to reduce malaria transmission by 17 and 31% compared to inert ITNs (Figure 5A), for moderately and highly attractive ITNs, respectively, and by 35 and 46% compared to deterrent ITNs (Figure 5B). Under the same conditions (62% spatiotemporal avoidance), an inert ITN is expected to reduce malaria transmission by 22% when compared to a deterrent ITN (Figure 5B). Compared to no nets (0% use), attractive LLINs (50% use) are predicted to achieve a 90% reduction in transmission potential under high resistance conditions (62% spatiotemporal avoidance), which is 14 percentage points (p.p.) more than deterrent ITNs (supplementary Figure 3D).

Figure 5 - Reduction in *Plasmodium* transmission potential ($RedTP$) induced (A) by attractive Insecticide-treated Nets (ITNs) when compared to inert ones and (B) by attractive and inert ITNs when compared to deterrent ITNs for varied levels of qualitative behavioral resistance (spatiotemporal avoidance expressed as 1 minus the efficacy of ITNs in reducing the number of bites, $1 - \Pi$) in the vector population. *Vector preference p_{ITN} of 0.6 indicates that 60% of malaria vectors choose an ITN-protected human if presented against an unprotected human. The reference inert ITN (panel A) induces a vector preference of 0.5 (vectors are equally likely to select an ITN-protected or unprotected human) and the reference deterrent ITN (panel B) induces a vector preference of 0.36 (only 36% of the vectors choose an ITN-protected human against an unprotected human). Lines become grey in the range [62; 100%], corresponding to spatiotemporal avoidance levels that were never observed in the field (see refs for parameter Π in Table 1).*

Uncertainty and sensitivity analyses

The uncertainty analysis revealed that, when all parameters varied within their plausible ranges, the mean reduction in transmission potential with attractive ITNs was estimated at 39.3% (95% Credible Interval: [11.9%, 71.4%]) relative to deterrent ITNs. The sensitivity analysis (Figure 6) identified the parameters with the greatest influence on model outcomes among those not expressly analyzed (Figure 6). Notably, we found that the reduction in transmission potential was greater for *Anopheles* populations with a higher baseline daily survival probability (S; Partial Rank Correlation Coefficients PRCC= -0.54 [95% CI: -0.63, -0.48], Partial Slope Coefficient PSC= -40.0 [95% CI: -44.7, -35.6]), shorter gonotrophic cycle duration (g in days; PRCC= 0.67 [0.32, 0.72], PSC= 5.76 [5.41, 6.19]), higher post-bite mortality probability in rooms with ITNs (μ_{2p}; PRCC= -0.55 [-0.63, -0.48], PSC= -15.6 [-17.7, -13.2]), and when parasites exhibited longer extrinsic incubation period duration (n in days; PRCC = -0.48 [-0.55, -0.41], PSC = -1.68 [: -1.91, -1.41]).

Fig 6: Results of a sensitivity analysis using Latin Hypercube Sampling, showing (A) Partial Rank Coefficients of Correlation (PRCC) of parameters and (B) simulation plots illustrating the impact on transmission potential of varying parameter values when comparing a moderately attractive ITN (p_{ITN} = 0.6) to a deterrent ITN (p_{ITN} = 0.36). *Intervals in panel A are 95% Confidence Interval of PRCC. In panel B, percent change in transmission potential (RedTP) is plotted against values for each model parameter (x-axis). Blue curves on panel B are loess regression. See Table 1 for parameters' definitions.*

Discussion

Recent evidence from both field and laboratory studies suggests that some ITNs may increase the attraction of *An. gambiae* mosquitoes to human hosts, potentially raising the bite exposure of ITN users [12,32]. However, the implications of such attractive ITNs on malaria transmission at the community level remained unclear. To address this, we developed a model to assess their impact under various levels of ITN usage and vector resistance, including both physiological and behavioral resistance.

Our model predicts that attractive ITNs consistently reduce malaria transmission across all simulated scenarios compared to inert or deterrent ITNs, even when lower ITN usage rates and

the presence of physiological or behavioral resistance limits the magnitude of the effect. Sensitivity analysis identified four additional key parameters that significantly influence the relative efficacy of attractive ITNs: daily survival probability (S), gonotrophic cycle duration (g), the extrinsic incubation period (n), and post-bite mortality in the room of an ITN-protected individual (μ_{2p}). The general pattern is that any factor reducing the average infectious life expectancy of mosquitoes, whether through increased mortality or more frequent encounters with ITNs, tends to enhance the relative efficacy of attractive ITNs. The corollary is that shorter extrinsic incubation periods reduce their relative efficacy, since parasites can reach transmissibility within a shorter mosquito lifespan.

In all cases, attractive ITNs performed better than inert ITNs, which in turn outperformed deterrent ITNs in reducing malaria transmission. Currently, most widely distributed ITNs have deterrent properties [12,26]. Our findings suggest that shifting from deterrent to inert (or attractive) ITNs could substantially enhance malaria control efforts, as also suggested by Killeen *et al.* [11]. We therefore urge national and international organizations responsible for ITN evaluation, selection, and distribution, in collaboration with manufacturers, to reassess whether deterrence should remain a desirable characteristic of ITNs.

Increasing reports of attractive effects in ITNs [12] should not be seen as a drawback but rather as an opportunity for innovation in malaria control. Our findings suggest that attractive ITNs could significantly enhance malaria prevention, opening new avenues for product development. Recent advances in mosquito chemical ecology have identified numerous host-derived attractants [33], while ITN manufacturers have developed techniques to incorporate dual compounds, such as insecticides and synergists [34]. Leveraging these innovations, next-generation ITNs could be designed to integrate both insecticidal and attractive properties, thereby optimizing vector control strategies. Such innovations could enhance the effectiveness of ITNs while maintaining their widespread acceptance and use in malaria-endemic regions. One key advantage of combining insecticides and attractants into ITNs is their potential to remain effective even in the absence of humans in the room or when old ITNs are repurposed for alternative uses [35]. This approach could extend the operational lifespan and functional utility of ITNs, further strengthening their role in malaria control. Nevertheless, it is important to recognize the potential adverse effects of attractive ITNs. By increasing host attractiveness, such nets could expose users to higher biting risk in situations of incomplete coverage or physical deterioration, which may reduce user satisfaction and compliance. These challenges underscore that, alongside technical optimization, the success of attractive ITNs would also depend on strong community engagement and communication strategies that emphasize their communal benefits while mitigating concerns about personal protection and nuisance biting [11,12].

While our model provides valuable insights into the potential impact of attractive ITNs on malaria transmission, it is important to acknowledge its limitations. The model relies on simplified assumptions about mosquitoes, parasites, and humans, which may limit its applicability to all real-world settings. In particular, we assume binary host choice, with mosquitoes selecting between ITN-protected and unprotected humans, without accounting for alternative hosts such as livestock. In areas where alternative hosts are abundant, mosquito feeding patterns could shift [36,37]. Our model also assumes spatio-temporal homogeneity, meaning that human and mosquito populations are evenly mixed in space and that model parameters remain constant over time (i.e., the model is static). This simplification neglects potential spatial heterogeneity, such as clustering of hosts, as well as temporal variability driven by seasonality, intervention effects, or evolutionary processes. Important model parameters such as daily survival probability, gonotrophic cycle duration, and the extrinsic incubation period are strongly driven by environmental factors, either directly (e.g., temperature and humidity [28,38,39]) or indirectly through their dependence on favorable ecological conditions determining the spatio-temporal distribution of the mosquito. It is therefore reasonable to expect the relative efficacy of attractive ITNs to vary across ecological settings.In variable real-world settings, such heterogeneities can influence both mosquito–human interactions and malaria transmission dynamics [8,40–42]. Another assumption is linked to the deterministic nature of our model, which does not account for stochastic variability in input parameters. Stochastic effects may be particularly relevant in small populations or low-

transmission settings, where random events can substantially influence outcomes. Relaxing these assumptions could alter quantitative predictions, potentially amplifying or dampening the effects of attractive ITNs in specific settings. However, our uncertainty analysis, varying all parameters across plausible ranges, indicates that attractive ITNs consistently reduce transmission potential more than deterrent ITNs under all simulations. This suggests that the relative benefit of attractive ITNs is likely robust, even if the absolute magnitude of their impact could vary in more complex scenarios. Incorporating these complexities into future modeling efforts could refine predictions and improve the relevance of our findings for diverse transmission settings.

As with any model that has not been validated using field data, these results should be interpreted with caution and considered to have theoretical value only. Validation of our model predictions would ideally require cluster-randomized controlled trials using ITNs whose attractiveness, inertness, or deterrence has been assessed beforehand in the vector population to ensure the interventions accurately reflect their intended behavioral properties.

Conclusion

Our findings predict that attractive ITNs could significantly reduce malaria transmission at the community level, even in the presence of physiological or behavioral resistance in vectors, relative to inert and deterrent ITNs. This challenges the conventional preference for deterrent ITNs, which prioritize personal protection, and expands the possibilities for designing next-generation ITNs. By combining insecticidal and attractant properties, innovative ITNs could enhance malaria control by remaining effective despite high resistance levels and repurposing. Advances in mosquito chemical ecology and ITN manufacturing present a unique opportunity to develop more efficient tools for malaria prevention. Future research should focus on better understanding the mechanisms underlying the attractant effect of some current ITNs, validating the theoretical predictions through field trials, and fostering collaborations with manufacturers to translate these insights into practical ITN prototypes.

Funding

The authors declare that they have received no specific funding for this study.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. The authors are recommenders for PCI Zoology. CP is recommender for PCI Infection.

Author contributions

Conceptualization: NM, CP; Methodology: NM; Software: NM; Formal analysis: NM; Investigation: NM, CP; Data curation: NM; Writing – original draft: NM; Writing – review and editing: NM, CP. Visualization: NM.

Data, scripts and code availability

Scripts, code and data are available online: https://doi.org/10.5281/zenodo.15121768 [43].

Supplementary information

Supplementary information is available online: https://doi.org/10.5281/zenodo.15121768 [43].

References

1.   World malaria report 2023. Geneva: World Health Organization; 2023. Available: https://iris.who.int/handle/10665/374472 [accessed 22 Feb 2024]

2.   Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature. 2015. doi:10.1038/nature15535

3.   Alonso P, Noor AM. The global fight against malaria is at crossroads. Lancet. 2017;390: 2532–2534. doi:10.1016/S0140-6736(17)33080-5

4.   Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, Corbel V. Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control? Trends Parasitol. 2011;27: 91–8. doi:10.1016/j.pt.2010.08.004

5.   Donnelly MJ, Corbel V, Weetman D, Wilding CS, Williamson MS, Black WC. Does kdr genotype predict insecticide-resistance phenotype in mosquitoes? Trends Parasitol. 2009;25: 213–219. doi:10.1016/j.pt.2009.02.007

6.   Carrasco D, Lefèvre T, Moiroux N, Pennetier C, Chandre F, Cohuet A. Behavioural adaptations of mosquito vectors to insecticide control. Curr Opin Insect Sci. 2019;34: 48–54. doi:10.1016/j.cois.2019.03.005

7.   Moiroux N, Gomez MB, Pennetier C, Elanga E, Djenontin A, Chandre F, et al. Changes in Anopheles funestus Biting Behavior Following Universal Coverage of Long-Lasting Insecticidal Nets in Benin. J Infect Dis. 2012;206: 1622–9. doi:10.1093/infdis/jis565

8.   Taconet P, Soma DD, Zogo B, Mouline K, Simard F, Koffi AA, et al. Physiological and behavioural resistance of malaria vectors in rural West-Africa: a data mining study to address their fine-scale spatiotemporal heterogeneity, drivers, and predictability. Peer Community Journal. 2024;4. doi:10.24072/pcjournal.367

9.   Chareonviriyaphap T, Roberts DR, Andre RG, Harlan HJ, Manguin S, Bangs MJ. Pesticide avoidance behavior in Anopheles albimanus, a malaria vector in the Americas. J Am Mosq Control Assoc. 1997;13: 171–183.

10. World Health Organization, WHO Pesticide Evaluation Scheme. Guidelines for laboratory and field-testing of long-lasting insecticidal nets. World Health Organization; 2013 p. 89. Report No.: WHO/HTM/NTD/WHOPES/2013.3. Available: https://iris.who.int/handle/10665/80270 [accessed 17 Sept 2025]

11. Killeen GF, Chitnis N, Moore SJ, Okumu FO. Target product profile choices for intra-domiciliary malaria vector control pesticide products: repel or kill? Malar J. 2011;10: 207. doi:10.1186/1475-2875-10-207

12. Moiroux N, Chandre F, Hougard J-M, Corbel V, Pennetier C. Remote Effect of Insecticide-Treated Nets and the Personal Protection against Malaria Mosquito Bites. PLOS ONE. 2017;12: e0170732. doi:10.1371/journal.pone.0170732

13. Glunt KD, Coetzee M, Huijben S, Koffi AA, Lynch PA, N'Guessan R, et al. Empirical and theoretical investigation into the potential impacts of insecticide resistance on the effectiveness of insecticide-treated bed nets. Evolutionary Applications. 2018;11: 431–441. doi:10.1111/eva.12574

14. Lines JD, Curtis CF, Myamba J, Njau R. Tests of repellent or insecticide impregnated curtains, bednets and anklets against malaria vectors in Tanzania. Geneva: World Health

Organization; 1985. Report No.: WHO/WBC/85.920. Available:
https://iris.who.int/handle/10665/59327

15.   Majori G, Sabatinelli G, Coluzzi M. Efficacy of permethrin-impregnated curtains for malaria
vector control. Medical and Veterinary Entomology. 1987;1: 185–192. doi:10.1111/j.1365-
2915.1987.tb00341.x

16.   Silver JB. Methods of Age-grading Adults and Estimation of Adult Survival Rates. Mosquito
Ecology. Dordrecht: Springer Netherlands; 2008. pp. 1161–1271. Available:
http://dx.doi.org/10.1007/978-1-4020-6666-5_13

17.   Afrane YA, Lawson BW, Githeko AK, Yan G. Effects of Microclimatic Changes Caused by
Land Use and Land Cover on Duration of Gonotrophic Cycles of Anopheles gambiae
(Diptera: Culicidae) in Western Kenya Highlands. Journal of Medical Entomology. 2005;42:
974–980. doi:10.1093/jmedent/42.6.974

18.   Carnevale P, Robert V, Manguin S, Corbel V, Fontenille D, Garros C, et al. Les anophèles :
biologie, transmission du Plasmodium et lutte antivectorielle. Marseille: IRD; 2009. Available:
http://www.documentation.ird.fr/hor/fdi:010047862 [accessed 15 Dec 2024]

19.   Detinova TS. Age-grouping methods in Diptera of medical importance with special reference
to some vectors of malaria. World Health Organization monograph series. 1962;47: 13–191.

20.   Churcher TS, Trape J-F, Cohuet A. Human-to-mosquito transmission efficiency increases as
malaria is controlled. Nat Commun. 2015;6: 6054. doi:10.1038/ncomms7054

21.   Russell TL, Govella NJ, Azizi S, Drakeley CJ, Kachur SP, Killeen GF. Increased proportions
of outdoor feeding among residual malaria vector populations following increased use of
insecticide-treated nets in rural Tanzania. Malar J. 2011;10: 80. doi:10.1186/1475-2875-10-
80

22.   Moiroux N, Damien GB, Egrot M, Djenontin A, Chandre F, Corbel V, et al. Human exposure
to early morning Anopheles funestus biting behavior and personal protection provided by
long-lasting insecticidal nets. PLoS ONE. 2014;9: e104967.
doi:10.1371/journal.pone.0104967

23.   Huho B, Briet O, Seyoum A, Sikaala C, Bayoh N, Gimnig J, et al. Consistently high estimates
for the proportion of human exposure to malaria vector populations occurring indoors in rural
Africa. Int J Epidemiol. 2013;42: 235–47. doi:10.1093/ije/dys214

24.   Cooke MK, Kahindi SC, Oriango RM, Owaga C, Ayoma E, Mabuka D, et al. 'A bite before
bed': exposure to malaria vectors outside the times of net use in the highlands of western
Kenya. Malaria Journal. 2015;14: 259. doi:10.1186/s12936-015-0766-4

25.   Malaria Atlas Project. In: MAP [Internet]. Available: https://malariaatlas.org/ [accessed 18
Dec 2024]

26.   Strode C, Donegan S, Garner P, Enayati AA, Hemingway J. The Impact of Pyrethroid
Resistance on the Efficacy of Insecticide-Treated Bed Nets against African Anopheline
Mosquitoes: Systematic Review and Meta-Analysis. PLoS Med. 2014;11: e1001619.
doi:10.1371/journal.pmed.1001619

27.   Hien DF d S, Dabiré KR, Roche B, Diabaté A, Yerbanga RS, Cohuet A, et al. Plant-Mediated
Effects on Mosquito Capacity to Transmit Human Malaria. PLOS Pathogens. 2016;12:
e1005773. doi:10.1371/journal.ppat.1005773

28.   Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, et al. Rethinking the extrinsic
incubation period of malaria parasites. Parasit Vectors. 2018;11: 178. doi:10.1186/s13071-
018-2761-4

29.   R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2022. Available: https://www.R-project.org/

30.   Chalom A, de Prado PIKL. pse: Parameter space exploration with latin hypercubes. 2017.
Available: https://cran.r-project.org/src/contrib/Archive/pse/ [accessed 16 Sept 2025]

31.   Geissbühler Y, Chaki P, Emidi B, Govella NJ, Shirima R, Mayagaya V, et al. Interdependence
of domestic malaria prevention measures and mosquito-human interactions in urban Dar es
Salaam, Tanzania. Malar J. 2007;6: 126. doi:10.1186/1475-2875-6-126

32.   Porciani A, Diop M, Moiroux N, Kadoke-Lambi T, Cohuet A, Chandre F, et al. Influence of
pyrethroïd-treated bed net on host seeking behavior of Anopheles gambiae s.s. carrying the
kdr allele. PLoS One. 2017;12. doi:10.1371/journal.pone.0164518

33.   Dormont L, Mulatier M, Carrasco D, Cohuet A. Mosquito Attractants. J Chem Ecol. 2021;47:
351–393. doi:10.1007/s10886-021-01261-2

34.   Vector Control Product List. In: WHO - Prequalification of Medical Products (IVDs, Medicines,
Vaccines and Immunization Devices, Vector Control) [Internet]. Available:
https://extranet.who.int/prequal/vector-control-products/prequalified-product-list   [accessed
31 Dec 2024]

35.   Hien AS, Hien H, Bacyè FY, Badolo H, Tiono A, Diallo CO, et al. What happens to old
insecticide-treated nets after households use in Burkina Faso? Malaria Journal. 2024;23:
350. doi:10.1186/s12936-024-05181-w

36.   Lefevre T, Gouagna L-C, Dabire KR, Elguero E, Fontenille D, Renaud F, et al. Beyond nature
and nurture: phenotypic plasticity in blood-feeding behavior of Anopheles gambiae s.s. when
humans are not readily accessible. American Journal of Tropical Medicine and Hygiene.
2009;81: 1023–1029. doi:10.4269/ajtmh.2009.09-0124

37.   Stone C, Gross K. Evolution of host preference in anthropophilic mosquitoes. Malaria
Journal. 2018;17: 257. doi:10.1186/s12936-018-2407-1

38.   Lyons CL, Coetzee M, Terblanche JS, Chown SL. Desiccation tolerance as a function of age,
sex, humidity and temperature in adults of the African malaria vectors Anopheles arabiensis
and Anopheles funestus. J Exp Biol. 2014;217: 3823–3833. doi:10.1242/jeb.104638

39.   Lardeux FJ, Tejerina RH, Quispe V, Chavez TK. A physiological time analysis of the duration
of the gonotrophic cycle of Anopheles pseudopunctipennis and its implications for malaria
transmission in Bolivia. Malaria Journal. 2008;7: 141. doi:10.1186/1475-2875-7-141

40.   Bousema T, Griffin JT, Sauerwein RW, Smith DL, Churcher TS, Takken W, et al. Hitting
Hotspots: Spatial Targeting of Malaria for Control and Elimination. PLOS Medicine. 2012;9:
e1001165. doi:10.1371/journal.pmed.1001165

41.   Kingsolver JG. Mosquito Host Choice and the Epidemiology of Malaria. The American
Naturalist. 1987;130: 811–827. doi:10.1086/284749

42.   Bationo CS, Gaudart J, Dieng S, Cissoko M, Taconet P, Ouedraogo B, et al. Spatio-temporal
analysis and prediction of malaria cases using remote sensing meteorological data in
Diébougou health district, Burkina Faso, 2016-2017. Sci Rep. 2021;11: 20027.
doi:10.1038/s41598-021-99457-9

43.   Moiroux N. Supplementary materials, as well as codes and data necessary to reproduce the
analyses presented in the study entitled "Potential of attractive insecticide-treated nets (ITNs)
in reducing malaria transmission: a modeling study." Zenodo; 2025.
doi:10.5281/zenodo.17142100