Leveraging questionnaire-based physical activity levels (PAL) to identify energy intake misreporting in the Goldberg method: A doubly-labeled water validation study

Heather K Neilson[1], Shervin Asgari[1], Karen A. Kopciuk[1,2,3], Janet A. Tooze[4], Farah Khandwala[1], Anita Koushik[5], Rémi Rabasa-Lhoret[6], Ilona Csizmadi[2*]

Author Affiliations:

[1]Department of Cancer Epidemiology and Prevention Research, Alberta Health Services – Cancer Care Alberta, Calgary, Alberta, Canada (HKN, SA, KAK, FK)

[2]Departments of Community Health Sciences and Oncology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada (KAK, IC)

[3]Department of Mathematics and Statistics, University of Calgary, Calgary, Alberta, Canada (KAK)

[4]Department of Biostatistics and Data Science, Wake Forest University School of Medicine, Wake Forest University, Winston-Salem, NC, USA (JAT)

[5]Department of Oncology, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec, Canada (AK)

[6]Departments of Medicine and Nutrition, Université de Montreal, and Montreal Clinical Research Institute, Montreal, Quebec, Canada (RRL)

Authors' Last Names: Neilson, Asgari, Kopciuk, Tooze, Khandwala, Koushik, Rabasa-Lhoret, Csizmadi

Abstract: 299 words

Manuscript: 4,098 words

ORCiD:

Ilona Csizmadi: 0000-0002-4895-1310

Heather K. Neilson: 0000-0002-6803-2800

Karen A. Kopciuk: 0000-0001-7499-1861

Anita Koushik: 0000-0001-5304-7660

Janet A. Tooze: 0000-0002-5160-0407

Rémi Rabasa-Lhoret: 0000-0003-4706-5170

Sources of Support: Funding for this study was provided to institutions by the Canadian Institutes of Health Research (Grant MOP-86632) and the Alberta Cancer Research Institute (Grant 24265).

Role of funding agencies: The funders had no role in the design of this study, the analysis, interpretation of results or drafting of the manuscript.

*Corresponding Author:

Dr Ilona Csizmadi, Departments of Community Health Sciences and Oncology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

Email: icsizmad@ucalgary.ca

Abbreviations:

BMR_{EST}: estimated basal metabolic rate; **C-DHQI**: Canadian Diet History Questionnaire I; **DLW**: doubly-labeled water; **EI**: energy intake; **NPV**: negative predictive values; **PAL**: physical activity level; **PPV**: positive predictive value; **rEI**: self-reported energy intake; **STAR-Q**: Sedentary Time and Activity Reporting Questionnaire; **TEE**: total energy expenditure; **TBW**: total body water

ABSTRACT

BACKGROUND: Dietary intake misreporting is widely acknowledged in nutritional

epidemiology. The Goldberg method has been proposed as an option for identifying under-

reporters of energy intake (EI) in large-scale epidemiologic studies. However, its implementation

remains limited by challenges associated with estimating physical activity levels (PAL), a critical

component.

OBJECTIVE: To quantify the accuracy of the Goldberg method to classify EI misreporting using

Sedentary Time and Activity Reporting Questionnaire (STAR-Q)-derived PAL_{STAR-Q} compared

with doubly-labeled water (DLW)-derived total energy expenditure (TEE_{DLW}).

DESIGN: Between 2009 and 2011, 99 weight-stable men and women (mean [SD]: 48 [8] years)

completed a two-week DLW protocol, the Canadian Diet History Questionnaire I, and the STAR-

Q, a comprehensive past-month activity questionnaire. TEE_{DLW} was the criterion measure of EI to

determine the sensitivity, specificity, positive predictive value (PPV), negative predictive value

(NPV) and accuracy (proportion correctly classified as under-reporters or acceptable reporters) of

Goldberg cut-points based on the PAL_{STAR-Q}. A global PAL of 1.55 frequently used in the original

Goldberg method was also studied.

RESULTS: The Goldberg method using PAL_{STAR-Q} classified 58% of men and women as under-

reporters, compared with 60% of men and 56% of women identified with TEE_{DLW}. Among men,

values for sensitivity, specificity, PPV, NPV and accuracy of the Goldberg method and PAL_{STAR-}

$_Q$ were estimated as: 88%, 87%, 91%, 81%, and 87%, respectively. Among women, these values

were: 79%, 69%, 77%, 72%, and 75%, respectively. Substituting a PAL of 1.55 for the PAL_{STAR-}

$_Q$ reduced the proportion of men and women classified as under-reporters to 35% and 19%,

respectively, lowered sensitivity to 54% and 33%, and increased specificity to 93% and 100%.

CONCLUSIONS: Population-specific validation sub-studies using the Goldberg method and

questionnaire-derived PALs can be informative for understanding EI misreporting to ultimately

improve estimates of diet-disease associations in large-scale epidemiologic studies lacking in

objective EI measures.

Keywords: energy intake, accuracy, misreporting, doubly-labeled water, Goldberg method,

physical activity level, adults, under-reporting, 24-hour movement paradigm

INTRODUCTION

Under-reporting of energy intake (EI) on dietary questonnaires is a widely recognized problem that distorts estimates of dietary inadequacies in nutritional surveys and potentially biases estimates of diet-disease associations in etiologic studies impeding progress in nutritional epidemiology (1). Statistical approaches that mitigate the effects of dietary intake misreporting have been suggested, but have not been routinely implemented, indicating a need for further study (2). Prerequisites for undertaking measurement error correction include having knowledge of *true* dietary intake and an understanding of the underlying systematic errors associated with self-report (3). Ideally, misreporting could be detected using 'gold standard' objective biomarkers of dietary intake; however, these methods are limited in number and often too labor-intensive and costly to implement in large-scale epidemiologic studies (4). A practical alternative approach specifically for EI, may be to classify individuals according to the plausibility of their reported EI (rEI), compared with a method validated to estimate energy requirements (5).

The Goldberg method (6) was originally proposed as a strategy for identifying EI misreporting in community-dwelling populations and is based on the principle that under long-term, weight-stable conditions, EI is approximately equal to total energy expenditure (TEE). In addition, the method leverages the well-established relations between EI, TEE, basal metabolic rate (BMR) and physical activity level (PAL), where EI/BMR approximates TEE/BMR and TEE/BMR approximates PAL (7). Further, the method proposes estimating cut-point boundaries, or 95% confidence intervals (95% CI), for the ratio EI:TEE which accounts for the variability of PAL and additional energy-related factors. From this we can infer that in weight-stable individuals, a habitual rEI below or above these boundaries represent under and over-reported EI.

While conceptually tenable, the method is logistically challenged by the difficulty associated with the estimation of PAL in free-living populations (8). As such, in the absence of population-specific objective measures a low activity PAL of 1.55 has frequently been assumed based on World Health Organization predictions of energy requirements (9) and the original Goldberg paper (10). Indeed, in a study of middle-aged men and women Tooze *et al*. (11) reported that the Goldberg method and a PAL of 1.55 had acceptable accuracy compared with doubly-labelled water (DLW), the 'gold standard' for estimating TEE (12, 13), although alternative PALs were not considered. In contrast, Black *et al*. (14) demonstrated that the proportion of misclassified men and women could be reduced by assigning low, medium, or high group-level PALs (e.g., 1.55, 1.75 or 1.95, respectively), based on DLW results. More recently, a large survey of Australian adults Grech et al (15) examined the associations between EI underreporting, diet quality and socioeconomic status using the Goldberg method with a PAL of 1.55 versus prediction equations that estimated individual TEE levels and reported comparable results. Nonetheless, based on historical DLW studies (16) and the recent Dietary Reference Intake (DRIs) for Energy report (median adult PAL = 1.68) (7), a global PAL of 1.55 may be unrealistically low for substantial segments of the population leading to active EI under-reporters misclassified as acceptable.

To our knowledge, the Goldberg method has not been evaluated using PALs derived from individual self-reports of physical activity despite their appeal given the potential to capture habitual behaviors of greatest relevance in chronic disease etiologic research. With that in mind, we appraised the accuracy of the Goldberg method for classifying EI reporting status using individual-level PALs derived from the Sedentary Time and Activity Reporting Questionnaire (STAR-Q), a comprehensive global activity questionnaire that we previously validated against DLW (17). In our primary analysis, we determined the accuracy of the Goldberg method with

76   STAR-Q-derived PALs compared with DLW-derived TEE_{DLW} to classify EI reporting status on a

77   food frequency questionnaire (FFQ). In secondary analyses we assumed the classic Goldberg PAL

78   of 1.55. In addition, in sensitivity analyses we studied the effect of modifying the FFQ variability

79   assumption in the estimation of Goldberg cut-points as previously studied (11).

81   **METHODS**

82   Participants in this analysis were those enrolled in the original STAR-Q DLW validation

83   study (17). The STAR-Q was conceptualized to capture 24-hour movement behaviours (18). It was

84   therefore uniquely designed to ascertain self-reported time spent in sleep, sedentary behavior, and

85   all types and intensities of physical activity (e.g., light, moderate and vigorous) within all domains

86   (e.g., occupation, leisure, household, and transportation) (17, 19).

88   **Subject Eligibility and Ethics**

89   Participant recruitment has previously been described (17). Briefly, urban residents living

in Calgary, Alberta, Canada were recruited from the community or from the Alberta Tomorrow

Project (ATP). The ATP is a large geographically-dispersed cohort established in 2001 in Alberta,

to study links between lifestyle behaviors and chronic disease risk (20, 21). ATP participants who

met eligibility criteria for the DLW validation study – body mass index (BMI) ≤ 35, weight stabile

(± 2.5 kg for at least 3 months), not planning to gain or lose weight, not pregnant or breastfeeding,

not diagnosed with metabolic disorders, and not taking medications that modify water balance –

were actively recruited by mail. Non-ATP participants were recruited by community-based

advertisements. The upper age limit for eligibility was 60 years; the lower age limit for the urban

Calgary sample was 30 years, and for ATP participants it was 35 years.

All participants provided written informed consent. Ethical approval was obtained from

the Alberta Cancer Research Ethics Committee of Alberta Health Services and the Conjoint Health

Research Ethics Board of the University of Calgary. All study procedures were in accordance with

the ethical standards outlined by the two governing bodies.

Doubly-labeled water study

Details of the 14-day DLW protocol have been published (17). Briefly, at baseline (Day

0), following an overnight fast, participants provided urine and saliva samples to determine

background isotope levels. Each subject was then orally dosed with 2.5g 10 atom % of ^{18}O per kg

total body water (TBW) and 0.18 g 99 atom % deuterium (^{2}H) per kg estimated TBW (Rotem Inc.,

Topsfield, Massachusetts). We collected post-dose saliva samples at 3 and 4 hours to measure

TBW from deuterium-isotope dilution, and second-void urine samples on Days 1, 8, and 14. On

Day 14 after an overnight fast, participants received 0.18 g 99 atom % ^{2}H per kg TBW. The post-

dose saliva sample collection protocol was followed as on Day 0. To measure the decline in

isotope enrichment, we batch analyzed samples in duplicate using the Isoprime Stable Isotope

Ratio Mass Spectrometer and the Multiflow-Bio module for Isoprime equipped with a Gilson

222XL Autosampler (GV Instruments, Manchester, United Kingdom). We performed data

processing using IonVantage (2012) software for Isoprime (Cheadle, United Kingdom).

We calculated TEE_{DLW} in kilocalories per day using a modified Weir equation (22) and an

assumed respiratory quotient of 0.85. We estimated BMR using the Schofield Equation (23) and

assumed 10% for the thermic effect of food (12), to derive individual level PALs from DLW as

follows:

$PAL_{DLW} = 0.9 TEE_{DLW} / BMR_{EST}$

Health and lifestyle measurements

Height and weight were measured (to the nearest 0.1 cm and 0.1 kg, respectively) on Days

0 and 14 of the DLW protocol using standard procedures (24). The TBF-310 Tanita Body

Composition Analyzer and Scale (TheCompetitiveEdge.com, Preston, Washington) was used to

measure body weight. BMI was determined as body weight divided by height-squared (kg/m^2).

The past-year Canadian Diet History Questionnaire I (C-DHQ I) was administered once,

on Day 0 of the DLW protocol to estimate reported rEI. The C-DHQ I is a comprehensive FFQ

that assesses usual dietary intake over the previous 12 months, modified for Canadian populations

from the U.S. National Cancer Institute's DHQ (25, 26). Several large epidemiologic studies use

the C-DHQ I in Canada including the ATP (20, 21, 27, 28).

The STAR-Q was completed on Day 14 of the DLW study to estimate energy expenditure

from all domains, types and intensities of past-month physical activity, sedentary behavior, and

sleep. STAR-Q-derived TEE and activity energy expenditure were highly reproducible over three

months, with intraclass correlation coefficients of 0.84 and 0.73, respectively (17). Compared with

DLW, the STAR-Q demonstrated moderate validity for ranking individuals by TEE and activity

energy expenditure (Spearman's rho=0.53 and 0.40, respectively) (17).

To derive TEE from the STAR-Q, we first assigned an activity code and metabolic

equivalents of task (MET) value (29) to each reported activity. If the total time reported for

physical activity, sedentary behavior, and sleep was less than 24 hours/day, then we assigned a

MET of 1.2 for unaccounted time. The reported frequency, duration and assigned MET values

were used to estimate energy expenditure for each activity in units of MET-hours/day, which were

then summed to derive the total average MET-hours/day. The STAR-Q-derived TEE (TEE_{STAR-Q})

was then estimated as follows:

$TEE_{STAR-Q} = [[0.9\ MET \times hours\ of\ sleep/d] + [MET\text{-}hours/d \times weight\ (kg)]]*1.1$

– where TEE_{STAR-Q} from reported hours of sleep per day on the STAR-Q was multiplied by 0.9

METs (29, 30) and then added to the total average MET-hours per day in awake time multiplied

by baseline body weight. We applied the energy equivalence of 1 MET = 1 kcal/kg/hr to estimate

the energy cost of activities and added 10% for the thermic effect of food (12).

The calculation for individual PALs derived from self-report was as follows:

$PAL_{STAR-Q} = 0.9\ TEE_{STAR-Q}/BMR_{EST}$

– where BMR_{EST} was estimated using the Schofield equation based on age, sex, weight and

height (23).

Classification of misreporting

For comparability with a previous DLW study, we applied methods described by Tooze *et*

al. (11) to classify misreporting. Each participant was classified as an under-reporter (UR),

acceptable reporter (AR), or over-reporter (OR) depending on his or her individual ratio [rEI:TEE]

relative to group-level cut-points described below (31). Participants with a ratio less than the lower

cut-point were deemed URs; participants with ratios greater than the upper cut-point were deemed

ORs; and participants with ratios between the lower and upper cut-points were deemed ARs. Cut-

points were calculated separately for TEE_{STAR-Q} and TEE_{DLW}.

In our primary analysis to evaluate the Goldberg method and PAL_{STAR-Q} for classifying

individual misreporting, we calculated cut-points for the log of $rEI:TEE_{STAR-Q}$ as follows:

$$e^{\left(\pm 2 \times \sqrt{CV_{EI}^2/d + CV_{BMR}^2 + CV_{PAL}^2}\right)}$$

– where CV_{EI} was the within-subject coefficient of variation for EI, d was the number of days of

dietary assessment, CV_{BMR} was the precision of estimated versus measured BMR, and CV_{PAL} was

the total variation in PAL. We assigned one day for the number of days of dietary assessment ($d=1$)

since the C-DHQ I was only administered once. The CV_{EI} was assumed to be 18.6% for men and

19.8% for women based on repeatability of the US DHQ in a previous study (11) and CV_{BMR} was

assumed to be 8.5% for men and women as previously suggested (31). The CV_{PALs} calculated in

our three-month STAR-Q repeatability study were 17.5% for men and 15.1% for women (17).

In secondary analyses, we applied the classic Goldberg method (10) assuming a constant

PAL of 1.55 for participants (using $[rEI:TEE_{1.55}]$ derived from $[rEI:(BMR_{EST}*1.55)]$). Goldberg

cut-points were calculated as described above for $rEI:TEE_{STAR-Q}$, except CV_{PAL} was assumed to be

15% for both men and women as previously reported (31).

To derive gold standard classifications of misreporting, we calculated DLW cut-points

based on the log of the ratio $[rEI:TEE_{DLW}]$ as:

$$e^{\left(\pm 2 \times \sqrt{CV_{EI}^2/d + CV_{DLW}^2}\right)}$$

– where CV_{EI} was the within-subject coefficient of variation for EI, d was the number of days of

dietary assessment, and CV_{DLW} was the within-subject coefficient of variation for TEE_{DLW}. We

assumed CV_{rEI} to be 18.6% for men and 19.8% for women and CV_{DLW} to be 5.1% for both men

and women as previously reported (11).

Finally in sensitivity analyses, for comparison with a previous study (11), we substituted

$d=\infty$ for $d=1$ for dietary assessment in each of the cut-point equations (above). This alternate

estimate of 'd' reflects the intended use of the past year C-DHQ I to ascertain habitual long-term

diet. Given a cut-point denominator of $d=\infty$, the CV^2_{EI} term was then eliminated.

Statistics

Means (SD) were determined for continuous sociodemographic characteristics,

frequencies and percentages for categorical characteristics, and medians (Q1, Q3) for EI and

energy expenditure. In all analyses, DLW classifications were assumed to be the 'gold standard'

or *true* classifications. All analyses were stratified by gender. To appraise the validity of Goldberg

classifications using PAL_{STAR-Q}, and $PAL_{1.55}$, we estimated the proportion of participants correctly

classified and misclassified in terms of UR, AR and OR. Sensitivity, specificity, positive predictive

value (PPV), negative predictive value (NPV), and the accuracy of UR and AR classifications

excluded the sole *true* OR (n=1) identified by TEE_{DLW}. Sensitivity was the proportion of true URs

correctly classified using the Goldberg method. Specificity was the proportion of true ARs

correctly classified using the Goldberg method. PPV was the proportion of Goldberg URs (i.e.,

classified as such using the Goldberg method) that were true URs, interpreted as the probability of

being a true UR if classified an UR by the Goldberg method. NPV was the proportion of Goldberg

ARs that were true ARs, interpreted as the probability of being a true AR if classified as an AR by

the Goldberg method. Accuracy of UR and AR was the proportion of [true URs + true ARs]

classified correctly with the Goldberg method.

To appraise the validity of PAL_{STAR-Q} for estimating the magnitude of misreporting (UR

and OR) in kilocalories per day, we compared the within-person differences of TEE_{STAR-Q}-rEI

versus TEE_{DLW}-rEI using Pearson correlation coefficients and the Wilcoxon signed-rank test.

Furthermore, we used graphical illustrations described by Tooze *et al*. (11) to examine the

magnitude of discrepancies relative to the gold standard ratio $rEI:TEE_{DLW}$. If an individual-level

$rEI:TEE_{STAR-Q}$ ratio fell between the Goldberg (TEE_{STAR-Q}) and TEE_{DLW} lower (or upper) cut-

points, we attributed the source of discrepancy to differences in cut-point calculations. Otherwise,

we attributed the discrepancy to differences in TEE estimation (STAR-Q vs DLW). In addition,

we estimated the contribution of these discrepancies to the resulting sensitivity and specificity

attributes of the Goldberg method (TEE_{STAR-Q}).

All analyses were done using SAS Institute Software, Gary, NC (Version 9.4). We

considered a *P*-value < 0.05 statistically significance.

RESULTS

Participants aged 30-60 years (n=106) were recruited to the original DLW study between

July 2009 and July 2010 (17). Seven participants were excluded from the present analysis for

reasons outlined in **Figure 1,** leaving 99 participants of whom 60% were women. Compared with

men, women were somewhat younger (mean: 46.2 vs 50.6 years of age), had a lower BMI (mean:

23.5 vs 25.9 kg/m^2), and were less likely to complete post-secondary education (45% vs 51%)

Table 1. Most of the participants were Caucasian and had paid or volunteer work. Median EI,

TEE_{DLW}, TEE_{STAR-Q}, and BMR_{EST} were lower for women (1676 kcal, 2604 kcal, 2889 kcal and

1362 kcal, respectively) compared with men (2036 kcal, 3306 kcal, 3620 kcal, and 1866 kcal,

respectively). Median values for PAL_{DLW} and PAL_{STAR-Q}, were 1.8 and 2.0 for men and, 1.9 and

2.1 , for women (Table 1), consistent with 'active' and 'very active' PALs (7).

Results from the primary, secondary and sensitivity analyses are presented in **Table 2**. In

the primary analysis, 60% of men and 56% of women were classified as URs by TEE_{DLW}, and

58% of men and women by TEE_{STAR-Q}. The TEE_{STAR-Q} misclassified fewer men (15%) than

women (25%), resulting in sensitivity, specificity, PPV, NPV and accuracy >80% among men,

whereas for women these estimates were <80%. In secondary analyses, $PAL_{1.55}$ resulted in fewer

participants classified as URs (35% and 19% for men and women, respectively) than with TEE_{DLW}

and TEE_{STAR-Q} which led to high specificity and PPVs (both 93% for men and 100% for women),

but low sensitivity and NPVs (all <60% for men and women).

In sensitivity analyses, where we assumed $d=\infty$ for the FFQ, a greater proportion of men

and women were misclassified than with $d=1$. Seven were deemed to be true OR in contrast to 1

true OR when $d=1$, attributed to narrower cut-points when $d=\infty$. Otherwise, when d=∞, results

were generally similar (or less favorable) to assumed $d=1$ except for specificity and PPV, which

were 100% among men.

Figures 2A and 2B illustrate individual-level agreement and magnitude of misreporting in

kilocalories per day (i.e., TEE_{STAR-Q}-rEI relative to TEE_{DLW}-rEI). Pearson correlation coefficients

were higher for men (r=0.76, *P*<0.0001; classification agreement in 34 of 40 men) than for women

(r=0.37, *P*=0.0042; classification agreement in 44 of 59 women). Nonetheless, the Wilcoxon

signed-rank test indicated statistically significant within-person differences between the two

methods (P = 0.0021 and P < 0.0001 for men and women, respectively). The magnitude of EI

under-reporting tended to be larger for misclassified URs (approximately 1,000 to 1,500 kcal/d; 3

men and 7 women) while the magnitude of EI under-reporting was typically smaller for

misclassified ARs (<1,000 kcal/d approximately; 2 men and 8 women).

In **Figures 3A and 3B**, we show whether discrepant classifications arose from different

cut-points or from different TEE estimation methods. Among 40 men (**Figure 3A**), five

misclassifications were due to TEE estimation (one classified as an OR), and one was due to cut-

point estimation. Among 59 women (**Figure 3B**), more misclassifications were due to differences

in TEE estimation than to cut-off estimation. Among *true* URs misclassified as ARs using the

PAL$_{STAR-Q}$ method (reflecting sensitivity), 2 of 3 men and 3 of 7 women were misclassified owing

to TEE estimation. Among true ARs misclassified as URs using the PAL$_{STAR-Q}$ method (reflecting

specificity), all misclassifications (2 men and 8 women) were due to differences in TEE estimation.

DISCUSSION

Using DLW, the 'gold-standard' reference biomarker for TEE and unbiased proxy for

habitual EI in weight-stable individuals (2), we determined the accuracy of STAR-Q-derived PALs

and the Goldberg method to classify the reporting status of FFQ-derived EI in community dwelling

middle-aged men and women. We found that the PAL$_{STAR-Q}$ generally improved the sensitivity

and overall accuracy of the Goldberg method for discriminating between URs and ARs compared

with the Goldberg method based on a global PAL of 1.55. We also identified gender-based

differences in the performance of the Goldberg-PAL$_{STAR-Q}$. In men, the method discriminated

between UR and AR with an overall high level of accuracy at 87%, while in women it exhibited

fair performance, with an overall classification accuracy of 75%.

A key determinant of the accuracy of the Goldberg method is the degree to which PALs

represent the individuals under study. Three previous DLW validation studies have explored

approaches to identify the optimal PAL for estimating EI misreporting (11, 14, 32). Two of these

studies examined EI misreporting on food records and/or diet histories and found under-reporting

in 14% (32) and 34% (14) of adult participants. Similar to our results these studies reported high

levels of specificity but relatively low sensitivity for identifying URs when a global PAL of 1.55

was assumed. Additionally, Black (14) leveraged the strength of a large database comprised of 21

DLW studies (n=429) to test the effect of various global PALs (from 1.55 to 1.95) and reported

incremental increases in sensitivity, with increasing PALs; however, this improvement coincided

with decreases in specificity. In a small sample of adults (n=14), Livingstone *et al* (32) tested the

performance of gender-specific PALs to detect URs based on individual heart-rate monitors

matched to low, medium and high activity categories of PAL (9). Compared with $PAL_{1.55}$,

sensitivity increased substantially from 50% to 100% with negligible decreases in specificity.

Subsequently, Tooze *et al* (11) reported favorable results for the Goldberg method with a $PAL_{1.55}$

to detect DLW-classified URs on 24-hour dietary recalls (~20%) and an FFQ (~50% UR) in the

Observing Protein and Energy Nutrition (OPEN) Study (n=451). Specificity was 88% for the FFQ

in both men and women. The sensitivity to detect URs was high for the FFQ at 96% when usual

intake was assumed for the FFQ ($d=\infty$), but substantially lower when OPEN study FFQ within-

person variability was applied (72% sensitivity for men and 62% for women).

The results reported by Livingstone *et al* (32) and Black (14) are consistent with our results

in supporting the premise that a global $PAL_{1.55}$ may be underestimating the activity levels of some

adult populations – and hence, more individualized PALs may improve the overall accuracy of the

Goldberg method to identify EI under-reporting. While our findings contrast with those of Tooze

et al (11), it is worth noting that the prevalence of true UR (50%) was lower than in our population,

which affects PPV and NPV values. In addition, participants in the OPEN study were on average

less active and slightly older than in the STAR-Q validation study. Median values for DLW-

derived TEE and PAL were 2,813 kcal and 1.69 for men and 2,283 kcal and 1.75 for women in

OPEN (33, 34) and 3,306 kcal and 1.8 for men and 2604 kcal and 1.9 for women in the STAR-Q

study, with ages ranging from 40 to 69 years in OPEN (33, 34) versus 30 to 60 years in the STAR-

Q study.

Statistical approaches have been explored with respect to handling URs once identified,

e.g., exclusion of misreporters, stratification by reporting status and adjustment for reporting status

in regression analyses (35-39). While the optimal approach may be study-specific, based on study

design, method of data collection and population characteristics, there is consensus that exclusion

of misreporters is undesirable given the loss of statistical power (36) and loss of valuable

information inherent in the characteristics of those excluded (35). As such, it is essential that

investigators understand the potential impact of measurement error on study results prior to

selecting the approach with the greatest potential to mitigate biasing effects of misreporting (39).

While the STAR-Q was designed to capture a comprehensive profile of daily activities

over a 24-hour period, 50% of true URs misclassified as ARs and all true ARs misclassified as

URs, were likely misclassified owing to errors in the estimation of TEE$_{STAR-Q}$ **(Figure 3a and 3b).**

Hence, both sensitivity and specificity which are important characteristics in the assessment of

exposure validity in epidemiologic studies (40), could be strengthened. For example, measuring

rather than estimating participant BMR (23), and using updated MET (energy expenditure cost)

values (41, 42), could improve future TEE$_{STAR-Q}$ estimation. In addition, the STAR-Q relies on the

reporting of activities during the previous month, a task not without cognitive challenges, though

less so than reporting past-year activities (19). Even less cognitively challenging is the Activities

Completed over Time in 24 hours (ACT24), an internet-based recall of previous day activities that

has been shown to estimate mean TEE and PAL values within 5% of DLW-based estimate and is

feasible to administer in epidemiologic studies (43, 44). The accuracy of short-term recall

combined with longer-term measures that reflect habitual lifestyle behaviors is an intriguing

approach that has been suggested for improving dietary intake measures (45) and may also have

potential to improve self reports of activities for TEE and PAL estimation. Moreover, new

wearable device-based technologies may in the foreseeable future more accurately estimate PALs

and TEE in large populations to complement the contextual data collected by traditional

questionnaires (46). Hence, it is anticipated that methodologies may be developed to combine

device-derived estimates of energy expenditure and body composition to validate self-reported EI

(47-49); though these approaches are still viewed as 'works in progress' (50).

A noteworthy strength of this study is the validation of the Goldberg-STAR-Q method

against DLW – and the unique use of individual-level PALs. Furthermore, as noted in the recent

report on DRIs for Energy, of 128 DLW studies representing 25 countries in the International

Atomic Energy Agency DLW Database, none were Canadian (7) highlighting a gap in this area

of Canadian research that this study begins to address. Nonetheless, our study has some limitations

that merit consideration. First, we used prediction equations for BMR, which can account for a

substantial proportion of TEE (51) and could be a source of error in the estimation of PAL_{STAR-Q}.

Second, we cannot be certain that some of the under-reporting on the past-year C-DHQ I did not

actually reflect undereating during the previous year, though we aimed to minimize this occurrence

by recruiting generally weight stable participants to the DLW study. Third, we caution that we

cannot assume that misreporting occurs proportionally across all nutrients similar to EI. Indeed,

while still relatively under-studied, evidence is accumulating to suggest that foods and nutrients

vary in the accuracy with which they are reported (3, 52-54). On a final note, our population was

largely Caucasian middle-aged adults; hence, our results may not be generalizable to older (55,

56) or more racially diverse populations and those with other characteristics that may affect

reporting and health status (57-60).

Overall, our DLW-validated results lend credibility to the thesis that detailed

comprehensive questionnaire-based activities may be employed with the Goldberg method to

explore EI misreporting in large-scale diet studies. Importantly, such strategies create opportunities

to carry out sensitivity analyses to better understand the impact of EI misreporting on diet-disease

associations in nutritional epidemiology.

STATEMENTS AND DECLARATIONS

Data availability: Data described in the manuscript, code book, and analytic code will be made available upon reasonable request to the corresponding author. The data are not publicly available due to privacy and/or ethical restrictions.

Conflicts of Interest: IC and KAK were awarded research funding paid to their institutions from the Canadian Institutes of Health Research (CIHR: Grant MOP-86632) and the Alberta Cancer Research Institute (Grant 24265) to support this investigator-initiated study. *Support during past 36 months*: IC has received honoraria from the American Institute of Cancer Research and Genome Quebec for her role as a grant reviewer. AK has been awarded research funding paid to her institution from CIHR to support investigator-initiated research unrelated to this study. AK has also received an honorarium from the Université Laval for her role as an epidemiology program evaluator. JAT discloses her role as Committee Member on the *Dietary Reference Intakes for Energy*, National Academies of Sciences, Engineering, and Medicine, 2023 (no remuneration received). All other co-authors have no conflicts or competing interests to declare.

Author contributions:

Concept and design: HKN, KAK and IC; *Acquisition of data and/or lab analysis*: HKN, KAK, RRL and IC; *Statistical analysis*: SA, FK and IC; *Interpretation of data*: All authors; *Drafting the manuscript*: HKN and IC; *Critical revision of manuscript and important intellectual contribution*: All authors; *Obtaining funding*: IC and KAK; *Administrative, technical or material support*: HKN, KAK and IC; *Supervision*: IC; *Primary responsibility for the final content*: IC.

FIGURE LEGENDS

Figure 1. Participant recruitment in the STAR-Q validation study (2009-2011), Alberta, Canada.

Figure 2. Correlation between the degree of energy intake misreporting based on Goldberg classifications (i.e., individual differences: TEE_{STAR-Q} – rEI) and the degree of energy intake misreporting based on doubly-labeled water classifications (i.e., individual differences: TEE_{DLW}-rEI) in kilocalories per day. **Figure 2A** shows the correlation among 40 men and **Figure 2B** shows the correlation among 59 women in the STAR-Q validation study (2009-2011), Alberta, Canada. AR, acceptable reporters; DLW, doubly labeled water; UR, under-reporters.

Figure 3. Sources of discrepancy between the Goldberg and doubly-labeled water methods for classifying energy intake misreporting. **Figure 3A** illustrates sources of discrepancy in men and **Figure 3B** illustrates sources of discrepancy in women. The gold standard doubly-labeled water classifications of misreporting are represented with shaded circles. The Goldberg classifications of misreporting, based on the ratio [$rEI:TEE_{STAR-Q}$], are represented with open circles (agreement between methods), squares (discrepant because of TEE estimation) or triangles (discrepant because of different cut-offs). The gray solid line represents the lower cut-off for gold standard doubly-labeled water classifications (0.68 for men; 0.66 for women). The black solid line represents the lower cut-off for Goldberg classifications (0.58 for men; 0.59 for women).

Table 1. Sociodemographic characteristics, energy expenditure and energy intake for 99 participants in the STAR-Q validation study (2009-2011), Alberta, Canada.

	Men, $n = 40$		Women, $n = 59$	
Sociodemographic characteristics	Mean (SD)	%	Mean (SD)	%
Age, years	50.6 (7.0)		46.2 (8.5)	
Body mass index, kg/m^2	25.9 (3.1)		23.5 (2.9)	
Education				
Less than post-secondary education		34		45
At least post-secondary education		51		45
Missing		15		10
Ethnicity				
Caucasian		73		85
Asian		3		5
Other		3		-
Missing		20		10
Paid or volunteer work		93		93
Energy intake and expenditure	Median [Q1, Q3]		Median [Q1, Q3]	
Energy intake, kcal/d	2036 [1627, 2634]		1676 [1317, 2033]	
Self-reported activity (STAR-Q), MET-h/d[1]	28.8 [24.6, 32.6]		30.6 [25.6, 39.5]	
TEE$_{DLW}$, kcal/d	3306 [2843, 3727]		2604 [2226, 2854]	
TEE$_{STAR-Q}$, kcal/d	3620 [3178, 4019]		2889 [2420, 3387]	
BMR$_{EST}$, kcal/d	1866 [1744, 1924]		1362 [1331. 1438]	
PAL (TEE$_{DLW}$:BMR$_{EST}$)	1.8 [1.6, 2.0]		1.9 [1.7, 2.1]	
PAL (TEE$_{STAR-Q}$:BMR$_{EST}$)	2.0 [1.8, 2.1]		2.1 [1.8, 2.4]	

[1] The sum of MET-h/d from all types of physical activity including light, moderate, and vigorous intensity and sedentary activities (sleep is excluded).

BMR_{EST}, estimated basal metabolic rate; MET, metabolic equivalent of task; PAL, physical activity level;

STAR-Q, Sedentary Time and Activity Reporting Questionnaire; TEE, total energy expenditure.

Table 2. Validity of the Goldberg method compared with the 'gold standard' doubly-labeled water method for classifying energy intake misreporting, for 40 men and 59 women[1] in the STAR-Q validation study (2009-2011), Alberta, Canada.

Analysis	Source of PAL	Gender	Cut-offs		UR^2	Mis-classified[2]	Sensitivity[1]	Specificity[1]	PPV[1]	NPV[1]	Accuracy of UR & AR[1]
			Lower	Upper	(%)	(%)	(%)	(%)	(%)	(%)	(%)
Primary & Secondary	$DLW_{d=1}$	M	0.68	1.47	60	-	-	-	-	-	-
		W	0.66	1.51	56	-	-	-	-	-	-
Primary (PAL_{STAR-Q})^3	STAR-Q	M	0.58	1.71	58	15	88	87	91	81	87
		W	0.59	1.69	58	25	79	69	77	72	75
Secondary ($PAL_{1.55}$)^3	Assumed 1.55	M	0.60	1.66	35	33	54	93	93	56	69
		W	0.59	1.69	19	37	33	100	100	54	63
Sensitivity ($d=\infty$) ^{4}	$DLW_{d=\infty}$	M	0.90	1.11	85	-	-	-	-	-	-
		W	0.90	1.11	83	-	-	-	-	-	-
	STAR-Q	M	0.68	1.48	60	35	71	100	100	38	75
		W	0.71	1.42	76	32	78	30	84	21	69

[1] Calculations of sensitivity, specificity, PPV, NPV, and accuracy - by definition - exclude true overreporters identified with DLW. In our Primary and Secondary analyses (where d=1 in the DLW cut-off formula), one participant was a true over-reporter (therefore n=98; 39 men, 59 women). In our Sensitivity analysis (where d=∞ in the DLW cut-off formula), seven participants were true over-reporters (therefore n=92; 36 men, 56 women). Doubly-labeled water classifications of misreporting, based on the ratio [rEI:TEE$_{DLW}$], served as the 'gold standard' (true) classification. Sensitivity is the proportion of true under-reporters who were correctly classified using the Goldberg method. Specificity

is the proportion of true acceptable reporters who were correctly classified using the Goldberg method. Positive predictive value (PPV) is the proportion of Goldberg under-reporters (i.e., classified as under-reporters using the Goldberg method) that were true under-reporters. Negative predictive value (NPV) is the proportion of Goldberg acceptable reporters (i.e., classified as acceptable reporters using the Goldberg method) that were true acceptable reporters. Accuracy of UR & AR is the proportion of [true under-reporters (UR) + true acceptable reporters (AR)] classified correctly using the Goldberg method.

[2] Calculations based on the entire study population (n=99; 40 men, 59 women), including overreporters. Doubly-labeled water classifications of misreporting, based on the ratio [rEI:TEE$_{DLW}$], served as the 'gold standard' (true) classification. UR was the proportion classified as an under-reporter. Misclassified was the proportion classified incorrectly by the Goldberg method.

[3] The formula for the Goldberg cut-offs included d=1 for the number of days of dietary assessment. We assigned d=1 because the C-DHQ I was administered once.

[4] The formula for Goldberg and DLW cut-offs included d=∞, representing an infinite number of days of dietary assessment. We assigned d=∞ for an extreme interpretation of 'd', as a sensitivity analysis, because the past year C-DHQ I is intended to measure habitual long-term diet.

AR, acceptable reporters; C-DHQ I, Canadian Diet History Questionnaire version I; DLW, doubly-labeled water; NPV, negative predictive value; PAL, Physical Activity Level; PPV, positive predictive value; rEI, self-reported energy intake; STAR-Q, Sedentary Time and Activity Reporting Questionnaire; TEE$_{DLW}$, total energy expenditure estimated from doubly labeled water; UR, underreporters.

Figure 1

r=0.76, *P*<0.0001

Figure 2A (men)

Figure 2B (women)

Figure 3A (men)

Figure 3B (women)

REFERENCES:

1.   Freedman LS, Commins JM, Moler JE, Arab L, Baer DJ, Kipnis V, et al. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake. Am J Epidemiol. 2014;180(2):172-88. doi: 10.1093/aje/kwu116.

2.   Ravelli MN, Schoeller DA. Traditional Self-Reported Dietary Instruments Are Prone to Inaccuracies and New Approaches Are Needed. Front Nutr. 2020;7(90):90. doi: 10.3389/fnut.2020.00090.

3.   Prentice RL. Intake Biomarkers for Nutrition and Health: Review and Discussion of Methodology Issues. Metabolites. 2024;14(5). doi: 10.3390/metabo14050276.

4.   Neuhouser ML, Prentice RL, Tinker LF, Lampe JW. Enhancing Capacity for Food and Nutrient Intake Assessment in Population Sciences Research. Annu Rev Public Health. 2022. doi: 10.1146/annurev-publhealth-071521-121621.

5.   Banna JC, McCrory MA, Fialkowski MK, Boushey C. Examining Plausibility of Self-Reported Energy Intake Data: Considerations for Method Selection. Frontiers in Nutrition. 2017;4(45). doi: 10.3389/fnut.2017.00045.

6.   Black AE, Goldberg GR, Jebb SA, Livingstone MB, Cole TJ, Prentice AM. Critical evaluation of energy intake data using fundamental principles of energy physiology: 2. Evaluating the results of published surveys. Eur J Clin Nutr. 1991;45(12):583-99.

7.   National Academies of Sciences, Engineering, and Medicine 2023. Dietary Reference Intakes for Energy. Washington (DC): National Academies Press (US) https://doi.org/10.17226/26818.

8.   Neilson HK, Robson PJ, Friedenreich CM, Csizmadi I. Estimating activity energy expenditure: how valid are physical activity questionnaires? Am J Clin Nutr. 2008;87(2):279-91 [PMID: 18258615]. doi: 10.1093/ajcn/87.2.279.

9.   Energy and protein requirements. Report of a joint FAO/WHO/UNU Expert Consultation. World Health Organ Tech Rep Ser. 1985;724:1-206.

10.   Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, Prentice AM. Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. Eur J Clin Nutr. 1991;45(12):569-81.

11.   Tooze JA, Krebs-Smith SM, Troiano RP, Subar AF. The accuracy of the Goldberg method for classifying misreporters of energy intake on a food frequency questionnaire and 24-h recalls: comparison with doubly labeled water. Eur J Clin Nutr. 2012;66(5):569-76. doi: 10.1038/ejcn.2011.198.

12.   Speakman JR, Pontzer H. Quantifying physical activity energy expenditure based on doubly labelled water and basal metabolism calorimetry: what are we actually measuring? Curr Opin Clin Nutr Metab Care. 2023;26(5):401-8. doi: 10.1097/mco.0000000000000937.

13.   Schoeller DA. Recent advances from application of doubly labeled water to measurement of human energy expenditure. J Nutr. 1999;129(10):1765-8. doi: 10.1093/jn/129.10.1765.

14.   Black AE. The sensitivity and specificity of the Goldberg cut-off for EI:BMR for identifying diet reports of poor validity. Eur J Clin Nutr. 2000;54(5):395-404. doi: 10.1038/sj.ejcn.1600971.

15. Grech A, Hasick M, Gemming L, Rangan A. Energy misreporting is more prevalent for those of lower socio-economic status and is associated with lower reported intake of discretionary foods. British Journal of Nutrition. 2021;125(11):1291-8. doi: 10.1017/S0007114520003621.

16. Black AE, Coward WA, Cole TJ, Prentice AM. Human energy expenditure in affluent societies: an analysis of 574 doubly-labelled water measurements. Eur J Clin Nutr. 1996;50(2):72-92.

17. Csizmadi I, Neilson HK, Kopciuk KA, Khandwala F, Liu A, Friedenreich CM, et al. The Sedentary Time and Activity Reporting Questionnaire (STAR-Q): reliability and validity against doubly-labeled water and 7-day activity diaries. Am J Epidemiol. 2014;180(4):424-35 - Editor's Choice. doi: 10.1093/aje/kwu150.

18. Šuc A, Einfalt L, Šarabon N, Kastelic K. Validity and reliability of self-reported methods for assessment of 24-h movement behaviours: a systematic review. The international journal of behavioral nutrition and physical activity. 2024;21(1):83 [PMID: 39095778]. doi: 10.1186/s12966-024-01632-4.

19. Neilson HK, Ullman R, Robson PJ, Friedenreich CM, Csizmadi I. Cognitive testing of the STAR-Q: insights in activity and sedentary time reporting. J Phys Act Health. 2013;10(3):379-89 [PMID: 22820674]. doi: 10.1123/jpah.10.3.379.

20. Robson PJ, Solbak NM, Haig TR, Whelan HK, Vena JE, Akawung AK, et al. Design, methods and demographics from phase I of Alberta's Tomorrow Project cohort: a prospective cohort profile. CMAJ Open. 2016;4(3):E515-e27. doi: 10.9778/cmajo.20160005.

21.   Ye M, Robson PJ, Eurich DT, Vena JE, Xu JY, Johnson JA. Cohort Profile: Alberta's Tomorrow Project. Int J Epidemiol. 2017;46(4):1097-8l. doi: 10.1093/ije/dyw256.

22.   Racette SB, Schoeller DA, Luke AH, Shay K, Hnilicka J, Kushner RF. Relative dilution spaces of 2H- and 18O-labeled water in humans. Am J Physiol. 1994;267(4 Pt 1):E585-90. doi: 10.1152/ajpendo.1994.267.4.E585.

23.   Schofield WN. Predicting basal metabolic rate, new standards and review of previous work. Hum Nutr Clin Nutr. 1985;39 Suppl 1:5-41.

24.   Douketis JD, Paradis G, Keller H, Martineau C. Canadian guidelines for body weight classification in adults: application in clinical practice to screen for overweight and obesity and to assess disease risk. CMAJ. 2005;172(8):995-8. doi: 10.1503/cmaj.045170.

25.   Csizmadi I, Kahle L, Ullman R, Dawe U, Zimmerman TP, Friedenreich CM, Bryant H, Subar AF. Adaptation and evaluation of the National Cancer Institute's Diet History Questionnaire and nutrient database for Canadian populations. Public Health Nutr. 2007;10(1):88-96 [PMID: 17212847]. doi: 10.1017/s1368980007184287.

26.   Csizmadi I, Boucher BA, Lo Siou G, Massarelli I, Rondeau I, Garriguet D, Koushik A, Elenko J, Subar AF. Using national dietary intake data to evaluate and adapt the US Diet History Questionnaire: the stepwise tailoring of an FFQ for Canadian use. Public Health Nutrition. 2016;19(18):3247-55. doi: 10.1017/S1368980016001506.

27.   Mickle AT, Brenner DR, Beattie T, Williamson T, Courneya KS, Friedenreich CM. The Dietary Inflammatory Index® and Alternative Healthy Eating Index 2010 in relation to leucocyte telomere length in postmenopausal women: a cross-sectional study. J Nutr Sci. 2019;8:e35. doi: 10.1017/jns.2019.32.

28.   Biel RK, Csizmadi I, Cook LS, Courneya KS, Magliocco AM, Friedenreich CM. Risk of endometrial cancer in relation to individual nutrients from diet and supplements. Public Health Nutr. 2011;14(11):1948-60. doi: 10.1017/s1368980011001066.

29.   Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, et al. Compendium of physical activities: an update of activity codes and MET intensities. Med Sci Sports Exerc. 2000;32(9 Suppl):S498-504. doi: 10.1097/00005768-200009001-00009.

30.   Ainsworth BE, Haskell WL, Leon AS, Jacobs DR, Jr., Montoye HJ, Sallis JF, Paffenbarger RS, Jr. Compendium of physical activities: classification of energy costs of human physical activities. Med Sci Sports Exerc. 1993;25(1):71-80. doi: 10.1249/00005768-199301000-00011.

31.   Black AE. Critical evaluation of energy intake using the Goldberg cut-off for energy intake:basal metabolic rate. A practical guide to its calculation, use and limitations. International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity. 2000;24(9):1119-30. doi: 10.1038/sj.ijo.0801376.

32.   Livingstone MB, Robson PJ, Black AE, Coward WA, Wallace JM, McKinley MC, Strain JJ, McKenna PG. An evaluation of the sensitivity and specificity of energy expenditure measured by heart rate and the Goldberg cut-off for energy intake: basal metabolic rate for identifying mis-reporting of energy intake by adults and children: a retrospective analysis. Eur J Clin Nutr. 2003;57(3):455-63. doi: 10.1038/sj.ejcn.1601563.

33.   Subar AF, Kipnis V, Troiano RP, Midthune D, Schoeller DA, Bingham S, et al. Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study. Am J Epidemiol. 2003;158(1):1-13. doi: 10.1093/aje/kwg092.

34.   Tooze JA, Schoeller DA, Subar AF, Kipnis V, Schatzkin A, Troiano RP. Total daily energy expenditure among middle-aged men and women: the OPEN Study. Am J Clin Nutr. 2007;86(2):382-7. doi: 10.1093/ajcn/86.2.382.

35.   Jessri M, Lou WY, L'Abbé MR. Evaluation of different methods to handle misreporting in obesity research: evidence from the Canadian national nutrition survey. Br J Nutr. 2016;115(1):147-59. doi: 10.1017/s0007114515004237.

36.   Tooze JA, Freedman LS, Carroll RJ, Midthune D, Kipnis V. The impact of stratification by implausible energy reporting status on estimates of diet-health relationships. Biometrical Journal. 2016;58(6):1538-51. doi: doi:10.1002/bimj.201500201.

37.   Solbak NM, Al Rajabi A, Akawung AK, Lo Siou G, Kirkpatrick SI, Robson PJ. Strategies to Address Misestimation of Energy Intake Based on Self-Report Dietary Consumption in Examining Associations Between Dietary Patterns and Cancer Risk. Nutrients. 2019;11(11 [PMID: 31683814]). doi: 10.3390/nu11112614.

38.   Lo Siou G, Akawung AK, Solbak NM, McDonald KL, Al Rajabi A, Whelan HK, Kirkpatrick SI. The effect of different methods to identify, and scenarios used to address energy intake misestimation on dietary patterns derived by cluster analysis. Nutr J. 2021;20(1):42 [PMID: 33964947]. doi: 10.1186/s12937-021-00696-3.

39.   Yamamoto N, Ejima K, Zoh RS, Brown AW. Bias in nutrition-health associations is not eliminated by excluding extreme reporters in empirical or simulation studies. Elife. 2023;12. doi: 10.7554/eLife.83616.

40.   Trevethan R. Sensitivity, Specificity, and Predictive Values: Foundations, Pliabilities, and Pitfalls in Research and Practice. Front Public Health. 2017;5:307. doi: 10.3389/fpubh.2017.00307.

41.   Herrmann SD, Willis EA, Ainsworth BE. The 2024 Compendium of Physical Activities and its expansion. J Sport Health Sci. 2024;13(1):1-2. doi: 10.1016/j.jshs.2023.09.008.

42.   Herrmann SD, Willis EA, Ainsworth BE, Barreira TV, Hastert M, Kracht CL, et al. 2024 Adult Compendium of Physical Activities: A third update of the energy costs of human activities. J Sport Health Sci. 2024;13(1):6-12. doi: 10.1016/j.jshs.2023.10.010.

43.   Matthews CE, Kozey Keadle S, Moore SC, Schoeller DS, Carroll RJ, Troiano RP, Sampson JN. Measurement of Active and Sedentary Behavior in Context of Large Epidemiologic Studies. Med Sci Sports Exerc. 2018;50(2):266-76. doi: 10.1249/MSS.0000000000001428.

44.   Matthews CE, Patel S, Saint-Maurice P, Loftfield E, Keadle SK, Chen KY, Brychta R, LaMunion S, Berrigan D. Physical Activity Levels (PAL) in United States Adults - 2019. Med Sci Sports Exerc. 2022. doi: 10.1249/mss.0000000000003102.

45.   Freedman LS, Midthune D, Arab L, Prentice RL, Subar AF, Willett W, Neuhouser ML, Tinker LF, Kipnis V. Combining a Food Frequency Questionnaire With 24-Hour Recalls to Increase the Precision of Estimation of Usual Dietary Intakes—Evidence From the Validation Studies Pooling Project. American Journal of Epidemiology. 2018;187(10):2227-32. doi: 10.1093/aje/kwy126.

46.   Sampson JN, Matthews CE, Freedman L, Carroll RJ, Kipnis V. Methods to Assess Measurement Error in Questionnaires of Sedentary Behavior. J Appl Stat. 2016;43(9):1706-21. doi: 10.1080/02664763.2015.1117593.

47.   Hibbing PR, Shook RP, Panda S, Manoogian ENC, Mashek DG, Chow LS. Predicting energy intake with an accelerometer-based intake-balance method. Br J Nutr. 2022:1-23. doi: 10.1017/s0007114522003312.

48. Hibbing PR, Welk GJ, Ries D, Yeh H-W, Shook RP. Criterion validity of wrist accelerometry for assessing energy intake via the intake-balance technique. International Journal of Behavioral Nutrition and Physical Activity. 2023;20(1):115. doi: 10.1186/s12966-023-01515-0.

49. Shook RP, Yeh H-W, Welk GJ, Davis AM, Ries D. Commercial Devices Provide Estimates of Energy Balance with Varying Degrees of Validity in Free-Living Adults. The Journal of Nutrition. 2021;152(2):630-8. doi: 10.1093/jn/nxab317.

50. Gebel K, Ding D. Using Commercially Available Measurement Devices for the Intake-Balance Method to Estimate Energy Intake: Work in Progress. The Journal of Nutrition. 2022;152(2):373-4. doi: https://doi.org/10.1093/jn/nxab393.

51. Manini TM. Energy expenditure and aging. Ageing Res Rev. 2010;9(1):1-11. doi: 10.1016/j.arr.2009.08.002.

52. Krebs-Smith SM, Graubard BI, Kahle LL, Subar AF, Cleveland LE, Ballard-Barbash R. Low energy reporters vs others: a comparison of reported food intakes. Eur J Clin Nutr. 2000;54(4):281-7. doi: 10.1038/sj.ejcn.1600936.

53. Lafay L, Mennen L, Basdevant A, Charles MA, Borys JM, Eschwège E, Romon M. Does energy intake underreporting involve all kinds of food or only specific food items? Results from the Fleurbaix Laventie Ville Santé (FLVS) study. International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity. 2000;24(11):1500-6. doi: 10.1038/sj.ijo.0801392.

54. Prentice RL, Aragaki AK, Zheng C, Manson JE, Tinker LF, Schoeller DA, et al. Energy intake is associated with dietary macronutrient densities: inversely with protein and

monounsaturated fat and positively with polyunsaturated fat and carbohydrate among postmenopausal females. Am J Clin Nutr. 2025. doi: 10.1016/j.ajcnut.2025.03.011.

55. Porter J, Ward LC, Nguo K, Ward A, Davidson Z, Gibson S, Prentice R, Neuhouser ML, Truby H. Development and validation of age-specific predictive equations for total energy expenditure and physical activity levels for older adults. The American Journal of Clinical Nutrition. 2024;119(5):1111-21. doi: 10.1016/j.ajcnut.2024.02.005.

56. Wong WW, Speakman JR, Ainslie PN, Anderson LJ, Arab L, Baddou I, et al. Decline in Isotope Dilution Space Ratio Above Age 60 Could Affect Energy Estimates Using the Doubly Labeled Water Method. J Nutr. 2024;154(12):3824-31. doi: 10.1016/j.tjnut.2024.10.016.

57. Prentice RL, Mossavar-Rahmani Y, Huang Y, Van Horn L, Beresford SA, Caan B, et al. Evaluation and comparison of food records, recalls, and frequencies for energy and protein assessment by using recovery biomarkers. Am J Epidemiol. 2011;174(5):591-603. doi: 10.1093/aje/kwr140.

58. Pryer JA, Vrijheid M, Nichols R, Kiggins M, Elliott P. Who are the 'low energy reporters' in the dietary and nutritional survey of British adults? Int J Epidemiol. 1997;26(1):146-54. doi: 10.1093/ije/26.1.146.

59. Prentice AM, Black AE, Coward WA, Cole TJ. Energy expenditure in overweight and obese adults in affluent societies: an analysis of 319 doubly-labelled water measurements. Eur J Clin Nutr. 1996;50(2):93-7.

60. Karelis AD, Lavoie ME, Fontaine J, Messier V, Strychar I, Rabasa-Lhoret R, Doucet E. Anthropometric, metabolic, dietary and psychosocial profiles of underreporters of energy intake: a doubly labeled water study among overweight/obese postmenopausal women--a

Montreal Ottawa New Emerging Team study. Eur J Clin Nutr. 2010;64(1):68-74. doi: 10.1038/ejcn.2009.119.