Identifying predictors of neuropathic pain medication prescribing, adherence, and discontinuation: a systematic review and meta-analysis

Mia E. Koponen [1,*], Dhaneesha N. S. Senaratne [1], Harry L. Hébert [1], Blair H. Smith [1], Lesley A. Colvin [1]

[1] Chronic Pain Research Group, Division of Population Health & Genomics, School of Medicine, University of Dundee, Dundee, UK

*Corresponding author: ME Koponen, mia.elena.koponen@outlook.com, Chronic Pain Research Group, University of Dundee, Level 7 Corridor E, Ninewells Hospital, Dundee, DD1 9SY, UK

Abstract

Background: Pharmacoepidemiological studies show that large proportions of people with neuropathic pain (NeuP) are not prescribed the medications recommended for NeuP, do not adhere well to these medications, and/or often discontinue the treatment within six months. Identifying what predicts these outcomes can inform a strategy to ensure that people with NeuP are prescribed recommended medication and continue their treatment when appropriate.

Methods: We carried out a systematic review and meta-analysis to identify predictors of pain medication prescribing, adherence, and discontinuation in adults with NeuP. Electronic searches were conducted in Embase, PubMed, Web of Science, and CINAHL Plus.

Results: We identified 69 relevant studies and divided them into non-mutually exclusive categories based on the outcomes they had investigated. There were 53 studies on prescribing, 14 on adherence, and 27 on discontinuation. Predictors associated with being prescribed recommended NeuP medications included having a mental health disorder, diabetes, and/or white ethnicity. Predictors associated with better adherence included being prescribed serotonin-norepinephrine reuptake inhibitors compared to gabapentinoids or tricyclic antidepressants; dose titration, and/or implementing a medicine reminder. Predictors associated with a higher likelihood of discontinuation included being prescribed tricyclic antidepressants rather than other antidepressants and/or having a combination of NeuP medications rather than monopharmacotherapy.

Discussion: There is a need to focus on improving NeuP medication prescribing for ethnic minorities and people without diabetes. Adherence may be improved by using dose titration and/or a medicine reminder. Prescribing monopharmacotherapy rather than combination pharmacotherapy for NeuP may result in better treatment persistence.

Systematic review protocol: PROSPERO CRD42023464307.

Keywords

Meta-analysis; neuropathic pain; pain medicine; pharmacoepidemiology; prescribing; systematic review; treatment adherence; treatment persistence

Introduction

Neuropathic pain (NeuP) is 'pain that arises as a direct consequence of a lesion or disease affecting the somatosensory system' and is associated with unpleasant sensations including those described as burning, pins and needles, electric shocks, and numbness.[1, 2] NeuP affects around 10% of the general population and has a substantial negative effect on people's quality of life even compared to those experiencing other types of chronic pain.[3–5] In addition, common analgesics such as non-steroidal anti-inflammatory drugs (NSAIDs) and opioids that are used to treat nociceptive pain may be ineffective in treating neuropathic pain.[6] According to the pharmacotherapy guidelines by the International Association for the Study of Pain (IASP) Neuropathic Pain Special Interest Group (NeuPSIG), standard first-line medications for NeuP are gabapentinoids, serotonin-norepinephrine reuptake inhibitors (SNRIs), or tricyclic antidepressants (TCAs).[6, 7] Second-line NeuP medications include topical capsaicin or lidocaine, and third-line medications include botulinum toxin A or opioids.[6, 7]

Despite the existing treatment guidelines and variety of choices, studies suggest that there are major inconsistencies in NeuP treatment. For example, in a retrospective cohort study conducted in the USA during 2001-2014, only 52.9% of 14,426 people with NeuP were prescribed a first-line medication in the year before or three years after diagnosis.[8] This suggests that a large proportion of people with NeuP are not being prescribed recommended treatment. Another retrospective cohort study conducted in the USA during 2001-2016 showed that in a sample of 57,495 people with NeuP, adherence to gabapentinoids and SNRIs was on average ~40-50% and ~50-60% respectively,[9] which suggests that adherence to NeuP medications is moderate or poor.[10] Moreover, in a retrospective cohort study conducted in Sweden during 2004-2009, 60% of 4,718 people with NeuP had discontinued their treatment five months after first prescription.[11] Other studies have shown that only a small proportion of people who discontinue their initial treatment switch to another medication.[12, 13]

Currently, it is not clear what predicts pharmacotherapy choice for NeuP, or adherence and persistence with treatment. To our knowledge, no published reviews have summarised the literature on the predictors of any of these outcomes. Identifying these predictors may inform strategies to ensure that people with NeuP are prescribed recommended medication and adhere to and persist with their treatment when appropriate. For example, understanding which patient characteristics or healthcare factors are associated with these outcomes could help clinicians target interventions, such as tailored patient education, closer follow-up, or system-level changes to reduce disparities and improve long-term treatment outcomes. Therefore, we carried out a systematic review and meta-analysis to identify predictors of pain medication prescribing, adherence, and discontinuation in adults with NeuP.

Methods

<u>Protocol and registration</u>

The protocol of this systematic review and meta-analysis was prospectively registered on the PROSPERO international prospective register of systematic reviews (Registration ID: CRD42023464307). This report follows the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) checklist.[14]

<u>Search strategy</u>

To identify relevant studies, Embase, PubMed, Web of Science, and CINAHL Plus were searched from inception to the 1st of August 2025. The search strategy combined keywords and medical subject headings for 1) neuropathic pain 2) pharmacological treatments and 3) prescribing, adherence, and discontinuation. The full search terms for all the databases are reported in Table S1. The retrieved records were exported to Covidence (Veritas Health Innovation, Melbourne, Australia) and screened independently by two reviewers (MK and DS). Discrepancies were resolved by discussion with the wider review team where necessary. The reference lists of the included studies were reviewed to identify further eligible studies not identified by the search strategy.

<u>Eligibility criteria</u>

Studies were included if they: (1) investigated potential predictors of pain medication prescribing, adherence, or discontinuation in adults (≥18 years old) with NeuP; (2) were a cohort study, case-control study, cross-sectional study, randomised controlled trial (RCT), or systematic review with meta-analysis; (3) were published in full-text in English. Studies were excluded if (1) they included participants who did not have NeuP, where these could not be analysed separately from those who had NeuP; (2) they included children (<18 years old) and did not present analysis on adults separately; (3) they were a case study, case series, review without a meta-analysis, protocol, editorial, or conference abstract; (4) they were an RCT that aimed to assess NeuP medication efficacy and the only relevant information for our review was discontinuation rates between trial medications; (5) the medications of our interest could not be analysed separately from other medications; and/or (6) they compared pain medications before and after surgery.

It should be noted that although the diagnosis of interest in this review was NeuP, we also included studies that defined NeuP using diagnostic codes for neuropathy, as diagnostic codes used to indicate NeuP often cannot separate painful and painless neuropathy. Our assumption was that relevant pain medicines were only prescribed for neuropathy in these studies when the neuropathy was painful - though we could not rule out the possibility of pain medicines given for other painful conditions or indications, in the presence of a painless neuropathy. We focused on peripheral neuropathic pain conditions, but did not exclude studies where central neuropathic pain conditions could not be separated from peripheral neuropathic pain conditions. The method of diagnosis used in each study is provided in Table S2. For the above discussed reason and because of the inconsistency of terms used for NeuP caused by diabetes, the following terms used in included papers,

were grouped under diabetic neuropathy (DN): diabetic peripheral neuropathic pain (DPNP), diabetic peripheral neuropathy (DPN), painful diabetic peripheral neuropathy (pDPN or PDPN), painful diabetic neuropathy (PDN), painful diabetic polyneuropathy, diabetic polyneuropathy.

The medication categories of interest were anticonvulsants, antidepressants, opioids, non-opioid analgesics, combinations of opioids and non-opioid analgesics, local anaesthetics, N-methyl-D-aspartate (NMDA) receptor antagonists, topical pain medications, botulinum toxin A, baclofen, and clonidine. The medications that were not of interest included but were not limited to muscle relaxants, corticosteroids, sedatives, and antimigraine medications. A summary of medications included and excluded from our review is provided in Table S3, while Table S4 provides medication included and excluded in each study. In addition, Table S5 shows medications considered recommended in the included studies. In our review, the word 'non-opioid analgesics' refers to non-steroidal anti-inflammatory drugs (NSAIDs), paracetamol, aspirin, and nefopam. This categorisation was used as some of the included studies categorised their medications as antidepressants, anticonvulsants, opioids, local anaesthetics, and non-opioid analgesics, without precisely defining the 'non-opioid analgesics' category.[15–18]

Data extraction

Data extraction was performed independently by two reviewers (MK and DS) and discrepancies were resolved by discussion. Study characteristics extracted included: author, year of publication, type of outcome(s) investigated, study design, country and time frame of data analysed, sample, size, proportion of females in the sample, mean age, NeuP diagnoses included in the study, and key findings.

Risk of bias assessment

Risk of bias assessment was performed independently by two reviewers (MK and DS) and discrepancies were resolved by discussion. The Risk Of Bias In Non-randomized Studies - of Exposures (ROBINS-E)[19] tool was used to assess cohort studies, cross-sectional studies, and case-control studies. The "exposure" in each study was defined as the potential predictors investigated (e.g. NeuP diagnosis, sex, household income). The revised Cochrane Risk of Bias tool for randomized trials (RoB 2)[20] was used to assess three RCTs. A secondary analysis of RCTs was assessed using A MeaSurement Tool to Assess systematic Reviews 2 (AMSTAR 2).[21]

Synthesis of results

The included studies were grouped into non-mutually exclusive categories based on the outcomes they investigated (prescribing, adherence, and discontinuation). The studies that investigated prescribing were further divided into non-mutually exclusive subgroups based on which medications were examined and how the analyses were conducted. For each category of studies, we listed all the potential predictors investigated and summarised the

evidence for each potential predictor. If the potential predictor definition and outcome measurement were similar or could be converted to be similar, meta-analyses were conducted using RStudio (v4.3.1). Statistical heterogeneity was described using the I^2 test. Random-effects models were used for all meta-analyses. The type of measure of effect estimated in each meta-analysis depended on the type of data provided in the included studies for the potential predictors. In the majority of cases, odds ratios (OR) and the associated 95% confidence intervals (CI) were calculated. In meta-analyses that were conducted using pre-calculated ORs, adjusted ORs were used rather than unadjusted ORs if possible. Finally, each potential predictor was categorised either to increase the likelihood of the investigated outcome, to not have an association with it, or to unclear or inconsistent results. Due to the extensive quantity of data generated, in this manuscript we have focused on describing the predictors that we considered to be the most useful to the readers and/or clinicians.

Results

<u>Study selection</u>

Figure 1 shows a flowchart of the study selection process. Database searches identified a total of 11,255 records. Sixty-nine studies were included in the review, and this included 6 studies identified from screening the reference lists of the eligible studies. The records excluded in full-text screening and the reasons for their exclusion are documented in Table S7.

<u>Study characteristics</u>

The characteristics of the included studies are summarised in Table 1. The majority of the included studies were retrospective cohort studies (44/69) conducted in North America and Europe (58/69). There were 53 studies on prescribing, 15 on adherence, and 27 on discontinuation. The division of the prescribing studies into different subcategories is shown in Table S7. Table S8 and Table S9 explain how adherence and discontinuation was assessed in each study. Most studies included people with diabetic neuropathy (DN) along with other aetiologies (55/69) – with 24/69 studies exclusively on DN.

<u>Risk of bias assessment</u>

The majority of the studies (65/69) were assessed using ROBINS-E. The results of this assessment are shown in Table S10. All of the studies assessed using ROBINS-E were deemed to have a low risk of bias in measuring exposures, post-exposure interventions, outcome measurement, and reporting of results (although the domain "post-exposure interventions" was generally irrelevant to our research question). Sixteen studies were deemed to have a high risk of bias in "confounding variables" due to adjusting for potential predictors that could affect each other. In the participant selection domain, most of the studies had "some concerns" as the exposure window of some of the investigated potential predictors would be lifelong (e.g. age or sex) and thus would precede the data collection time frame. However, these factors were unlikely to influence our interpretation of these studies. In the missing data domain, in many of the studies with a high risk of bias, no information was provided about the handling of missing data. Three RCTs were assessed using Rob 2. The results are shown in Table S11. In these studies, the only finding of a high risk of bias was in the missing data domain in one of the studies[22] as there was no information about missing data. One study, a non-systematic analysis of three RCTs, [23] was assessed using AMSTAR 2 and was concluded to have a critically low (poor) overall rating. The results for each domain are shown in Table S12.

<u>Summary of results</u>

The analyses of all potential predictors, including meta-analysis results, are provided Tables S13 to S20. A summary of these results is provided in Table S21, which categorises all potential predictors to either to increase the likelihood of the investigated outcome, to not have an association with it, or to unclear or inconsistent results. The key findings are summarised in Table 2, which provides examples of association strength between the predictor and the outcome. Other relevant results for the key predictors are described below.

It should be noted that studies that investigated potential predictors of NeuP medication prescribing often did not report their analysis such that the participants were exclusively prescribed the medication that was analysed and no other pain medications. For example, if potential predictors of being prescribed recommended medication were analysed, it was not reported whether these people received other pain medications.

<u>Receiving recommended NeuP medication</u>

The key predictors for receiving recommended NeuP medication were having a mental health disorder (in particular anxiety and/or depression), diabetes, white ethnicity, and/or purely neuropathic pain rather than mixed pain.

Mental health disorders as predictors for receiving recommended medication were investigated in four studies. Lakkad and colleagues (2023) found that people with a mental health disorder (a variety of diagnoses included) were more likely to receive recommended NeuP medication (OR: 3.57, 95% CI: 3.12 to 4.09).[24] Boulanger and colleagues (2009) (N=27,779) found that people with anxiety and/or depression were more likely to receive recommended medication (OR: 1.74, 95% CI: 1.63 to 1.86).[25] Callaghan and colleagues (2019) and Chahine and Al Souheil (2021) investigated anxiety and depression separately, and meta-analyses based on their results found that people with depression were more likely to receive recommended medication (OR: 1.62, 95% CI: 1.33 to 1.98), while the result for anxiety was no significant (OR: 0.84, 95% CI: 0.20 to 3.59).[8, 26] It is worth noting that the result from Chahine and Al Souheil (2021) on anxiety was the only non-significant result, and their study was much smaller (N=360) than the study by Callaghan and colleagues (2019) (N=14,426).

Eight studies investigated whether having diabetes or diabetic neuropathy changed the likelihood of receiving recommended NeuP medication. A meta-analysis based on three studies investigating diabetes found that people with diabetes were more likely to receive recommended medication (OR: 1.41, 95% CI: 1.22 to 1.62).[8, 24, 26] The results relating to DN compared to other NeuP aetiologies were mixed, although the results generally showed that either people with DN were more likely to receive recommended medication or that there was no difference between the aetiologies.[15, 16, 26–29] Only people with TN were found to be more likely to receive recommended medication than people with DN.[15]

Ethnicity as a potential predictor for receiving recommended medication was investigated in three studies. Lakkad and colleagues (2023) (N=7,116) found that people with white ethnicity were more likely to receive recommended medication compared to people with

black ethnicity (OR: 1.76, 95% CI: 1.43 to 1.75) or any other ethnicity (OR: 1.37, 95% CI: 1.07 to 1.75).[24] In contrast, Reynolds and colleagues (2020a) (N=202) found no differences based between these categories.[22] Callaghan and colleagues (2019) found that people with white ethnicity were more likely receive recommended medication compared to people with Asian (OR: 1.64, 95% CI: 1.22 to 2.17) or Hispanic ethnicity (OR: 1.32, 95% CI: 1.12 to 1.52), but found no difference between white and black ethnicity (OR: 1.11, 95% CI: 0.97 to 1.27).[8] A meta-analysis contrasting white and non-white groups found that people with ethnicity were more likely to receive recommended medication (OR: 1.40, 95% CI: 1.18 to 1.66).

Receiving any pain medication

The key predictors for receiving any pain medication were higher pain severity, having anxiety and/or depression, and/or younger age (among people aged >50 years).

Sadosky and colleagues (2013) (N=112) investigated how pain severity measured using the Brief Pain Inventory (BPI) influenced the likelihood of receiving any pain medication.[30] They found that the proportion of people with medication increased as pain severity increased (54.5% in mild pain, 84.2% in moderate pain, 96.9% in severe pain). In this small sample, the difference between severe pain and mild pain was very large (OR: 25.83, 95% CI: 2.97 to 225.21). Another small study investigated the differences in pain intensity measured using a 0-10 numerical rating scale (NRS).[31] In this study, Pillay and colleagues (2015) (N=130) found that the median pain intensity in the group with medication was significantly higher than in the group without medication [Median NRS (IQR): 10 (2-10) vs 6 (2-10)].[31]

Two studies investigated whether having a mental health disorder changed the likelihood of receiving any pain medication. Boulanger and colleagues (2009) (N=27,779) found that people with anxiety and/or depression were more likely to receive any pain medication (OR: 3.66, 95% CI: 3.14 to 4.26).[25] Mbrah and colleagues (2022) investigated these separately and found similar results (adjusted prevalence ratio for receiving medication, for depression: 1.47, 95 % CI: 1.43 to 1.52; for anxiety: 1.20, 95% CI: 118 to 1.23).[32]

The differences in the likelihood of receiving any pain medication based on age were investigated by Mbrah and colleagues (2022) and Pillay and colleagues (2015). Mbrah and colleagues (2022) (N=69,160) investigated people in nursing homes who were over 50 years old. They compared different age categories and found that younger people were more likely to receive medication (50-64 years vs +85, prevalence ratio: 1.45, 95% CI: 1.38 to 1.52).[32] Pillay and colleagues (2015) (N=130) found no difference in age between the groups that received vs did not receive medication [mean (SD) age in years: 45.4 (9.3) vs 46.2 (9.8), p=0.63].[31] The differing results may be due to differences in the ages analysed or due to the small sample size of the latter study.

Adherence

The key predictors for having better adherence to NeuP medications were taking SNRIs rather than gabapentinoids or TCAs, having dose titration, brand-name medication, a

medicine reminder, lower medication cost to the patient (USA), diabetic neuropathy rather than postherpetic neuralgia, older age (in below 75-year-olds), and/or white ethnicity. Most of the adherence studies measured the outcome using medication possession ratio (MPR) or proportion of days covered (PDC) [See Table S6].

Six studies investigated the differences in adherence between NeuP medications.[9, 13, 33–36] Reynolds and colleagues (2020b) had the largest sample (N=57,495) and longest duration of follow-up (12 years for this data).[9] They found that the mean MPR for gabapentinoids was around 40-50% and for SRNIs 50-60%.[9] Three other studies compared SNRIs and gabapentinoids and found better adherence to SNRIs.[13, 33, 34] Gharibian and colleagues (2013) found a large difference in adherence between venlafaxine and gabapentin (OR: 2.83, 95% CI: 1.45 to 5.76).[33] Two studies compared adherence between SNRIs and TCAs. Oladapo and colleagues (2013) found that mean MPR in people taking duloxetine was 85.6% while the mean MPR in people taking TCAs was 76.0% (n=652 vs n=931, p<0.0001).[34] Similarly, Gharibian and colleagues (2013) found that the proportion of people with ≥ 80% MPR was 69.4% in people taking venlafaxine and 44.4% in people taking TCAs (n=49 vs n=1208, p<0.01).[33]

The difference in adherence between groups receiving or not receiving dose titration was investigated in two studies. Meta-analysis of these studies found that people who received any dose titration within 45 days of initiation their medication had better adherence compared to people with no dose titration (OR: 2.20, 95% CI: 1.51 to 3.21).[37, 38]

Five studies examined adherence based on age. Wu and colleagues (2009) found that people aged 55-64 years had better adherence than people aged 18-44 years (OR: 2.12, 95% CI: 1.30 to 3.46), while there was no difference between people aged 45-54 years and 18-44 years (OR: 1.63, 95% CI: 0.99 to 2.71).[39] The other studies that investigated age supported that increasing age was associated with better adherence.[9, 37, 38, 40] Wu and colleagues (2009) were the only ones to examine adherence in people aged 65 years and older.[39] They found that people aged 65-74 years had better adherence than people aged 75-84 years (OR: 1.33, 95% CI: 1.01 to 1.79).[39] Thus, increasing age seems to be associated with better adherence until the age of 75 years, after which it may decline.

Two studies from Spain found that people with brand-name gabapentin or pregabalin had better adherence compared to people with generic name gabapentin or pregabalin (Mean MPR MD: 0.05, 95% CI: 0.03 to 0.07).[18, 41] In contrast, a study from the USA found that lower medication cost to patient was associated with better adherence (IRR: 1.09, 95% CI: 1.03 to 1.14 when +$50 for 30-day supply).[9]

Discontinuation

The key predictors for being more likely to discontinue NeuP medications were taking TCAs rather than other antidepressants, having generic-name medication rather than brand-name medication, combination pharmacotherapy for NeuP rather than monopharmacotherapy, not experiencing clinically significant pain relief from the medication, experiencing moderate or severe adverse effects from the medication, having postherpetic neuralgia rather than diabetic neuropathy, and/or pain-related worrying.

Only two eligible studies investigated the clinical effectiveness of NeuP medications or their adverse effects as potential predictors of discontinuation. Sicras-Mainar and colleagues (2019)[41] (N=4,860) found that people who did not experience clinically meaningful pain relief measured using a visual analogue scale (≥4 out of 0-10 points scale) were more likely to discontinue pregabalin (OR: 1.2, 95% CI: 1.0 to 1.4). Shaparin and colleagues (2015)[23] (N= 556) found that people with PHN taking gabapentin were more likely to discontinue if they experienced moderate (OR: 8.64, 95% CI: 4.42 to 16.89) or severe adverse events (OR: 5.37, 95% CI: 2.01 to 14.00).

The differences in therapy duration between medications were investigated by eleven studies. Most of these compared individual medications (e.g. pregabalin vs duloxetine). Two studies provided useful comparisons between different medication classes. Hall and colleagues (2008)[15] found that people taking TCAs were more likely to not have stable pharmacotherapy after one year compared to people using other types of antidepressants (OR: 1.56, 95% CI: 1.01 to 2.42). Similarly, Gore and colleagues (2007a)[42] found that people taking TCAs had fewer mean days with pharmacotherapy compared to people with SNRIs (MD -12, 95% CI: -21 to -3) or SSRIs (MD -35, 95% CI: -40 to -30). Heatmaps showing the pairwise meta-analyses between all the investigated medications in these studies are shown in Figures S1 and S2.

Four studies compared pharmacotherapy durations between people with PHN and people with DN. Our meta-analysis found that people with PHN were more likely to discontinue sooner than people with DN (MD in treatment days: -51, 95% CI: -4 to -98). Three of the included studies[15, 16, 43] found a mean difference in treatment days to be around -20 to -30 days, while Wang and colleagues (2020)[44] found a mean difference of -133 days (95% CI: 126 to 139).

Discussion

Our systematic review and meta-analysis aimed to understand the predictors of NeuP pharmacotherapy prescribing, adherence, and discontinuation. In the 69 included studies, prescribing was analysed in 53 studies, adherence in 14 studies, and discontinuation in 27 studies. They key predictors for receiving recommended medication were having diabetes and/or white ethnicity. People with SNRIs, dose titration and/or medicine reminder were more likely to have better adherence. Pain-related worrying and/or combination pharmacotherapy were associated with shorter treatment durations.

The association between diabetes and receiving recommended medication is not unexpected. There are many potential contributing factors to this. Diabetes is the most common cause for peripheral neuropathic pain and is widely recognized in the clinical and research communities.[45] People with diabetes usually have regular appointments for diabetes management, which increases the chance of neuropathic symptoms being detected and opportunities for medication review.

Similarly, ethnic differences in prescription medicines have been noted in Western countries for a long time.[46, 47] Socioeconomic status is often considered to be the key explanatory variable between health care and ethnicity.[48] However, in many cases disparities by ethnicity cannot be explained by socioeconomic status or other factors.[48] In a key study included in our review, Callaghan and colleagues (2019) found that people with white ethnicity were more likely to receive recommended medication compared to people with Asian or Hispanic ethnicity despite controlling for education, household income, or insurance type.[8]

In the current IASP NeuPSIG pharmacotherapy guidelines (which are based on results from RCTs) SNRIs, TCAs, and gabapentinoids are recommended as first-line treatments.[6] The number needed to treat (low indicating better efficacy) is lower for TCAs (4.6, 95% CI: 3.2 to 7.7) compared to that of SNRIs (7.4, 95% CI: 5.6 to 10.9) or gabapentinoids (8.9, 95% CI: 7.4 to 1.11).[6] The number needed to harm (low indicating more risk) is lowest for SNRIs (13.9, 95% CI: 10.9 to 19.0), a little bit higher for TCAs (17.1, 95% CI: 11.4 to 33.6), and considerably higher for gabapentinoids (26.2, 95% CI: 20.4 to 36.5).[6] It is also noteworthy that TCAs are not recommended for older adults due their anticholinergic and sedative side effects. Interestingly, the studies included in our review (mostly epidemiological), found that adherence and persistence with TCAs was lower compared to SNRIs.[15, 33, 34, 42] In addition, adherence to SNRIs was better compared to adherence to gabapentinoids.[9, 13, 33–35] Reynolds and colleagues (2020b) were a key study for this due to their large sample (n=57,495).[9] Since these results come from epidemiological studies, it is possible that these results are confounded by variables that are not evident. On the other hand, RCTs cannot provide real life evidence, in particular for outcomes like adherence or persistence.

Dose titration and combination pharmacotherapy, which were identified as key predictors for adherence and discontinuation respectively, have been discussed in the IASP NeuPSIG pharmacotherapy guideline.[6] The studies included in our review found that people who received any dose titration within 45 days of medication initiation were more likely to adherence to pregabalin or mirogabalin (OR: 2.40, 95% CI: 1.51 to 3.21).[37, 38] Dose titration has been taken in account in the guideline, as the key recommendations

table suggests that systemic drugs should be initiated at low doses and titrated slowly.[6] Regarding combination pharmacotherapy, the guideline concluded that there was insufficient evidence to recommend any combination. This included one systematic review and meta-analysis which found no difference in efficacy or safety between combinations and single medications.[49] In our review, one study (n=7,566) found that people with combination pharmacotherapy were more likely to discontinue compared to people with monopharmacotherapy.[50]

In addition to the key predictors identified for NeuP medication discontinuation, we noted that in most studies the majority of people discontinued their treatment within half a year regardless of having predictors for longer treatment duration. None of the included studies investigated potential predictors for the minority that continued treatment for a long time. Yang and colleagues (2015)[13] and Dworkin and colleagues (2012)[12] found that only a small proportion of people who discontinued their initial treatment switched to another one. It is possible that NeuP pharmacotherapy is discontinued due to the resolution of pain, for example in people with PHN,[51] but this is unlikely to be the case for people with diabetic neuropathic pain, whose pain is highly likely to persist and exacerbate.[52]

Limitations

It should be noted that we investigated associations, meaning that the predictors identified in our review do not provide information about causality. Although some studies provided results adjusted for many confounding variables, unidentified confounding variables may be present in these results.

Although this review identified a good number of studies, investigating many different potential predictors, the overlap of potential predictors investigated between the studies was limited. Thus, the conclusions about each potential predictor were generally based on a small number of studies. Only two eligible studies investigated the clinical effectiveness of NeuP medications and their adverse effects as potential predictors of discontinuation. It is interesting that our review did not identify any other studies specifically investigating adverse events or clinical effectiveness as potential predictors of discontinuation although they are key aspects of successful drug development as well as pharmacoepidemiology.

Studies that investigated potential predictors of NeuP medication prescribing often did not report their analysis such that the participants were exclusively prescribed the medication that was analysed and no other pain medications. For example, if potential predictors of being prescribed recommended medication were analysed, it was not reported whether these people received other pain medications. Analysing the predictors of prescribing this way provides useful binary categorisation between people with recommended medication and people without recommended medication. However, this simple categorisation misses other treatments which could be informative about real life prescribing. In future studies, it would be worth exploring whether there are differences between people who exclusively receive recommended medications and people with recommended medication and additional pain medications.

Although the diagnosis of interest in this review was NeuP, most of the included studies were not able to differentiate between painful and painless neuropathy based on the

information provided in the databases they investigated. Studies often defined their sample using International Classification of Diseases (ICD) 9 or 10 codes and pain medication claims, but it was only in ICD-11 that a specific code for neuropathic pain was developed.[53] The lack of information about the presence and level of pain in the included studies creates a major limitation in identifying NeuP pharmacotherapy predictors in those with the most need for treatment. Future studies could be improved by seeking additional data sources that allow classifying the sample into mild, moderate, and severe pain categories.

Another limitation in our work is that our analysis focused on finding an overall association for each potential predictor and thus we did not seek associations within the potential predictors. It is possible that some predictors are context-specific (e.g. specific to a type of NeuP diagnosis or medication) and more detailed conclusions may arise later with more studies. Analysing within-predictor associations was not considered worthwhile in this review due to the small number of studies per each potential predictor.

We made a pragmatic decision to exclude RCTs that are primarily designed to assess the efficacy of NeuP medications from our review. The medications given in these trials could be compared as the potential predictors for discontinuation. However, as our review aims to summarise all the potential predictors for NeuP medication prescribing, adherence, and discontinuation; including these trials in our review would focus disproportionately on one type of predictor (>200 RCTs [7]). We do not believe that the inclusion of this group of studies would materially change the conclusions derived from our review.

Conclusion

Our review has identified useful predictors of pain medication prescribing, adherence, and discontinuation in adults with NeuP. The results show that there is a need to focus on improving treatment allocation for people without diabetes and to non-white populations. Adherence may be improved by using dose titration and/or a medicine reminder. Monopharmacotherapy rather than combination pharmacotherapy for NeuP may result in better treatment persistence. These, and the other factors we identified should be considered in the development of future NeuP pharmacotherapy guidelines and in policy making. Future research should investigate how treatment effectiveness and adverse effects are associated with adherence and persistence.

Details of authors' contributions

Study conception and design: MK, LC, BS, HH

Acquisition of data: MK, DS

Analysis and interpretation of data: MK, DS, BS, HH, LC

Drafting of final manuscript: MK

Revision and critical appraisal of the manuscript: MK, BS, HH, DS, LC

Final approval of the submitted manuscript: MK, BS, HH, DS, LC

Acknowledgments

We would like to acknowledge Mr Scott McGregor (University of Dundee Library Academic and Cultural Services) for his assistance in developing the search strategy.

Declaration of interest

This systematic review was conducted as part of MK's PhD studentship funded by The National Institute of Academic Anaesthesia [British Journal of Anaesthesia / Royal College of Anaesthetists, WKR0-2022-0028]. DS is a fellow on the Multimorbidity Doctoral Training Programme for Health Professionals, which is supported by the Wellcome Trust [223499/Z/21/Z]. LC receives research support funding, on behalf of her institution, from the Advanced Pain Discovery Platform (funded by UK Research and Innovation, Versus Arthritis, Eli Lilly), the Scottish Government (Chief Scientist Office), The Wellcome Trust, and the National Institute of Academic Anaesthesia. She is Vice Chair of Scottish Intercollegiate Guidelines Network (SIGN) Council and is currently chairing the SIGN Guideline Development Group for Management of Chronic Pain. BHS has received research support funding, on behalf of his institution, from UK Research and Innovation (Medical Research Council), Versus Arthritis, the Scottish Government (Chief Scientist Office), Eli Lilly, The Wellcome Trust, and the National Institute of Academic Anaesthesia. HLH has received funding to support research from The National Institute of Academic Anaesthesia [British Journal of Anaesthesia / Royal College of Anaesthetists].

Funding

This systematic review was conducted as part of MK's PhD studentship funded by The National Institute of Academic Anaesthesia [British Journal of Anaesthesia / Royal College of Anaesthetists, WKR0-2022-0028].

References

1. Bouhassira D, Attal N, Alchaar H, Boureau F, Brochet B, Bruxelle J, et al. Comparison of pain syndromes associated with nervous or somatic lesions and development of a new neuropathic pain diagnostic questionnaire (DN4). *Pain* 2005; **114**: 29–36

2. Treede R-D, Jensen TS, Campbell JN, Cruccu G, Dostrovsky JO, Griffin JW, et al. Neuropathic pain: redefinition and a grading system for clinical and research purposes. *Neurology* 2008; **70**: 1630–5

3. Baskozos G, Hébert HL, Pascal MM, Themistocleous AC, Macfarlane GJ, Wynick D, et al. Epidemiology of neuropathic pain: an analysis of prevalence and associated factors in UK Biobank. *Pain Rep* 2023; **8**: e1066

4. van Hecke O, Austin SK, Khan RA, Smith BH, Torrance N. Neuropathic pain in the general population: A systematic review of epidemiological studies. *PAIN* 2014; **155**: 654

5. Langley PC, Van Litsenburg C, Cappelleri JC, Carroll D. The burden associated with neuropathic pain in Western Europe. *J Med Econ* 2013; **16**: 85–95

6. Soliman N, Moisset X, Ferraro MC, Andrade DC de, Baron R, Belton J, et al. Pharmacotherapy and non-invasive neuromodulation for neuropathic pain: a systematic review and meta-analysis. *Lancet Neurol* Elsevier; 2025; **24**: 413–28

7. Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015; **14**: 162–73

8. Callaghan BC, Reynolds E, Banerjee M, Kerber KA, Skolarus LE, Burke JF. Longitudinal pattern of pain medication utilization in peripheral neuropathy patients. *Pain* 2019; **160**: 592–9

9. Reynolds EL, Burke JF, Banerjee M, Kerber KA, Skolarus LE, Magliocco B, et al. Association of out-of-pocket costs on adherence to common neurologic medications. *Neurology* 2020; **94**

10. Burnier M. Is There a Threshold for Medication Adherence? Lessons Learnt From Electronic Monitoring of Drug Adherence. *Front Pharmacol* Frontiers; 2019; **9**

11. Gustavsson A, Bjorkman J, Ljungcrantz C, Rhodin A, Rivano-Fischer M, Sjolund K-F, et al. Pharmacological Treatment Patterns in Neuropathic Pain—Lessons from Swedish Administrative Registries. *Pain Med* 2013; **14**: 1072–80

12. Dworkin RH, Panarites CJ, Armstrong EP, Malone DC, Pham SV. Is treatment of postherpetic neuralgia in the community consistent with evidence-based recommendations? *Pain* 2012; **153**: 869–75

13. Yang M, Qian C, Liu Y. Suboptimal Treatment of Diabetic Peripheral Neuropathic Pain in the United States. *Pain Med* 2015; **16**: 2075–83

14. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* British Medical Journal Publishing Group; 2021; **372**: n160

15. Hall GC, Carroll D, McQuay HJ. Primary care incidence and treatment of four neuropathic pain conditions: A descriptive study, 2002–2005. *BMC Fam Pract* 2008; **9**: 26

16. Hall GC, Morant SV, Carroll D, Gabriel ZL, McQuay HJ. An observational descriptive study of the epidemiology and treatment of neuropathic pain in a UK general population. *BMC Fam Pract* 2013; **14**: 28

17. Reed C, Hong J, Novick D, Lenox-Smith A, Happich M. Incidence of diabetic peripheral neuropathic pain in primary care – a retrospective cohort study using the United Kingdom General Practice Research Database. *Pragmatic Obs Res* Dove Press; 2013; **4**: 27–37

18. Sicras-Mainar A, Rejas J, Navarro-Artieda R. Comparative effectiveness and costs of generic and brand-name gabapentin and venlafaxine in patients with neuropathic pain or generalized anxiety disorder in Spain. *Clin Outcomes Res* 2015; 299

19. Higgins JPT, Morgan RL, Rooney AA, Taylor KW, Thayer KA, Silva RA, et al. A tool to assess risk of bias in non-randomized follow-up studies of exposure effects (ROBINS-E). *Environ Int* 2024; **186**: 108602

20. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* British Medical Journal Publishing Group; 2019; **366**: l4898

21. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* British Medical Journal Publishing Group; 2017; **358**: j4008

22. Reynolds EL, Burke JF, Banerjee M, Callaghan BC. Randomized controlled trial of a clinical decision support system for painful polyneuropathy. *Muscle Nerve* 2020; **61**: 640–4

23. Shaparin N, Slattum PW, Bucior I, Nalamachu S. Relationships Among Adverse Events, Disease Characteristics, and Demographics in Treatment of Postherpetic Neuralgia With Gastroretentive Gabapentin. *Clin J Pain* 2015; **31**: 983–91

24. Lakkad M, Martin B, Li C, Harrington S, Dayer L, Painter JT. Factors Associated With Guideline-Concordant Pharmacological Treatment for Neuropathic Pain Among Breast Cancer Survivors. *Clin Breast Cancer* 2023; **23**: 598–619

25. Boulanger L, Zhao Y, Foster TS, Fraser K, Bledsoe SL, Russell MW. Impact of comorbid depression or anxiety on patterns of treatment and economic outcomes among patients with diabetic peripheral neuropathic pain. *Curr Med Res Opin* 2009; **25**: 1763–73

26. Chahine B, Al Souheil F. Dispensing patterns of drugs used for neuropathic pain and adherence to NeuPSIG guideline: an observational study. *Egypt J Neurol Psychiatry Neurosurg* 2021; **57**: 142

27. Udall M, Kudel I, Cappelleri JC, Sadosky A, King-Concialdi K, Parsons B, et al. Epidemiology of physician-diagnosed neuropathic pain in Brazil. *J Pain Res* 2019; **Volume 12**: 243–53

28. Butler S, Eek D, Ring L, Gordon A, Karlsten R. The utility/futility of medications for neuropathic pain – an observational study. *Scand J Pain* De Gruyter; 2019; **19**: 327–35

29. Margolis J, Princic N, Smith D, Abraham L, Cappelleri J, Shah S, et al. Development of a novel algorithm to determine adherence to chronic pain treatment guidelines using administrative claims. *J Pain Res* 2017; **Volume 10**: 327–39

30. Sadosky A, Schaefer C, Mann R, Bergstrom F, Baik R, Parsons B, et al. Burden of illness associated with painful diabetic peripheral neuropathy among adults seeking treatment in the US: results from a

retrospective chart review and cross-sectional survey. *Diabetes Metab Syndr Obes Targets Ther* 2013;
79

31. Pillay P, Wadley AL, Cherry CL, Karstaedt AS, Kamerman PR. Pharmacological treatment of painful HIV-associated sensory neuropathy. *S Afr Med J* 2015; **105**: 769–72

32. Mbrah AK, Nunes AP, Hume AL, Zhao D, Jesdale BM, Bova C, et al. Prevalence and treatment of neuropathic pain diagnoses among U.S. nursing home residents. *Pain* 2022; **163**: 1370–7

33. Gharibian D, Polzin JK, Rho JP. Compliance and Persistence of Antidepressants Versus Anticonvulsants in Patients With Neuropathic Pain During the First Year of Therapy. *Clin J Pain* 2013; **29**: 377–81

34. Oladapo AO, Barner JC, Rascati KL, Strassels SA. A Retrospective Database Analysis of Neuropathic Pain and Oral Antidiabetic Medication Use and Adherence Among Texas Adults With Type 2 Diabetes Enrolled in Medicaid. *Clin Ther* 2012; **34**: 605–13

35. Zhao Y, Sun P, Watson P. Medication adherence and healthcare costs among patients with diabetic peripheral neuropathic pain initiating duloxetine versus pregabalin. *Curr Med Res Opin* 2011; **27**: 785–92

36. Giannopoulos S, Kosmidou M, Sarmas I, Markoula S, Pelidou S-H, Lagos G, et al. Patient Compliance With SSRIs and Gabapentin in Painful Diabetic Neuropathy. *Clin J Pain* 2007; **23**: 267–9

37. Kato K, Kodama S, Shiosakai K, Kimura T. Relationship between the dose titration and adherence of mirogabalin in patients with peripheral neuropathic pain depending on renal function: a nationwide electronic medical record database study. *Expert Opin Pharmacother* 2023; **24**: 267–82

38. Yeh Y-C, Cappelleri JC, Marston XL, Shelbaya A. Effects of dose titration on adherence and treatment duration of pregabalin among patients with neuropathic pain: A MarketScan database study. Mallhi TH, editor. *PLOS ONE* 2021; **16**: e0242467

39. Wu N, Chen S, Boulanger L, Fraser K, Bledsoe SL, Zhao Y. Duloxetine compliance and its association with healthcare costs among patients with diabetic peripheral neuropathic pain. *J Med Econ* 2009; **12**: 192–202

40. Chen S-Y, Wu N, Boulanger L, Fraser KA, Zhao Y. The Relationship Between Average Daily Dose, Medication Adherence, and Health-Care Costs among Diabetic Peripheral Neuropathic Pain Patients Initiated on Duloxetine Therapy: Duloxetine Average Daily Dose, Adherence and Costs. *Pain Pract* 2010; **10**: 530–9

41. Sicras-Mainar A, Rejas-Gutiérrez J, Pérez-Paramo M, Sánchez-Alvarez L, Navarro-Artieda R, Darbà J. Consequences on economic outcomes of generic versus brand-name drugs used in routine clinical practice: the case of treating peripheral neuropathic pain or generalized anxiety disorder with pregabalin. *Expert Rev Pharmacoecon Outcomes Res* 2019; **19**: 45–57

42. Gore M, Dukes E, Rowbotham DJ, Tai K, Leslie D. Clinical characteristics and pain management among patients with painful peripheral neuropathic disorders in general practice settings. *Eur J Pain* 2007; **11**: 652–64

43. Hall GC, Carroll D, Parry D, McQuay HJ. Epidemiology and treatment of neuropathic pain: The UK primary care perspective. *Pain* 2006; **122**: 156–62

44. Wang Y-F, Chen Y-T, Tsai C-W, Yen Y-C, Chen Y-C, Shia B-C, et al. Persistence of pregabalin treatment in Taiwan: a nation-wide population-based study. *J Headache Pain* 2020; **21**: 54

45. Lehmann HC, Wunderlich G, Fink GR, Sommer C. Diagnosis of peripheral neuropathy. *Neurol Res Pract* 2020; **2**: 20

46. Morgan S, Hanley G, Cunningham C, Quan Q. Ethnic differences in the use of prescription drugs: a cross-sectional analysis of linked survey and administrative data. *Open Med Peer-Rev Indep Open-Access J* Open Med; 2011; **5**

47. Fiscella K, Franks P, Doescher MP, Saver BG. Disparities in Health Care by Race, Ethnicity, and Language Among the Insured: Findings From a National Sample. *Med Care* 2002; **40**: 52

48. Mayberry RM, Mili F, Ofili E. Racial and ethnic differences in access to medical care. *Med Care Res Rev MCRR* Med Care Res Rev; 2000; **57 Suppl 1**

49. Balanaser M, Carley M, Baron R, Finnerup N, Moore R, Rowbotham M, et al. Combination pharmacotherapy for the treatment of neuropathic pain in adults: systematic review and meta-analysis. *Pain* 2023; **164**

50. Kuo K-L, Brixner D, Lipman AG, Goodman M, Hung M, Oderda GM. Single- Versus Multiple-Drug Pharmacotherapy in the Management of Diabetic Painful Neuropathy. *J Pain Palliat Care Pharmacother* 2016; **30**: 184–94

51. Reda H, Greene K, Rice FL, Rowbotham MC, Petersen KL. Natural history of herpes zoster: Late follow-up of 3.9 years (n = 43) and 7.7 years (n = 10). *PAIN®* 2013; **154**: 2227–33

52. Boulton AJM, Armstrong WD, Scarpello JHB, Ward JD. The natural history of painful diabetic neuropathy—a 4-year study. *Postgrad Med J* 1983; **59**: 556–9

53. Scholz J, Finnerup NB, Attal N, Aziz Q, Baron R, Bennett MI, et al. The IASP classification of chronic pain for ICD-11: chronic neuropathic pain. *Pain* 2019; **160**: 53–9

54. Adhikari S, Bista D, Shrestha R, Woods D. Prescribing Patterns and Off-Label Use of Gabapentinoid Agents at Dhulikhel Hospital, Nepal: A Cross-Sectional Study. *J Pain Res* 2024; **17**: 4377–91

55. Aggarwal P, Mishra PN, Mathur VN, Velivela KC, Khan S, Deshmukh P, et al. Drug Usability Survey (DUS) of Gabapentinoid and Its Combinations Among Indian Patients With Neuropathic Pain: Results From a Real-World, Multicenter, Retrospective Survey at Neurology Clinics. *Cureus* Cureus; 2025; **17**

56. Anderson SG, Malipatil NS, Roberts H, Dunn G, Heald AH. Socioeconomic deprivation independently predicts symptomatic painful diabetic neuropathy in type 1 diabetes. *Prim Care Diabetes* 2014; **8**: 65–9

57. Anderson SG, Narayanan RP, Dunn G, Heald AH. Socioeconomic deprivation independently predicts symptomatic painful diabetic neuropathy in people with type 2 diabetes. 2016; **20**

58. Banks C, A. Bowman L, Merrey J, Waldfogel JM. Characterization of Outpatient Gabapentinoid Prescribing for Pain. *J Pain Palliat Care Pharmacother* 2023; **37**: 143–7

59. Berger A, Dukes E, McCarberg B, Liss M, Oster G. Change in opioid use after the initiation of gabapentin therapy in patients with postherpetic neuralgia. *Clin Ther* 2003; **25**: 2809–21

60. Chen S, Wu N, Fraser K, Boulanger L, Zhao Y. Opioid use and healthcare costs among patients with DPNP initiating duloxetine versus other treatments. *Curr Med Res Opin* 2010; **26**: 2507–16

61. Chen S, Wu N, Boulanger L, Fraser K, Zhao Z, Zhao Y. Factors associated with pain medication selection among patients diagnosed with diabetic peripheral neuropathic pain: a retrospective study. *J Med Econ* 2011; **14**: 411–20

62. Dieleman JP, Kerklaan J, Huygen FJPM, Bouma PAD, Sturkenboom MCJM. Incidence rates and treatment of neuropathic pain conditions in the general population ⬚. *Pain* 2008; **137**: 681–8

63. Dinesh Babu U, Pradeep Kumar B. T, Gomathi G. Drug utilization patterns for peripheral neuropathy in a neurology outpatient department at a tertiary care center: A cross-sectional observational study. *Natl J Physiol Pharm Pharmacol* 2024; **14**: 940–940

64. Dragic L, Webb T, Chandler M, Harrington SB, McDade E, Dayer L, et al. Comparing Effectiveness of Gabapentin and Pregabalin in Treatment of Neuropathic Pain: A Retrospective Cohort of Palliative Care Outpatients. *J Pain Palliat Care Pharmacother* 2020; **34**: 192–6

65. Gewandter JS, Kleckner AS, Marshall JH, Brown JS, Curtis LH, Bautista J, et al. Chemotherapy-induced peripheral neuropathy (CIPN) and its treatment: an NIH Collaboratory study of claims data. *Support Care Cancer Off J Multinatl Assoc Support Care Cancer* 2020; **28**: 2553–62

66. Gore M, Sadosky A, Tai K-S, Stacey B. A Retrospective Evaluation of the Use of Gabapentin and Pregabalin in Patients with Postherpetic Neuralgia in Usual-Care Settings. *Clin Ther* 2007; **29**: 1655–70

67. Gore M, Zlateva G, Tai K, Chandran AB, Leslie D. Retrospective Evaluation of Clinical Characteristics, Pharmacotherapy and Healthcare Resource Use among Patients Prescribed Pregabalin or Duloxetine for Diabetic Peripheral Neuropathy in Usual Care. *Pain Pract* 2011; **11**: 167–79

68. Gore M, Tai K, Zlateva G, Bala Chandran A, Leslie D. Clinical Characteristics, Pharmacotherapy, and Healthcare Resource Use among Patients with Diabetic Neuropathy Newly Prescribed Pregabalin or Gabapentin. *Pain Pract* 2011; **11**: 528–39

69. Goswami S, Ramachandran S, Sharma M, Barnard M. Gabapentin and Opioids Utilization in Patients With Diabetic Neuropathy Enrolled in Medicare (2012–2016): A Cohort Study. *J Pharm Pract* 2023; **36**: 1085–94

70. Han G, Han Y, Yu L, Zhao Y, Yu Z. Patterns and Trends in Pharmacological Treatment for Outpatients with Postherpetic Neuralgia in Six Major Areas of China, 2015–2019. *Healthcare* Multidisciplinary Digital Publishing Institute; 2023; **11**: 764

71. Jacob L, Kaiser M, Kostev K. Incidence of antiepileptic drug therapy and factors associated with their prescribing in outpatients with diabetic polyneuropathy. *Prim Care Diabetes* 2021; **15**: 535–40

72. Jha KK, Zaman ZA. An Observational Study on Effectiveness and Safety of Amitriptyline, Duloxetine, and Pregabalin in Painful Diabetic Neuropathy. *IJPCR* 2024; **16**

73. Johnson P, Becker L, Halpern R, Sweeney M. Real-World Treatment of Post-herpetic Neuralgia with Gabapentin or Pregabalin. *Clin Drug Investig* 2013; **33**: 35–44

74. Johnston SS, Udall M, Alvir J, McMorrow D, Fowler R, Mullins D. Characteristics, Treatment, and Health Care Expenditures of Medicare Supplemental-Insured Patients with Painful Diabetic Peripheral Neuropathy, Post-Herpetic Neuralgia, or Fibromyalgia. *Pain Med* 2014; **15**: 562–76

75. Knoerl R, Sohn MB, Foust M, Francar L, O'Rourke MA, Morrow GM, et al. Exploring Analgesic Use Patterns Among Cancer Survivors With Chronic Chemotherapy-Induced Peripheral Neuropathy. *Oncol Nurs Forum* 2024; **51**: 445–50

76. Koopman JSHA, Huygen FrankJ, Dieleman JP, De Mos M, Sturkenboom MCJM. Pharmacological Treatment of Neuropathic Facial Pain in the Dutch General Population. *J Pain* 2010; **11**: 264–72

77. Lin CC, Callaghan BC, Burke JF, Kerber KA, Bicket MC, Esper GJ, et al. Prescription Opioid Initiation for Neuropathy, Headache, and Low Back Pain: A US Population-based Medicare Study. *J Pain* 2023; **24**: 2268–82

78. Marcianò G, Siniscalchi A, Di Gennaro G, Rania V, Vocca C, Palleria C, et al. Assessing Gender Differences in Neuropathic Pain Management: Findings from a Real-Life Clinical Cross-Sectional Observational Study. *J Clin Med* Multidisciplinary Digital Publishing Institute; 2024; **13**: 5682

79. Mittal M, Pasnoor M, Mummaneni RB, Khan S, McVey A, Saperstein D, et al. Retrospective Chart Review of Duloxetine and Pregabalin in the Treatment of Painful Neuropathy. *Int J Neurosci* 2011; **121**: 521–7

80. Muñoz-Vendrell A, Valín-Villanueva P, Tena-Cucala R, Campoy S, Martínez-Yélamos S, Huerta-Villanueva M. Second-line pharmacological treatment strategies for trigeminal neuralgia: A retrospective comparison of lacosamide, gabapentin and baclofen. *Headache* 2025;

81. Nyqvist L, Åkerstedt J, Thoreson O. Current trends in the medical treatment of neuropathic low back pain: a Swedish registry-based study of 1.7 million people. *BMC Musculoskelet Disord* 2024; **25**: 486

82. Patil PR, Wolfe J, Said Q, Thomas J, Martin BC. Opioid Use in the Management of Diabetic Peripheral Neuropathy (DPN) in a Large Commercially Insured Population. *Clin J Pain* 2015; **31**: 414–24

83. Pérez C, Navarro A, Saldaña MT, Masramón X, Pérez M, Rejas J. Clinical and Resource Utilization Patterns in Patients with Refractory Neuropathic Pain Prescribed Pregabalin for the First Time in Routine Medical Practice in Primary Care Settings in Spain. *Pain Med* 2013; **14**: 1954–63

84. Sanchez RJ, Mardekian J, Clair AG, Cappelleri JC. Therapeutic and Subtherapeutic Dosing of Pregabalin: Medication Adherence, Healthcare Resource Utilization, and Costs. *Peripher Neuropathy*

85. Sutema IAMP, Jaya MKA, Bakta IM. Medicine reminder to improve treatment compliance on geriatric patients with diabetic neuropathy at Sanglah Central Hospital, Bali-Indonesia. *Bali Med J* 2018; **7**: 516

86. Toth C, Brady S, Hatfield M. The importance of catastrophizing for successful pharmacological treatment of peripheral neuropathic pain. *J Pain Res* 2014; 327

87. Winterbottom LM, Fong AM, Benkstein KL, Liang B, Snodgrass LS, Parks-Huitron H. Impact of a Clinical Pharmacy Consult Service on Guideline Adherence and Management of Gabapentin for Neuropathic Pain. *J Manag Care Pharm* 2006; **12**: 61–9

88. Wu N, Chen S, Hallett LA, Boulanger L, Fraser KA, Patel CK, et al. Opioid Utilization and Health-Care Costs among Patients with Diabetic Peripheral Neuropathic Pain Treated with Duloxetine vs. Other Therapies. *Pain Pract* 2011; **11**: 48–56

89. Zhao Y, Liu J, Zhao Y, Thethi T, Fonseca V, Shi L. Predictors of Duloxetine versus Other Treatments among Veterans with Diabetic Peripheral Neuropathic Pain: A Retrospective Study. *Pain Pract* 2012; **12**: 366–73

Figure legends

Figure 1. Flowchart of the study selection process. Out of 7,778 titles and abstracts, 69 studies were included in our review.

Tables

Table 1. Characteristics of the included studies. Age is reported as mean unless stated otherwise. F: female, NR: not reported, NeuP: neuropathic pain, DN: diabetic neuropathy, PHN: postherpetic neuralgia, CTS: carpal tunnel syndrome, TN: trigeminal neuralgia, PLP: phantom limb pain, CIPN: chemotherapy-induced peripheral neuropathic pain, PSN: post-surgical neuropathic pain, PTN: post-traumatic neuropathic pain, HIV-SN: human immunodeficiency virus-associated sensory neuropathy, SCI: spinal cord injury, COX: cyclooxygenase, TCA: tricyclic antidepressant, SNRI: serotonin and norepinephrine reuptake inhibitor, SSRI: selective serotonin reuptake inhibitor.

Study	Outcomes	Study design	Country & data time frame	Sample characteristics	NeuP diagnoses	Key findings
Adhikari and colleagues (2024)[54]	Prescribing	Cross-sectional	Nepal 2024	Size: 23 F: NR Age: NR	Peripheral NeuP PHN DN	There was no difference in the likelihood of being prescribed gabapentin or pregabalin between NeuP diagnoses.
Aggarwal and colleagues (2025)[55]	Prescribing	Cross-sectional	India 2023-2024	Size: 2,251 F: 36.3% Age: 51.4	Any NeuP	Compared to people with moderate or severe pain, people with mild pain were more likely to be prescribed gabapentin rather than pregabalin or combinations of gabapentinoid with SNRI.
Anderson and colleagues (2014)[56]	Prescribing	Cross-sectional	UK NR	Size: 1,621 F: 42.2% Age: 47.6	DN	Factors associated with being prescribed pharmacological treatment included: living in a region with higher deprivation, female gender, higher body mass index, older age.
Anderson and colleagues (2015)[57]	Prescribing	Cross-sectional	UK NR	Size: 15,387 F: NR Age: NR	DN	Factors associated with being prescribed pharmacological treatment included: living in a region with higher deprivation, female gender, higher body mass index, older age.
Banks and colleagues (2023)[58]	Discontinuation	Retrospective cohort	USA 2017-2021	Size: 1,221 F: 58.6% Age: 58 (median age)	DN	No significant difference in duration of therapy between people with pregabalin compared to people with gabapentin.
Berger and colleagues (2003)[59]	Prescribing	Retrospective cohort	USA 1999-2002	Size: 45 F: 73.3% Age: 73	PHN	Having several gabapentin prescriptions was associated with a higher likelihood of opioid use.

Boulanger and colleagues (2009)[25]	Prescribing	Retrospective cohort	USA 2004-2006	Size: 27,779 F: 51.5% Age: 64.8	DN	People who had anxiety and/or depression were more likely to be prescribed any pain medication, antidepressants, anticonvulsants, or opioids.
Butler and colleagues (2018)[28]	Prescribing	Prospective cohort	Canada 2010-2013	Size: 210 F: 42.2% Age: 55.2	DN PTN	People with PTN were more likely to use opioids, antiepileptics, TCAs, and other antidepressants, compared to people with DN. People with DN were more likely to use COX inhibitors.
Callaghan and colleagues (2019)[8]	Prescribing	Retrospective cohort	USA 2001-2014	Size: 14,426 F: 47.6% Age: 43.1	Peripheral NeuP	No factors were associated with both a high probability of opioid prescription initiation and chronic prescription, *and* a low probability of guideline-recommended medication prescriptions. Factors associated with a higher probability of being prescribed any pain medication included female gender, younger age, white ethnicity, having diabetes, anxiety, or depression.
Chahine and Al Souheil (2021)[26]	Prescribing	Cross-sectional	Lebanon 2019	Size: 360 F: 55% Age: 50.2	Several	Factors associated with being prescribed/dispensed SNRIs, TCAs, or gabapentinoids included: younger age, low alcohol intake, and DN diagnosis. Factors associated with being prescribed opioids included smoking and high alcohol intake.
Chen and colleagues (2010a)[60]	Prescribing Discontinuation	Retrospective cohort	USA 2004-2006	Size: 234 F: 57.3% Age: 55	DN	People who were prescribed duloxetine were less likely to use opioids compared to those who were prescribed other recommended NeuP medications.
Chen and colleagues (2010b)[40]	Adherence Discontinuation	Retrospective cohort	USA 2003-2007	Size: 1,351 F: 59% Age: 54.7	DN	30 mg/day duloxetine was associated with lower adherence and shorter therapy duration compared to 60 mg/day duloxetine.
Chen and colleagues (2011)[61]	Prescribing	Retrospective case-control	USA 2004-2007	Size: 11,060 F: 54% Age: 55.4	DN	People who had lower back pain or had previously used duloxetine or pregabalin were more likely to initiate duloxetine compared to other medications.
Dieleman and colleagues (2008)[62]	Prescribing	Retrospective cohort	Netherlands 1996-2003	Size: NR F: NR Age: NR	Mononeuropathy CTS DN PHN Facial NeuP	People with CTS were less likely to be prescribed any pain medication compared to those with other NeuP diagnosis. People with PHN were more likely to be prescribed more than one medication compared to those with other NeuP diagnosis. Different NeuP diagnoses affected the probability of being prescribed specific pain medications.

Author	Topic	Study design	Country / Years	Demographics	Condition	Findings
Dinesh Babu and colleagues (2024)[63]	Prescribing	Cross-sectional	India NR	Size: 13 F: NR Age: NR	Peripheral NeuP	Four people with DN were prescribed duloxetine, while none were prescribed pregabalin and amitriptyline combination.
Dragic and colleagues (2020)[64]	Discontinuation	Retrospective cohort	USA 2014-2017	Size: 252 F: 51.2% Age: 52.9	NR	A higher proportion of people prescribed gabapentin reported discontinuations due to adverse events compared to people prescribed pregabalin.
Dworkin and colleagues (2012)[12]	Discontinuation	Retrospective case-control	USA 2005-2006	Size: 252 F: 51.2% Age: 52.9	PHN	People with PHN had longer therapy duration with first-line and third-line medications compared to second-line medications.
Gewandter and colleagues (2020)[65]	Prescribing	Retrospective cohort	USA 2006-2016	Size: 401,114 F: 61.6% Age: 57	CIPN	People treated with neurotoxic chemotherapy where more likely to receive gabapentin or pregabalin for CIPN compared to people treated with non-neurotoxic chemotherapy.
Gharibian and colleagues (2013)[33]	Adherence Discontinuation	Retrospective cohort	USA 2007-2008	Size: 1,817 F: 57.5% Age: 59.4	NR	No difference in adherence or treatment persistence was observed between people who were prescribed antidepressants and people who were prescribed anticonvulsants. People prescribed venlafaxine had higher adherence than people prescribed gabapentin or TCAs.
Giannopoulos and colleagues (2007)[36]	Adherence	RCT	Greece NR	Size: 101 F: NR Age: 57	DN	People with DN had better adherence with SSRIs than gabapentin in a 6-month-long trial.
Gore and colleagues (2007a)[42]	Prescribing Discontinuation	Retrospective cohort	UK 2001	Size: 30,999 F: 59.8% Age: 54.9	Peripheral NeuP	People with pure NeuP were more likely to be prescribed TCAs, second-generation antidepressants, and antiepileptics, while people with mixed NeuP were more likely to be prescribed NSAIDs and opioids. People with SSRIs or SNRIs had longer therapy duration compared to people with other pain medications.
Gore and colleagues (2007b)[66]	Prescribing	Retrospective case-control	USA 2005-2006	Size: 251 F: 59% Age: 54.6	PHN	People who were younger, had depression, or other NeuP diagnoses were more likely to be prescribed pregabalin than gabapentin. Those who were prescribed pregabalin had less opioid use compared to those who were prescribed gabapentin.

Gore and colleagues (2011a)[67]	Prescribing	Retrospective cohort	USA 2004-2006	Size: 1,426 F: 60.3% Age: 55.9	DN	After being prescribed pregabalin, people had fewer anticonvulsant and opioid prescriptions and more SNRI prescriptions. After being prescribed duloxetine, people had fewer SNRI and opioid prescriptions and more anticonvulsant prescriptions.
Gore and colleagues (2011b)[68]	Prescribing	Retrospective cohort	USA 2004-2006	Size: 2,356 F: 53.4% Age: 56.7	DN	People who were prescribed pregabalin as well as those who were prescribed gabapentin had higher proportions of people with most pain medications after the prescription compared to before the prescription.
Goswami and colleagues (2023)[69]	Prescribing	Retrospective case-control	USA 2012-2016	Size: 22,037 F: 55.2% Age: NR	DN	Factors associated with being prescribed concomitant gabapentin and opioids included: female gender, younger age, having anxiety, and having other comorbidities.
Gustavsson and colleagues (2013)[11]	Prescribing Discontinuation	Retrospective cohort	Sweden 2004-2009	Size: 4,718 F: 60.6% Age: 56.2	Pure NeuP Mixed pain	Over 3 years, people with mixed pain had a faster rate of pain medication discontinuation compared to people with pure NeuP.
Hall and colleagues (2006)[43]	Prescribing Discontinuation	Retrospective cohort	UK 1992-2002	Size: 25,824 F: 62% Age: NR	PHN TN PLP DN	The most common first-line treatment in people with DN, PLP, or PHN was amitriptyline; the most common first-line treatment in people with TN was carbamazepine.
Hall and colleagues (2008)[15]	Prescribing Discontinuation	Retrospective cohort	UK 2002-2005	Size: 5,286 F: NR Age: NR	PHN TN PLP DN	People with PHN were less likely to be prescribed antidepressants or antiepileptics as first-line treatment compared to people with other NeuP diagnoses. People with PLP were more likely to be prescribed opioids as first-line treatment compared to people with other NeuP diagnoses.
Hall and colleagues (2013)[16]	Prescribing Discontinuation	Retrospective cohort	UK 2005-2010	Size: 102,386 F: NR Age: NR	PHN DN PLP NeuP back pain	People with DN were more likely to be prescribed antidepressants or anticonvulsants as first-line treatment compared to people with other NeuP diagnoses. People with neuropathic back pain were the most likely to be prescribed tramadol.
Han and colleagues (2023)[70]	Prescribing	Cross-sectional	China 2015-2019	Size: NR F: NR Age: NR	PHN	People with gabapentin were more likely to be prescribed concomitant tramadol compared to people with pregabalin.

Jacob and colleagues (2021)[71]	Prescribing	Retrospective cohort	Germany 2008-2017	Size: 48,324 F: 44.4% Age: 65.8	DN	Factors associated with being prescribed antiepileptics included: female gender, older age, having private insurance, or having type 1 diabetes.
Jha and Zaman (2024)[72]	Prescribing	Cross-sectional	India	Size: 75 F: 53.3% Age: 61	DN	People who were prescribed amitriptyline had significantly higher body mass index compared to people who were prescribed duloxetine or pregabalin.
Johnson and colleagues (2013)[73]	Prescribing Discontinuation	Retrospective cohort	USA 2005-2010	Size: 1,645 F: 59.5% Age: 63	PHN	People who had <150 mg/day of pregabalin or <1800 mg/day of gabapentin were more likely to have a shorter duration of therapy compared to those with ≥150 mg/day of pregabalin day or ≥1800 mg/day of gabapentin.
Johnston and colleagues (2014)[74]	Prescribing	Retrospective cohort	USA 2008-2009	Size: 30,428 F: 53.2% Age: 75.6	DN PHN	Compared to people with DN, people with PHN were more likely to have gabapentin, pregabalin, and topical pain medications. Compared to people with PHN, people with DN were more likely to have non-opioid analgesics.
Kato and colleagues (2023)[37]	Adherence Discontinuation	Retrospective cohort	Japan 2019-2021	Size: 4,138 F: 50.9% Age: 67.1	Peripheral NeuP	Mirogabalin dose titration was associated with better adherence and persistence compared to fixed-dose mirogabalin.
Knoerl and colleagues (2024)[75]	Prescribing	Cross-sectional	USA NR	Size: 142 F: 35.2% Age: 62.9	CIPN	There was no difference in the likelihood of being prescribed opioids based on pain severity.
Koopman and colleagues (2010)[76]	Prescribing	Retrospective cohort	Netherlands 1996-2006	Size: 203 F: 62.1% Age: 54.3	Facial NeuP	Compared to people with TN, people with PHN were more likely to be prescribed medication according to the guidelines by the European Federation of Neurological Societies (EFNS). The most common medications as the first choice were antidepressants for people with PHN while the most common medications as the first choice were anticonvulsants for people with TN.
Kuo and colleagues (2016)[50]	Prescribing Discontinuation	Retrospective cohort	USA Puerto Rico 2006-2013	Size: 7,566 F: 54.7% Age: 64.4	DN	People who were prescribed combination pharmacotherapy were more likely to discontinue their treatment compared to those who were prescribed monopharmacotherapy.

Lakkad and colleagues (2023)[24]	Prescribing	Retrospective case-control	USA 2007-2015	Size: 7,116 F: 100% Age: 74.8	NeuP related to breast cancer treatment	Factors associated with being prescribed recommended medication as the first choice rather than opioids as the first choice included: white ethnicity, mental health disorders, diabetes, and dementia.
Lin and colleagues (2023)[77]	Prescribing	Retrospective cohort	USA 2010-2017	Size: 20,402 F: 50.7% Age: 74.8	Any NeuP	People who were diagnosed for neuropathic pain by a neurologist or podiatrist were less likely to be prescribed an opioid compared to people who were diagnosed by a primary care physician. People who were diagnosed for neuropathic pain by a pain management physician or emergency medicine physician were more likely to be prescribed an opioid compared to people who were diagnosed by a primary care physician.
Marcianò and colleagues (2024)[78]	Prescribing	Cross-sectional	Italy 2021-2024	Size: 174 F: 63.2% Age: 61.5	Any NeuP	There was no difference in prescribing of variety of medications based on sex, age, or body mass index.
Margolis and colleagues (2017)[29]	Prescribing	Retrospective cohort	USA 2005-2014	Size: 14,655 F: 54.4% Age: 63.2	DN PHN	Compared to people with PHN, people with DN were more likely to be prescribed recommended medication as the first choice.
Mittal et al. (2011)[79]	Prescribing Discontinuation	Retrospective cohort	USA	Size: 143 F: 50% Age: 56	Any NeuP	There was no difference in the number of people discontinuing pregabalin or duloxetine.
Mbrah and colleagues (2022)[32]	Prescribing	Retrospective case-control	USA 2016	Size: 69,160 F: 68.6% Age: NR	Any NeuP	Factors associated with being prescribed any pain medication included: female gender, anxiety, depression, multiple sclerosis, arthritis, and seizure disorder. Factors associated with not being prescribed any pain medication included: older age, cognitive impairment, Alzheimer's disease or dementia, cancer, coronary artery disease, fracture, and urinary tract disease.
Muñoz-Vendrell and colleagues (2025)[80]	Prescribing Discontinuation	Retrospective cohort	Spain 2015-2023	Size: 49 F: 53.1% Age: 62.1	TN	There was no difference in discontinuation between people who used gabapentin, baclofen, or lacosamide as a second-choice treatment after carbamazepine discontinuation.
Nygvist and colleagues (2024)[81]	Prescribing	Retrospective cohort	Sweden 2017-2021	Size: >1.7M F: 49.6% Age: NR	NeuP back pain	Factors associated with being prescribed recommended medication included: female gender, more recent year of NeuP diagnosis, and certain types of NeuP back pain.

Oladapo and colleagues (2012)[34]	Adherence	Retrospective cohort	USA 2003-2009	Size: 4,277 F: 72.4% Age: 53.2	DN	People were more adherent with duloxetine than TCAs, with TCAs than gabapentin, and with gabapentin than pregabalin.
Patil and colleagues (2015)[82]	Prescribing	Retrospective cohort	USA 1998-2008	Size: 666 F: 32.9% Age: 58.6	DN	People were more likely to be prescribed opioids as the first choice if they had diabetes-related organ damage or nephropathy.
Pérez and colleagues (2013)[83]	Prescribing	Prospective case-control	Spain 2005-2006	Size: 1,354 F: 49.8% Age: 59.3	DN PHN TN	Higher pain intensity, disability, depression severity, and worse health-related quality of life were associated with a higher likelihood of being prescribed pregabalin.
Pillay and colleagues (2015)[31]	Prescribing	Retrospective case-control	South Africa 2012-2014	Size: 130 F: 71.5% Age: 45.7	HIV-SN	People who had higher pain severity or lower CD4 T-cell count were more likely to be prescribed any pain medication.
Reed and colleagues (2013)[17]	Prescribing	Retrospective cohort	UK 2002-2011	Size: 6,779 F: 45.5% Age: 65.5	DN	People who were diagnosed with DN during 2002-2005 were more likely to be prescribed NSAIDs or opioids compared to people who were diagnosed with DN after 2005. People diagnosed with DN after 2005 were more likely to be prescribed anticonvulsants or SNRIs compared to people diagnosed with DN before 2005.
Reynolds and colleagues (2020a)[22]	Prescribing	RCT	USA 2016-2017	Size: 202 F: NR Age: NR	Peripheral NeuP	Factors associated with being prescribed recommended treatment included: male gender and being treated by a junior doctor.
Reynolds and colleagues (2020b)[9]	Prescribing Adherence	Retrospective cohort	USA 2001-2016	Size: 57,495 F: 54.2% Age: 64.7	Peripheral NeuP	Adherence to gabapentin was higher than adherence to pregabalin. Adherence to venlafaxine was higher than adherence to duloxetine. Adherence to SNRIs tended to be higher than adherence to gabapentinoids. Higher cost of medication to the patient was associated with poorer adherence.
Sadosky and colleagues (2013)[30]	Prescribing	Cross-sectional	USA 2011	Size: 112 F: 52.5% Age: 61.1	DN	Higher pain intensity was associated with more pain medication prescriptions compared to lower pain intensity.

Study	Outcome	Design	Country/Years	Sample	Condition	Findings
Sanchez and colleagues (2012)[84]	Adherence Discontinuation	Retrospective cohort	USA 2007-2009	Size: 6,459 F: 51.6% Age: 64.4	PHN DN	Compared to people prescribed <300 mg/day of pregabalin, people prescribed ≥300 mg/day of pregabalin had better adherence and persistence. Compared to people with PHN, people with DN had better adherence and persistence.
Shaparin and colleagues (2015)[23]	Discontinuation	Analysis of RCTs	USA Russia Argentina 2006-2012	Size: 556 F: 60.3% Age: 66.7	PHN	People were more likely to discontinue gabapentin if they had moderate or severe adverse events (e.g. dizziness, somnolence, nausea, or headache).
Sicras-Mainar and colleagues (2015)[18]	Prescribing Adherence Discontinuation	Retrospective cohort	Spain 2007-2013	Size: 1,369 F: 61.1% Age: 64.6	Peripheral NeuP	Compared to generic gabapentin, brand-name gabapentin was associated with better adherence and persistence.
Sicras-Mainar and colleagues (2019)[41]	Prescribing Adherence Discontinuation	Retrospective cohort	Spain 2014-2017	Size: 4,860 F: 61.3% Age: 59.9	Peripheral NeuP	Compared to generic pregabalin, brand-name pregabalin was associated with better adherence and persistence. Older age and being female were associated with better pregabalin treatment persistence.
Sutema and colleagues (2018)[85]	Adherence	RCT	Indonesia 2016-2017	Size: 62 F: 40.3% Age: 63.3	DN	In an RCT of people with DN who were 60 years or older, having a medicine reminder improved adherence.
Toth and colleagues (2014)[86]	Discontinuation	Prospective cohort	Canada NR	Size: 62 F: 58% Age: 51.9	Peripheral NeuP	Pain-related worrying was associated with a higher likelihood of discontinuation.
Udall and colleagues (2019)[27]	Prescribing	Cross-sectional	Brazil NR	Size: 209 F: 77.6% Age: 53	DN PHN CNP PTN PSN NeuP back pain	Compared to people with other NeuP diagnoses, people with PHN were the most likely to be prescribed any pain medication, antiepileptics, or SNRIs.
Wang and colleagues (2020)[44]	Discontinuation	Retrospective cohort	Taiwan 2012-2017	Size: 48,288 F: NR Age: NR	DN PHN	Compared to people with PHN, people with DN were more likely to persist with pregabalin.
Winterbottom and colleagues (2006)[87]	Discontinuation	Retrospective cohort	USA 2002-2003	Size: 124 F: NR Age: NR	Any NeuP	Compared to usual practice, prescribing gabapentin through a clinical pharmacy consultation process was not associated with any difference in treatment persistence.

Wu and colleagues (2009)[39]	Prescribing Adherence	Retrospective case-control	USA 2003-2007	Size: 2,354 F: 57.8% Age: 61.8	DN	People who were prescribed >30mg/day duloxetine had better adherence than those who were prescribed ≤30mg/day duloxetine.
Wu and colleagues (2011)[88]	Prescribing Discontinuation	Retrospective cohort	USA 2004-2006	Size: 499 F: 55.3% Age: 54.8	DN	People who were prescribed duloxetine were less likely to use opioids compared to those who were prescribed other recommended NeuP medications.
Yang and colleagues (2015)[13]	Adherence Discontinuation	Retrospective cohort	USA 2005-2012	Size: 8,523 F: NR Age: NR	DN	People who were prescribed duloxetine had better adherence than those who were prescribed gabapentin or pregabalin. At 180 days, most people had discontinued their treatment in all groups.
Yeh and colleagues (2021)[38]	Adherence Discontinuation	Retrospective cohort	USA 2009-2014	Size: 5,186 F: 47.6% Age: 53	DN PHN NeuP related to SCI	The characteristics associated with better pregabalin adherence included: dose titration, older age, male gender, previous gabapentin or TCA use, and concomitant lidocaine use.
Zhao and colleagues (2011)[35]	Prescribing Adherence Discontinuation	Retrospective cohort	USA 2005-2007	Size: 2,573 F: 57.7% Age: 55.7	DN	People who were prescribed pregabalin were more likely to discontinue their treatment earlier and be less adherent compared to those who were prescribed duloxetine.
Zhao and colleagues (2012)[89]	Prescribing	Retrospective case-control	USA 2004-2008	Size: 2,694 F: 2.1% Age: 64.3	DN	Factors associated with being prescribed duloxetine rather than other recommended NeuP medications included: previous gabapentin use, previous TCAs use, previous venlafaxine use, previous opioid use, anxiety, substance abuse, cerebrovascular or peripheral vascular disease, and self-reported severe pain.

Table 2. Summary of key predictors. For each predictor, an example result is reported. OR: odds ratio, PR: prevalence ratio, IRR: incidence rate ratio, MPR: medication possession ratio, PDC: proportion of days covered, RR: relative risk, HR: hazard ratio.

Outcome	Key predictors	Example result
Receiving recommended NeuP medication	White ethnicity	OR: 1.40 (95% CI: 1.18 to 1.66)[8, 24]
	Diabetes	OR: 1.41 (95% CI: 1.23 to 1.62)[8, 24, 26]
	Mental health disorder (especially depression and/or anxiety)	OR: 3.57 (95% CI: 3.12 to 4.09)[24]
	Pure NeuP (compared to mixed pain)	OR: 1.37 (95% CI: 1.30 to 1.45)[42]
Receiving any pain medication	Younger age (50-64 years vs +85 years)	PR: 1.45 (95% CI: 1.38 to 1.52)[32]
	Higher pain severity (BPI severe vs mild)	OR: 25.83 (95% CI: 2.97 to 225.21)[30]
	Anxiety and/or depression	OR: 3.66 (95% CI: 3.14 to 4.26)[25]
Adherence	Older age (55-64 years vs 18-44 years)	OR: 2.11 (95% CI: 1.50 to 2.97)[39]
	Younger age (65-74 years vs 75-84 years)	OR: 1.33 (95% CI: 1.01 to 1.79)[39]
	White ethnicity	IRR: 1.14 (95% 1.08 to 1.21)[9]
	Lower medication cost to patient in the USA (compared to +$50 for 30-day supply)	IRR: 1.09 (95% CI: 1.03 to 1.14)[9]
	SNRIs (compared to gabapentinoids)	Difference in mean MPR: >0.10[9]
	SNRIs (compared to tricyclic antidepressants)	Difference in mean MPR: 9.6 %[34]
	Dose titration	OR: 2.40 (95% CI: 1.51 to 3.21)[37, 38]
	Brand-name medication (compared to generic-name medication)	MD in MPR: 0.05 (95% CI: 0.03 to 0.07)[18, 41]
	Medicine reminder	RR: 2.38 (95% CI: 1.58 to 3.61)[85]
	Diabetic neuropathy (compared to postherpetic neuralgia)	MD in PDC: 0.13 (95% CI: 0.10 to 0.16)[84]
Discontinuation	Tricyclic antidepressants (compared to other antidepressants)	OR: 1.56 (95% CI: 1.01 to 2.42)[15]
	Generic rather than brand name medication	HR: 1.43 (95% CI: 1.18 to 1.72)[41]
	Combination pharmacotherapy	HR: 1.30 (95% CI: 1.13 to 1.51)[50]
	Not experiencing clinically significant pain relief	OR: 1.2 (95% CI: 1.0 to 1.4)[41]
	Experiencing moderate adverse events	OR: 8.64 (95% CI: 4.42 to 16.89)[23]
	Experiencing severe adverse events	OR: 5.37 (95% CI: 2.01 to 14.00)[23]
	Postherpetic neuralgia (compared to diabetic neuropathy)	MD in treatment days: 51 (95% CI: 4 to 98)[15, 16, 43, 44]
	Pain-related worrying (CSQ catastrophising subscale score)	Pearson's correlation: 0.36[86]

Identification

Records from databases **(n = 11,255)**
Embase (n = 5,048)
PubMed (n = 3,033)
Web of Science (n = 2,577)
CINAHL (n = 597)

Records from reference lists of included studies **(n = 7)**

Duplicates removed **(n = 3,484)**

Screening

Titles/abstracts screened **(n = 7,778)**

Records excluded **(n = 7,531)**

Records assessed for eligibility **(n = 247)**

Records excluded **(n = 178)**
No relevant outcomes (n=24)
Sample not specific to neuropathic pain (n = 66)
No predictors (n = 35)
Irrelevant medications/treatments (n=7)
Wrong study design (n=1)
Hypothetical prescribing study (n=3)
Prescribing study in a surgical context (n=1)
RCT: discontinuation between mediations (n = 6)
Carpal tunnel syndrome only (n=3)
Not a full text article (n = 30)
Not in English (n = 1)
No data to support relevant statements (n = 1)

Included

Studies included in review **(n = 69)**