Genomic surveillance reveals age-structured SARS-CoV-2 transmission across demographics and settings

Gage K. Moreno*[†1], Taylor Brock-Fisher*[1,2], Lydia A. Krasilnikova*[1,2,3], Stephen F. Schaffner*[1,2,4], Meagan Burns[5], Carolyn E. Casiello[5], Katelyn S. Messer[1], Brittany Petros[1,6,7], Ivan Specht[1], Katherine C. DeRuff[1], Katherine J. Siddle[8], Christine Loreth[1], Nicholas A. Fitzgerald[1], Heather M. Rooke[1], Stacey B. Gabriel[1], Sandra Smole[5], Shirlee Wohl[5,9], Daniel J. Park[1], Lawrence C. Madoff[5], Catherine M. Brown*[5], Bronwyn L. MacInnis*[†1,10], Pardis C. Sabeti*[1,3,10]

Broad Institute of Harvard and MIT, Cambridge, MA, USA

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA

Howard Hughes Medical Institute, Chevy Chase, MD, USA

Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA

Massachusetts Department of Public Health, Boston, MA, USA

Harvard-MIT Program in Health Sciences and Technology, Cambridge, MA, USA

Harvard/MIT MD-PhD Program, Boston, MA, USA

Department of Molecular Microbiology and Immunology, Brown University, Providence, RI, USA

Brigham and Women's Hospital, Boston, Massachusetts, USA

Massachusetts Consortium for Pathogen Readiness, Boston, MA, USA

* These authors contributed equally

† Corresponding authors: gmoreno@broadinstitute.org (G.K.M.) and bronwyn@broadinstitute.org (B.L.M.)

SUMMARY

Understanding respiratory virus transmission requires integrating pathogen genomic data with detailed epidemiological context. We analyzed >85,000 SARS-CoV-2 genomes linked to epidemiological data collected in Massachusetts from November 2021 to January 2023. Collected in 666 settings, including schools, colleges, and skilled nursing facilities, this dataset enabled direct comparison of within-setting transmission to community-wide risk. Elevated transmission was concentrated in specific age-defined subpopulations–older adolescents in schools, undergraduates in colleges, and residents in nursing facilities–with little additional risk among other groups, notably including staff. Young adults, particularly in colleges, played a disproportionate role in spread, and new viral lineages consistently expanded first within this group. Viral spread followed reproducible urban-to-rural gradients, and vaccination was associated with reduced onward transmission. We further estimated sequencing thresholds required for timely detection of emerging variants. Together, these findings reveal consistent demographic and spatial structure in SARS-CoV-2 transmission that can inform future viral respiratory pathogen genomic surveillance.

KEYWORDS

Genomic surveillance; genomic epidemiology; surveillance; SARS-CoV-2; COVID-19; transmission; genomics; vaccination; age-structured transmission

INTRODUCTION

Genomic surveillance transformed understanding of SARS-CoV-2 evolution and spread during the COVID-19 pandemic. Most large-scale surveillance efforts were designed primarily to detect

and track emerging variants of concern, often with limited linked epidemiological context. In

contrast, studies integrating viral genome data with detailed metadata have enabled more

comprehensive analyses of the pandemic, yielding important insights into risk factors for

infection[1,2,3] and outbreak dynamics[4–7]. Nevertheless, many questions remain about the

epidemiological drivers that shaped SARS-CoV-2 transmission, and addressing these questions

using linked genomic and epidemiological data is likely to yield broadly relevant insights into the

spread of respiratory viruses.

The Broad Institute's clinical laboratory provided large-scale, PCR-based COVID-19 diagnostic

testing throughout the acute phase of the pandemic, with a focus in its home state of

Massachusetts, USA. The laboratory served public testing programs as well as educational,

medical, congregate living, and other settings. The Broad Institute also sequenced a subset of

positive samples through the US CDC's National SARS-CoV-2 Strain Surveillance (NS3)

program. By early 2023, these efforts had generated more than 37 million diagnostic tests and

175,000 SARS-CoV-2 genomes, representing ~5% of all tests and genomes produced in the

United States.[8,9] In parallel, the Massachusetts Department of Public Health (MDPH) and

municipal public health departments collected extensive epidemiological data on COVID-19

cases statewide. Combining these efforts, we assembled a dataset of over 130,000 SARS-CoV-

2 genomes collected in Massachusetts from late 2021 through early 2023, spanning the late

Delta wave and successive Omicron subvariant waves. Over 85,000 genomes were linked to

epidemiological data, including age, gender, vaccination status, municipality of residence, and

testing setting.

This dataset enabled unusually detailed analyses of SARS-CoV-2 transmission across

institutional and community environments. Metadata describing the testing setting, including

schools, colleges, and skilled nursing facilities (SNFs), allowed systematic comparison of

transmission dynamics across varied settings and among sub-populations within them.

Municipality-level residence information provided fine-scale geographic resolution. Demographic

and vaccination data, available for a substantial fraction of cases, added further individual-level

context. Together, these features created an opportunity to examine how age, setting,

geography, and vaccination status shaped transmission patterns at scale.

In this study, we integrated genomic, geospatial, and epidemiological data to examine how

SARS-CoV-2 transmission varied across age groups, institutional environments, and viral

lineages. We analyzed how new variants were introduced and propagated, how transmission

risk differed across demographic and contextual settings, and how these dynamics shifted

across epidemic waves. We also assessed associations between vaccination status and both

infection risk and onward transmission, and estimated the sampling intensity required for timely

detection of emerging lineages. Together, these analyses define demographic and spatial

structure in viral spread and provide insights to guide future public health genomic surveillance

and response strategies for viral respiratory pathogens.

RESULTS

Over 85,000 SARS-CoV-2 genomes paired to individualized

epidemiological data

We analyzed 134,785 high-quality (≥80% complete) SARS-CoV-2 genomes generated by the

Broad Institute from residual positive diagnostic samples collected from Massachusetts

residents between November 1, 2021, and January 17, 2023. These genomes comprised 492

named Pango lineages, which fell into seven dominant lineages: Delta (B.1.1.617.2*, where the

asterisk indicates the inclusion of sublineages[10]) and the BA.1*, BA.2*, BA.4*, BA.5*, BQ.1*, and

XBB* Omicron subvariants (**Fig 1A, 1B**). We identified 434 infections involving multiple viral

lineages and 85 additional cases consistent with reinfection within 90 days by a different lineage

(see Methods) (**Fig S1**).

The samples were collected at 666 facilities statewide, which we classified into seven sectors

(**Fig 1C, Fig S2, S3**): 29 public testing facilities (accounting for 35% of genomes), 101 colleges

(19%), 52 hospitals and clinics (17%), 301 skilled nursing, rehabilitation, and long-term care

facilities collectively referred to as SNFs, (5%), 44 primary and secondary schools (1%), 23

other workplaces (<1%), and 117 testing facilities of unknown sector (22%). We obtained

deidentified epidemiological data, including age, sex, city of residence, and vaccination

information, for 85,125 total genomes from 81,952 unique individuals. Unless otherwise

indicated, we used this set of genomes with accompanying epidemiological data as the core

dataset for downstream analyses.

Sequenced samples broadly reflected the distribution of contributing testing facilities (**Fig S2-**

S4) and tracked statewide case count trends, with a mean sampling fraction of 10.5% of all

confirmed cases reported to MDPH per week during the study period (**Fig 1D**). Several sectors

were enriched for specific demographic groups, including children in primary and secondary

schools, young adults in colleges, and older adults in SNFs (**Fig S2D**). In contrast, free public

testing was available statewide regardless of age or symptom status and therefore served as

the sector most representative of the overall population (**Fig S4**).

Distinct transmission patterns in different demographic groups

The diversity of environments and demographics represented by the testing sectors in our

dataset allowed us to examine how these factors shaped SARS-CoV-2 transmission. We

focused on settings with clear implications for shared exposure, including SNFs, colleges,

primary and secondary schools, and a mixed group of other workplaces that included hospital

staff, first responders, and warehouse staff. For each setting, we asked how much additional

risk of infection was attributable to the shared facility environment compared to broader

community risk.

We assessed within-facility transmission by quantifying the enrichment of closely related viral

genomes within each setting compared to the surrounding community. That is, for each virus

from a facility, we calculated how often other viruses from either the same facility or from the

surrounding community were closely related to it, where the community was represented by

public testing data from individuals in the same municipality. Increased within-facility viral

relatedness therefore served as a proxy for the probability that transmission occurred within the

facility rather than between unrelated individuals in the same municipality. For this purpose,

closely related viruses were defined as those differing by ≤2 substitutions and collected within

10 days of one another, consistent with previous studies.[11–13]

Using this framework, we observed marked differences in within-facility transmission between

settings. Closely related viruses were strongly enriched in SNFs (4.9x, 95% CI: 4.3–5.5x) and

substantially enriched in colleges (3.3x, CI: 3.1–3.6x). In contrast, enrichment in schools was

modest (1.4x, CI: 1.2–1.7x) and was not statistically significant in other workplaces (1.1x, CI:

0.9–1.3x) (**Fig 2A**). These patterns indicated that the contribution of within-facility transmission

varied considerably by setting; notably, average viral relatedness in SNFs was nearly twice that

observed in any other setting. These results were robust to differences in sampling density

across sectors, as subsampling analyses yielded consistent enrichment patterns (**Fig S5**), and

were not driven by a small number of highly sampled facilities (**Fig S6**).

We also observed that in SNFs, schools, and colleges, enrichment of closely related viruses

was largely confined to specific age groups. In SNFs, the probability that a virus had a close

relative within the same facility increased steadily with age, beginning around 50 years old.

Among the oldest individuals, viruses were 60 percentage points more likely to be closely

related to another virus from the same facility than to one from the surrounding community (**Fig**

2B). Although our SNF dataset likely included some younger patients in rehabilitation and other

care settings—so age does not perfectly distinguish staff from residents—the proportion of true

residents increases with age. The strongest enrichment of closely related viruses therefore most

likely reflects resident-to-resident transmission.

Examining within-facility and community relatedness separately provided additional insight into

transmission in SNFs. Relatedness between viruses in SNFs and those in the community varied

little by age, whereas relatedness among viruses in SNFs increased sharply with age (**Fig. S7**).

This pattern is consistent with repeated viral introductions from the community into SNFs

followed by relatively short transmission chains within facilities. Prolonged outbreaks from single

introductions would instead be expected to generate longer transmission chains linking

residents to community viruses, producing declining relatedness with age. These observations

are not, however, informative about the frequency of introductions or overall incidence among

staff and residents.

In schools, staff-aged individuals (defined here as >23 years old) showed the lowest within-

facility enrichment of closely related viruses, lower even than in SNFs: their viruses were only

2% more likely to have a close relative in the school than in the broader community. Younger

students (5-14 years old) also showed only modest enrichment (3%), with some variation by

age. In contrast, enrichment increased sharply among older adolescents, peaking at ~20%

among 18-year-olds. Because both younger students and staff show little evidence of increased

transmission, these findings suggest that the school environment itself conferred little additional

risk. Instead, elevated transmission among older adolescents likely reflects age-specific social

interaction patterns, whether inside or outside school.

Colleges showed a similar age-structured pattern, with enrichment of related viruses largely

confined to undergraduates (typically 18 to 22 year olds). The strongest signal was among

those 18–20 years old, for whom enrichment peaked at ~20%. We found little evidence of

increased viral relatedness among other age groups within colleges, including those typical of

post-graduate students, faculty or staff. Together, the school and college analyses suggest that

elevated SARS-CoV-2 transmission was not a general feature of these environments but was

concentrated within a narrow age range. Consistent with this interpretation, public testing data

also showed increased transmission among individuals in their late teens to early twenties (**Fig**

S8), suggesting this age group contributed disproportionately to overall transmission, although

the proportion who were college students is unknown.

The importance of young adults in colleges in SARS-CoV-2 transmission was further evident in

examining lineage dynamics. We previously observed that BA.1 cases rose earlier and more

rapidly in colleges than statewide[14]. Extending this analysis to major lineages for which we had

sufficient data, BA.1*, BA.2*, BA.2.12.1* reached 50% frequency earlier in college settings than

in other sectors, with lead times ranging from 4 to 13 days. BA.5* was an exception, it showed

similar timing across all sectors, but rose when colleges were not in session (**Fig 2C**).

To disentangle effects of age from college settings, we compared 18- to 22-year-olds in college

and public testing to all-age public testing for BA.1*, BA.2*, and BA.2.12.1* (omitting BA.5* for

lack of data). BA.1* reached 50% frequency earliest among 18- to 22-year-olds in colleges,

whereas BA.2* and BA.2.12.1* reached 50% frequency earlier among 18- to 22-year-olds in

both colleges and public testing than in all-age public testing (**Fig S9**). These findings suggest

that both the college environment and the 18–22 age group contributed to early lineage

expansion, supporting the potential role of young adults as a sentinel population for genomic

surveillance.

We considered the possibility that asymptomatic surveillance testing conducted in many

colleges contributed to the initial rise of emerging variants there, since such testing could enrich

for early detection of infections[14,15]. To test this, we grouped together the other sectors known to

have been conducting predominantly asymptomatic surveillance testing (schools, workplaces,

and SNFs) to increase statistical power. We then compared the rise of variants in this combined

group with colleges, as well as with the remaining sector categories, (hospitals and clinics, and

public testing), which performed mixtures of symptomatic and asymptomatic testing (**Fig S10**).

As before, all variants except for BA.5* (which emerged when college was not in session)

reached 50% frequency earliest in colleges. The consistent early rise of multiple lineages in

colleges, coupled with the absence of this pattern in other sectors conducting asymptomatic

surveillance, suggests that the early signal in college testing was not simply driven by

asymptomatic testing. Instead, features of the college population and environment likely

contributed to rapid early expansion, reinforcing the potential of colleges as settings for early

viral detection.

204   High resolution temporal and spatial mapping of viral spread

Our dataset captured the introduction of six major Omicron sub-lineages at high geographic

resolution, enabling detailed analysis of how new variants were introduced, established, and

spread. The dataset had broad coverage of the state, including genomes from ≥1% of all

confirmed cases in 87% (304 of 351) of municipalities (**Fig 3A**). To quantify viral introductions

and movement, we reconstructed phylogenetic trees using two complementary sampling

strategies: one normalized to case counts and one based on equal sample sizes per week per

municipality. We defined an introduction into the state as a virus detected in our dataset whose

immediate ancestral node was inferred to be outside Massachusetts. Similarly, we identified

intrastate movement when the inferred locations of ancestral and descendant nodes shifted

between municipalities within the state.

New introductions into the state were disproportionately likely to occur into more urban areas.

That is, even when municipalities were equally sampled, the rate of introductions was

significantly and independently associated with both the size and the density of the

municipality's population, suggesting greater per-capita linkage to out-of-state travel in urban

areas (**Fig S11**). Movement of viruses between municipalities was similarly associated with both

the size and density of the source population. As a result, urban municipalities acted as both

recipients of new introductions from outside the state and sources of intrastate spread (**Fig 3B**).

Consistent with this, we found that BA.1* and BA.2* (the only lineages with enough data) rose to

high frequency initially in the primary urban center, followed by the surrounding suburbs and

then by smaller independent urban areas, with more rural areas trailing (**Fig 3C**).

We estimated the speed of viral movement throughout the state by looking at how genetic

distance varied at different geographic distances. For pairs of viruses collected within 10 days of

each other, the number of genetic differences between them should be proportional, on

average, to the number of intervening transmissions since the pair's common ancestor. Closely

related viruses (≤2 substitutions) occurred predominantly at short physical distances (<~15 km)

(**Fig 3D**). By the time a pair of viruses were separated by 3-5 substitutions (corresponding to ~4

generations or 20 days of transmission since their common ancestor), they were almost as likely

to be found >100 km apart as within the same municipality. A substantial component of this

spread involved movement between major urban centers. For example, the relatedness peak in

Fig 3D around 125 km, which is most pronounced for a genetic distance of 3–5 substitutions,

reflects the distance between two major, well-sampled cities (Boston and Springfield); similar

patterns were observed for other metropolitan pairs. In contrast, pairs involving at least one

virus from a more rural municipality consistently showed lower relatedness across distances,

consistent with viruses primarily moving between urban centers rather than from urban to rural

municipalities. When genetic distance exceeded ~12 substitutions, geographic structure

became undetectable, suggesting that descendant viruses typically diffused broadly throughout

the state within roughly two months.

Although viruses in rural areas were less likely to be closely related to those from distant

municipalities, they were more likely to be closely related to other viruses from the same

community. In developing suburbs and rural municipalities, viruses from the same municipality

were 11.3x as likely to be closely related as viruses from different municipalities, compared to

only 5.6x in the inner core, regional urban centers, and maturing suburbs (**Fig 3E**). Greater

relatedness in less populous towns may reflect both the greater isolation from intrastate

movement and smaller effective viral population sizes in those settings. These different aspects

of the genomic data, then, create a consistent picture, in which new variants typically arrived

first in urban areas, spread rapidly to adjacent suburbs and then to other regional urban centers,

and reached rural areas more slowly and less completely.

Vaccination was associated with decreased infection and transmission

Although this opportunistically collected dataset was not designed to directly assess the effects

of vaccination on infection or transmission, we leveraged the available data to explore these

relationships and observed modest but informative associations. In one cohort, the timing of the

COVID-19 vaccine rollout allowed us to examine the qualitative impact of vaccination on

infection. Children aged 5–11 years old first became eligible for vaccination in the US on

October 29, 2021, two days before the start of our study, with administration of first doses

ramping up in mid-November (**Fig S12A**) and second doses following in mid-December.

Comparing the periods immediately before and after widespread vaccination (November vs. late

December/early January), we found that the proportion of infected individuals in the 5- to11-

year-old cohort declined by approximately 50% in both school-based and public testing data

(Fig. 4A,B), with the decline evident for both Delta and BA.1* **(Fig. S12B)**. The concordance

between school-based and public testing suggests that vaccination provided substantial short-

term protection against infection.This protection may have contributed to the relatively low

within-school transmission among younger students noted above, although it is unlikely to be

the dominant factor, since similarly low values were observed among 12- to 14-year-olds.

We asked whether vaccination altered SARS-CoV-2 transmission between individuals using two

complementary genomic approaches. Both analyses suggested an association between booster

vaccination and reduced onward spread. The first approach distinguished likely transmitters

from non-transmitters by proxy, classifying individuals whose viruses had closely related

genomes elsewhere in the dataset as "linked", i.e. as evidence of a short transmission chain

between them, and those without such relatives as "unlinked". Unlinked individuals were

significantly more likely to have received a booster than linked individuals (33.2% vs. 19.1%; **Fig

4C**), indicating a weaker connection between boosted individuals and sampled transmission

chains. The second approach identified likely transmitters by detecting within-host variants that

later appeared as consensus variants in other individuals, consistent with onward

transmission[16]. In this analysis, boosted individuals were 0.65x (0.50–0.84) as likely as

unvaccinated individuals to initiate at least one transmission event (**Fig 4D**). In contrast, we

observed no significant differences between unvaccinated individuals and those who had

completed a primary vaccine series without a booster in either analysis.

We next asked whether the association between booster vaccination and reduced onward

spread reflected recency of vaccination rather than number of vaccine doses received. To test

this, we compared unboosted individuals who were first vaccinated early in the initial rollout

(February-May 2021) with those first vaccinated later (November 2021- January 2023).

Individuals vaccinated later showed a 0.85x (0.54x–1.33x) likelihood of initiating transmission

relative to unvaccinated individuals (**Fig S13**). Although this difference was not statistically

significant, the trend is consistent with previous observations of waning vaccination protection

against SARS-CoV-2 transmission[17]. We also observed lower viral loads (based on diagnostic

Ct values) with increasing number of vaccine doses, and a modest but statistically significant

increase in viral load with time since the most recent vaccine dose[18] (**Fig S14**). Together, these

results suggest that the increased probability of transmission with time since vaccination may be

partly mediated by biological factors related to viral load. However, both the number and timing

of vaccine doses are also likely correlated with behavioral factors that influence transmission

probability, such as risk tolerance or the ability to self-isolate.

Sampling rates for detection of novel lineages

During the pandemic, practical questions arose about the rate of viral sequencing needed to

effectively surveil emerging variants of concern. Using our dataset of all 130,000 genomes, we

first asked how sampling rate influences the speed of detecting new variants. Using both a

simple binomial model and the tool PhyloSamp[19], we modeled the detection of lineages with the

observed growth rates of BA.1*, BA.2*, BA.2.12.1*, BA.5*, and BQ.1* (the lineages with

sufficient data). Specifically, we assessed how the weekly sampling rates affected the timing of

first detection relative to the date each variant reached 1% prevalence in the dataset, a

commonly used surveillance threshold[20,21]. Sequencing 100 genomes per week resulted in

detection before the 1% threshold more than 50% of the time, while 50 genomes/week typically

delayed detection by only a few days. The probability of detection increased with increasing

sequencing rate up to approximately 500 genomes per week. At this rate, four of the five

lineages (BA.2*, BA.2.12.1*, BA.5*, and BQ.1*) had >95% probability of being detected by the

target date, with little additional benefit from higher sampling (**Fig 5A, Fig S15**). In contrast,

BA.1* rose so rapidly in frequency that sampling intensity had little effect on detection timing;

increased sampling would not have enabled earlier detection.

We next estimated the sampling rate required for a more data-intensive surveillance goal:

identifying previously unknown lineages that are increasing in frequency and therefore represent

potential variants of concern. To do this, we repeatedly subsampled our dataset of 130,000

genomes for the same five lineages at increasing weekly sampling rates. For each subsample,

we fit logistic regression models beginning on the day the lineage reached 3% frequency and

extending over varying time windows to determine the earliest date at which the growth rate

became significantly positive (95% confidence). The 3% threshold minimizes the highly

stochastic fluctuations that occur prior to the onset of exponential growth. Based on these

empirical data, sequencing 300 genomes per week was sufficient to determine within 14 days

that BA.1* was increasing in frequency **(Fig. 5B)**. In contrast, for slower-growing variants such

as BA.5*, confident detection of growth within 14 days was effectively unattainable; even

detection within 21 days would have required sequencing at least 400 genomes per week **(Fig.**

5C).

DISCUSSION

Large-scale genomic surveillance provided valuable insights into the emergence and spread of

SARS-CoV-2 variants during the COVID-19 pandemic, but deeper epidemiological analyses

were often hindered by the lack of detailed metadata linked to viral genomes. Here, we leverage

large-scale, high-resolution SARS-CoV-2 genomic surveillance data paired with epidemiological

metadata across multiple variant waves to examine spatial patterns of viral spread, differences

in transmission dynamics across demographic groups and settings, and factors associated with

the risk of infection and onward transmission. We also provide practical estimates of the

sampling rate required for timely detection of known variants of concern and for identifying

previously unknown variants that are increasing in frequency.

Within schools, colleges, and SNFs, elevated SARS-CoV-2 transmission relative to public

testing was concentrated in specific age-defined subpopulations rather than reflecting a general

feature of these institutional settings. Notably, in all three settings, transmission involving staff-

aged individuals was only modestly higher than, or indistinguishable from, transmission in the

surrounding community. We also found no evidence of substantially elevated transmission in a

heterogeneous group of other workplaces, nor among children under the age of 15 in schools.

Although sample sizes for schools were relatively small in our dataset, the essential findings,

particularly the importance of young adults and the lower transmission risk among staff, were

consistent across sectors. Together, these results suggest that increased infection risk in these

settings was largely driven by characteristics of specific subpopulations rather than by the

institutional settings themselves.

The demographic groups that showed elevated transmission likely reflect different processes in

their respective contexts. SNFs were the only setting that showed a consistent enrichment of

within-facility transmission. The substantial increase among older age groups suggests a unique

risk to residents in these facilities, while the more modest increase among individuals under 50

could indicate increased risk among staff, although it may also reflect younger in-patients in

rehabilitation or other care settings.

In schools, the sharp increase in transmission was largely confined to 15- to 18-year-olds,

suggesting that it reflects distinctive social interaction patterns in this age group, only some of

which occur within the school environment. It was somewhat surprising that we did not observe

stronger evidence for school-associated transmission across a broader age range, given the

well-established role of younger children in the spread of endemic respiratory viruses, the many

opportunities for transmission within schools and school-adjacent environments, and the

additional shared risk associated with siblings attending the same school. A particularly notable

and reassuring finding was how limited the apparent workplace risk for teachers was, at least

under the conditions present in our data.

We observed similar age-structured transmission among the undergraduate-aged population in

college settings. The particularly sharp increase between ages 18 and 20 may reflect the

additional effect of shared living environments, as on campus college dormitories are more

common in this age group than among older students who typically live off campus. Whatever

the underlying causes, evidence for elevated transmission among young adults comes from

both college and school populations, groups that typically have limited overlap in their social

networks.

Our findings here and elsewhere[14] suggest that individuals aged 18 to 22 served as leading

indicators for the establishment of new SARS-CoV-2 lineages; emerging variants expanded

within young adult populations before becoming widespread across the rest of the state, with

the effect strongest among college students. Because Massachusetts has a high concentration

of colleges and other post secondary institutions, we considered whether an overrepresentation

of young adults, particularly residential college students, might influence this pattern. We were

unable to readily identify data directly comparing the proportion of residential college students in

Massachusetts with that of the U.S. overall. However, U.S. Census Bureau data indicate that

18- to 22-year-olds constitute a very similar share of the total population in Massachusetts and

nationwide, approximately 6.5% in both cases[22,23]. (We note that the Census counts college

students where they live during the academic year, whether on or off campus, even if they are

originally from another state). In any case, our observation reflects the relative transmission

rate within this demographic group rather than its share of the population, a relationship likely to

hold in other contexts.

Although a genomic epidemiology dataset of this scale and resolution would ideally enable more

extensive evaluation of public health interventions on transmission and infection risk, the timing

and nature of interventions during our study period largely precluded such analysis. By the start

of the study, many earlier widespread measures, including masking and distancing mandates

and broad school closures, had already ended, leaving vaccination as the primary public health

intervention for most of the period. Early in the study, some masking and other mitigation

measures were intermittently in place, but these coincided with rapid shifts in the immune

landscape as Delta was replaced by the BA.1 Omicron surge, along with extensive travel and

holiday-related school and college closures. As a result, we were unable to reliably isolate the

effects of specific policy changes on transmission.

Nevertheless, we observed a short-term reduction in infections among 5- to 11-year-olds

coinciding with vaccine rollout, detectable in both school-based surveillance and broader public

testing. We were unable to estimate the effect of vaccination on infection more broadly because

we did not have vaccination data for uninfected people to use as a comparator. Assessing

vaccination effects on transmission was more tractable, as these could be inferred from viral

genetic relationships combined with vaccination status metadata within our dataset. Individuals

with more recent and additional vaccine doses had a lower probability of initiating transmission,

although causal interpretation remains limited by potential confounding factors, such as

behavioral differences between groups.

The arrival of multiple new lineages during our study period, each effectively representing a new

outbreak, allowed us to examine patterns in where lineages were introduced, how they

subsequently spread, and how they behaved across different populations and settings. New

lineages were introduced primarily through major urban areas, from which they spread quickly

to nearby suburbs and regional centers, and more gradually to less populated areas, becoming

broadly distributed statewide after approximately two months. These patterns suggest that

effective strategies to reduce transmission may differ by geographic density, for example,

prioritizing efforts to limit new introductions into urban centers while focusing on interrupting

local transmission chains in rural areas.

Our analyses also clarify how sequencing coverage influences the timeliness of variant

detection. Increasing the fraction of cases sequenced reduces the delay to detecting emerging

lineages, but with diminishing returns at higher coverage levels. These results highlight a

practical tradeoff between surveillance effort and detection speed, suggesting that sustained

sequencing at moderate coverage can capture much of the achievable early-warning benefit.

Despite the constraints inherent to opportunistic genomic surveillance data collected during a

rapidly evolving pandemic, this study illustrates the power of integrating genomic and

epidemiological information to resolve fine-scale transmission patterns. By identifying which

groups were most likely to drive transmission within and across settings, and when emerging

variants were most likely to become detectable, our findings provide actionable insights for the

design of future genomic surveillance systems, particularly through targeted sampling of high-

transmission demographic groups.

422 Limitations of study

While offering important novel findings, this study was subject to common limitations of genomic

epidemiology, including opportunistic, observational datasets, siloed data streams, and the

broader constraints of a rapidly evolving public health emergency. In particular, our sampling

framework was shaped by the realities of a large-scale COVID-19 testing program and was not

designed to resolve details of transmission dynamics. Although epidemiological metadata, rarely

available at this scale, enabled comparisons across demographic groups and settings, it did not

permit definitive linkage of infections to individual behaviors or specific exposure events, limiting

causal inference. These constraints also restricted our ability to distinguish the behavioral and

biological drivers underlying the observed patterns. For example, exposure risk and testing

practices may vary in correlated ways between facility types, and vaccination status may itself

be associated with differences in risk-taking behavior, not captured in this study. In addition,

clinical data were available for only a subset of individuals and are often complex and

subjective, limiting the scope of related analyses, while vaccination status was unavailable for

uninfected individuals, precluding more detailed assessment of vaccine-associated effects. As a

result, analyses requiring deeper integration of clinical or behavioral data could not be pursued

robustly.

439 Acknowledgements

We thank members of the Massachusetts Department of Public Health and Broad Clinical Labs

(formerly Broad's Clinical Research Sequencing Platform) for their efforts and support of this

work. We are also grateful to the laboratories that shared SARS-CoV-2 sequence data publicly

in GenBank. We thank Miguel Paredes for helpful feedback on the draft manuscript and

analyses.

Funding: Samples analyzed were from CDC-funded COVID-19 diagnostic testing and genomic

surveillance programs (75D30121C10501 and 75D30122C13720 to the Broad Institute Clinical

Research Sequencing Platform, LLC) and through the U.S. CDC Pathogen Genomic Centers of

Excellence (INTF5104H78W22195346 to P.C.S. and B.L.M.). This work was also funded by a

CDC Broad Agency Announcement (75D30120C09605 to B.L.M.), a Howard Hughes Medical

Institute Investigator Award to P.C.S., the National Institute of Allergy and Infectious Diseases

(U19AI110818, to P.C.S.), the Rockefeller Foundation (2021HTH013 to B.L.M. and P.C.S.), the

National Human Genome Research Institute (T32HG010464-06 to G.K.M.), the National

Institute of General Medical Sciences (T32GM007753 and T32GM144273 to B.P.), and the

Broad MD-PhD Fellowship (to B.P.).

Competing interests: P.C.S. is a co-founder of, shareholder in Delve Bio; she was formerly a

co-founder of and shareholder in Sherlock Biosciences, Inc and a Board member of and

shareholder in Danaher Corporation.

Author contributions

Conceptualization: G.K.M., T.B.-F., L.A.K., S.F.S., C.M.B., B.L.M., P.C.S.

Methodology: G.K.M., T.B.-F., L.A.K., S.F.S., M.B., B.P., I.S., S.W., D.J.P.

Software: G.K.M., T.B.-F., L.A.K., S.F.S., M.B., C.E.C., K.S.M., B.P., I.S., S.W., D.J.P.

Formal analysis: G.K.M., T.B.-F., L.A.K., S.F.S., M.B., C.E.C., K.S.M., B.P., I.S., S.W., D.J.P.

Investigation: G.K.M., T.B.-F., L.A.K., M.B., C.E.C., K.S.M., B.P., I.S., K.C.D., K.J.S., C.L.,

N.A.F., H.M.R.

Resources: S.B.G., S.S., S.W., L.C.M., C.M.B., B.L.M., P.C.S.

Data curation: G.K.M., T.B.-F., L.A.K., M.B., C.E.C., K.S.M., N.A.F., H.M.R.

Writing – original draft: G.K.M., T.B.-F., L.A.K., S.F.S., B.L.M.

Writing – review & editing: G.K.M., T.B.-F., L.A.K., S.F.S., B.L.M., P.C.S.

Supervision: G.K.M., S.F.S., D.J.P., L.C.M., C.M.B., B.L.M., P.C.S.

Project administration: G.K.M., T.B.-F., L.A.K., C.M.B., B.L.M.

Funding acquisition: C.M.B., L.C.M., P.C.S., B.L.M.

All authors reviewed and approved the manuscript.

Declaration of interests

B.L.M. and P.C.S hold patents related to genomic sequencing technologies. P.C.S. is a co-founder of, shareholder in Delve Bio and Lyra Labs; she was formerly a co-founder of and shareholder in Sherlock Biosciences, Inc and a Board member of and shareholder in Danaher Corporation.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used ChatGPT in order to refine language in the writing process. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the published article.

Figure titles and legends

Figure 1. Study spans multiple viral lineages across diverse settings and demographic groups. A) Number of SARS-CoV-2 genomes sequenced per week attributed to each lineage in the complete dataset. The asterisk denotes that the lineage label (e.g., BA.1*) includes all sublineages. B) Multinomial regression modeling logistic growth of Delta and Omicron sublineages in the complete dataset. Shaded regions indicate 95% confidence intervals; points represent daily lineage frequencies from sequencing data. C) Number of sequenced genomes stratified by testing sector. D) Blue dashed line indicates the percentage of confirmed cases

sequenced per day in this study shown as a 7-day rolling average (left y-axis). Gray shaded

region shows SARS-CoV-2 case counts reported to Massachusetts Department of Public Health

from November 1, 2021 to January 17, 2023 (7-day rolling average; right y-axis). The horizontal

black line marks the United States Center for Disease Control and Prevention's national target

of sequencing ≥5% of confirmed cases.

Figure 2. Heterogeneous transmission across facility types and age-defined

subpopulations. A) Bootstrap distributions of fold-enrichment (x-axis) of closely related viruses

within a facility relative to relatedness between the facility and surrounding municipality. The

dashed line indicates the mean enrichment for each testing sector. B) Age-stratified enrichment

of closely related viruses within each sector, defined as the difference in probability of a close

viral relative within the same facility versus in public testing data from the individual's

municipality of residence. Shown for age groups with >30 samples (3-year rolling average;

bootstrapped 95% confidence bands). C) Logistic regression of BA.1* rise over time (top) and

estimated date at which lineage frequency reached 50% for BA.1*, BA.2*, BA.2.12.1*, and

BA.5* across sectors (bottom, with 95% confidence intervals).

Figure 3. Viral movement within Massachusetts driven by larger and denser populations.

A) Percentage of confirmed cases sequenced in each municipality during the study period. B)

Mean number of inferred introductions into each municipality, averaged across 10 phylogenetic

trees and all six major lineages, stratified by municipality type. C) Logistic regression of BA.1*

(top) and BA.2* (bottom) lineage frequency over time, showing estimated date at which lineage

reached 50% frequency in municipalities of different urbanicity classes (95% confidence

intervals). D) Proportion of virus pairs within specified genetic distance (as a fraction of all pairs)

across geographic distance between municipalities of residence. Closely related pairs (≤2

substitutions) are shown in blue. E) Relative enrichment of closely related virus pairs across

different municipal settings. "Rural" includes rural municipalities and developing suburbs;

"urban" includes maturing suburbs, regional urban centers, and the inner urban core. The

dashed line indicates background relatedness for each category.

Figure 4. Booster vaccination is associated with reduced onward SARS-CoV-2

transmission. A) Proportion of individuals in school-based testing who were 5–11 years old,

comparing November 2021 with late December 2021/early January 2022. B) Fraction of the

Massachusetts population represented in the public testing dataset by age during November

2021 (blue) versus late December 2021/early January 2022 (gray). Plots are normalized to have

equal areas. C) Proportion of linked (coral) and unlinked (dark purple) cases across vaccination

categories in the complete dataset; linked individuals have closely related viral genomes

elsewhere in the dataset and are therefore more likely to have been involved in transmission

chains. D) Proportion of individuals with a detected descendant virus as, identified through

shared intrahost single nucleotide variants (iSNVs), stratified by vaccination status in the public

testing dataset.

Figure 5. Genomic sampling intensity determines the speed of variant detection. A)

Estimated timing of detection of a rising variant relative to the date it reaches 1% frequency in

the dataset, across different weekly sampling rates (95% credible intervals). B) Subsampling

analysis of BA.1* empirical data at N genomes per week. Logistic regression models were fit

beginning one day after BA.1* reached 3% frequency in the dataset. Shown is the probability

that the 95% confidence interval of the estimated growth rate exceeds 0, indicating strong

evidence for logistic growth. Red colors denote probabilities >50%. C) Same as in (B), applied

to BA.5*.

538   STAR Methods

539   RESOURCE AVAILABILITY

540   Lead Contact

541   Further information and requests for resources and reagents should be directed to and will be

542   fulfilled by the lead contact, Gage K. Moreno (gmoreno@broadinstitute.org).

543   Materials availability

544   This study did not generate new unique reagents.

545   Data and code availability:

-    Raw reads for all samples (including those that did not produce a successful genome)

were deposited in NCBI SRA. All NCBI data were deposited under BioProject

PRJNA715749.

-    All code used for sequence data processing, genome assembly, and phylogenetic

analysis is publicly available either via the Dockstore Tool Registry Service

(dockstore.org/organizations/BroadInstitute/collections/pgs) or on GitHub

(github.com/AndrewLangvt/genomic_analyses/blob/main/workflows/wf_viral_refbased_a

ssembly.wdl).

-      For any additional information required to reanalyze the data reported in this paper,

contact the lead author.

556 EXPERIMENTAL MODEL AND SUBJECT DETAILS

Ethical approvals

As previously described[5], the research project (Protocol #1603078) was reviewed and approved

by the Massachusetts Department of Public Health (MADPH) Institutional Review Board and

covered by a reliance agreement at the Broad Institute. An additional non-human subjects

research determination was made by the Harvard Longwood Campus Institutional Review

Board. A summary of all available metadata categories and the number of individuals within

each category is provided in **Table 1**.

564 METHOD DETAILS

SARS-CoV-2 RT-qPCR

For specimens submitted for testing under Emergency Use Authorization to the Broad Institute's

CLIA-certified laboratory, Clinical Research Sequencing Platform (now Broad Clinical Labs),

total RNA was extracted from inactivated anterior nasal (AN) swabs using the Thermo Fisher

MagMAX Viral RNA Isolation kit and presence of virus was confirmed by RT-qPCR assay

detecting the N1 and N2 SARS-CoV-2 gene regions. Ct values for the N1 gene were used to

compare viral titers between individuals; samples for which the RP positive control gene had a

Ct>32 were excluded from the analyses to prevent biasing from poorly collected samples. A

specimen was determined positive for SARS-CoV-2 if the initial cycle threshold of the N1 probe

was less than or equal to Ct 40. As seen in previous studies[24,25], clinical diagnostic PCR cycle

threshold (Ct) values were significantly lower for Delta than for the Omicron sublineages, with

little variation among the latter (**Fig S14**).

SARS-CoV-2 sequencing

Candidate samples were re-extracted from the source material for Illumina sequencing.

Libraries were prepared using the NEBNext ARTIC SARS-CoV-2 FS Library Prep Kit and

transitioned through ARTIC, VarSkip Short v1, v1a, or v2a amplicon primers to ensure genomic

coverage of emerging variants. During library preparation, some volumes were adjusted from

manufacturer recommendations to accommodate 384-well plate reactions and high-throughput

automated processing. Libraries were sequenced on Novaseq SP flowcells with 75-nucleotide

paired-end reads.

Genome assembly and analysis

For genomes generated at the Broad Institute, we conducted all analyses using viral-ngs 2.1.28

on the Terra platform (app.terra.bio). All of the workflows named below are publicly available via

the Dockstore Tool Registry Service

(dockstore.org/organizations/BroadInstitute/collections/pgs). Briefly, samples were

demultiplexed, reads were filtered for known sequencing contaminants, and SARS-CoV-2 reads

were assembled using a reference-based assembly approach with the SARS-CoV-2 isolate

Wuhan-Hu-1 reference genome NC_045512.2 (using workflow sarscov2_illumina_full.wdl;

publicly available on GitHub). We processed all raw read data using a reference-based

consensus calling method with the same NC_045512.2 reference genome. Assembled

genomes meeting the CDC criteria for submission to public repositories (unambiguous length

≥24,000 nt and successful gene annotation) were deposited in NCBI Genbank and GISAID

immediately upon completion. Raw reads for all samples (including those that did not produce a

successful genome) were deposited in NCBI SRA. All NCBI data were deposited under

BioProject PRJNA715749.

We used LoFreq version 2.1.5 to call intrahost single nucleotide variants (iSNVs) with default

parameters (minimum read depth ≥10, strand bias <85%, and default iSNV quality scoring)[26].

Variants with frequency ≥3% and in positions with read depth ≥100 were used for downstream

analysis. Minor variants arising in primer binding sites were masked[27]. All non-fixed consensus

mutations (<97% and >50%) were flipped to their corresponding minor variant (e.g. a mutation

at C11803T at 65%, would now show up as a T11803C at 35%). An additional filtering step was

applied to account for sequencing bias[28,29]. Briefly, a binomial distribution was applied that took

into account the total number of variant reads, sequencing depth, and sequencing error rate.

Variants with significance (≤0.05) after applying a Bernoulli correction were kept for downstream

analysis[29].

Identification of reinfections and coinfections

Among the 81,952 unique individuals, 2,714 had more than one genome sequence available

within a 90-day window, corresponding to the timeframe during which repeated tests per

individual could be tracked in the Massachusetts public health database (**Fig S1**). For

individuals with multiple sequenced samples, we recorded the pairwise genetic distances

between consensus genomes and the number of days between sample collections. Each

genome was assigned a Pango lineage using pangolin (v4.1.3). Samples collected more than

14 days apart that were assigned to different Pango lineages were classified as reinfections,

according to the CDC definition[30]. Individuals with multiple samples collected within 14 days

were compared for lineage consistency; those showing identical lineage assignments were

considered to have the same infection, while those with discordant lineages were classified as

potential multiple infections within a short interval. Pairwise distances and collection intervals

were visualized and colored by Pango lineage to confirm lineage differences and reinfection

events (**Fig S1**).

To detect simultaneous infection with multiple viral lineages, we first assigned a consensus

Pango lineage to each genome using pangolin (v4.1.3). We then compiled a list of major

lineage-defining mutations for all major lineages (BA.1, BA.2, BA.2.12.1, BA.5, BQ.1, XBB).

Samples were classified as co-infections if two or more lineage-defining mutations from a

different lineage were present as iSNVs. Only samples with sufficient genome coverage (>90%

completeness) and sequencing depth (>100x mean coverage) were considered to minimize

false positives. The majority of identified co-infections involved Delta and BA.1 lineages,

consistent with the high transmission period during the early BA.1 wave.

Phylogenetic tree construction

All genomes paired with metadata during the study period (November 1, 2021, to January 17,

2023) were used for phylogenetic inference (n=94,404). Prior to phylogenetic inference,

genomes were aligned using Nextclade CLI v1.3.0. Genomes were excluded if they 1)

contained any mixed bases, 2) were missing >10% of the SARS-CoV-2 genome, 3) lacked

adequate metadata, or 4) were denoted as "bad" during Nextclade QC. The remaining

(n=91,080) genomes were masked using a light masking scheme[31].

To understand the number of introductions into each municipality, we generated 60 variant-

specific (10 per variant) maximum likelihood phylogenetic trees using publicly available SARS-

CoV-2 genomes from Genbank (retrieved January 23, 2024). Datasets were filtered to only

include genomes collected prior to the date that each major lineage reached 50% frequency in

MA. The resulting dataset was further subsampled so that 1% of confirmed infections were

sequenced within each geographic resolution (country, state, municipality), as the geographic

distribution of cases in Massachusetts was skewed towards highly urban areas. We constructed

a maximum-likelihood (ML) phylogenetic tree using FastTree[32] with 100 bootstraps and

otherwise default parameters.

Under this case-normalized sampling strategy, replicate tree sizes varied modestly by lineage:

BA.1 trees contained 72,356–72,746 genomes; BA.2.12.1 trees contained 8,721–8,803

genomes; BA.5 trees contained 39,368–39,717 genomes; BQ.1 trees contained 6,605–6,772

genomes; and XBB trees contained 1,893–1,933 genomes.

We generated a set of temporally balanced subsampled phylogenetic trees. First, using the

augur filter tool, we randomly sampled 2,000 genomes per month from the USA and from the

global dataset, respectively, prioritizing sequences with close genetic similarity to samples in our

study. Second, we included 5 genomes per week from each Massachusetts municipality with an

average sequencing rate of at least five genomes per week.

Under this temporally balanced strategy, final tree sizes were smaller and more uniform across

lineages: BA.1 trees contained 9,601 genomes (5,840 from Massachusetts); BA.2 trees

contained 4,211 genomes (1,129 from Massachusetts); BA.2.12.1 trees contained 4,206

genomes (468 from Massachusetts); BA.5 trees contained 3,616 genomes (542 from

Massachusetts); BQ.1 trees contained 3,609 genomes (147 from Massachusetts); and XBB

trees contained 3,985 genomes (75 from Massachusetts).

Using custom scripts, the resulting phylogenies were processed using the refine, ancestral,

traits, and export steps of the Nextstrain pipeline[33].

Viral introductions

To identify viral introductions into Massachusetts municipalities, we assigned a trait to each

genome in the phylogeny of either "out-of-state" or the name of the municipality. We used

Nextstrain's ancestral inference[33] to infer the state of that trait for each internal node of the tree.

We defined an introduction by traversing the phylogenetic tree using baltic[34], identifying

branches in which an ancestral node was inferred to be out of state with ≥70% confidence and

gave rise to a descendant node inferred to be a Massachusetts municipality with ≥70%

confidence.

For viral movements occurring between Massachusetts municipalities, we applied the same

logic, defining a viral movement between municipalities as instances of an ancestral node and

descendant node being different municipalities, both with confidence ≥70%. We combined the

movement data between origin and destination municipalities with associated variables such as

case counts, percent positivity, and population size and density for both origin and destination.

Movements across the 10 replicate phylogenies for each variant were deduplicated on the

origin, destination, date, and downstream genomes. The remaining movements were then

counted based on origin, destination, and date. We used a Generalized Linear Model (GLM)

with a Gaussian family and identity link function to examine the relationship between movement

counts and various predictors. The model was specified as:

Movement_count $= \beta 0 + \beta 1(\text{predictors}) + \epsilon$

Model fitting was performed using the Iteratively Reweighted Least Squares (IRLS) method in

the statsmodels library v0.14.2 [35] in python. Model performance was assessed using the Akaike

Information Criterion (AIC) and log-likelihood.

Normalized age distribution pre- and post-vaccine rollout, ages 5–11

The age distribution was calculated for all public testing samples with submittable genomes and

recorded subject ages during two periods, November 1, 2021 – November 30, 2021 and

December 15, 2021 – January 13, 2022. The statewide population by age was interpolated

using a cubic spline from US Census estimated data for 2021 (

https://www.census.gov/data/tables/time-series/demo/popest/2020s-state-detail.html). The two

distributions are normalized to have equal areas.

Pairwise analysis

First, all possible pairs of genomes within the same lineage, collected within 10 days of each

other in the public testing sector were identified. Then, the genetic distance between all possible

pairs, defined as the Hamming distance between genomes only including positions at which

both genomes had confident reads, was calculated using the ape[36] package v5.7.1 in R.

Additionally, we similarly calculated all pairs among all genomes collected at SNFs, other

workplaces, schools, and colleges.

Geographic analysis

For all resulting genome pairs, we calculated geographic distance (calculated as haversine

distance) between the interior points of each pair of municipalities with the R package

geosphere[37] V 1.5.18. Geographic distance between pairs was then associated with genetic

distance and plotted. BA.1* was excluded from the geographic distance versus genetic distance

analysis because it showed too little geographic structure to be informative about viral spread,

perhaps because of its rapid, nearly clonal rise or an unusually large number of introductions

(**Fig S16**). Given we observed geographic structure for genomes differing by up to 12

substitutions, while more distantly related viruses were approximately uniformly distributed

throughout the state, we can take this as the amount of divergence required for pairs to be

distributed randomly throughout the state. As a simple model to translate this into time, we can

treat the two viruses being compared as descending from a recent common ancestor, with each

descendant lineage accumulating ~0.5 mutations per generation. Thus a difference of 12

substitutions would accumulate in ~12 generations or, assuming ~5 days per generation, ~60

days.

Identification of closely related pairs

The identification of closely related cases was performed in Julia (v1.9). To identify all pairs of

closely related cases within our dataset, first a random case was selected and all cases

occurring within 10 days of this case were identified. Then the genetic distance from the initial

randomly selected case to all of these temporally linked cases was calculated, and cases with a

genetic distance of 2 or fewer SNPs were recorded as being closely related. This process was

repeated until all cases had been compared against all other temporally linked cases. Resulting

cases which had any number of additional closely related cases were then recorded as being

"linked" while cases with no additional closely related cases were "unlinked."

To compute the percent enrichment of closely related viruses within the same testing facility

compared to within the same municipality, we calculated, per virus, the percent of contemporary

(occurring within 10 days) same-facility cases which were closely related and subtracted the

percent of contemporary (occurring within 10 days) same-municipality cases collected in public

testing which were closely related. To additionally get a relative percent change, the likelihood

of close relatives in each facility was calculated on a sector level (that is, summing the number

of related cases for all cases within a given sector to avoid dividing by zero) and then divided by

the percent of contemporary closely related in public testing in the community. Due to high

variability in the likelihood of close relatives among public testing for specific age groups in each

sector, we only calculated this on average for sectors, not within specific ages in each sector.

We then averaged all cases per sector to report summary findings. To describe these patterns

over age groups we computed overlapping, 3-year rolling averages per sector. Age groups

which had fewer than 25 viruses for comparison were dropped from the analysis.

Detection of putative transmission events

An individual was defined as a transmitter when an index and contact met the following criteria:

1) the index and contact consensus sequences differed by no more than two consensus-level

changes, 2) the consensus level changes in the contact were detected at sub-consensus levels

in the index, 3) the contact appeared downstream of the index in a divergence phylogeny, 4) the

collection date of the index was ≤6 days before that of the contact, and 5) if two genomes were

identical, the defining mutations were at ≤90% frequency in the index, only seen <5 times in the

community, and rose to 100% frequency in the contact.

Estimation of sequencing rates to detect novel lineages

A multinomial regression model was applied to estimate the daily frequencies of variants BA.1*,

BA.2*, BA.2.12.1*, BA.5*, and BQ.1*. Using the estimated variant frequencies, a simple model

was used to estimate the earliest possible detection dates. This model incorporates a binomial

draw, accounting for both sampling rates and the estimated frequencies of each variant,

outputting the number of variant genomes that would be detected on a given day. This model

was used to calculate the probability of detecting a variant genome by running it 1,000 times per

genome sequencing rate (10, 50, 100, 200, up to 1,000 genomes per week). The resulting

probabilities were used to estimate a 95% credible interval of the first date of detection for each

variant under each sequencing rate.

We additionally applied the vartrack_prob_detect_cont function from PhyloSamp[19] to verify

these modeling results. For this function we set the coefficient of detection ratio to 1 (i.e.

assuming no difference in the probability of detection between lineages), the initial prevalence to

0.0001, and the percent of samples which generate high quality genomes to 100% (to reflect

that the modelling approach we employed relied on the number of high quality genomes). This

function was then run with logistic growth rates of 0.075, 0.125, 0.175, and 0.225 and sampling

rates ranging from 7 genomes per week to 1000 genomes per week.

Estimation of sequencing rates to detect growth of novel lineages

To estimate how many genomes would be needed and when growth of variants would be

detected after reaching 3% frequency, the dataset was subsampled 100 times at rates of 10, 50,

100, 200, 300, and so on, up to 1,000 genomes per week. For each subsample, the date when

the lineage reached 3% frequency was recorded. A logistic regression was generated starting

from day 1 (after reaching 3% frequency) through 30 days, capturing the day when the growth

rate became positive with 95% probability. The probability was plotted that a positive 95%

growth rate could be detected within each sampling rate N days after the lineage reached 3%

frequency.

Statistical comparison of date when each lineage reached 50% frequency

A logistic growth model was used to analyze the growth trajectories of Omicron sublineages

(BA.1*, BA.2*, BA.2.12.1*, and BA.5*) in each sector. For each sector, the logistic growth model

was fit using nonlinear least squares (NLS) regression to estimate the parameters growth rate

and lineage frequency. The model for each lineage was constrained to only consider data

starting from the first day where the proportion of genomes belonging to the lineage exceeded

0. The fitted models were used to generate predicted frequency of each lineage over time.

The tsum.test() function within BSDA[38] was used to compare the growth rates of the variants

between sectors. Specifically, the significance was tested of the differences between sectors in

the timing of when each variant reached 50% frequency. The tsum.test() function was used

while accounting for standard errors and sample sizes. A 95% confidence level was applied to

assess the significance of differences in the growth rates and 50% threshold timing.

Statistical analysis

All statistical analyses were performed in R (The R Core team, 2017), python 3, or Julia (v1.9).

Pairwise ranksum tests for comparisons of Ct values were performed using the ranksum

function in scipy[39]. All multiple comparisons were corrected using multipletests in statsmodels[35]

using the Benjamini-Hochberg correction. All additional statistical details including sample

numbers can be found in the figure legends.

Supplemental information titles and legends

Figure S1. Genomic signatures of repeated and mixed SARS-CoV-2 infection. Within our

dataset, we were able to link genomes back to an individual within 90 days of their first genome.

A) Number of mixed lineage infections. X-axis: lineage of the consensus genome. Y-axis:

lineage detected at sub-consensus frequencies. B) Number of mutations separating genomes

collected from the same individual vs time between their collection dates. Colors indicate the

same (gray) or different (blue) Pango lineages for the two genomes. Dashed lines delimit

candidate region for evolution in persistently infected individuals (>14 days and >2 mutations).

C) Histogram showing the number of genomes per individual among those with more than one

genome in the dataset. D) Histogram of the number of days between sequential genomes from

the same individual; the dotted line denotes the median interval (4 days). E) Kernel density plot

relating time between sample collection to the number of mutations observed; samples

belonging to the same lineage cluster below 2 mutations and within 14 days, consistent with a

single infection. F) Monthly prevalence of individuals with detected mixed infections, overlaid on

a multinomial regression of circulating lineage frequencies.

Figure S2. Temporal and demographic distribution of sequenced genomes. A–C)

Temporal distribution of the number of genomes collected grouped by A) age, B) municipality

type, and C) gender. (D–E) Number of genomes collected within each facility type by D) age

and E) gender.

Figure S3. Overview of testing sectors and sampling intensity. Left, table summarizing the

testing sectors included in the dataset, showing the total number of genomes and the number of

unique facilities per sector. Right, weekly histogram of genomes collected from each testing

sector; the orange line indicates the total number of unique facilities contributing data in each

week.

Figure S4. Genomes collected from public testing by age, as a fraction of the

Massachusetts census population of that age.

Figure S5. Robustness of sector-specific relatedness to subsampling. Data were randomly

subsampled per sector such that the sample size was the same in each sector and then

bootstrap samples of fold-enrichment (x-axis) of closely related viruses within a facility versus

between facility and municipality were plotted; dashed line: testing sector mean.

Figure S6. Per facility enrichment of closely related viruses. Scatterplot of the average

enrichment in closely related viruses per facility compared to the square root of the total number

of contemporary viral pairs in each facility, which is proportional to the total number of genomes

each facility contributed to the complete dataset.

Figure S7. Age-specific likelihood of close viral relatedness across settings. Likelihood of

identifying close viral relatives among cases within the same facility (colored lines) and among

cases tested at public facilities within the same municipality (grey), stratified by age and sector.

Curves show 3-year rolling averages for age groups with more than 30 samples; shaded

regions denote bootstrapped 95% confidence intervals.

Figure S8. Age-specific relatedness within public testing facilities. Likelihood of identifying

close viral relatives within the same testing facility as a function of age among public testing

sites. Curves show 3-year rolling averages for age groups with more than 30 samples; shaded

regions represent bootstrapped 95% confidence intervals.

Figure S9. Lineage growth dynamics in college-aged individuals. Logistic regressions of

four SARS-CoV-2 lineages prevalence among 18- to 22-year-olds in college testing settings

(teal) and in public testing (dark blue), compared with all public testing data (grey). Estimated

dates at which each lineage reached 50% frequency are shown below, with 95% confidence

intervals where available; for BA.5*, insufficient data precluded estimation of a 95% confidence

interval for the 50% date.

Figure S10. Lineage growth dynamics across testing settings. Logistic regression fits and

estimated dates at which lineage frequency reached 50% (with 95% confidence intervals) for

BA.1, BA.2, BA.2.12.1, and BA.5, shown from top left to bottom right. Results are stratified by

testing setting: public testing (blue), colleges (yellow), hospitals and clinics (red), and schools,

employers, and skilled nursing facilities (green).

Figure S11. Viral introductions scale with municipality population size and density

across lineages. Datasets were restricted to genomes collected prior to the date at which each

major lineage reached 50% frequency in Massachusetts and were further subsampled to

sequence 1% of confirmed infections within each geographic level (country, state, municipality)

to account for urban–rural sampling biases. A) Scatter plots of the number of detected

introductions per municipality as a function of municipality population, shown separately for

each lineage; solid lines indicate linear regression fits with shaded 95% confidence intervals. B)

Scatter plots of the number of detected introductions per municipality as a function of

municipality population density, shown separately for each lineage; solid lines indicate linear

regression fits with shaded 95% confidence intervals. C) Scatter plots of the number of detected

introductions per municipality as a function of municipality population (left) and population

density (right), averaged across all lineages; solid lines indicate linear regression fits with

shaded 95% confidence intervals. Blue lines show the regression fits including Boston. Gray

lines show regression fits removing Boston. The positive trends remain after removing Boston,

although their strength is reduced.

Figure S12. Vaccination uptake and infection timing in children. A) Cumulative percentage

of individuals aged 5–11 years receiving their first (black) and second (grey) COVID-19 vaccine

doses following vaccine availability. B) Biweekly histograms showing the proportion of the

Massachusetts population infected with Delta (top) and BA.1* (bottom) over time, stratified by

age.

Figure S13. Transmission patterns by demographic and clinical features. A) Proportion of

males and females who initiated at least one transmission event. B) Proportion of males and

females who were linked versus unlinked. C) Proportion of individuals initiating a transmission

event stratified by symptom status; symptom data were restricted to reports prior to January 1,

2022, and no significant difference was observed (p = 0.1). D) Proportion of symptomatic and

asymptomatic individuals who were linked versus unlinked. E) Proportion of individuals initiating

a transmission event by vaccination timing among vaccinated individuals; "early" vaccination

corresponds to Feb 1 - May 1, 2021, while "late" vaccination corresponds to vaccination after

booster availability (from Nov, 2021 - Jan 2023). F) Proportion of individuals initiating a

transmission event by age group.

Figure S14. Factors associated with SARS-CoV-2 viral load (Ct). A) Heatmap of pairwise

comparisons of lineage-specific Ct values; colored cells indicate significant differences after

multiple-testing correction. The asterisk denotes the lineage label (e.g., BA.1*) and includes all

sublineages following that designation, according to WHO/Nextstrain nomenclature. B)

Temporal trends in mean lineage Ct values, showing lower Cts during lineage emergence and

higher Cts during decline; shaded regions denote 95% confidence intervals. C) Violin plot

comparing Ct values in symptomatic and asymptomatic cases. D) Ct values decrease with

increasing time since last vaccination (-0.002 Ct per day; $p < 2 \times 10^{-16}$). E) Ct values decrease

modestly with age (-0.001 Ct per year). F) Small but statistically significant differences in Ct by

sex. G) Ct values increase with number of vaccine doses ($+0.14$ Ct per dose; $p < 2 \times 10^{-16}$).

Figure S15. Sampling requirements for early lineage detection. Application of PhyloSamp to

estimate the number of high-quality genomes required to detect emerging lineages before they

reach 1% prevalence. Curves are shown for growth rates comparable to BA.1 (fast-growing),

BA.5 (slow-growing), and intermediate rates. The x-axis (plotted in reverse for comparability to

Fig. 4A) shows the empirical probability of detecting a lineage before 1% prevalence, assuming

logistic growth from an initial prevalence of 0.0001. The y-axis indicates the number of high-

quality genomes generated per week.

Figure S16. Reduced geographic structure during BA.1 emergence. Relationship between

genetic and geographic distance, shown as the density of the fraction of sequence pairs at a

given SNP difference, during the exponential phase of BA.1* emergence (top; cases detected

before December 21, 2021) and during the subsequent plateau and decline phase (bottom).

References

1. Lopes, R., Pham, K., Klaassen, F., Chitwood, M.H., Hahn, A.M., Redmond, S., Swartwood,

N.A., Salomon, J.A., Menzies, N.A., Cohen, T., et al. (2024). Combining genomic data and

infection estimates to characterize the complex dynamics of SARS-CoV-2 Omicron variants

in the US. Cell Rep *43*, 114451.

2. Tran-Kiem, C., Paredes, M.I., Perofsky, A.C., Frisbie, L.A., Xie, H., Kong, K., Weixler, A.,

Greninger, A.L., Roychoudhury, P., Peterson, J.M., et al. (2025). Fine-scale patterns of

SARS-CoV-2 spread from identical pathogen sequences. Nature, 1–10.

3. Hamilton, W.L., Tonkin-Hill, G., Smith, E.R., Aggarwal, D., Houldcroft, C.J., Warne, B.,

Meredith, L.W., Hosmillo, M., Jahun, A.S., Curran, M.D., et al. (2021). Genomic

epidemiology of COVID-19 in care homes in the east of England. eLife *10*, e64618.

4. Lindsey, B.B., Villabona-Arenas, C.J., Campbell, F., Keeley, A.J., Parker, M.D., Shah, D.R.,

Parsons, H., Zhang, P., Kakkar, N., Gallis, M., et al. (2022). Characterising within-hospital

SARS-CoV-2 transmission events using epidemiological and viral genomic data across two

pandemic waves. Nature Communications *13*, 671.

5. Siddle, K.J., Krasilnikova, L.A., Moreno, G.K., Schaffner, S.F., Vostok, J., Fitzgerald, N.A.,

Lemieux, J.E., Barkas, N., Loreth, C., Specht, I., et al. (2022). Transmission from

vaccinated individuals in a large SARS-CoV-2 Delta variant outbreak. Cell *185*, 485–

492.e10.

6. Oltean, H.N., Black, A., Lunn, S.M., Smith, N., Templeton, A., Bevers, E., Kibiger, L.,

Sixberry, M., Bickel, J.B., Hughes, J.P., et al. (2024). Changing genomic epidemiology of

COVID-19 in long-term care facilities during the 2020–2022 pandemic, Washington State.

BMC Public Health *24*, 182.

7. Page, A.J., Mather, A.E., Le-Viet, T., Meader, E.J., Alikhan, N.-F., Kay, G.L., de Oliveira

Martins, L., Aydin, A., Baker, D.J., Trotter, A.J., et al. (2021). Large-scale sequencing of

SARS-CoV-2 genomes from one region allows detailed epidemiology and enables local

outbreak management. Microb Genom *7*. https://doi.org/10.1099/mgen.0.000589.

8. Total COVID-19 tests Our World in Data. https://ourworldindata.org/grapher/full-list-total-

tests-for-covid-19?country=~USA.

9. CDC (2020). COVID Data Tracker. Centers for Disease Control and Prevention.

https://covid.cdc.gov/covid-data-tracker.

10. Tracking SARS-CoV-2 variants https://www.who.int/activities/tracking-SARS-CoV-2-

variants.

11. Tran-Kiem, C., Paredes, M.I., Perofsky, A.C., Frisbie, L.A., Xie, H., Kong, K., Weixler, A.,

Greninger, A.L., Roychoudhury, P., Peterson, J.M., et al. (2025). Fine-scale patterns of

SARS-CoV-2 spread from identical pathogen sequences. Nature, 1–10.

12. Braun, K.M., Moreno, G.K., Wagner, C., Accola, M.A., Rehrauer, W.M., Baker, D.A., Koelle,

K., O'Connor, D.H., Bedford, T., Friedrich, T.C., et al. (2021). Acute SARS-CoV-2 infections

harbor limited within-host diversity and transmit via tight transmission bottlenecks. PLoS

Pathog. *17*, e1009849.

13. Bendall, E.E., Callear, A.P., Getz, A., Goforth, K., Edwards, D., Monto, A.S., Martin, E.T.,

and Lauring, A.S. (2023). Rapid transmission and tight bottlenecks constrain the evolution

of highly transmissible SARS-CoV-2 variants. Nature Communications *14*, 272.

14. Petros, B.A., Turcinovic, J., Welch, N.L., White, L.F., Kolaczyk, E.D., Bauer, M.R., Cleary,

M., Dobbins, S.T., Doucette-Stamm, L., Gore, M., et al. (2023). Early Introduction and Rise

of the Omicron Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Variant

in Highly Vaccinated University Populations. Clin. Infect. Dis. *76*, e400–e408.

15. Petros, B.A., Paull, J.S., Tomkins-Tinch, C.H., Loftness, B.C., DeRuff, K.C., Nair, P.,

Gionet, G.L., Benz, A., Brock-Fisher, T., Hughes, M., et al. (2022). Multimodal surveillance

of SARS-CoV-2 at a university enables development of a robust outbreak response

framework. Med (New York, N.Y.) *3*. https://doi.org/10.1016/j.medj.2022.09.003.

16. Lythgoe, K.A., Hall, M., Ferretti, L., de Cesare, M., MacIntyre-Cockett, G., Trebes, A.,

Andersson, M., Otecko, N., Wise, E.L., Moore, N., et al. (2021). SARS-CoV-2 within-host

diversity and transmission. Science *372*. https://doi.org/10.1126/science.abg0821.

17. Eyre, D.W., Taylor, D., Purver, M., Chapman, D., Fowler, T., Pouwels, K.B., Sarah Walker,

A., and Peto, T.E.A. (2022). Effect of Covid-19 Vaccination on Transmission of Alpha and

Delta Variants. N. Engl. J. Med. https://doi.org/10.1056/NEJMoa2116597.

18. Woodbridge, Y., Amit, S., Huppert, A., and Kopelman, N.M. (2022). Viral load dynamics of

SARS-CoV-2 Delta and Omicron variants following multiple vaccine doses and previous

infection. Nature Communications *13*, 6706.

19.   Wohl, S., Lee, E.C., DiPrete, B.L., and Lessler, J. (2023). Sample size calculations for

pathogen variant surveillance in the presence of biological and systematic biases. Cell Rep

Med *4*, 101022.

20.   Brito, A.F., Semenova, E., Dudas, G., Hassler, G.W., Kalinich, C.C., Kraemer, M.U.G., Ho,

J., Tegally, H., Githinji, G., Agoti, C.N., et al. (2022). Global disparities in SARS-CoV-2

genomic surveillance. Nat. Commun. *13*, 7003.

21.   Vavrek, D., Speroni, L., Curnow, K.J., Oberholzer, M., Moeder, V., and Febbo, P.G. (2021).

Genomic surveillance at scale is required to detect newly emerging strains at an early

timepoint. bioRxiv. https://doi.org/10.1101/2021.01.12.21249613.

22.   Digest of Education Statistics,

23.   US Census Bureau (2026). State Population Totals and Components of Change: 2020-

2025.

24.   Chaguza, C., Coppi, A., Earnest, R., Ferguson, D., Kerantzas, N., Warner, F., Young, H.P.,

Breban, M.I., Billig, K., Koch, R.T., et al. (2022). Rapid emergence of SARS-CoV-2

Omicron variant is associated with an infection advantage over Delta in vaccinated

persons. Med (New York, N.Y.) *3*. https://doi.org/10.1016/j.medj.2022.03.010.

25.   Li, K., Chaguza, C., Stamp, J., Chew, Y.T., Chen, N.F.G., Ferguson, D., Pandya, S.,

Kerantzas, N., Schulz, W., Yale SARS-CoV-2 Genomic Surveillance Initiative, et al. (2024).

Genome-wide association study between SARS-CoV-2 single nucleotide polymorphisms

and virus copies during infections. PLoS Comput Biol *20*, e1012469.

26.   Wilm, A., Aw, P.P.K., Bertrand, D., Yeo, G.H.T., Ong, S.H., Wong, C.H., Khor, C.C., Petric,

R., Hibberd, M.L., and Nagarajan, N. (2012). LoFreq: a sequence-quality aware, ultra-

sensitive variant caller for uncovering cell-population heterogeneity from high-throughput

sequencing datasets. Nucleic Acids Res. *40*, 11189–11201.

27. VarSkip: VarSkip multiplex PCR designs for SARS-CoV-2 sequencing (Github).

28. VCFgenie: Program for reproducibly filtering VCF files and eliminating false positive

variants (Github).

29. Nelson, C.W., Ardern, Z., Goldberg, T.L., Meng, C., Kuo, C.-H., Ludwig, C., Kolokotronis,

S.-O., and Wei, X. (2020). Dynamically evolving novel overlapping gene as a factor in the

SARS-CoV-2 pandemic. Elife *9*. https://doi.org/10.7554/eLife.59633.

30. Muehleman, D., Gruner, B., Hogan, V., Fanning, P., Garrett, C., Meyer, J., Lanter, K.,

Purves, S., DeMarcus, L., Thervil, J., et al. (2024). Using SARS-CoV-2 Sequencing Data to

Identify Reinfection Cases in the Global Emerging Infections Surveillance Program, United

States. Emerg Infect Dis *30*, 53–61.

31. Roemer, C. exclude_sites_light.txt at main · corneliusroemer/ncov-simplest (Github).

32. Price, M.N., Dehal, P.S., and Arkin, A.P. (2009). FastTree: computing large minimum

evolution trees with profiles instead of a distance matrix. Mol. Biol. Evol. *26*, 1641–1650.

33. Hadfield, J., Megill, C., Bell, S.M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P.,

Bedford, T., and Neher, R.A. (2018). Nextstrain: real-time tracking of pathogen evolution.

Bioinformatics *34*, 4121–4123.

34. GitHub - evogytis/baltic: baltic - backronymed adaptable lightweight tree import code for

molecular phylogeny manipulation, analysis and visualisation. Development is back on the

evogytis/baltic branch (i.e. here) GitHub. https://github.com/evogytis/baltic.

35. statsmodels 0.14.4 https://www.statsmodels.org/stable/index.html.

36. Analyses of Phylogenetics and Evolution [R package ape version 5.8] (2024).

37. Hijmans, R.J. (2024). Spherical Trigonometry [R package geosphere version 1.5-20].

38. GitHub - alanarnholt/BSDA: R package for "Basic Statistics and Data Analysis" web page:

https://alanarnholt.github.io/BSDA/ GitHub. https://github.com/alanarnholt/BSDA.

39. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental

algorithms for scientific computing in Python. Nat Methods *17*, 261–272.

1009