LDSC++: Improving linkage disequilibrium score regression estimation of heritability and genetic correlation for multivariate GWAS analysis

Johan Zvrskovec[1,2(0000-0002-8862-0874)], Alexandra Gillett[1,2(0000-0002-5069-3197)], Michel Nivard[5,6,7], Jonathan R I Coleman[1,2(0000-0002-6759-0944)], Christopher Hübel[1,3,4(0000-0002-1267-8287)], Raquel Iniesta[1,2], Gerome Breen[1,2(0000-0003-2053-1792)]

1. Institute of Psychiatry, Psychology and Neuroscience, King's College London, Denmark Hill, Camberwell, London, UK
2. National Institute for Health and Care Research (NIHR) Maudsley Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, London, UK
3. Clinic for Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, German Red Cross Hospitals Berlin Westend, Berlin, Germany
4. National Centre for Register-based Research, Aarhus University, Aarhus, Denmark
5. Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, UK
6. Population Health Sciences, University of Bristol, Bristol, UK
7. Dept of biological psychology, Vrije Universiteit Amsterdam, the Netherlands

Abstract

Introduction: Linkage disequilibrium (LD) score regression is widely used for estimating common variant heritability and genetic correlations from genome-wide association study (GWAS) summary statistics. We hypothesise that segmented regression (also known as piecewise regression) improves on previous LD score regression implementations, when estimating both genetic covariance and its standard error.

Methods: We present novel extensions to LD score regression (LDSC++) improving I.) handling of varying numbers of shared genetic variants across trait pairs and reference panels, II.) estimation of genetic covariance and its variance, and III.) handling of imputation quality. We propose supporting statistical tests that use our novel extensions to improve sensitivity, and are further aimed at comparing parameter estimates that are highly correlated, such as those obtained from the same trait but from different methods. We validate LDSC++ first on real-world individual level data from the Genetic Links to Anxiety and Depression study and the United

Kingdom National Institute of Health and Social Care Research BioResource (N: 14,190 - 20,144),

second on simulated data with different degrees of shared QTL, and third on a battery of publicly

available GWASs of ten diverse traits of varying statistical power and heritability.

Results: Using variance-component method (GCTA-GREML) estimates for reference, LDSC++

extensions were found to yield heritability estimates with a bias of about -10% to -20% while

standard LD score regression yielded a bias of -30%, and heritability variability estimates with a

bias of -1% to -7% while standard LD score regression yielded a bias of 8%. For ten external trait

GWASs, LDSC++ was shown to recover 5% to 8% larger heritabilities with 4% smaller variability

on average compared to standard LD score regression. Weighting by imputation quality in the

model, rather than excluding genetic variants of low imputation quality, contributed to retaining

information. Our supporting statistical tests enabled us to detect statistically significant differences

in genetic covariance and its standard error while considering the varying number of shared

genetic variants across bivariate trait pairs.

Conclusion: LDSC++ was confirmed to produce less biassed estimates of genetic covariance

and its variability in our GLAD+ sample compared to standard LD score regression, using GCTA-

REML as reference. This performance was supported by results from external trait GWASs of

varying character, also implying an important performance of our extended weighting schemes.

Our proposed extensions to LD score regression, among which genome-wide parameters are

constructed as aggregates of heterogeneous local parameters, may prove important for large-

scale multivariate studies such as genomic structural equation models or local genetic covariance

analyses.

1 | INTRODUCTION

Genome-wide association study (GWAS) summary statistics are readily available to researchers worldwide compared to individual level genotyped or sequenced data, which typically require considerable cost and time to access due to data sharing restrictions[1,2]. This prevents the use of optimal methods that use individual level genotype data to estimate components of variance, such as GCTA-GREML[3,4], BOLT-REML[5,6] and REML in LDAK[7]. Popular alternatives[8] are Linkage Disequilibrium (LD) score regression[9] (LDSR) and SumHer[10], which instead use GWAS summary statistics and an LD reference panel to obtain common genetic variant covariance (covG) and heritability (h^2), which are standardised to estimate genetic correlations[11] (r_g). LDSR was later re-implemented for calculation of large-scale bivariate trait pair analyses to obtain covG matrices and variance estimates of the elements of these for genomic structural equation modelling in Genomic SEM[12].

However, available implementations of LDSR do not: 1.) explicitly account for varying numbers of genetic variants between association statistics from contributing GWASs, 2.) consider local association and LD patterns in GWASs, or 3.) adjust for imputation quality differences. In bivariate LDSR, genetic variant overlap between two traits and the LD score panel is usually drastically smaller compared to the regressions performed on single traits (supplementary tab. S6), due to differing patterns of missingness across datasets. This limits the power to carry out many downstream multivariate GWAS analyses, such as in Genomic SEM[12], and related methods such as mtCOJO[13,4], MTAG[14], and METAL[15]. These methods rely on the covG estimates from LDSR, and/or use the LDSR intercept to adjust associations (genomic control) for spurious associations due to confounding from population stratification or sample overlap. In comparison, the variance-component REML method in LDAK[7] is both assuming a variable contribution to heritability across the genome and includes an information score weighting[16] to differentially treat variants of varying genotyping or imputation quality. It may be of interest to bring this functionality

over to an LDSR setting to be applicable to summary level data rather than individual level data.

Standard LDSR further assumes there to exist a heteroskedastic relationship between genome-

wide covG and its standard error, mainly due to the expected larger variability of variants in high

LD regions. Despite a weighting correction to account for genome-wide covG heteroskedasticity,

it also served as justification[9] for employing resampling to estimate standard errors that are robust

to bias from heteroskedasticity. It may therefore be of interest to investigate whether this

assumption can be relaxed when estimating covG standard errors, to allow for sampling across

the genome.

Here we present an enhanced version of LDSR, LDSC++, extended to improve on 1.) the

handling of varying numbers of shared genetic variants across trait pairs and reference panels,

2.) the estimation of genetic covariance and its standard error, and 3.) the handling of imputation

quality of imputed genotypes or genetic associations. Finally, through our extensions we aim to

allow for tractable statistical tests of covG, its standard error, or differences between them, also

accounting for high correlations between compared variables. We validate the performance of our

extensions to LDSR, first using simulations, second on real-world data in the form of GWASs

performed on selected measurements in the Genetic Links to Anxiety and Depression study and

the United Kingdom National Institute of Health and Social Care Research BioResource, and third

on a selection of ten GWAS summary statistics representing a wide variety of traits. We

hypothesise the accuracy of estimating both covG and its standard error may be improved and

the average measured variability of covG estimates across a sample of multiple diverse traits will

be reduced when applying our extensions in LDSC++ individually or in combination, as compared

to standard LDSR.

2 | MATERIALS AND METHODS

To enable more precise covG estimation from GWAS summary statistics, we extend the LDSR

method based on the multivariate implementation included in Genomic SEM. We have named

this extension LDSC++ (supplementary info. 1.8). The extensions to LDSR are:

❖ **Extension 1 - Variable block-count:** Allowing the number of blocks to vary when

estimating covG and covG standard error, in contrast to a fixed number of blocks.

❖ **Extension 2 - Extended block definitions to allow for variable block-count:** Block

definitions to be used while allowing the number of blocks to vary with Extension 1. These

are based on either a set number of variants or a recombination distance in cM.

❖ **Extension 3 - Variable block-count sampling:** A novel approach to sample across

genome blocks to estimate genome-wide covG and covG variance, rather than performing

block jackknife resampling.

❖ **Extension 4 - Extended weighting scheme for imputation quality:** An addition to the

standard LDSR weighting scheme to also control for imputation quality.

❖ **Extension 5 - Adjusted weighting scheme to correct for the correlation between LD**

score and association statistic: We adjusted the standard weighting scheme in LDSR

to correct for correlation between the regression variables, to be less influenced by

extreme LD values due to noise.

LDSC++ alongside standard LDSR (emulated in our LDSC++ code) were validated in both

real-world data and simulated traits, using REML covG estimates for reference.

An individual level sample from cohorts part of the National Institute of Health Research

(NIHR) BioResource was used for assessing genetic associations with phenotypes and for

simulation. This selection included participants of European ancestry from the Genetic Links to

Anxiety and Depression study (GLAD)[17], the United Kingdom Eating Disorders Genetics Initiative

(EDGI)[18,19], the NIHR BioResource Inflammatory Bowel Disease BioResource panel and the

NIHR BioResource General Population panel, which together form a collection of cohorts

colloquially referred to as GLAD+ (N=36,746).

We selected the traits height, weight, neuroticism measured by the Eysenck Personality

Questionnaire-Revised Short form (EPQR-S)[20], and negative affect measured by the PID-5 Brief

Form[21,22], in the GLAD+ cohort. Additionally, we simulated two phenotypes (further referred to as

SIM1 and SIM2 respectively) created to have 20% heritability and a genetic correlation of 0.4 with

each other, in four pairs with fractions of 100%, 75%, 50%, and 25% shared quantitative trait loci

(QTL) respectively. A simulation was performed in GCTA[4] for each phenotype and fraction

combination, using 30,000 QTL and GLAD+ as an individual level sample.

We used GCTA-GREML[3] implemented in GCTA[4] to estimate variance component covG

for our individual level GLAD+ samples. Association tests were performed in GCTA fastGWA[23].

The covariates included in the association analysis were: genotyping array type/version, tissue

source, genotyping batch, and the first seven principal component scores of participant genotypes

as computed by the genetic principal component eigenvalues of the 1000 Genome Project[24]

Phase 3 (1kG3) reference panel projected on the genetic variants of GLAD+. Additionally, ten

external GWAS summary statistics covering a range of traits of various statistical power, h^2, and

genome coverage in terms of the number of variants included were selected for validating

methods (Tab. 1). All summary statistics were documented to be based on samples with

predominantly European ancestry.

To match GCTA-GREML, we used GCTA[4,25] to generate the GLAD+ LD scores, using a

window of 1 Mb and a prior filter on minor allele frequency (MAF) > 0.01. For our external GWAS

summary statistics, we instead created LD scores based on the European ancestry subset of the

1kG3[26], not including Finnish ancestry. For this second LD score library, we used a 1 cM window

and 250 blocks setting in LDSC[9,27].

GWAS summary statistics were first harmonised using our *Supermunge* routine part of

the *shru*[28] package for the R language [https://github.com/tnggroup/tng_fork_shru_jz2024],

based on the original LDSC munge routine[9], the Genomic SEM routines for munging and

preparation for latent factor GWAS[12], and the MungeSumstats routine[29] (supplementary info.

1.4.4).

To account for heteroskedastic relationships for comparisons of covG inferential

uncertainty (sampling variance and thus standard errors) across traits of varying heritabilities, we

applied the coefficient of variation (CV) of covG rather than comparing direct estimates

(supplementary info. 1.5.1).

We formulated simple parametric statistical tests specifically accounting for the novel

varying number of blocks in the LDSR sampling and resampling methods and non-independent

variables (supplementary info. 1.9), supported by paired samples Wilcoxon signed-rank tests[30]

(supplementary info. 1.9.12). Test results were corrected for multiple comparisons using the

Benjamini & Hochberg[31] method (q-values, also called false discovery rate; FDR), additionally

and unconventionally taking the effective number of comparisons from correlated variables[32–34]

into account (supplementary info. 1.9.13).

174 **Table 1. Selected external GWAS summary statistics**

175 A selection of ten diverse publicly available GWAS summary statistics. The table describes the dataset

176 reference, the number of cases or total number of participants, and for binary traits; the number of

177 controls, the sample prevalence estimate, and the population prevalence estimate used throughout the

178 study. 'INFO range' describes the range of the provided imputation quality information score.

Trait	Reference	Year	N case	N control	Sample prev.	Population prev.	INFO range
Anxiety disorder (ANXD)	Purves et al.[35]	2019	25,453	58,113	0.305	0.16[36]	0.8-1.0
Bipolar disorder (BIPD)	Stahl et al.[37]	2019	20,352	31,358	0.394	0.025[38]	0.1-1.6
BMI (BMI)	Pulit et al.[39]	2019	806,834				0.3-1.0
MDD (MDD)	Wray et al.[40]	2018	16,823	25,632	0.396	0.146[41]	0.6-1.9
Broad depression (MDDB)	Howard et al.[42]	2019	170,756	329,443	0.341	0.302[42]	-
MDD symptoms (MDDS)	Wray et al.[40]	2018	14,260	15,480	0.479	0.146[41]	0.9-1.0
Educational attainment (EDUA)	Lee et al.[43]	2018	766,345				-
Insomnia (INSO)	Jansen et al.[44]	2019	109,402	277,131	0.283	0.69[45]	0.9-1.0
Neuroticism (NEUR)	Nagel et al.[46]	2018	449,484				0.9-1.0
Schizophrenia (SCHI)	Pardiñas et al.[47]	2018	40,675	64,643	0.386	0.0072[48]	-

3 | RESULTS

We estimated REML h^2 and r_g of our four selected GLAD+ traits and their bivariate combinations. The REML h^2 of the simulated traits SIM1 and SIM2 were estimated in the range from 13% to 15%, and the REML r_g between traits in a pair to 0.43, 0.24, 0.19, and 0.2 for the QTL sharing fractions 100%, 75%, 50%, and 25% respectively. Deviations of LDSR method results from GCTA-REML estimates were obtained, run on both the selected GLAD+ traits (Tab. 2, supplementary tab. S7A, S7B) and the simulated traits (supplementary tab. S8A, S8B). Most deviations between covG and covG variance estimates by our LDSR method extensions compared to standard LDSR were seen to be statistically significant, supporting our extensions to yield empirically distinct results. As results were quite different comparing results from diagonal and off-diagonal elements of covG matrices, we decided to stratify them accordingly.

Only considering h^2 estimates (diagonal elements in the covG or covG SE matrix), the top performing LDSR method combination was variable block-count sampling using the recombination distance block-definition, which yielded average relative h^2 differences as small as -11% and average relative h^2 CV differences of -1% compared to GCTA-GREML in GLAD+. The same pattern was seen for the simulated traits only using the 100% QTL sharing fraction, while for lesser sharing fractions the performance among methods was mixed with the standard LDSR block jackknife or LDSC++ variable block-count sampling with recombination distance blocks yielding the on average most accurate and less biassed estimates. In the GLAD+ data, all LDSR methods under-estimated diagonal covG with about -30% for block jackknife resampling and -10% to -20% for variable block-count sampling. Block jackknife resampling was also consistently seen to over-estimate diagonal variability estimates from 5% to 9%, while variable block-count sampling seemed to under-estimate diagonal variability estimates with about -1% to -7%. The influence of our weighting extensions were seen to only have tiny effects on the differences in the

preferred direction for the GLAD+ traits, primarily detectable for h^2 estimates and when using

block jackknife resampling and imputation quality weighting.

Off-diagonal deviations were markedly larger by about a factor of two to four for average

relative covG estimates and a factor of ten to sixty for average relative covG CV estimates,

compared to the corresponding diagonal matrix element results for the GLAD+ traits. A similar

but attenuated pattern was seen for the simulated traits. As for the diagonal elements, the LDSR

methods under-estimated off-diagonal covG estimates with about -60% for block jackknife

resampling and -40% to -50% for variable block-count sampling in the GLAD+ traits. Off-diagonal

covG CVs were consistently and grossly over-estimated by all LDSR methods. For block jackknife

resampling, the recombination distance block definition brought the off-diagonal covG CV

difference to about 100% compared to 120% when using number-of-variants defined blocks.

Variable block count sampling produced off-diagonal covG CV estimates with deviations from

about 60% with the recombination distance block definition to 80% with the number-of-variants

block definition.

We applied the different LDSR method combinations in our selection of ten external trait

GWASs (Tab. 1) for which samples we did not have individual level data (supplementary tab.

S9A, S9B). Default block-size settings resulted in block-counts in the range of 249 - 405 for the

number-of-variants block definition and 338 - 364 for the recombination distance block definition,

without INFO filter. Estimates from method combinations using our extensions were first assessed

independently in reference to suitable baselines (supplementary tab. S9A). Compared to standard

LDSR, applying our block definitions and sampling extensions yielded 4% to 8% larger h^2

estimates, 3% smaller h^2 CV estimates using number-of-variants blocks, but 7% larger h^2 CV

estimates using recombination distance blocks. The weighting extensions made a relatively small

contribution to average differences in estimates compared to the block and sampling extensions,

but specific traits showed statistically significant results. Applying imputation score weighting

resulted in an approximate gain of 4% to 5% (q<0.05) in BIPD h^2 and a 1% to 7% (q<0.02)

reduction in BIPD bivariate covG CV with specific traits across block and sampling methods.

Similarly, applying alternative correlation correction weighting yielded an approximate gain of 3%

to 4% (q<0.05) in BMI h^2 and multiple significant bivariate BMI covG CV reductions in the range

of 1% to 10% (q<0.05). Applying all extensions in concert and comparing to standard LDSR

yielded 5% to 8% larger h^2 estimates, 4% smaller h^2 CV estimates using number-of-variants

blocks, but 6% larger h^2 CV estimates using recombination distance blocks.

Table 2. LDSR estimate deviation from a REML ideal measured over four GLAD+ traits

The table shows deviations of LD score regression (LDSR) method estimates of genetic covariance (covG) and covG SE (converted to coefficient of variation, CV) from the chosen ideal; estimates from GCTA-GREML using individual level data. We used the LDSC++ implementation of LDSR applied to our selection of four real-world traits from the GLAD+ sample: height, weight, neuroticism measured by the Eysenck Personality Questionnaire-Revised Short form (EPQR-S), and negative affect measured by the PID-5 Brief Form. Deviation was measured as average relative difference (Avg. Δ) and root-mean-square error (RMSE). LDSR method combinations were defined by their configuration in regard to block definition, (re)sampling method, alternative correlation correction weighting, and imputation quality weighting. Estimates are stratified in diagonal (heritability) and off-diagonal (bivariate genetic covariance) estimates from a genetic covariance matrix. Only significant covG values (q<0.05) were considered.

Block definition	(Re)sampling method	Alternative correlation correction weighting	Imputation quality weighting	covG				covG CV			
				Diagonal		Off-diagonal		Diagonal		Off-diagonal	
				Avg. Δ (%)	RMSE	Avg. Δ (%)	RMSE	Avg. Δ (%)	RMSE	Avg. Δ (%)	RMSE
Fixed 200 blocks	Block jackknife resampling	□	□	-32.7	0.0754	-61.0	0.0994	7.6	0.0756	118.0	0.1283
Number-of-variants blocks	Block jackknife resampling	□	□	-32.7	0.0754	-61.0	0.0994	7.9	0.0788	120.0	0.1300
		☑	□	-33.0	0.0759	-61.3	0.0998	8.7	0.0769	122.4	0.1327
		□	☑	-33.0	0.0756	-61.3	0.0998	6.4	0.0769	116.5	0.1231
		☑	☑	-33.2	0.0760	-61.5	0.1002	7.3	0.0750	118.8	0.1256
	Variable block-count sampling	□	□	-21.7	0.0542	-50.1	0.0836	-5.8	0.1182	79.4	0.0783
		☑	□	-22.0	0.0549	-50.4	0.0841	-5.6	0.1173	80.5	0.0795
		□	☑	-21.0	0.0523	-52.3	0.0861	-6.7	0.1185	81.4	0.0807
		☑	☑	-21.2	0.0529	-52.6	0.0867	-6.5	0.1175	82.8	0.0821
Recombination distance blocks	Block jackknife resampling	□	□	-31.7	0.0722	-59.8	0.0972	5.1	0.0727	101.8	0.1155
		☑	□	-31.9	0.0726	-60.0	0.0976	5.9	0.0707	103.6	0.1176
		□	☑	-31.9	0.0721	-60.0	0.0975	5.7	0.0703	103.1	0.1177
		☑	☑	-32.1	0.0726	-60.2	0.0979	6.5	0.0683	105.0	0.1198
	Variable block-count sampling	□	□	-11.1	0.0475	-38.5	0.0638	-1.6	0.1132	59.7	0.0925
		☑	□	-11.4	0.0480	-38.5	0.0643	-1.1	0.1127	60.9	0.0944
		□	☑	-11.2	0.0482	-46.4	0.0784	-1.4	0.1134	99.0	0.0962
		☑	☑	-11.5	0.0487	-39.3	0.0647	-0.8	0.1129	62.9	0.0910

4 | DISCUSSION

We have adapted the LDSR routine from Genomic SEM into a modified implementation called

LDSC++, which performs LDSR with extensions specifically aimed at improving covG estimation

in large-scale multivariate analyses with varying genomic overlaps among trait GWAS dataset

and reference panels. For this purpose, we created a variable block-count sampling routine in

LDSC++ aimed to be applied with a variable number of blocks to discriminate between situations

with a varying number of participating genetic variants, which further makes it possible to use this

information in statistical tests. Variable block-count sampling performs segmented (also called

piecewise) regression using one of our two block definitions, which putatively makes the

aggregate genome-wide parameter estimates (averages of segmented regression parameters)

robust to heteroskedasticity and outliers. The most obvious and important assumption for both

the variable block-count sampling method and the statistical tests is that the estimated genome-

wide covG can be assumed to have the same variance across the genome, or rather can be

estimated as an aggregate of piecewise estimates. While resampling techniques[49,50] traditionally

have been used to obtain robust covG standard errors in LDSR, a sufficiently better model from

the segmented regression is in our project theorised to make robust standard errors superfluous[51].

LDSC++ further takes advantage of imputation quality information to weight the LDSR, instead of

relying on filters which otherwise reduce the overall information content in the analysis. Variable

block-count sampling may be interpreted as applying a slightly different h^2 model, allowing for a

different contribution to h^2 across the genome similar to the LDAK h^2 model[7,16,52], in contrast to

the standard LDSR heritability model[9,16,52]. We additionally proposed statistical tests in which the

varying number of blocks from our extensions can be applied to translate statistical power from a

large number of genetic variants in a GWAS into statistical power for LDSR covG and covG

variance, and thus making these tests more sensitive and differentiating in collections of bivariate

trait pairs proportional to the variability in genetic variant overlap.

Our LDSR extensions were shown to yield more accurate results as validated by GCTA-

GREML than standard LDSR settings, both when using each extension independently and when

applied in concert, yielding average relative deviations from about -10% to -20% for h^2 and -1%

to -6% for h^2 CV compared to the corresponding deviations of -30% and 8% respectively for

standard LDSR. This performance was seen in estimates made in the individual level GLAD+

sample and replicated to a lesser extent in simulated data using GCTA-GREML. The performance

increase when using the combined extensions could be attributed to allowing the number of

blocks to vary and using our new block definitions or using our variable block-count sampling, and

only a small fraction could be attributed to the weighting extensions in this setting. Applying our

extensions to a set of ten trait GWASs external to our individual level sample, supported the above

validations through convergent results primarily for h^2 estimates rather than off-diagonal covG

estimates and when using number-of-variants block definitions. In this diverse selection of

GWASs of varying power and coverage our combined LDSC++ extensions yielded on average

up to 8% larger and 4% less variable h^2 estimates compared to standard LDSR. Larger h^2

estimates were convergent with the previous individual level data validation validations, but only

the number-of-variants block definition were seen to produce smaller average h^2 estimates

compared to standard LDSR, while using the recombination distance block definition instead

yielded larger average h^2 estimates - contrary to the previous validation results. This may indicate

a sensitivity of the recombination distance block definition to missingness patterns or other factors

that were not represented by our four GLAD+ trait GWASs. The external GWASs also allowed us

to explore other imputation score ranges and highly polygenic trait GWASs of high power, which

implicated putative improved estimates when applying the imputation score weighting to the BIPD

GWAS and when applying the alternate correlation correction weighting to the BMI GWAS. Our

proposed explanation for these results is that the BIPD GWAS has a wide distribution of

imputation quality INFO scores in contrast to other included datasets which the corresponding

weighting extension would be able to leverage, and the BMI GWAS is relatively well-powered and

represents a highly polygenic trait for which the alternative correlation correction weighting may

be able to correct bias due to tiny LD scores to a higher degree.

The major strength in our simulations came from using a relatively large sample size of

36,746 individuals from the GLAD+ cohort, in contrast to the Bulik-Sullivan study which simulated

associations based on a sample of 2,062 individuals. We additionally simulated whole genome

associations, where the Bulik-Sullivan study only simulated chromosome 2. Unfortunately, we did

not manage to perform sufficiently many repetitions in our simulations for them to appear

trustworthy, with an effective number of GCTA simulations of 2.5 for any randomly sampled QTL

compared to 100 repetitions done in a previous study by Bulik-Sullivan[53]. Our effort to investigate

the influence of effect QTL overlap on REML and LDSR estimates was interesting, but proved

practically challenging enough to be one of the reasons for why we did not manage to repeat the

simulations to a higher degree.

The discrepancy between validation results from diagonal and off-diagonal covG elements

was identified as a cause for concern about the LDSR method in general. Because of this we

were inclined to trust off-diagonal validation results less. Off-diagonal covG estimates from

bivariate LDSR are however the foundation for genetic correlation analyses, and also provide the

majority and arguably the more important information for analyses performed on covG such as

Genomic SEM, underscoring the importance of accurate off-diagonal covG estimation and the

need to critically investigate bivariate LDSR further.

Our results provide support for the accuracy of statistical models where genome-wide h^2

and covG parameters are constructed as aggregates of heterogeneous local parameters. The

recombination distance block definition was part of the method combination displaying the

strongest performance in our individual level real-world and simulated GCTA-GREML validations.

In the external GWASs, the recombination distance block definition also recovered some of the

largest covG estimates on average compared to the number-of-variants block definition and

standard LDSR. We would suggest that these properties are promising and may prompt further

research into block definitions for LDSR that are consistent across traits and bivariate trait pairs

in the way our recombination distance definition was made or how local genetic correlation

methods such as LAVA[54] define consistent genomic blocks. The arbitrary definition of blocks is a

general limitation of LDSR. Creating more optimised block definitions is indeed an active research

area[55,56]. Reconciling the recombination based block definition approach with the utility of the

number-of-variants block definition may be important to advance LDSR and similar methods

further.

5 | CONFLICT OF INTEREST

6 | ACKNOWLEDGEMENTS

This study/research is funded by the National Institute for Health and Care Research (NIHR)

Maudsley Biomedical Research Centre (BRC). The views expressed are those of the author(s)

and not necessarily those of the NIHR or the Department of Health and Social Care.

Christopher Hübel acknowledges funding by Lundbeckfonden (R276-2018-4581).

Alexandra Gillett is supported by the Medical Research Council (MR/X009815/1).

The authors acknowledge use of the research computing facility at King's College London, King's

Computational   Research,   Engineering   and   Technology   Environment   (CREATE)

(https://doi.org/10.18742/rnvf-m076)[57].

7 | DATA AVAILABILITY

All data produced in the present study are available upon reasonable request to the authors. While some data is contained in the manuscript, an attempt has been made to additionally present results in table format in the accompanying supplementary tables. GWAS summary statistics of our traits used for validation are available upon reasonable request. LD scores based on the GLAD+ sample are available upon reasonable request. The analysis code will be made available at https://github.com/tnggroup/jz-phd-c2c3-2024. The software package 'shru' will be made available on https://github.com/johanzvrskovec/shru.

8 | ETHICS APPROVAL STATEMENT

The London—Fulham Research Ethics Committee approved the GLAD Study on 21st August 2018 (REC reference: 18/LO/1218). The London—Fulham Research Ethics Committee approved the EDGI UK Study on 29th July 2019 (REC reference: 19/LO/1254). The East of England—Cambridge Central Committee approved the NIHR BioResource as a Research Tissue Bank (REC reference: 17/EE/0025). The South West—Central Bristol Research Ethics Committee approved the COVID-19 Psychiatry and Neurological Genetics study on 27th April 2020 (REC reference: 20/SW/0078).

9 | REFERENCES

1. Abdellaoui, A. & Verweij, K. J. H. Dissecting polygenic signals from genome-wide association studies on human behaviour. *Nat. Hum. Behav.* 1–9 (2021) doi:10.1038/s41562-021-01110-y.

2. Pasaniuc, B. & Price, A. L. Dissecting the genetics of complex traits using summary association statistics. *Nat. Rev. Genet.* **18**, 117–127 (2017).

3. Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nat. Genet.* **42**, 565–569 (2010).

4. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: A tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).

5. Loh, P.-R. *et al.* Contrasting genetic architectures of schizophrenia and other complex diseases using fast variance-components analysis. *Nat. Genet.* **47**, 1385–1392 (2015).

6. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat. Genet.* **47**, 284–290 (2015).

7. Speed, D., Hemani, G., Johnson, M. R. & Balding, D. J. Improved Heritability Estimation from Genome-wide SNPs. *Am. J. Hum. Genet.* **91**, 1011–1021 (2012).

8. Barry, C.-J. S. *et al.* How to estimate heritability: a guide for genetic epidemiologists. *Int. J. Epidemiol.* **52**, 624–632 (2023).

9. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

10. Speed, D. & Balding, D. J. SumHer better estimates the SNP heritability of complex traits from summary statistics. *Nat. Genet.* **51**, 277–284 (2019).

11. van Rheenen, W., Peyrot, W. J., Schork, A. J., Lee, S. H. & Wray, N. R. Genetic correlations of polygenic disease traits: from theory to practice. *Nat. Rev. Genet.* **20**, 567–581 (2019).

12. Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into the

multivariate genetic architecture of complex traits. *Nat. Hum. Behav.* **3**, 513–525 (2019).

13. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from

GWAS summary data. *Nat. Commun.* **9**, 224 (2018).

14. Turley, P. *et al.* Multi-trait analysis of genome-wide association summary statistics using

MTAG. *Nat. Genet.* **50**, 229–237 (2018).

15. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of

genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).

16. Speed, D., Cai, N., Johnson, M. R., Nejentsev, S. & Balding, D. J. Reevaluation of SNP

heritability in complex human traits. *Nat. Genet.* **49**, 986–992 (2017).

17. Davies, M. R. *et al.* The Genetic Links to Anxiety and Depression (GLAD) Study: Online

recruitment into the largest recontactable study of depression and anxiety. *Behav. Res.*

Ther. (2019) doi:10.1016/j.brat.2019.103503.

18. Monssen, D. *et al.* The United Kingdom Eating Disorders Genetics Initiative. *Int. J. Eat.*

Disord. **57**, 1145–1159 (2024).

19. Bulik, C. M. *et al.* The Eating Disorders Genetics Initiative (EDGI): study protocol. *BMC*

Psychiatry **21**, 234 (2021).

20. Eysenck, H. J. & Eysenck, M. W. Personality and Individual Differences. *Plenum Press N.*

Y. (1985) doi:10.1007/978-1-4613-2413-3.

21. Krueger, R. F., Derringer, J., Markon, K. E., Watson, D. & Skodol, A. E. Initial construction

of a maladaptive personality trait model and inventory for DSM-5. *Psychol. Med.* **42**, 1879–

1890 (2012).

22. Maples, J. L. *et al.* Supplemental Material for Testing Whether the DSM-5 Personality

Disorder Trait Model Can Be Measured With a Reduced Set of Items: An Item Response

Theory Investigation of the Personality Inventory for DSM-5. *Psychol. Assess.* **27**, 1195–

1210 (2015).

23. Jiang, L. *et al.* A resource-efficient tool for mixed model association analysis of large-scale

data. *Nat. Genet.* (2019) doi:10.1038/s41588-019-0530-8.

24. Altshuler, D. M. *et al.* An integrated map of genetic variation from 1,092 human genomes.

Nat. 2012 4917422 **491**, 56–65 (2012).

25. Yang, J. *et al.* Genetic variance estimation with imputed variants finds negligible missing

heritability for human height and body mass index. *Nat. Genet.* **47**, 1114–1120 (2015).

26. Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).

27. Bulik-Sullivan, B. K. LDSC (LD SCore). (2015).

28. Zvrskovec, J. shru. https://github.com/johanzvrskovec/shru (2023).

29. Murphy, A. E. & Skene, N. G. MungeSumstats: A Bioconductor package for the

standardisation and quality control of many GWAS summary statistics. *bioRxiv*

2021.06.21.449239 (2021) doi:10.1101/2021.06.21.449239.

30. Wilcoxon, F. Individual Comparisons by Ranking Methods. *Biom. Bull.* **1**, 80–83 (1945).

31. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300

(1995).

32. Gao, X., Starmer, J. & Martin, E. R. A multiple testing correction method for genetic

association studies using correlated single nucleotide polymorphisms. *Genet. Epidemiol.* **32**,

361–369 (2008).

33. Nyholt, D. R. A Simple Correction for Multiple Testing for Single-Nucleotide Polymorphisms

in Linkage Disequilibrium with Each Other. *Am. J. Hum. Genet.* **74**, 765–769 (2004).

34. Cheverud, J. M. A simple correction for multiple comparisons in interval mapping genome

scans. *Heredity* **87**, 52–58 (2001).

35. Purves, K. L. *et al.* A major role for common genetic variation in anxiety disorders. *Mol.*

Psychiatry (2019) doi:10.1038/s41380-019-0559-1.

36. Kessler, R. C. *et al.* The global burden of mental disorders: An update from the WHO World

Mental Health (WMH) Surveys. *Epidemiol. Psichiatr. Soc.* **18**, 23–33 (2009).

37. Stahl, E. A. *et al.* Genome-wide association study identifies 30 loci associated with bipolar

disorder. *Nat. Genet.* **51**, 793–803 (2019).

38. Kessler, R. C., Petukhova, M., Sampson, N. A., Zaslavsky, A. M. & Wittchen, H.-U. Twelve-

month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the

United States. *Int. J. Methods Psychiatr. Res.* **21**, 169–184 (2012).

39. Pulit, S. L. *et al.* Meta-analysis of genome-wide association studies for body fat distribution

in 694 649 individuals of European ancestry. *Hum. Mol. Genet.* **28**, 166–174 (2019).

40. Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine the

genetic architecture of major depression. *Nat. Genet.* **50**, 668–681 (2018).

41. Kessler, R. C. & Bromet, E. J. The Epidemiology of Depression Across Cultures. *Annu. Rev.*

Public Health **34**, 119–138 (2013).

42. Howard, D. M. *et al.* Genome-wide meta-analysis of depression identifies 102 independent

variants and highlights the importance of the prefrontal brain regions. *Nat. Neurosci.* **22**,

343–352 (2019).

43. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association

study of educational attainment in 1.1 million individuals. *Nat. Genet. 2018 508* **50**, 1112–

1121 (2018).

44. Jansen, P. R. *et al.* Genome-wide analysis of insomnia in 1,331,010 individuals identifies

new risk loci and functional pathways. *Nat. Genet.* **51**, 394–403 (2019).

45. Morphy, H., Dunn, K. M., Lewis, M., Boardman, H. F. & Croft, P. R. Epidemiology of

Insomnia: a Longitudinal Study in a UK Population. *Sleep* **30**, 274–280 (2007).

46. Nagel, M. *et al.* Meta-analysis of genome-wide association studies for neuroticism in

449,484 individuals identifies novel genetic loci and pathways. *Nat. Genet. 2018 507* **50**,

920–927 (2018).

47. Pardiñas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-intolerant

genes and in regions under strong background selection. *Nat. Genet.* **50**, 381–389 (2018).

48. McGrath, J., Saha, S., Chant, D. & Welham, J. Schizophrenia: A Concise Overview of

Incidence, Prevalence, and Mortality. *Epidemiol. Rev.* **30**, 67–76 (2008).

49. Friedl, H. & Stampfer, E. Jackknife Resampling. *Encycl. Environmetrics* **2**, (2001).

50. Efron, B. Nonparametric estimates of standard error: The jackknife, the bootstrap and other

methods. *Biometrika* **68**, 589–599 (1981).

51. King, G. & Roberts, M. E. How Robust Standard Errors Expose Methodological Problems

They Do Not Fix, and What to Do About It. *Polit. Anal.* **23**, 159–179 (2015).

52. Speed, D., Holmes, J. & Balding, D. J. Evaluating and improving heritability models using

summary statistics. *Nat. Genet.* **52**, 458–462 (2020).

53. Bulik-Sullivan, B. Relationship between LD Score and Haseman-Elston Regression. 018283

Preprint at https://doi.org/10.1101/018283 (2015).

54. Werme, J., Sluis, S. van der, Posthuma, D. & Leeuw, C. A. de. LAVA: An integrated

framework for local genetic correlation analysis. *bioRxiv* (2021)

doi:10.1101/2020.12.31.424652.

55. Privé, F. Optimal linkage disequilibrium splitting. *Bioinformatics* **38**, 255–256 (2022).

56. Muggeo, V. M. R. Estimating regression models with unknown break-points. *Stat. Med.* **22**,

3055–3071 (2003).

57. King's College London. King's Computational Research, Engineering and Technology

Environment (CREATE). (2022) doi:https://doi.org/10.18742/rnvf-m076.

493