1 **ORIGINAL RESEARCH**
2
3 **Effects of Sacubitril/valsartan on Hypertensive Heart Disease: the REVERSE-LVH**
4 **Randomized Phase 2 Trial**
5
6 Brief Title: Myocardial Fibrosis and Hypertrophy Regression
7

8 Vivian Lee[a], Mayank Dalakoti[b], Qishi Zheng[c], Desiree-Faye Toh[a], Redha Boubertakh[a],
9 Jennifer A Bryant[d], Tar-Choon Aw[e], Chi-Hang Lee[b], A Mark Richards[f], Javed Butler[g,h],
10 Javier Díez[i,j], Roger Foo[b,k], Stuart A Cook[a,d,l], Carolyn SP Lam[a,d,l], Thu-Thao Le[a,m], Calvin
11 WL Chin*[a,d,m]
12
13 These authors contributed equally: Vivian Lee and Mayank Dalakoti
14 These authors jointly supervised the work: Thu-Thao Le and Calvin WL Chin
15 *Corresponding author
16
Affiliations
[a]National Heart Research Institute of Singapore, National Heart Centre Singapore, Singapore,
Singapore
[b]Department of Cardiology, National University Heart Centre Singapore, National University
Health System, Singapore, Singapore
[c]Cochrane Singapore, Singapore, Singapore
[d]Department of Cardiology, National Heart Centre Singapore, Singapore, Singapore
[e]Department of Laboratory Medicine, Changi General Hospital, Singapore, Singapore
[f]Christchurch Heart Institute, University of Otago, Christchurch, New Zealand
[g]Baylor Scott and White Research Institute, Dallas, Texas, United States
[h]Department of Medicine, University of Mississippi School of Medicine, Mississippi, United
States
[i]Centre of Applied Medical Research (CIMA) and School of Medicine, University of
Navarra, Pamplona, Spain
[j]Centre for Network Biomedical Research of Cardiovascular Diseases (CIBERCV), Carlos III
Institute of Health, Madrid, Spain
[k]Cardiovascular Metabolic Disease Translational Research Program, Yong Loo Lin School of
Medicine, National University of Singapore, Singapore, Singapore
[l]Cardiovascular & Metabolic Disorders SRP, Duke-NUS Medical School, Singapore,
Singapore
[m]Cardiovascular Medicine Academic Clinical Program (ACP), Duke-NUS Medical School,
Singapore, Singapore
39
Address for Correspondence
Dr Calvin Chin
Department of Cardiology
5 Hospital Drive, Singapore 169609
Email: cchin03m@gmail.com
Phone: (65) 6704 8905
Fax: (65) 6844 9030
47

ABSTRACT

Diffuse interstitial fibrosis is associated with adverse outcomes in hypertensive heart disease

and may be reversible. Sacubitril/valsartan could offer greater anti-fibrotic effects than

valsartan alone. In the REVERSE-LVH phase 2 open-labelled trial (clinicaltrials.gov NCT:

03553810; funded by the National Medical Research Council of Singapore), 78 patients with

essential hypertension and left ventricular hypertrophy (LVH) were randomized 1:1 to

sacubitril/valsartan or valsartan for 52 weeks. Primary endpoint was a change in interstitial

volume, assessed using cardiovascular magnetic resonance. Despite similar 24-hour systolic

blood pressure at 52 weeks (125±11 vs. 126±11 mmHg; P=0.379), sacubitril/valsartan

resulted in a greater absolute reduction in interstitial volume compared to valsartan (−5.2±5.4

vs. −2.5±3.1 mL; P=0.006). Secondary endpoints showed significant differences favoring

sacubitril/valsartan in LV mass, left atrial volume, estimated LV filling pressure, and

improved cardiac circulating biomarkers (N-terminal pro-B-type natriuretic peptide and high-

sensitivity troponin T). Other markers of cardiac volumes, function and mechanics were

similar between the two treatment arms. Here we show the potential myocardial benefits of

sacubitril/valsartan beyond blood pressure control, though larger studies are needed to

confirm their clinical relevance.

KEYWORDS

Diffuse interstitial fibrosis; cardiovascular magnetic resonance; hypertensive heart disease

70 INTRODUCTION

About 30% of the burden related to hypertension occurred in individuals with well-controlled

blood pressures (BP)[1]. The updated definition of heart failure (HF) stages had identified more

asymptomatic individuals in a pre-clinical stage[2]. This suggests that despite substantial

benefits associated with BP lowering, there remains opportunities to improve risk

stratification and develop therapeutic options beyond managing peripheral BP to

slow/prevent the development of symptomatic HF.

In hypertensive heat disease (HHD), left ventricular hypertrophy (LVH) occurs in response to

hemodynamic and neuro-hormonal stresses triggered by prolonged exposure to elevated BP[3].

Its pathophysiology is accompanied by an expansion in extracellular volume (ECV) due to

accumulation of extracellular matrix components and collagen deposition throughout the

myocardium – diffuse interstitial fibrosis. This type of fibrosis typically develops gradually

and is initially reversible[4]. Myocardial fibrosis significantly contributes to cardiac

dysfunction, arrhythmias, and reduced coronary perfusion, ultimately leading to HF and

adverse cardiovascular outcomes[4,5]. Therefore in addition to peripheral BP, diffuse interstitial

fibrosis is a potential marker for enhancing risk stratification and directing targeted therapies

to prevent the progression of hypertensive LVH to advanced HF stages.

Angiotensin receptor–neprilysin inhibitor (ARNI) represents a novel drug class that inhibits

the renin-angiotensin-aldosterone system (RAAS) and enhances natriuretic peptides. This

dual agent combines a neprilysin inhibitor (sacubitril) and an angiotensin receptor blocker

(ARB; valsartan). Sacubitril/valsartan has shown superior benefits over conventional ARB or

angiotensin-converting enzyme inhibitors (ACEI) monotherapy in lowering BP and

improving outcomes in those with HF and reduced LV ejection fraction (HFrEF)[6,7]. Pre-

clinical studies of mouse models have demonstrated that sacubitril/valsartan is effective in

reducing myocardial fibrosis and improving cardiac function by attenuating angiotensin II

and transforming growth factor b1 fibrotic pathways[8,9].

Using cardiovascular magnetic resonance (CMR) to quantify interstitial volume as a measure

of diffuse interstitial fibrosis non-invasively[10], the aim of the study was to compare

sacubitril/valsartan versus valsartan in reducing diffuse interstitial fibrosis in patients with

hypertensive LVH. The hypothesis was that 52 weeks of sacubitril/valsartan therapy would

result in greater regression of diffuse interstitial fibrosis compared to valsartan, independent

of BP control.

RESULTS

Study Participants

A total of 80 patients were recruited, and 78 individuals were successfully randomized to

sacubitril/valsartan (n=39) and valsartan (n=39) (Figure 1). The mean age at baseline was

58±11 years, and 32 (41%) were males. The baseline 24h mean SBP and DBP were 137±14

and 81±11 mmHg, respectively. The baseline clinical, renal function and CMR

characteristics were well-balanced between the two treatment groups (Table 1). There was no

change in body weight at the start and end of study.

The median dose prescribed in the sacubitril/valsartan group was 200 [100, 200] mg twice a

day and 160 [160, 320] mg once a day in the valsartan group, respectively. A total of 50

participants required additional anti-hypertensive medications (sacubitril/valsartan group,

n=26 and valsartan group, n=24; P=0.571; Supplemental Table 2). The overall treatment

adherence rate achieved in the trial was 97% (sacubitril/valsartan group, 97%; valsartan

group, 96%).

Primary Trial Endpoint

The baseline interstitial volume in the sacubitril/valsartan group was 29.2±12.8 mL compared

to 28.0±11.3 mL in the valsartan group. After 52 weeks, sacubitril/valsartan resulted in

greater regression in diffuse interstitial fibrosis compared to valsartan alone (-5.2±5.4 versus

-2.5±3.1 mL, P=0.006), with an absolute between-group difference of -2.8 (95% CI: -4.8 to -

0.8) mL. Sacubitril/valsartan resulted in an 18% reduction in interstitial volume from

baseline, compared to an 8.9% reduction with valsartan. The reduction in diffuse interstitial

fibrosis remained greater in the sacubitril/valsartan group compared to valsartan alone after

accounting for baseline interstitial volume, prior ACEI or ARB use, and baseline 24h SBP

(Figure 2).

Exploratory subgroup analysis suggested treatment heterogeneity in favor of

sacubitril/valsartan in individuals with 24h median SBP >135 mmHg, with an absolute

between-group difference of -5.2 (95% CI: -8.0 to -2.5) mL compared to -0.5 (95% CI: -3.2

to 2.2) mL in those with 24h SBP ≤135 mmHg (P=0.016 for interaction). There was no

significant treatment heterogeneity in the other subgroups stratified by age, sex, LV ejection

fraction, baseline interstitial volume, prior ACEI or ARB use and body mass index

(Supplemental Table 3).

Secondary Trial Endpoints

Compared to valsartan alone, sacubitril/valsartan resulted in greater reduction in LV mass (-

21.9±25.4 versus -11.0±9.0 g, P=0.014) and myocyte volume (-16.0±19.5 versus -8.0±7.1

mL, P=0.019). Sacubitril/valsartan also led to a greater decrease in LA volume (-18.3±16.6

versus -8.6±13.6 mL, P=0.006) and LV filling pressures (-1.8±1.5 versus -0.9±1.1 mmHg,

P=0.003) compared to valsartan, suggestive of improved diastolic function.

There were no significant differences in changes in cardiac volumes, systolic function, and

other measures of myocardial mechanics between the treatment groups. Although

sacubitril/valsartan was associated with a statistically significant improvement in global

longitudinal strain (GLS) compared to valsartan (-0.9±2.3 versus 0.2±1.9, P=0.027), the

effect size was small and the isolated improvement in GLS was not associated with

improvement in the other strain components (Table 2). Similar findings were observed when

relevant CMR metrics were indexed to body surface area (Supplemental Table 4). Compared

to valsartan, sacubitril/valsartan was associated with a significant reduction from baseline to

52 weeks in NT-proBNP (ETR=0.72; 95% CI: 0.54–0.91; P=0.006) and hsTnT (ETR=0.83;

95% CI: 0.69–0.99; P=0.005).

Treatment Effects on Blood Pressure Control

Participants in both groups achieved similarly lowered 24h mean BP at all timepoints

assessed in the trial (Figure 3). At the end of 52 weeks, the 24h mean SBP was 125±11

mmHg with sacubitril/valsartan versus 126±11 mmHg with valsartan (P=0.762 for

comparison). This corresponded to a decrease of 13.3±13.1 and 10.6±13.4 mmHg from

baseline in the sacubitril/valsartan and valsartan groups, respectively. Treatment difference

between the two groups was not significant (-2.7 [95% CI: -8.7 to 3.3] mmHg, P=0.379).

Similarly, the 24h mean DBP at the end of study was 75±9 mmHg with sacubitril/valsartan

versus 75±12 mmHg with valsartan (P=0.943 for comparison), corresponding to a decrease

of 7.1±7.1 and 5.6±9.6 mmHg from baseline in the two respective treatment groups. The

treatment difference in 24h DBP reduction between the two groups was also not significant (-

1.5 [95% CI: -5.4 to 2.3] mmHg, P=0.426).

Adverse Events

Treatment with sacubitril/valsartan and valsartan were very well-tolerated in this trial. One

patient developed nocturnal cough with sacubitril/valsartan three months after randomization.

Symptoms improved upon dose reduction, and the participant continued with the trial. Renal

function remained stable throughout the trial (Table 3). There were no major adverse

cardiovascular events such as hospitalizations for HF, myocardial infarction, strokes and

deaths reported during the trial.

DISCUSSION

The main finding of REVERSE-LVH was that sacubitril/valsartan resulted in a reduction in

diffuse interstitial fibrosis in hypertensive patients with LVH following 52 weeks of

treatment compared to valsartan alone. This was accompanied by a favorable change in LV

mass, circulating cardiac biomarkers and diastolic function (LA size and estimated LV filling

pressure). There was no significant difference in systolic function improvement between the

two treatment groups.

The use of CMR in REVERSE-LVH has allowed reliable quantification of fibrosis without

invasive endomyocardial biopsies, a procedure that was required in previous studies on anti-

fibrotic therapies. Novel CMR mapping techniques have dramatically advanced tissue

characterization non-invasively, allowing us to investigate novel anti-fibrotic therapies in

diverse populations.

ECV fraction represents the ratio of interstitium to the total myocardial volume. Although it

is widely used to measure diffuse interstitial fibrosis, its utility may be limited where

treatments affect both myocyte and interstitial compartments. In such cases, therapies that

reduce both compartments may lead to an unchanged or even increased ECV fraction. For

instance, in REVERSE-LVH, despite reductions in interstitial and myocyte volumes, both

treatment groups showed a slight increase in ECV fraction, likely due to a greater reduction

in myocyte volume relative to interstitial volume. Instead, deeper mechanistic insights can be

accomplished by separately assessing the interstitial and myocyte compartments, particularly

for novel treatments or interventions with potential effects on both compartments.

Myocardial fibrosis is a hallmark of HF, and not all BP medications have anti-fibrotic

properties. Among conventional anti-hypertensive therapies, ACEIs or ARBs that target the

RAAS have shown the most promise in regressing biopsy-confirmed myocardial fibrosis in

HHD, independent of BP lowering effects. Similarly, mineralocorticoid receptor antagonists

(MRAs) like spironolactone and eplerenone reduce serum markers of collagen turnover and

improve diastolic function on echocardiography[16–18].

Sacubitril/valsartan, a first-in-class ARNI used primarily for HFrEF, is also an effective anti-

hypertensive medication. While sacubitril/valsartan is approved for hypertension in some

countries, its effectiveness in regressing myocardial fibrosis in patients with hypertension has

not been established. Through the REVERSE-LVH trial using CMR to assess interstitial

volume, sacubitril/valsartan demonstrated a two-fold greater reduction in diffuse interstitial

fibrosis compared to valsartan alone. Both treatment groups achieved similar BP throughout

the trial, an important feature to isolate the anti-fibrotic effects of sacubitril/valsartan from

any potential influence of BP control. It is also noteworthy that the other classes of anti-

hypertensive medications prescribed were similar between the two treatment groups.

Sacubitril/valsartan also led to greater LV mass regression compared to valsartan alone,

findings that align with results from a previous trial comparing sacubitril/valsartan and

olmesartan. Overall, these results reinforced the beneficial impact of sacubitril/valsartan on

reverse remodeling in patients with HHD.

PARAGON-HF (Prospective Comparison of ARNI with ARB Global Outcomes in HF with

Preserved Ejection Fraction) reported that sacubitril/valsartan did not result in significantly

lower rates of HF hospitalizations and cardiovascular deaths in patients with HF and

preserved ejection fraction (HFpEF) compared to valsartan[22]. Considering these results, it is

important to discuss the relevance of REVERSE-LVH after PARAGON-HF and possible

implications for future HFpEF trials.

HFpEF represents a syndrome characterized by phenotypic heterogeneity, underpinned by

complex underlying mechanisms, and contributed by various coexisting comorbidities. This

heterogeneity is exemplified by the observation that despite similar E/e' (echocardiographic

marker of LV filling pressure), the prevalence of LVH ranges between 14 and 50% in recent

HFpEF trials (21% in PARAGON-HF). This observation hints at the presence of additional

factors contributing to elevated LV filling pressure (such as chronic lung disease, atrial

fibrillation and/or chronic kidney disease). Consequently, it is plausible that not all these

patients had derived the anticipated benefits from the HF medications being investigated.

To enhance the effectiveness of future HFpEF trials, one key consideration lies in the

strategic inclusion of patients with common phenotypes driving the underlying

pathophysiology. This approach will enable more precise selection of pharmaceutical

therapies tailored at specific mechanistic phenotypes. For instance, pirfenidone, a small-

molecule anti-fibrotic agent without hemodynamic effects, reduced myocardial fibrosis in

patients with HFpEF[26]. This reduction in interstitial space without significance changes in

myocyte compartment explains for the overall decrease in ECV fraction. In contrast,

sacubitril/valsartan reduces both myocardial fibrosis and hypertrophy because of its effects

on hemodynamics, natriuretic and RAAS system.

The effectiveness of hypertension treatment is often assessed by the BP targets achieved,

with the fundamental goal to minimize end-organ complications such as HF. However,

cardiac remodeling in the hypertensive heart is complex and the peripheral BP may not

adequately reflect the abnormalities in the myocardium. A moderate correlation was

demonstrated between interstitial volume and 24h SBP ($r=0.36$, $P<0.001$), worse with office

SBP ($r=0.21$; $P<0.01$)[27]. This poses a challenge in defining the optimal BP targets in

hypertensive patients with LVH, particularly if BP are adequately controlled.

Because the apparent amelioration of diffuse interstitial fibrosis is associated with improved

outcomes, the interstitium is an attractive target for drug repurposing and development.

Sacubitril/valsartan resulted in a greater reduction in diffuse interstitial fibrosis compared to

valsartan alone after 52 weeks of treatment. The observed association with an improvement

in diastolic function and circulating biomarkers was encouraging.

The findings from REVERSE-LVH strengthened the case for considering sacubitril/valsartan

in selected individuals with hypertensive LVH, especially those who failed to achieve

adequate reduction in diffuse interstitial fibrosis with ACEIs or ARBs. Individuals with 24h

SBP greater than 135mmHg (median SBP in this study) may have further reduction in diffuse

interstitial fibrosis with sacubitril/valsartan but this finding from the post-hoc subgroup

analysis was exploratory and should be investigated in future trials.

It was by chance that a higher proportion of female participants (59%) were recruited.

Despite the challenges posed by COVID-19 between 2020 and 2021, the trial had very low

dropout rates (5%) and exceptionally high treatment adherence rates, surpassing the 80%

threshold widely used to define "good adherence". Both treatment arms achieved similar BP

reduction at the end of the trial. These observations highlighted the meticulous conduct of the

trial.

The study had some limitations. Two participants (one in each treatment group) were given

MRAs, which may confound regression of diffuse interstitial fibrosis. This was inevitable

because their BPs were not optimally controlled despite being prescribed the maximum

tolerated doses of trial medications and other classes of anti-hypertensives. Although

REVERSE-LVH was adequately powered as a phase 2 trial, the effect size observed is small

and warrants further investigation in larger studies powered for clinical outcomes. Prior

exposure to ACEIs and ARBs in the participants may have introduced confounding, although

no significant treatment interactions were observed.

In conclusion, compared to valsartan alone, sacubitril/valsartan appears to exert beneficial

myocardial effects beyond blood pressure reduction in patients with hypertensive LVH.

However, larger and longer-term studies are necessary to determine the clinical significance

of the observed effect sizes.

METHODS

Trial Design

The Role of ARNI in the Ventricular Remodeling in Hypertensive LVH (REVERSE-LVH)

was a prospective, randomized, open-label, blinded endpoint (PROBE) phase 2 trial

(ClinicalTrials.gov identifier: NCT03553810). The original design and methods of the trial

has been published[31] (Supplemental note 1). All study procedures were conducted at the

National Heart Centre Singapore (NHCS). This was an investigator-initiated trial. All trial-

related expenses, including investigations and medications were funded by the National

Medical Research Council. Novartis did not participate in the design of the trial, analysis and

interpretation of the data.

REVERSE-LVH was reviewed and approved by the SingHealth Centralized Institutional

Review Board (CIRB ref: 2018/2182) and the Health Sciences Authority. All participants

provided written informed consent.

Trial Participants and Recruitment

Participants were recruited between June 2019 and June 2023. These participants were

screened and recruited from the REMODEL cohort, an ongoing non-interventional

prospective study to examine the role of CMR in patients with hypertension

(ClinicalTrials.gov Identifier: NCT02670031). Eligible participants were adults at least 21

years old with essential hypertension and LVH, diagnosed using Asian specific thresholds on

CMR[32].

Essential hypertension was defined as systolic BP (SBP) ≥140 mmHg and/or diastolic BP

(DBP) ≥90 mmHg at time of diagnosis, on at least one medication for BP control. Individuals

with known secondary causes of hypertension, inherited cardiomyopathies, atrial fibrillation,

and history of cardiovascular complications (such as myocardial infarction, HF and stroke)

were excluded. Other exclusion criteria included intolerance to ARBs, contraindications for

CMR, stage IV or V chronic renal disease, presence of disease with limited life expectancy

(<3 years), pregnancy and/or breast-feeding.

Randomization, Treatment and Blinding

Study participants were randomized to either sacubitril/valsartan or valsartan in a 1:1 fashion.

Randomization was performed using sealed and sequentially numbered envelopes, prepared

by the Singapore Clinical Research Institute. Crossover of treatment allocation was not

allowed, and the treatment duration was 52 weeks. Treatment allocation was not concealed

from patients. Treatment assignment and trial endpoints were concealed from the core-lab

staff analyzing the CMR images.

Sacubitril/valsartan was initiated at 50-100mg twice a day, uptitrated to maximum 200mg

twice a day; valsartan was initiated at 80-160mg once a day, uptitrated to maximum 320 mg

once a day. Medications in both treatment groups were uptitrated as tolerated to achieve a

target SBP of <140 mmHg according to contemporary guidelines at the time of study

conception[33,34]. Additional non-RAAS inhibiting anti-hypertensive agents were prescribed

where necessary. Participants who were on ACEI or ARB at the time of enrollment

underwent a washout period of two weeks before commencing trial medication to eliminate

residual RAAS-intervening effects, during which alternative medicines or uptitration of

existing concomitant anti-hypertensive agents were prescribed for BP control.

Study medications were dispensed by NHCS pharmacists after each study visit. Treatment

adherence was assessed by pill counting by the pharmacists.

Study Procedures

Baseline demographic and clinical information were obtained from participants upon

randomization. Participants were seen at the clinic with 24h ambulatory BP at weeks 4, 8, 26

and 52. Renal function tests were performed during these visits as part of safety monitoring.

Phone calls were made to the participants at weeks 12 and 38. Additional un-scheduled visits

were arranged if deemed necessary by the study team (Supplemental Table 1). If withdrawal

of participation occurred after 12 weeks of treatment allocation, final visit procedures

(including CMR to assess endpoints) would be performed at the time of withdrawal. At each

clinic visit, adverse events related to study treatment were assessed by the study team. If

present, they were recorded and reported to the local ethics board and the Health Sciences

Authority within the required timeline.

Cardiac volumes, function, myocardial mass, and interstitial volumes were assessed at

baseline and after 52 weeks of treatment using standardized imaging CMR protocols on the

Siemens Aera 1.5T scanner (Siemens Healthineers, Erlangen, Germany). Balanced steady-

state free precession cine images were acquired in the long-axis 2-, 3- and 4-chamber views;

and short axis cines extending from the mitral valve annulus to the apex (acquired voxel size

of $1.6 \times 1.3 \times 8.0$ mm^3 and 30 phases per cardiac cycle). T1 mapping was performed with the

Modified Look-Locker Inversion-recovery sequence, where native T1 maps were acquired

using a heartbeat scheme of 5(3)3, and post-contrast T1 maps were acquired 15 minutes after

administration of 0.1 mmol/kg gadobutrol (Gadovist; Bayer Pharma AG, Germany) using a

heartbeat scheme of 4(1)3(1)2. Details of the imaging protocol were published in the trial

protocol[11].

All CMR images were de-identified and analyzed by trained personnel at the National Heart

Research Institute Singapore CMR core-lab using CVI42 software (Circle Cardiovascular

Imaging, Calgary, Canada) according to the published analysis protocols[32,35].

Primary Trial Endpoint

The primary endpoint was the difference in absolute change in interstitial volume from

baseline to post-treatment between the two treatment groups. Interstitial volume, reflecting

diffuse interstitial fibrosis in the absence of infiltrative diseases, was derived as the product of

ECV fraction × myocardial volume. This approach of quantifying interstitial volume was

previously validated against histological fibrosis[10] and demonstrated prognostic value in

patients with HHD[36].

ECV fraction was measured using the T1 mapping module in CVI42. Hematocrit for

calculating ECV fraction was taken on the day of CMR. Myocardial volume (mL) was

calculated as myocardial mass (g) divided by the specific density of the myocardium (1.05

g/mL).

Secondary Trial Endpoints

Secondary endpoints were absolute change in CMR markers of cardiac remodeling that

included LV morphology, function (including multi-directional LV strain) and mechanics.

LV concentricity was defined by the ratio between LV mass and end-diastolic volume

(EDV). Myocyte volume was derived as the product of (1-ECV fraction) and myocardial

volume. Maximal left atrial volume was estimated based on the biplane area-length method

using the 2- and 4-chamber views, at end atrial diastole[32].

Several CMR measures of myocardial mechanics were assessed. The Remodeling Index (RI)

was used as a marker of LV wall stress[35,37]. The RI was calculated as $\frac{\sqrt[3]{EDV}}{t}$, where t is the

maximal wall thickness (cm) across the 16 myocardial segments. The ventricular-arterial

coupling ratio (VCR) was used to describe the relationship between LV and arterial system. It

was calculated as the ratio of LV end-systolic volume to stroke volume[38]. LV filling

pressures were estimated using a CMR model consisting of left atrial (LA) volume and LV

mass, which had been validated with invasively measured pulmonary wedge pressure and

demonstrated to have prognostic implications[39].

Blood samples were collected on the first and final study visit; and stored at −80°C.

Biochemical analyses were conducted in a single freeze-thaw cycle across two assay runs at a

College of American Pathologists-accredited laboratory (Changi General Hospital,

Singapore). Serum NT-proBNP (N-terminal pro-B-type natriuretic peptide; proBNP II STAT,

Roche Diagnostics, Germany) and hsTnT (high-sensitivity cardiac troponin T; Trop T hs

STAT; Roche Diagnostics, Germany) were measured using electrochemiluminescence

immunoassay on the Cobas E801 analyzer (Roche Diagnostics Asia Pacific, Singapore). The

lab-reported lower detection limits for NT-proBNP and hsTnT were 11 pg/mL and 3 pg/mL,

respectively. All biochemical concentrations below detection thresholds were assigned a

value of half the detection limit.

Sample Size Estimation

An estimated sample size of 70 (35 per treatment group) would provide effective power of 80% at a 5% significance level (two-sided) to detect superiority of sacubitril/valsartan over valsartan for the primary endpoint, assuming an absolute minimum difference of 3.5 mL/m^2 in mean interstitial volume change between the two groups, a standard deviation (SD) of 5.8mL/m^2 (data subsequently published[17]) and a moderate correlation of 0.60 between interstitial volume at baseline and week 52. This effect size was based on an estimate of the magnitude of myocardial fibrosis regression that could be expected to translate into improved cardiac function and clinical outcomes[8]. The study plan was to recruit 80 participants, allowing a withdrawal rate of up to 15%.

Statistical Analysis

The endpoint comparison was based on intention-to-treat analysis, i.e., all participants were analyzed as part of the group to which they had been randomized. Continuous variables were presented as mean ± SD and categorical variables were expressed as count (percentage). Skewed data were presented as medians and inter-quartile range. Comparisons between treatment groups were performed using independent samples t-tests for continuous variables and χ^2 test for dichotomous variables.

For primary endpoint, analysis-of-covariance (ANCOVA) was performed to adjust for baseline covariates. Treatment group at randomization was included as fixed effect. Baseline interstitial volume, 24h SBP and prior ACEI/ARB use were included as covariates. Results were reported as adjusted mean difference with 95% confidence intervals (CI). Post-hoc subgroup analyses of the primary endpoint were performed using two-way analysis of variance (ANOVA) to test heterogeneity of treatment effects in the following subgroups: age,

sex, body mass index, whether washout was performed, LV ejection fraction, interstitial

volume and 24h SBP. Continuous covariates were categorized by median values. The

hypothesis testing on secondary endpoints was considered exploratory. The analyses were

presented with effect estimates and unadjusted 95% CI. Adjustment for multiple comparisons

was not performed.

The change in NT-proBNP and hsTnT were reported as the ratio of geometric mean values at

baseline and 52 weeks. For treatment comparisons, the effective treatment ratio (ETR) was

calculated as the ratio of geometric mean changes in biomarkers between the two groups. An

ETR closer to 0 indicates greater effectiveness of sacubitril/valsartan in reducing biomarker

level compared to valsartan.

Statistical significance was set at two-sided P value <0.05. All analyses were performed with

SPSS, version 24 (SPSS; IBM, INC, Armonk, NY) and R, version 4.3.3 (R foundation for

statistical computing, Vienna, Austria). Graphs were created using GraphPad Prism 8.1.2

(GraphPad Software, Inc, San Diego, CA). Statistical analyses were performed by team

members who were not involved in image analysis and participants' follow up.

DATA AVAILABILITY

De-identified data for this study is available for non-commercial academic purposes. Request

for access can be made with the corresponding author and should include a clear research

plan with statistical considerations. Requests will be reviewed by the Data Access Committee

at the National Heart Centre Singapore within 6 weeks. The data required for the approved,

specified purposes will be provided after completion of a data sharing agreement that include

instructions on publications, reporting and usage policy. Source data are provided with paper.

REFERENCES

1. Goldberg LR, Jessup M. Stage B heart failure: management of asymptomatic left

ventricular systolic dysfunction. *Circulation*. **113**, 2851-2860 (2006).

2. Mohebi R, Wang D, Lau ES, et al. Effect of 2022 ACC/AHA/HFSA criteria on stages

of heart failure in a pooled community cohort. *J Am Coll Cardiol*. **81**, 2231-2242

(2023).

3. Messerli FH, Rimoldi SF, Bangalore S. The transition from hypertension

to heart failure: contemporary update. *JACC Heart Fail*. **5**, 543-551 (2017).

4. González A, López B, Ravassa S, et al. Myocardial Interstitial fibrosis in hypertensive

heart disease: from mechanisms to clinical management. *Hypertens*. **81**, 218-228

(2024).

5. González A, Schelbert EB, Díez J, Butler J. Myocardial interstitial fibrosis in heart

failure. *J Am Coll Cardiol*. **71**, 1696-1706 (2018).

6. Ruilope LM, Dukat A, Böhm M, Lacourcière Y, Gong J, Lefkowitz MP. Blood-

pressure reduction with LCZ696, a novel dual-acting inhibitor of the angiotensin II

receptor and neprilysin: a randomised, double-blind, placebo-controlled, active

comparator study. *Lancetl*. **375**, 1255-1266 (2010).

7. McMurray JJV, Packer M, Desai AS, et al. Angiotensin–neprilysin inhibition versus

enalapril in heart failure. *N Engl J Med*. **371**, 993-1004 (2014).

8. Suematsu Y, Miura S, Goto M, et al. LCZ696, an angiotensin receptor-neprilysin

inhibitor, improves cardiac function with the attenuation of fibrosis in heart failure with

reduced ejection fraction in streptozotocin-induced diabetic mice. *Eur J Heart Fail*. **18**,

386-393 (2016).

9.   Burke RM, Lighthouse JK, Mickelsen DM, Small EM. Sacubitril/Valsartan decreases

cardiac fibrosis in left ventricle pressure overload by restoring pkg signaling in cardiac

fibroblasts. *Circ Heart Fail*. **12**:e005565 (2019).

10. Chin CWL, Everett RJ, Kwiecinski J, et al. Myocardial fibrosis and cardiac

decompensation in aortic stenosis. *JACC Cardiovasc Imaging*. **10**, 1320-1333 (2017).

11. Brilla CG, Funck RC, Rupp H. Lisinopril-mediated regression of myocardial fibrosis in

patients with hypertensive heart disease. *Circulation*. **102**, 1388-1393 (2000).

12. Díez J, Querejeta R, López B, González A, Larman M, Martínez Ubago JL. Losartan-

dependent regression of myocardial fibrosis is associated with reduction of left

ventricular chamber stiffness in hypertensive patients. *Circulation*. **105**, 2512-2517

(2002).

13. López B, Querejeta R, González A, Sánchez E, Larman M, Díez J. Effects of loop

diuretics on myocardial fibrosis and collagen type I turnover in chronic heart failure. *J

Am Coll Cardiol*. **43**, 2028-2035 (2004).

14. Izawa H, Murohara T, Nagata K, et al. Mineralocorticoid receptor antagonism

ameliorates left ventricular diastolic dysfunction and myocardial fibrosis in mildly

symptomatic patients with idiopathic dilated cardiomyopathy: a pilot study.

Circulation. **112**, 2940-2945 (2005).

15. López B, Querejeta R, Varo N, et al. Usefulness of serum carboxy-terminal propeptide

of procollagen type I in assessment of the cardioreparative ability of antihypertensive

treatment in hypertensive patients. *Circulation*. **104,** 286-291 (2001).

16. Mak GJ, Ledwidge MT, Watson CJ, et al. Natural history of markers of collagen

turnover in patients with early diastolic dysfunction and impact of eplerenone. *J Am

Coll Cardiol*. **54**, 1674-1682 (2009).

17. Kosmala W, Przewlocka-Kosmala M, Szczepanik-Osadnik H, Mysiak A, O'Moore-

Sullivan T, Marwick TH. A randomized study of the beneficial effects of aldosterone

antagonism on LV function, structure, and fibrosis markers in metabolic syndrome.

JACC Cardiovasc Imaging. **4**, 1239-1249 (2011).

18. Kosmala W, Przewlocka-Kosmala M, Szczepanik-Osadnik H, Mysiak A, Marwick TH.

Fibrosis and cardiac function in obesity: a randomised controlled trial of aldosterone

blockade. *Heart*. **99**, 320-326 (2013).

19. Bavishi C, Messerli FH, Kadosh B, Ruilope LM, Kario K. Role of neprilysin inhibitor

combinations in hypertension: insights from hypertension and heart failure trials. *Eur*

Heart J. **36**, 1967-1973 (2015).

20. Loo G, Puar T, Foo R, et al. Unique characteristics of Asians with hypertension: what

is known and what can be done? *J Hypertens*. **42**, 1482-1489 (2024).

21. Schmieder RE, Wagner F, Mayr M, et al. The effect of sacubitril/valsartan compared to

olmesartan on cardiovascular remodelling in subjects with essential hypertension: the

results of a randomized, double-blind, active-controlled study. *Eur Heart J*. **38**, 3308-

3317 (2017).

22. Solomon SD, McMurray JJV, Anand IS, et al. Angiotensin–neprilysin inhibition in

heart failure with preserved ejection fraction. *N Engl J Med*. **381**, 1609-1620 (2019).

23.   Shah AM, Cikes M, Prasad N, et al. Echocardiographic features of patients with

heart failure and preserved left ventricular ejection fraction. *J Am Coll Cardiol*. **74,**

2858-2873 (2019).

24.   Palazzuoli A, Caravita S, Paolillo S, et al. Current gaps in HFpEF trials: Time to

reconsider patients' selection and to target phenotypes. *Prog Cardiovasc Dis*. **67**, 89-97

(2021).

25.   Sweeney M, Corden B, Cook SA. Targeting cardiac fibrosis in heart failure with
539   preserved ejection fraction: mirage or miracle? *EMBO Mol Med*. **12**, e10865 (2020).

540   26.   Lewis GA, Dodd S, Clayton D, et al. Pirfenidone in heart failure with preserved
541   ejection fraction: a randomized phase 2 trial. *Nat Med*. **27**, 1477-1482 (2021).

542   27.   Goh VJ, Le TT, Bryant J, et al. Novel Index of maladaptive myocardial remodeling in
543   hypertension. *Circ Cardiovasc Imaging*. **10**, e006840 (2017).

544   28.   Schelbert EB, Fonarow GC, Bonow RO, Butler J, Gheorghiade M. Therapeutic targets
545   in heart failure: refocusing on the myocardial interstitium. *J Am Coll Cardiol*. **63**, 2188-
546   2198 (2014).

547   29.   Schelbert EB, Butler J, Diez J. Why clinicians should care about the cardiac
548   interstitium. *JACC Cardiovasc Imaging*. **12**, 2305-2318 (2019).

549   30.   Haynes RB, Taylor DW, Sackett DL, Gibson ES, Bernholz CD, Mukherjee J. Can
550   simple clinical measurements detect patient noncompliance? *Hypertens*. **2**, 757-764
551   (1980).

552   31.   Lee V, Zheng Q, Toh DF, et al. Sacubitril/valsartan versus valsartan in regressing
553   myocardial fibrosis in hypertension: a prospective, randomized, open-label, blinded
554   endpoint clinical trial protocol. *Front Cardiovasc Med*. **10**, 1248468 (2023).

555   32.   Le TT, Tan RS, De Deyn M, et al. Cardiovascular magnetic resonance reference ranges
556   for the heart and aorta in Chinese at 3T. *J Cardiovasc Magn Reson*. **18**, 21 (2016).

557   33.   Weber MA, Schiffrin EL, White WB, et al. Clinical practice guidelines for the
558   management of hypertension in the community: a statement by the American Society of
559   Hypertension and the International Society of Hypertension. *J Clin Hypertens*. **16**, 14-
560   26 (2014).

561   34.   Ministry of Health, Singapore. *MOH clinical practice guidelines on hypertension*
562   *1/2017*. Ministry of Health, Singapore; 2017.

https://www.moh.gov.sg/docs/librariesprovider4/guidelines/cpg_hypertension-booklet--

-nov-2017.pdf.

35. Goh VJ, Le TT, Bryant J, et al. Novel Index of maladaptive myocardial remodeling in

hypertension. *Circ Cardiovasc Imaging*. **10**, e006840 (2017).

36. Iyer NR, Le TT, Kui MSL, et al. Markers of focal and diffuse nonischemic myocardial

fibrosis are associated with adverse cardiac remodeling and prognosis in patients with

hypertension: the REMODEL study. *Hypertens*. **79**, 1804-1813 (2022;).

37. Le TT, Lim V, Ibrahim R, et al. The remodelling index risk stratifies patients with

hypertensive left ventricular hypertrophy. *Eur Heart J Cardiovasc Imaging*. **22**, 670-

679 (2021).

38. Holm H, Magnusson M, Jujić A, Bozec E, Girerd N. How to calculate ventricular–

arterial coupling? *Eur J Heart Fail*. **24**, 600-602 (2022).

39. Garg P, Gosling R, Swoboda P, et al. Cardiac magnetic resonance identifies raised left

ventricular filling pressure: prognostic implications. *Eur Heart J*. **43**, 2511-2522

(2022).

ACKNOWLEDGEMENTS

The authors wish to thank all the study participants, the radiographers at the Department of

Cardiovascular Magnetic Resonance, the pharmacists at the Department of Pharmacy and the

research coordinators at the Clinical Trial Research Office, National Heart Centre Singapore

for their assistance in the study. REVERSE-LVH is funded by the National Medical Research

Council of Singapore (MOH-CTGIIT17may-0001).

AUTHOR CONTRIBUTIONS

CWLC, TTL, SAC, AMR and CHL contributed to the conception and design of the study.

QZ designed the statistical analysis plan. VL and DFT led the data collection and

management. VL, MD, and QZ performed the data analysis. JAB and TTL led the CMR

image analysis. TA analyzed the circulating biomarkers. JB, JD, RF and CSPL provided

important inputs to the overall trial conduct. VL and MD wrote the first draft of the

manuscript. CWLC secured funding for the trial. All authors made critical revisions for

important intellectual content, contributed to the article, and approved the submitted version.

COMPETING INTERESTS

Prof. Lam has received consulting fees from Novartis. Prof. Butler is a consultant to

Novartis. The other authors do not have any relevant conflicts of interest to disclose.

Table 1. Baseline Characteristics of All Randomized Participants

	All Participants (n=78)	Sacubitril/ valsartan (n=39)	Valsartan (n=39)
Baseline Clinical Characteristics			
Age, y	58±11	56±11	59±11
Males, n (%)	32 (41)	18 (46)	14 (36)
Chinese ethnicity, n (%)	71 (91)	35 (90)	36 (92)
Weight, kg	71±18	74±19	68±16
Body mass index, kg/m^2	27.2±5.6	27.7±6.2	26.7±5.1
Dyslipidemia, n (%)	33 (42)	18 (46)	15 (38)
Diabetes mellitus, n (%)	14 (18)	9 (23)	5 (13)
Smoking, n (%)	6 (8)	3 (8)	3 (8)
24h systolic blood pressure, mmHg	137±14	138±17	137±10
24h diastolic blood pressure, mmHg	81±11	82±11	80±11
Baseline Biochemical Characteristics			
Potassium, mmol/L	3.8 [3.6, 4.0]	3.8 [3.6, 4.0]	3.7 [3.6, 4.0]
Creatinine, µmol/L	72 [52, 86]	73 [51, 88]	72 [54, 85]
NT-proBNP, pg/mL	53 [28, 141]	42 [28, 101]	60 [31, 275]
High sensitivity cTnT, pg/mL	9.8 [6.5, 14.5]	9.5 [6.6, 15.0]	10.9 [6.0, 14.5]
Estimated GFR, mL/min/1.73 m^{2a}	90 [72, 106]	94 [74, 108]	88 [70, 102]
Baseline Cardiovascular Magnetic Resonance Characteristics			
Indexed LV mass, g/m^2	64±19	64±19	64±18
Indexed LV end diastolic volume, mL/m^2	78±15	76±16	79±15
Indexed LV end systolic volume, mL/m^2	32±12	32±13	32±11
Indexed LV stroke volume, mL/m^2	46±7	45±8	47±7
LV ejection fraction	60±9	59±10	60±7
Indexed RV end diastolic volume, mL/m^2	74±13	73±13	74±13
Indexed RV end systolic volume, mL/m^2	28±10	28±9	27±11
Indexed RV stroke volume, mL/m^2	46±7	45±8	47±7
RV ejection fraction	63±9	62±8	64±9
Indexed left atrial volume, mL/m^2	55±15	53±12	56±18
Indexed right atrial area, cm^2/m^2	12.0±2.1	11.7±1.8	12.4±2.2
Remodeling Index	5.33±0.99	5.22±0.80	5.44±1.15
Ventricular-arterial coupling ratio	0.71±0.29	0.74±0.36	0.69±0.22
Estimated LV filling pressure, mmHg	15.6±2.8	15.7±3.0	15.5±2.6
Maximal wall thickness, mm	10.0±2.4	10.1±2.1	9.9±2.7
Mean wall thickness, mm	7.5±1.6	7.7±1.6	7.4±1.6
LV mass/end-diastolic volume ratio	0.83±0.17	0.84±0.14	0.81±0.19

Global longitudinal strain, %	-16.5±3.3	-16.0±3.2	-17.0±3.2
Global circumferential strain, %	-18.7±3.2	-18.2±3.5	-19.1±2.8
Global radial strain, %	32.4±8.6	31.2±9.2	33.6±7.8
Extracellular volume fraction, %	26.4±2.6	26.1±2.4	26.8±2.8
Indexed myocyte volume, mL/m^2	45.0±13.1	45.4±13.7	44.5±12.6
Non ischemic late gadolinium enhancement, n (%)	29 (37)	15 (38)	14 (36)

Values were presented as mean ± standard deviations or median [interquartile range] unless

otherwise indicated. [a]Estimated GFR calculated using MDRD equation = 175 x [serum

creatinine (μmol/L)/88.4]$^{-1.154}$ ´ (age) $^{-0.203}$ ´ (0.742 if female). Left ventricular hypertrophy

was defined based on age- and sex-specific reference ranges established in the local

population (females: 47g/m^2 and males: 65g/m^2). Abnormal global longitudinal strain in the

population was defined as < -14.0 to -15.0% in males and < -17.0 to -18.0% in females[20].

GFR = glomerular filtration rate; LV = left ventricular; RV = right ventricular; NT-proBNP =

N-terminal pro-B-type natriuretic peptide; cTnT = cardiac troponin T

Table 2. Secondary Endpoints After 52 Weeks of Treatment

	Change from Baseline to Post-treatment		Treatment Difference	
	Sacubitril/valsartan	Valsartan	95% Confidence Interval	P Value
LV mass, g	-21.9±25.4	-11.0±9.0	-19.5 to -2.3	0.014
LV end diastolic volume, mL	-12.4±18.7	-7.1±16.4	-13.2 to 2.7	0.194
LV end systolic volume, mL	-8.0±18.3	-5.2±15.7	-10.5 to 5.0	0.482
LV stroke volume, mL	-4.4±9.3	-1.9±9.4	-6.7 to 1.7	0.243
LV ejection fraction	1.2±6.5	2.1±6.2	-3.8 to 1.9	0.511
RV end diastolic volume, mL	-6.8±11.5	-4.8±16.5	-8.4 to 4.5	0.545
RV end systolic volume, mL	-2.2±10.5	-2.4±13.0	-5.1 to 5.6	0.926
RV stroke volume, mL	-4.6±8.6	-2.4±12.5	-7.1 to 2.7	0.368
RV ejection fraction	-0.3±6.3	0.1±8.3	-3.7 to 2.9	0.811
Left atrial volume, mL	-18.3±16.6	-8.6±13.6	-16.6 to -2.9	0.006
Right atrial area, cm^2	-1.5±2.5	-1.3±2.2	-1.3 to 0.9	0.694
LV mass/end-diastolic volume ratio	-0.08±0.09	-0.04±0.09	-0.08 to 0.00	0.072
Remodeling Index	0.46±0.56	0.38±0.50	-0.17 to 0.31	0.544
Ventricular-arterial coupling ratio	-0.07±0.26	-0.04±0.23	-0.14 to 0.09	0.655
Estimated LV filling pressure, mmHg	-1.8±1.5	-0.9±1.1	-1.5 to -0.3	0.003
Maximal wall thickness, mm	-1.1±1.0	-0.7±0.7	-0.8 to 0.0	0.065

Global longitudinal strain, %	-0.9±2.3[a]	0.2±1.9	-2.0 to -0.1	0.027
Global circumferential strain, %	-0.6±2.2[a]	-0.03±1.9[a]	-1.5 to 0.4	0.253
Global radial strain, %	1.7±6.5	-0.2±5.8	-0.9 to 4.7	0.189
Extracellular volume fraction, %	0.3±2.0	0.3±2.7	-1.1 to 1.1	0.962
Myocyte volume, mL	-16.0±19.5	-8.0±7.1	-14.6 to -1.4	0.019

612 [a]Negative value signifies an improvement in strain.

613 LV = left ventricular; RV = right ventricular; NT-proBNP = N-terminal pro-B-type natriuretic peptide; cTnT = cardiac troponin T.

614 Values were presented in mean ± standard deviations. Treatment difference was calculated using Student's t-test (two-sided) and presented in

615 95% confidence intervals and corresponding P values.

616 **Table 3. Renal Function Between Treatment Groups Throughout Study Trial**

	Week	Sacubitril/ valsartan	Valsartan	P value
	0	3.8 [3.6, 4.0]	3.7 [3.6, 4.0]	0.800
	4	3.9 [3.7, 4.3]	4.0 [3.7, 4.3]	0.778
Potassium, mmol/L	8	4.0 [3.8, 4.2]	4.0 [3.9, 4.2]	0.766
	26	4.1 [3.7, 4.5]	4.1 [3.9, 4.3]	0.765
	52	3.9 [3.6, 4.2]	3.8 [3.5, 4.2]	0.602
	0	94 [74, 108]	88 [70, 102]	0.420
	4	82 [68, 108]	84 [74, 100]	0.996
Estimated GFR, mL/min/1.73m^2	8	86 [71, 102]	82 [63, 110]	0.909
	26	89 [65, 107]	83 [69, 96]	0.897
	52	89 [63, 104]	83 [69, 100]	0.796

618 Values were presented in median [inter-quartile range]. Comparisons between treatment

619 groups were performed with Mann-Whitney U (two-sided) test.

620 GFR = glomerular filtration rate.

 FIGURE LEGENDS

Figure 1. Participant Recruitment, Randomization and Treatment

A total of 78 participants were randomized in the trial. One patient in the sacubitril/valsartan

group withdrew at week 4 due to concerns of medication side effects. One patient in the

valsartan group withdrew at week 26 due to difficulties with trial commitment. This

participant had final visit investigations (including cardiovascular magnetic resonance)

performed at time of withdrawal, according to study protocol. Primary and secondary

endpoints were analyzed as intention-to-treat.

Figure 2. Effects of Sacubitril/valsartan Compared to Valsartan on Primary Endpoint

The primary endpoint in 78 participants. The effect estimates were presented in mean and

95% confidence intervals, with corresponding P values from one-way ANCOVA testing

before and after adjustment for baseline covariates. Abbreviations: ACEI angiotensin-

converting enzyme inhibitor; ARB, angiotensin receptor blocker.

Figure 3. Blood Pressure Control Between Sacubitril/valsartan and Valsartan

The mean blood pressure assessed at all timepoints during the trial were similar between the

two treatment groups (n=78). Data were presented in mean and 95% confidence interval; and

P values from Student's t-test (two-sided) comparison.

Systolic Blood Pressure