M-PreSS: A Model Pre-training Approach for Study Screening in Systematic Reviews

Zhaozhen Xu [*1,2], Philippa Davies[2], Louise AC Millard[1,2], Lam Teng[4], Georgios Markozannes[4], Eduardo AP Seleiro[4], Julian PT Higgins[2], Richard M Martin[2,3], Konstantinos K Tsilidis[4], Doris SM Chan[4], Tom R Gaunt[1,2], and Yi Liu [†1,2]

[1]MRC Integrative Epidemiology Unit, University of Bristol, UK

[2]Bristol Medical School, Department of Population Health Sciences, University of Bristol, UK

[3]National Institute for Health Research (NIHR) Bristol Biomedical Research Centre, University Hospitals Bristol and Weston NHS Foundation Trust and University of Bristol, Bristol, UK

[4]Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, UK

Abstract

Background

Conducting a systematic review is labour-intensive and time-consuming, especially during the study screening process. Previous research has introduced traditional machine learning models (e.g. Support Vector Machines) to automate the study screening process, but it is difficult to generalise across topics. Therefore, recent research has explored the use of existing large language models (LLMs), such as ChatGPT/GPT-4, for study screening. However, the lack of transparency in training data and consistency in output results make applying such large, commercial LLMs challenging in the context of systematic reviews where transparency in methods is particularly important.

Results

We introduce an approach to fine-tune an open-source biomedical language model (Blue-BERT) using a Siamese neural network [1] so that it screens the scientific literature databases on multiple research topics. We evaluate different training approaches in seven COVID-19 systematic reviews. The results indicate good generalisation among topics with an average recall/sensitivity of 0.86 (minimum: 0.67, maximum: 1.00) and an average false positive rate of 6.48% (minimum: 1.38%, maximum: 11.41%). Furthermore, adding study selec-

---

*zhaozhen.xu@bristol.ac.uk

†yi6240.liu@bristol.ac.uk

tion criteria to the topic definition can improve the model performance (Area Under the Precision-Recall Curve [PRAUC]) by 2.74%, and adding more related review topics during training can increase the performance by 15.82%.

Conclusions

Our results indicate that fine-tuning BlueBERT with study screening datasets can outperform ChatGPT/GPT-4 in two out of three COVID-19 review topics reported in the literature, whilst maintaining the ability for researchers to continue updating or extending the search for related evidence and significantly reducing the computational resource requirements.

1 Introduction

Systematic literature reviews are widely used to support evidence-based research and inform policy and practice in many domains, including biomedical and clinical sciences. Compared to a narrative literature review, which gives a summary or overview of a topic, a systematic review aims to provide a transparent, thorough, and unbiased evaluation of all the existing evidence for a specific research question, including the assessment of the risk of bias, and often the quantitative summation of study results using a meta-analysis. Guidelines are provided for conducting systematic reviews, which aim to make the processes transparent and repeatable. The same review team might want to update the review with new evidence or conduct different reviews under a general research theme, where reproducibility and efficiency can be improved by the use of computational tools.[2]

Automation in systematic reviews has been widely studied due to the potential for computerising the repeatable systematic processes involved. The process to automate usually breaks down into three tasks: study screening, key information extraction, and risk of bias assessment to evaluate the validity of the study results. In this study, we focus on the first of these key processes: study screening.

Study screening (also known as study selection) aims to identify the primary studies for a review topic. To include all related studies, the traditional approach is to conduct an exhaustive search in relevant bibliographic databases using pre-defined searching strategies, followed by manual screening of the retrieved study records according to the eligibility criteria. In practice, a search from literature databases can return thousands of candidate studies for each systematic review topic. Then, a minimum of two reviewers typically independently screen the study records and

agree on a final decision on whether a study should be included or excluded in the systematic review.[3] The screening typically comprises two phases. In the initial phase, the titles and abstracts are screened to remove obviously irrelevant references, which significantly reduces the number of references for the next phase. Then, the full-text papers are screened to make the decisions on inclusion in the review.

A significant challenge in study selection is the substantial time and human resources needed for the initial screening for relevant studies. Previous study[4] surveyed 33 experienced systematic review reviewers and found that it can take a full-time systematic reviewer an average of 33 days for the entire screening process. Hence, automating the study screening process with machine learning (ML) has the potential to increase the efficiency of conducting systematic reviews.

Existing tools[5] and models[6,7] apply traditional ML and natural language processing (NLP) approaches to identify primary studies and have shown promise in using ML models to semi-automate study screening in the fields of software engineering, medicine, and biology.[6,7] However, a specific ML model needs to be trained for each review topic to obtain optimal prediction results, and cannot then be directly applied to new topics without new labelled data and further training. Another challenge of applying ML to study screening is class imbalance. After screening, only a small percentage of candidate studies are included in the review, which limits the amount of 'positive' examples in the training data. This can cause a bias towards the majority class during training, which limits the performance of the learnt model on identifying relevant studies. Previous research[7] observed that up-sampling of included studies resulted in poor predictions and increasing computation time in study screening, while down-sampling of excluded studies was a better strategy for imbalanced data but resulted in reducing training data.

Recent developments in ML, particularly in large language models (LLMs), such as BERT[8] and GPTs,[9,10] have resulted in models that understand key aspects of human language and learn general knowledge by pre-training on large amounts of text, such as books, Wikipedia, and academic literature. Such models therefore have the potential to be used in various tasks related to the processing of natural language (e.g. information retrieval,[11] text classification[12]) and for public health research problems in the biomedical domain.[13,14] Furthermore, LLMs exhibit a good generalisation to unseen data without any training (i.e. zero-shot learning) and are increasingly used for automating systematic reviews.[15]

In study screening, some studies[16–18] have shown the potential to use an existing language

model to perform zero-shot classification on different topics. However, they use pre-trained models without tailoring them for study screening. Thus, the recall and false positive rate of those models can potentially be increased by fine-tuning the model with existing study selection datasets.

Another study from Guo et al.[19] demonstrates that using ChatGPT and GPT-4 can provide promising results as a complementary tool to human reviewers in screening clinical reviews. However, GPT-4 misclassified 22.8% of the relevant studies (average recall: 0.77) as irrelevant. In addition, responses from ChatGPT can be inconsistent across sessions, limiting the reproducibility of this approach. The results of previous studies highlight the need to investigate whether fine-tuning an open-source LLM with systematic review datasets can further improve the performance of the LLM on study screening. Our work builds on Guo et al.'s by fine-tuning a different LLM using the same data and comparing the performance with their results.

In this study, we introduce M-PreSS, a model pre-training approach for study screening, for different systematic review topics. This approach aims to train a general study model that can be used among multiple topics and potentially extended to new/unseen topics. A general model is not expected to excel in all topics but demonstrates robust performance across multiple diverse topics. For instance, a general model may perform well on cancer-related topics but not across all disease categories.

Our model is based on BlueBERT,[20] a biomedical version of the original BERT network. BERT is a well-established language model pre-trained with Wikipedia and BookCorpus, and demonstrates strong performance across various language understanding tasks.[8] Based on the original BERT, BlueBERT is further trained with PubMed abstracts and clinical notes, which results in a good understanding of biomedicine text and performance in multiple biomedical natural language processing tasks, such as sentence similarity and named entity recognition. By fine-tuning BlueBERT with existing study screening datasets, the model may then be more effectively directed at identifying relevant studies on a given systematic review topic.

One previous research[21] fine-tuned BERT to classify included and excluded studies using study titles and abstracts. Similar to traditional ML approaches, each systematic review topic requires its own model to be trained. Training a topic-specific screening model in this way relies on a sufficient number of human-labelled records, which would not typically be the case in real-world usage.

Our study leverages a Siamese network to fine-tune BlueBERT so that one general model can be used across topics. The Siamese network is an architecture that consists of two identical subnetworks for comparing the relationship and similarity between two inputs.[1] In our study, the Siamese network allows the model to take both review topic and reference as inputs, and make different inclusion/exclusion decisions for a reference depending on the topic provided. Siamese networks have been shown to be robust to imbalanced data for classification task.[22] Other research[23] has also demonstrated that Siamese structures achieve higher recall for positive examples (i.e. included studies) compared to cross-encoders, which concatenate inputs for training. Therefore, it is applicable for study screening, where there are few positive examples and identifying all relevant studies is important (i.e. high recall).

Combining the power of LLM and Siamese networks, M-PreSS can be generalised to multiple topics for a research focus (e.g. efficacy studies of COVID-19), which can be useful for research teams consistently looking for different types of evidence for a particular sub-domain. More details of the M-PreSS architecture are explained in the Method: General Model for Study Screening Section.

The Experiments Section presents the methodology for training and evaluating the M-PreSS model. When training our model, we leverage screening datasets from multiple sources. The idea is for the model to capture the general skill for study screening from diverse topics so that it can improve the overall performance among these topics, and can be potentially generalised to unseen topics. We train our model with various data sizes and topic homogeneity and evaluate its performance on open datasets to understand the impact of different training data on model performance. Moreover, to further explore the model's generalisability, we evaluate it on unseen topics. The results are demonstrated in the Results and Discussions Section.

2   Method: General Model for Study Screening

We call M-PreSS a general model as it aims to use one model for multiple topics instead of customising topic-specific models. Figure 1 illustrates the architecture of our model. M-Press produces a priority score for a candidate study based on the review topic. During the initial study screening process, reviewers can utilise the trained model to rank studies and generate machine predictions on inclusion/exclusion, which support their decision-making.

Figure 1: Architecture of M-PreSS. Blue: Input ($Systematic\ Review\ Topic, Candidate\ Study$) pair. Green: M-PreSS. Orange: Output priority score. Note that the trained study screening models are identical, which means they share the same setup and weights.

M-PreSS takes ($Systematic\ Review\ Topic, Candidate\ Study$) pairs as inputs, where the number of candidate studies per topic differs across the dataset. The training size is determined by the total number of candidate studies across all topics.

The systematic review topic is the review title. The candidate study includes the title and abstract. Some keywords, such as 'Query', 'Title', and 'Abstract', were provided to assist the language model in capturing the information in the topic and study. The input is constructed as below.

```
Query: {Systematic review title}.
```
```
Title: {Study title}. Abstract: {Study Abstract}
```

Then, the trained study model is able to embed the inputs into dense vector representations (i.e. embeddings). These embeddings can capture the semantic meaning and key features of the topic and study.

The output of M-PreSS is a priority score, which can be used to rank the candidate studies for later review. The priority score is the cosine similarity between the systematic review topic and the candidate study embeddings. During the screening, the reviewers can pay less attention to studies with extremely high or low priority scores, thus increasing their efficiency.

Examples of input and output data from the model can be found in supplementary material Table 6.

To train the study screening model, we employ a Siamese BERT structure using the Sentence-

BERT[24] framework. Sentence-BERT leverages the Siamese architecture to capture the text similarity between two input sentences. The embeddings of sentences with similar meanings are close to each other in the embedding space.

In this study, we utilise the same Siamese structure to capture the relationship between the given systematic review topic and a candidate study. However, instead of the sentence-level inputs used in Sentence-BERT, we take the entire topic and candidate study as document-level inputs. We train the study screening model by fine-tuning BlueBERT using existing study screening records from previously completed systematic reviews. The aim is that a relevant study receives a higher cosine similarity to the topic than an irrelevant one during training.

Once trained, a topic and candidate study can be input and a priority score is assigned based on their relationship. Candidate studies relevant to the review topic are supposed to receive a higher priority score than irrelevant ones. The priority score for a given candidate study varies with the review topic, enabling one general model to determine whether a study should be included for various topics.

2.1 Binary Decision Threshold

When M-Press is used to assist human reviewers, it is also helpful for the model to propose a decision on whether a candidate study should be included in a specific review and a boundary for a reviewer to stop looking for relevant studies. Therefore, we introduce a decision threshold to give a binary output on inclusion/exclusion. If a $(Systematic\ Review\ Topic, Candidate\ Study)$ pair has a priority score higher than the defined threshold, it is recommended for inclusion. A specific threshold is calculated for each topic. As the inclusion/exclusion decision is local to a specific topic and the general priority score cannot be compared between topics, we calculated the topic-specific threshold based on the priority score and the ground truth for each topic to maximise the performance. Furthermore, due to the variability of text, such as language usage and semantic structures, an embedding distance that signifies relevance in one topic might not indicate the same level of relevance in another topic. Therefore, using a topic-specific threshold can accommodate this variability and increase classification accuracy.

The threshold is determined by ranking studies based on their priority score for the topic. Each score in this ordered list serves as a possible threshold. We can identify the optimal threshold from all the possible thresholds using the human labelling.

Multiple strategies can be used to identify the optimal thresholds, such as a threshold that results in the best accuracy metric on the training data. However, in the study screening task for a systematic review, the aim is to include all relevant studies. Therefore, we calculate the threshold that provides the lowest false positive rate (FPR) at 100% recall, encouraging the model to include all relevant studies without too many false positives for a human to examine.

More specifically, we rank the candidate studies for a topic by the priority score, and calculate the recall using each priority score as the threshold.

$$Recall = \frac{TP}{TP + FN} \tag{1}$$

Where TP is the true positive and FN is the false negative from the confusion matrix. Then we identify the threshold where $Recall = 100\%$, and the false positive rate (FPR) is minimised.

$$FPR = \frac{FP}{FP + TN} \tag{2}$$

Where FP represents false positives and TN is the true negatives from the confusion matrix. The threshold is defined as the average of the priority score yielding the lowest FPR at 100% recall and the score of the next lower-ranked study. The process of identifying the best threshold is referred to as the 'best FPR at 100% recall' strategy and demonstrated in algorithm 1.

3 Experiments

To understand the performance and generalisability of our model, we train the model on various data sources and different setups.

3.1 Datasets

We use existing systematic review screening records from different sources to train our model. Each review topic contains a list of study titles, abstracts, human decisions of inclusion or exclusion, and selection criteria.

Algorithm 1 Find the best threshold with the lowest FPR at 100% recall. *value* is the priority score predicted by the model. *label* is the true label of an inclusion/exclusion decision for a study. The algorithm aims to find the threshold with the best performance for the given evaluation metrics.

Require: score: List of numerical values $\epsilon[0, 2]$
Require: label: List of binary labels, 1: positive, 0: negative
Ensure: threshold, max_performance
  1: Initialize $threshold \leftarrow -1$
  2: Initialize $max\_performance \leftarrow 0$
  3: Initialize $ncorrect \leftarrow 0$
  4: $positive\_true \leftarrow$ Number of positive examples in `label`
  5: Sort `score` in ascending order $\rightarrow$ `sorted_value`
  6: Sort `label` based on `value` sorting $\rightarrow$ `sorted_label`
  7: **for** $i = 0$ to (len(sorted_label) $- 2$) **do**
  8:   **if** `sorted_label`[i] $== 1$ **then**
  9:     Increment $ncorrect$ by 1
 10:   **end if**
 11:   **if** $ncorrect > 0$ **then**
 12:     $recall \leftarrow ncorrect/positive\_true$
 13:     **if** $recall > max\_performance$ **then**
 14:       $max\_performance \leftarrow recall$
 15:       $threshold \leftarrow ($`sorted_value`$[i] + $`sorted_value`$[i+1])/2$
 16:     **end if**
 17:   **end if**
 18: **end for**
 19: **return** $threshold, max\_performance$

Guo et al.[19] (GUO)   Guo et al. provided screening records for ten clinical systematic reviews. All the topics were related to efficacy studies; out of them, seven studies are related to COVID-19. However, only three COVID-19 topics have inclusion/exclusion criteria and were evaluated in their study.

Intervention Effectiveness   We curated study screening records for six systematic reviews addressing questions of the effectiveness of various medical interventions from our previous research[25–31] and adapted query criteria from the review objectives. This dataset enriches our training set on intervention studies and can be used to understand how model generalises to this study type.

SYNERGY[32]   This is an open study screening dataset consisting of 26 systematic reviews. In this study, we removed three reviews for computer science, which resulted in 23 review topics left among medicine, psychology, and biology domains. SYNERGY did not provide selection criteria. Therefore, we collected the selection criteria for each topic from the published systematic

review.

The systematic review topics and screening records were separated into three categories: COVID-19, effectiveness studies (ES), and SYNERGY, where ES includes three non-COVID-19 studies from GUO and all the topics from the Intervention Effectiveness Dataset. The varied heterogeneity within and across these three datasets enables evaluation of how the model performs with more versus less diverse training data. We split each of these three datasets into training (70%), validation (10%) and test (20%) sets, stratifying by review topic. Table 1 shows the basic information for each training set. An overview for all topics included is listed in the supplementary materials (Table 7).

Table 1: Overview of training sets for the general study screening model. We removed three computer science systematic review topics from the 26 SYNERGY topics. Average inclusion percentage is the macro average of the inclusion percentages for all topics in the dataset.

Dataset	Number of Topics	Number of Inclusion	Number of Records	Average Inclusion Percentage (Minimum, Maximum)
COVID-19	7	923	27, 169	5.05 (0.71, 10.98)
ES	9	609	26, 473	3.54 (0.45, 7.95)
SYNERGY	23	1, 842	106, 315	4.70 (0.16, 21.96)

3.2 Implementation Details

We first trained the model by fine-tuning BlueBERT on the COVID-19 training set and extended the training process with more diverse data from ES and SYNERGY.

During training, we applied a maximum sequence length of 512 tokens (roughly 350400 words) due to the limitation of BERT, which means a topic or study with more than 512 tokens was truncated. Then, we optimised the model using Online Contrastive Loss and AdamW optimiser[33] with a margin of 0.5 and a learning rate of $2e-5$. The aim was to increase the distance between irrelevant studies and systematic review topics in the embedding space while decreasing the distance for the relevant studies during training. We trained the model for 6 epochs with a batch size of 6 for each dataset. The model's hyperparameters, including learning rate, margin, epochs and batch size, were tuned empirically with training and validation sets, and the model was evaluated on the test set. Four NVIDIA GeForce RTX 2080 Ti GPUs were used to train the model.

The binary decision thresholds are calculated on the training set and evaluated on the test

 set.

3.3 Exploring the impact of including selection criteria in topic descriptions

BERT and its derivative models are able to capture the semantic meaning and contextual information. Therefore, adding more contextual information on the topic could potentially improve the study screening performance. To validate this assumption, we added selection criteria to the topic and compared it with the model trained without criteria using the COVID-19 dataset.

Selection criteria are the pre-specified eligibility criteria for determining which studies are included or excluded from the review. They are specified based on the review question and include aspects such as the study participants, settings, treatments of interest, and appropriate study designs. The input for the systematic review topics is structured as follows.

`Query: {Systematic review title}. Criteria: {Selection cri-`
`teria}`

We extracted and curated three types of criteria for the COVID-19 dataset: free text criteria, structured criteria, and inclusion/exclusion criteria. Free text criteria are the descriptions of the study selection criteria, which we extracted from the methods section in the published systematic review papers. In the real world however, reviewers tend to have structured criteria that contain information about participants, interventions, comparators, outcomes, and study designs. Thus, we built the structured criteria for all topics in the COVID-19 dataset. In Guo et al.'s study,[19] the criteria were defined as a list of inclusion criteria and a list of exclusion criteria. We curated our criteria following the same framework and compared the performance between different types of criteria.

The criteria used for the COVID-19 dataset can be found in supplementary material Table 8.

3.4 Exploring the impact of differences in dataset size and topic homogeneity on predictive performance

We trained the models on the three selected COVID-19 efficiency studies (the same as Guo et al.[19]), all COVID-19 topics, COVID-19 topics with ES topics, and COVID-19 topics with ES and SYNERGY topics, respectively. The trained models were evaluated on all topics in

the COVID-19 dataset. By comparing the performance of the study screening models trained with different numbers of topics and topics from various domains, we can understand whether adding more training data and diverse topics can improve the performance on a target theme (COVID-19).

To investigate how the data homogeneity affects the model performance, we calculated a measure of textual similarity between the selected COVID-19 topics and other datasets. Firstly, the topics in each dataset were embedded with Sentence-BERT.[24] The average of the selected topic embeddings in the COVID-19 dataset was considered the centroid of the dataset. Then, we compared the cosine similarity between the centroid and other topics. The similarity between the selected COVID-19 topics and other datasets was calculated as follows.

$$Similarity = \frac{1}{n} \sum_{i=1}^{n} CosineSimilarity(\mathrm{C}, \mathrm{T}_i) \tag{3}$$

Where n is the number of topics in a dataset, C is the centroid of the 3 selected COVID-19 topics, and T is the topic embedding.

3.5 Evaluating the models performance on unseen topics: A Leave-One-Topic-Out analysis

The model's ability to generalise to topics not used in training was assessed using the seven topics from the COVID-19 dataset. We used a leave-one-topic-out cross validation approach, where the dataset is split by topic. For each cross-validation iteration, six topics are used to fine-tune BlueBERT and the trained model was evaluated on the remaining topic. If the trained models achieved good performance among target datasets, this would indicate that M-PreSS has a good generalisation of COVID-19 efficacy studies and could be used for new reviews in this area.

3.6 Evaluation

We evaluated our models trained with various strategies on the COVID-19 dataset. Table 2 gives an overview of the topics included in the dataset. Seven systematic reviews were included in the dataset; most of them have fewer than 10% of the studies labelled as 'included', hence, imbalanced.

Table 2: Overview of the COVID-19 dataset. A lower inclusion percentage indicates greater imbalance within the topic.

Abbreviation	Topic	Inclusion Count	Exclusion Count	Inclusion Percentage
NMABS	Clinical efficacy and safety of SARS-CoV-2-neutralizing monoclonal antibody in patients with COVID-19: A living systematic review and meta-analysis	297	2, 801	9.59%
CSR	Efficacy and safety of corticosteroid regimens for the treatment of hospitalized COVID-19 patients: a meta-analysis	140	8, 865	1.55%
IVM	Efficacy and safety of ivermectin for the treatment of COVID-19: a systematic review and meta-analysis	35	279	11.15%
SSRI	Efficacy and safety of selective serotonin reuptake inhibitors in COVID-19 management: a systematic review and meta-analysis	29	3, 989	0.72%
CQHCQ	Efficacy of chloroquine and hydroxychloroquine for the treatment of hospitalized COVID-19 patients: a meta-analysis	311	8, 148	3.68%
LPVR	Efficacy of lopinavir-ritonavir combination therapy for the treatment of hospitalized COVID-19 patients: a meta-analysis	91	1, 456	5.88%
PMH	Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis	422	13, 280	3.08%

Due to the imbalance of the dataset, using accuracy metric can be misleading because it does not sufficiently reflect the model's effectiveness across all classes. In our datasets, where the number of included studies significantly outnumbers the exclusions, a high accuracy score can be achieved by predicting the exclusion class for all instances. Therefore, we evaluated different trained study screening models with PRAUC (Area under the Precision-Recall Curve), which focuses on the performance relative to the inclusion (minority) class. A perfect prediction would have a PRAUC equal to 1. The PRAUC of a random model would equal the inclusion percentage, meaning that a model with a PRAUC greater than the inclusion percentage can discriminate between inclusion and exclusion to some extent.

To further evaluate the performance of our model, we compared our trained COVID-19 study screening models with the results reported by Guo et al. In Guo et al.'s study, the GPT model was required to give a decision on inclusion/exclusion for a candidate study. Therefore, we applied a binary threshold to the COVID-19 model for comparison. The thresholds were calculated respectively for each review topic using the best FPR at 100% recall strategy (explained in the

 Section).

When evaluating the performance of inclusion/exclusion prediction, we calculated the recall (equation 1), FPR (equation 2), and true negative rate (TNR, also known as specificity) as follows.

$$TNR = \frac{TN}{TN + FP} \tag{4}$$

Recall indicates how well the model captured all the included studies, which is a key metric for study screening. FPR is the percentage of exclusion examples predicted as inclusion by the model, which indicates the extra irrelevant studies needed to review for the reviewers. TNR (also known as specificity) indicates how well the model correctly identifies exclusion examples from all actual excluded studies.

4 Results and Discussions

4.1 Exploring the impact of including selection criteria in topic descriptions

Figure 2 compares the performance of the study screening models trained with and without criteria. As shown in the figure, the PRAUC of all the models is better than the baseline, which is a random classifier. Furthermore, the model trained with criteria achieved a comparable or better performance than the model without criteria, for all except one topic (CSR). This shows that extra contextual information provided to the model is beneficial for distinguishing included references from excluded ones. Therefore, criteria were added to the inputs during training for all the models in this study.

Our model trained with criteria (free text) achieved an average PRAUC of 0.61 among the COVID-19 review topics and had the best performance of 0.84 PRAUC on IVM. Furthermore, the PRAUC among all topics was higher than the inclusion percentages, which shows that our model is better than a random classifier and can discriminate between inclusion and exclusion studies.

One limitation of using BlueBERT is that input sequence length has a limitation of 512 tokens, typically representing around 350400 words. For any input that is more than 512 tokens, the model will truncate the input text. Therefore, using Siamese architecture can separate the selection criteria and study information into two identical language models, hence minimising

Figure 2: Comparing the models trained with and without selection criteria on the COVID-19 dataset. The criteria were manually extracted from the published systematic reviews. The models were trained on the training set and evaluated on the test set. The baseline for PRAUC were the inclusion percentage.

the loss of information from the review topics and studies.

When pre-training the study selection model with a larger number of topics, the free text criteria are more accessible from the published paper than manually curated criteria. However, to further understand the role of selection criteria, we compared the free text criteria with other curated criteria.

The results, provided in Table 3, show that free text achieved the highest PRAUC among four out of seven topics, followed by structured criteria (three out of seven) and inclusion/exclusion criteria (two out of seven). The structured and inclusion/exclusion criteria had a comparable or higher recall for all topics than free text. Structured criteria achieved better recall on SSRI, LPVR, and PMH, whereas inclusion/exclusion criteria outperformed structured criteria on NM-BAS, CSR, and CQHCQ. Nevertheless, the structured criteria performed better in terms of average recall and average FPR, which shows that the model (with our choice of classification threshold) is better at including all the relevant studies while adding fewer irrelevant ones.

In Table 4, we compared the model trained with inclusion/exclusion criteria with Guo et al.'s results using ChatGPT/GPT-4. The thresholds that achieved the lowest FPR at 100% recall

Table 3: Performance of the models trained with different types of criteria. The criteria for each topic can be found at ??. Free text criteria are the study selection description extracted from the published systematic review. Structured criteria are the curated selection objectives, which include participants, interventions, comparators, outcomes, and study designs. Inclusion/Exclusion criteria are the criteria curated in an inclusion and exclusion format.

Topic	Free Text			Structured			Inclusion/Exclusion		
	PRAUC	Recall	FPR(%)	PRAUC	Recall	FPR(%)	PRAUC	Recall	FPR(%)
NMABS	0.80	0.93	10.16	0.74	0.95	11.41	0.76	0.97	13.55
CSR	0.53	0.86	1.24	0.77	0.86	2.31	0.53	0.93	6.77
IVM	0.84	1.00	19.64	0.59	1.00	7.14	0.80	1.00	3.57
SSRI	0.29	0.67	0.88	0.33	0.67	1.38	0.46	0.50	0.63
CQHCQ	0.60	0.79	5.40	0.60	0.84	5.58	0.51	0.86	8.65
LPVR	0.48	0.63	5.82	0.45	0.74	10.62	0.47	0.68	7.53
PMH	0.69	0.89	6.40	0.70	0.95	6.93	0.70	0.91	4.97
Average	0.61	0.83	7.08	0.60	0.86	6.48	0.60	0.83	6.52

were applied to obtain inclusion/exclusion predictions. Note that Guo et al. evaluated the GPT model on the entire study screening dataset without training or fine-tuning the model. However, by fine-tuning BlueBERT with 80% screening records from the COVID-19 dataset, we evaluated our model only on the sampled data (i.e. test set) from the original dataset.

Table 4: Model performance compared to Guo et al.[19] We trained the COVID-19 model with the training set of all seven topics from the COVID-19 datasets. The inclusion/exclusion selection criteria were included in the input during the training. A threshold for each topic was calculated to achieve the lowest FPR at 100% recall in the training set. Note that Guo et al. evaluated the performance with all the records in each topic. However, we trained our model with 80% of the records. Hence, our models were evaluated on the test set (20% of the records) for each topic.

Topic	Guo et al.			Ours: COVID-19 Model (best FPR at 100% recall)		
	Recall	TNR(%)	FPR (%)	Recall	TNR(%)	FPR(%)
IVM	0.69	75.63	24.37	1.00	96.43	3.57
SSRI	0.97	94.91	5.09	0.50	99.37	0.63
LPVR	0.59	86.20	13.80	0.68	92.47	7.53

Although BlueBERT is a smaller language model compared to the GPT models, the results in Table 4 show that fine-tuning BlueBERT with study screening data can obtain comparable results to a large general model, such as ChatGPT, on the three topics reported in Guo et al.'s study. For IVM and LPVR, our model managed to identify more relevant studies without including additional false positive studies compared to Guo et al. 's results from GPT-4. Moreover, our model achieved 100% recall on IVM, which means it managed to identify all the relevant studies from the candidates.

4.2 Exploring the impact of differences in dataset size and topic homogeneity on predictive performance

The evaluation results of training on different datasets, with different dataset sizes and topic homogeneity, are illustrated in Figure 3. To maintain consistency between datasets, we applied free text criteria for all the topics in this experiment. The models were trained with training sets from specific topics and evaluated on the test sets from the COVID-19 dataset. Following Guo et al.'s study, the selected COVID-19 model was trained on IVM, SSRI, and LPVR. Thus, we did not evaluate the selected COVID-19 topics model on other unseen topics.

Figure 3: Comparing the performance of models trained on various data sizes (see Table 1). The models were trained on the COVID-19 training set and evaluated on the test set. We measured the cosine similarity between selected COVID-19 topics and other topics for training the models. The selected COVID-19 topics model was only trained on IVM, SSRI, AND LPVR. Therefore, we only evaluated it on these three topics.

The results in Figure 3 show that the model (orange) trained with all COVID-19 topics outperformed the model (blue) trained with selected topics on the IVM and LPVR test set, showing that adding in domain data can improve the model performance. However, the model performance decreased on SSRI when more COVID-19 topics were added during training. This may be because that free text criteria resulted in lower performance than other criteria formats for SSRI but performs better for IVM and LPVR (Table 3).

We quantified the data homogeneity by calculating the cosine similarity between the selected

COVID-19 and added topics. The similarity dropped from 0.8 to 0.55 and 0.48 when adding ES and SYNERGY datasets. The improvements were inconsistent among all the topics when increasing the size of study screening data from a broader but related domain (e.g. ES and SYNERGY dataset), indicating that the homogeneity of the data matters during scaling.

Furthermore, we compared our model trained on three selected COVID-19 topics with the results from Guo et al. using the same criteria. Similar to the COVID-19 model, we used the best FPR at 100% recall threshold to give a binary prediction for each candidate study. With limited topics, our model still improved the recall for IVM and LPVR (shown in Table 5), demonstrating that fine-tuning with study screening datasets is an effective approach for researchers with constrained computational resources to improve the performance of general LLMs.

Table 5: Performance of the model trained with three selected COVID-19 topics compared to the results reported by Guo et al.[19] We used the same topics and inclusion/exclusion criteria as Guo et al. for training the model.

Topic	Guo et al.			Selected COVID-19 Model (best FPR at 100% recall)		
	Recall	TNR(%)	FPR (%)	Recall	TNR(%)	FPR(%)
IVM	0.69	75.63	24.37	1.00	92.86	7.14
SSRI	0.97	94.91	5.09	0.17	99.12	0.88
LPVR	0.59	86.20	13.80	0.79	86.64	13.36

4.3 Evaluating the models performance on unseen topics: A Leave-One-Topic-Out analysis

The results so far demonstrated in this study show that our model has a good generalisation among different topics, for the set of topics used in model training. Another advantage of utilising an LLM is that the model performance can be generalised and transferred to unseen data or topics after training. Therefore, we performed a leave-one-topic-out validation for all COVID-19 topics to evaluate the transferability of knowledge learned from given study screening records to a new topic. The results are illustrated in figure 4.

The results show that the model performance on unseen topics decreased compared to the results trained with the entire COVID-19 dataset (see free text results in Table 3), as predicting without showing the model examples from the target topic was more challenging. However, the model still discriminated included studies from exclusions to some extent, achieving a larger PRAUC than the topic baseline (i.e. inclusion percentage - see Table 2).

Performance of our model decreased least for LPVR (PRAUC 0.38 vs 0.48) and IVM (PRAUC

Figure 4: Precision-Recall curves for leave-one-topic-out validation. The performance on each topic is based on zero-shot learning.

0.64 vs 0.84), better than PMH (PRAUC 0.04 vs 0.69) which had the largest decrease. This may be because these LPVR and IVM contain the least number of records, such that removing these examples from the training data does not dramatically affect the model's performance.

5 Conclusion

In this study, we applied a Siamese architecture to fine-tune BlueBERT for study screening. Our results show that our model has good generalisation among different topics, where these examples from these topics are used in training. This training method can also be applied to other large language model frameworks, such as Llama [1]. Furthermore, adding selection criteria during training can increase prediction performance. The models we trained achieved better recall and lower FPR compared to Guo et al.'s study using ChatGPT on two out of three COVID-19 topics. This shows that even with limited data and a smaller language model, fine-

---

[1]Llama: https://huggingface.co/meta-llama

tuning can achieve comparable or better performance. We experimented with various types of selection criteria and found that the criteria curated in the format of participants, interventions, comparators, outcomes, and study designs achieved the best average recall and FPR among the COVID-19 dataset. We also observed that adding more systematic review topics with high homogeneity to the model can potentially increase the overall performance across topics.

Our method shows potential in performing study screening for systematic review topics where the topics are related to a theme or focus. For example, we applied our method to automate the screening of Global Cancer Update Programme (CUP Global) systematic reviews about cancer incidence topics for a variety of cancer sites,[34] and achieved good results where the recall for all topics are consistently above 90% while the FPRs are under 10% for the majority of the topics. Our developed model can also be used for other study screening processes in a similar setting. In addition, this method could be applied in a more semi-automated manner to assist expert decision-making, where the developed model can be used under different thresholds to achieve multiple inclusion/exclusion strategies, or being combined with more rule-based approaches for weighted scoring.

However, transferring the screening knowledge learned from existing topics to new topics with zero-shot learning remains challenging for our model and BlueBERT. In future research, few-shot learning (labels a small amount of data) or active learning could be applied to fine-tune the pre-trained study screening model for better performance on new review topics. For instance, with active learning, human reviewers can review the top K studies prioritised by the pre-trained model. Then, the decisions could be fed back to tuning the study screening model to improve the accuracy. This process can be repeated until the model achieves a better prediction.

6 Declarations

6.1 List of abbreviations

CQHCQ: Efficacy of chloroquine and hydroxychloroquine for the treatment of hospitalized COVID-19 patients: a meta-analysis

CUP Global: Global Cancer Update Programme

ES: Effectiveness studies

IVM: Efficacy and safety of ivermectin for the treatment of COVID-19: a systematic review and

meta-analysis

FPR: False positive rate

LLM: Large Language Model

LPVR: Efficacy of lopinavir-ritonavir combination therapy for the treatment of hospitalized COVID-19 patients: a meta-analysis

ML: Machine learning

NLP: Natural language processing

NMABS: Clinical efficacy and safety of SARS-CoV-2-neutralizing monoclonal antibody in patients with COVID-19: A living systematic review and meta-analysis

PRAUC: Area under the Precision-Recall curve

SSRI: Efficacy and safety of selective serotonin reuptake inhibitors in COVID-19 management: a systematic review and meta-analysis

TNR: True negative rate

WCRF: World Cancer Research Fund

6.2   Data and Code Availability

The code will be available on GitHub (https://github.com/automation-in-systematic-reviews/M-PreSS) when published. The open dataset can be found at:

- Guo et al.: https://data.mendeley.com/datasets/np79tmhkh5/1

- SYNERGY: https://github.com/asreview/synergy-dataset

6.3   Consent for publication

Not applicable.

6.4   Competing Interests

The authors declare that they have no competing interests.

6.5   Acknowledgements

yse and judge global research on how diet, nutrition, physical activity and body weight affect cancer risk and survival. Findings and evidence judgements are used to develop public health recommendations for cancer prevention and recommendations for future research.

6.6 Funding

Funding for this work (Automating WCRF/AICR systematic review processes, CUP_2021_002) was obtained from World Cancer Research Fund (WCRF), as part of World Cancer Research Fund Internationals Global Cancer Update Programme. This research was carried out in the Medical Research Council Integrative Epidemiology Unit (MC_UU_00032/3).

RMM is a National Institute for Health Research Senior Investigator (NIHR202411). RMM is supported by a Cancer Research UK 25 (C18281/A29019) programme grant (the Integrative Cancer Epidemiology Programme). RMM is also supported by the NIHR Bristol Biomedical Research Centre which is funded by the NIHR (BRC-1215-20011) and is a partnership between University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. Department of Health and Social Care disclaimer: The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

6.7 Author's Contributions

Zhaozhen Xu: Conceptualization, Methodology, Investigation, Software, Visualization, Writing - Original draft; Philippa Davies: Resources, Investigation, Writing - review & editing; Louise AC Millard: Conceptualization, Methodology, Writing - review & editing; Lam Teng: Conceptualization, Writing - review & editing; Georgios Markozannes: Conceptualization, Writing - review & editing; Eduardo AP Seleiro: Conceptualization, Writing - review & editing; Julian PT Higgins: Conceptualization, Resources, Writing - review & editing; Richard M Martin: Funding acquisition, Conceptualization, Supervision, Writing - review & editing; Konstantinos K Tsilidis: Conceptualization, Writing - review & editing; Doris SM Chan: Conceptualization, Writing - review & editing; Tom R Gaunt: Funding acquisition, Conceptualization, Methodology, Supervision, Writing - review & editing; Yi Liu: Funding acquisition, Conceptualization, Methodology, Project administration, Supervision, Writing - review & editing.

References

[1] Bromley, J.; Guyon, I.; LeCun, Y.; Säckinger, E.; Shah, R. Advances in neural information processing systems 1993, 6.

[2] Beller, E.; Clark, J.; Tsafnat, G.; Adams, C.; Diehl, H.; Lund, H.; Ouzzani, M.; Thayer, K.; Thomas, J.; Turner, T.; others Systematic reviews 2018, 7, 1–7.

[3] Lefebvre, C.; Glanville, J.; Briscoe, S.; Featherstone, R.; Littlewood, A.; Metzendorf, M.-I.; Noel-Storr, A.; Paynter, R.; Rader, T.; Thomas, J.; Wieland, L. S. In Cochrane Handbook for Systematic Reviews of Interventions version 6.5; Higgins, J. P., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., Welch, V. A., Eds.; Cochrane, 2024.

[4] Haddaway, N. R.; Westgate, M. J. Conservation Biology 2019, 33, 434–443.

[5] Ouzzani, M.; Hammady, H.; Fedorowicz, Z.; Elmagarmid, A. RayyanA web and mobile app for systematic reviews. Systematic Reviews, 5 (1), Article 210. 2016.

[6] Van De Schoot, R.; De Bruin, J.; Schram, R.; Zahedi, P.; De Boer, J.; Weijdema, F.; Kramer, B.; Huijts, M.; Hoogerwerf, M.; Ferdinands, G.; others Nature machine intelligence 2021, 3, 125–133.

[7] Kebede, M. M.; Le Cornet, C.; Fortner, R. T. Research Synthesis Methods 2023, 14, 156–172.

[8] Devlin, J.; Chang, M.-W.; Lee, K.; Toutanova, K. BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding. Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, Volume 1 (Long and Short Papers). 2019; pp 4171–4186.

[9] Radford, A.; Wu, J.; Child, R.; Luan, D.; Amodei, D.; Sutskever, I.; others OpenAI blog 2019, 1, 9.

[10] Brown, T.; Mann, B.; Ryder, N.; Subbiah, M.; Kaplan, J. D.; Dhariwal, P.; Neelakantan, A.; Shyam, P.; Sastry, G.; Askell, A.; others Advances in neural information processing systems 2020, 33, 1877–1901.

[11] Ma, X.; Wang, L.; Yang, N.; Wei, F.; Lin, J. Fine-tuning llama for multi-stage text retrieval. Proceedings of the 47th International ACM SIGIR Conference on Research and Development in Information Retrieval. 2024; pp 2421–2425.

[12] Xu, Z.; Cristianini, N. QBERT: Generalist model for processing questions. International symposium on intelligent data analysis. 2023; pp 472–483.

[13] Liu, Y.; Elsworth, B. L.; Gaunt, T. R. Bioinformatics 2023, 39, btad169.

[14] Liu, Y.; Gaunt, T. R. Bioinformatics 2024, 40, btae519.

[15] Luo, X.; Chen, F.; Zhu, D.; Wang, L.; Wang, Z.; Liu, H.; Lyu, M.; Wang, Y.; Wang, Q.; Chen, Y. Journal of Medical Internet Research 2024, 26, e56780.

[16] Moreno-Garcia, C. F.; Jayne, C.; Elyan, E.; Aceves-Martins, M. Decision Analytics Journal 2023, 6, 100162.

[17] Issaiy, M.; Ghanaati, H.; Kolahi, S.; Shakiba, M.; Jalali, A. H.; Zarei, D.; Kazemian, S.; Avanaki, M. A.; Firouznia, K. BMC Medical Research Methodology 2024, 24, 78.

[18] Akinseloyin, O.; Jiang, X.; Palade, V. Journal of the American Medical Informatics Association 2024, 31, 1939–1952.

[19] Guo, E.; Gupta, M.; Deng, J.; Park, Y.-J.; Paget, M.; Naugler, C. Journal of Medical Internet Research 2024, 26, e48996.

[20] Peng, Y.; Yan, S.; Lu, Z. Transfer Learning in Biomedical Natural Language Processing: An Evaluation of BERT and ELMo on Ten Benchmarking Datasets. Proceedings of the 2019 Workshop on Biomedical Natural Language Processing (BioNLP 2019). 2019; pp 58–65.

[21] Hasny, M.; Vasile, A.-P.; Gianni, M.; Bannach-Brown, A.; Nasser, M.; Mackay, M.; Donovan, D.; Šorli, J.; Domocos, I.; Dulloo, M.; others BERT for complex systematic review screening to support the future of medical research. International Conference on Artificial Intelligence in Medicine. 2023; pp 173–182.

[22] Sun, D.; Wu, Z.; Wang, Y.; Lv, Q.; Hu, B. Risk Prediction for Imbalanced Data in Cyber Security : A Siamese Network-based Deep Learning Classification Framework. 2019 International Joint Conference on Neural Networks (IJCNN). 2019; pp 1–8.

[23] Ye, Y.; Simpson, E.; Rodriguez, R. S. arXiv preprint arXiv:2409.15090 2024,

[24] Reimers, N.; Gurevych, I. Sentence-bert: Sentence embeddings using siamese bert-networks. 2019; Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing and the 9th International Joint Conference on Natural Language Processing (EMNLP-IJCNLP).

[25] Pickard, A.; Davies, P.; Birnie, K.; Beringer, R. British journal of anaesthesia 2014, 112, 982–990.

[26] López-López, J. A.; Sterne, J. A.; Thom, H. H.; Higgins, J. P.; Hingorani, A. D.; Okoli, G. N.; Davies, P. A.; Bodalia, P. N.; Bryden, P. A.; Welton, N. J.; others bmj 2017, 359.

[27] Sterne, J. A.; Bodalia, P. N.; Bryden, P. A.; Davies, P. A.; López-López, J. A.; Okoli, G. N.; Thom, H. H.; Caldwell, D. M.; Dias, S.; Eaton, D.; others Health Technology Assessment Volume 2017, 21.

[28] Ijaz, S.; Davies, P.; Kessler, D.; Lewis, G.; Wiles, N.; others Cochrane database of systematic reviews 2018,

[29] Jackson, J.; Davies, P.; Leggett, N.; Nugawela, M.; Scott, L.; Leach, V.; Richards, A.; Blacker, A.; Abrams, P.; Sharma, J.; others BJS open 2019, 3, 11–23.

[30] Davies, P.; Ijaz, S.; Kessler, D.; Lewis, G.; Wiles, N.; others Cochrane Database of Systematic Reviews 2019,

[31] de Araújo Pio, C. S.; Chaves, G. S.; Davies, P.; Taylor, R. S.; Grace, S. L. Cochrane Database of Systematic Reviews 2019,

[32] De Bruin, J.; Ma, Y.; Ferdinands, G.; Teijema, J.; Van de Schoot, R. SYNERGY - Open machine learning dataset on study selection in systematic reviews. 2023; https://doi.org/10.34894/HE6NAQ.

[33] Loshchilov, I.; Hutter, F. Decoupled Weight Decay Regularization. International Conference on Learning Representations.

[34] Xu, Z.; Teng, L.; Millard, L.; Higgins, J. P.; Martin, R. M.; Markozannes, G.; Tsilidis, K. K.; Chan, D. S.; Gaunt, T. R.; Liu, Y. Automating Screening of Studies for Systematic Reviews

Using a Large Language Model. Abstracts accepted for the 2nd Global Evidence Summit 2024. Prague, Czech Republic.

Supplementary Materials

Table 6: Input and Output Examples for M-PreSS

Systematic Review Topic	Candidate Study	Priority Score
Query: Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis	Title: COVID-19 and Adolescent Depression and Suicide Risk Screening Outcomes Abstract: BACKGROUND: Mental health concerns increased during the coronavirus disease 2019 pandemic, but previous studies have not examined depression screening in pediatric primary care. We aimed to describe changes in screening, depressive symptoms, and suicide risk among adolescents during the coronavirus disease 2019 pandemic. METHODS: In a repeat cross-sectional analysis of electronic health record data from a large pediatric primary care network, we compared the percentage of primary care visits where adolescents aged 12 to 21 were screened for depression, screened positive for depressive symptoms, or screened positive for suicide risk between June and December 2019 (prepandemic) and June and December 2020 (pandemic). Changes were examined overall, by month, and by sex, race and ethnicity, insurance type, and income. Modified Poisson regression was used to calculate prevalence ratios (PRs) for the prepandemic to pandemic changes. RESULTS: Depression screening at primary care visits declined from 77.6% to 75.8% during the pandemic period (PR: 0.98, 95% confidence interval [CI]: 0.90-1.06). The percentage of adolescents screening positive for depressive symptoms increased from 5.0% to 6.2% (PR: 1.24, 95% CI: 1.15-1.34), with greater increases among female, non-Hispanic Black, and non-Hispanic white adolescents. Positive suicide risk screens increased from 6.1% to 7.1% (PR: 1.16, 95% CI: 1.08-1.26), with a 34% relative increase in reporting recent suicidal thoughts among female adolescents (PR: 1.34, 95% CI: 1.18-1.52). CONCLUSIONS: Results suggest that depression and suicide concerns have increased during the pandemic, especially among female adolescents. Results underscore the importance of consistent depression and suicidality screening.	0.4994128
Query: Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis	Title: Hopes and fears: emerging adults in mood and anxiety disorder treatment predict outcomes of the COVID-19 pandemic Abstract: Active patients at The First Episode Mood and Anxiety Program (FEMAP) were asked to participate in a study monitoring their symptoms during early COVID-19 closures. Participants were asked to predict both possible positive and feared outcomes of the pandemic. Responses were analyzed with a descriptive qualitative approach using thematic analysis. Primary feared outcomes included death, prolonged pandemic impacts, and concerns about employment, mental health, and education. Prevalent positive predictions included a new perspective, improved relationships, personal growth, and environmental improvement. These findings suggest the complex relationship between fear and hope for young people in the early stages of the pandemic.	0.4899408

Table 7: Topics Overview. Topics used for training M-PreSS.

Dataset	Topic
SYNERGY	A meta-analysis on the role older adults with cancer favour in treatment decision making
SYNERGY	A Systematic Review Comparing Experimental Design of Animal and Human Methotrexate Efficacy Studies for Rheumatoid Arthritis: Lessons for the Translational Value of Animal Studies
SYNERGY	Bayesian PTSD-Trajectory Analysis with Informed Priors Based on a Systematic Literature Search and Expert Elicitation
SYNERGY	Cerebral small vessel disease and the risk of dementia: A systematic review and meta_analysis of population_based evidence
SYNERGY	Comparability of Objective Structured Clinical Examinations (OSCEs) and Written Tests for Assessing Medical School Students Competencies: A Scoping Review
SYNERGY	Comparative effectiveness of common therapies for Wilson disease: A systematic review and meta_analysis of controlled studies
SYNERGY	Comparative efficacy and safety of long-acting oral opioids for chronic non-cancer pain: a systematic review
SYNERGY	Comparative efficacy and safety of skeletal muscle relaxants for spasticity and musculoskeletal conditions: a systematic review
SYNERGY	Coronary heart disease, heart failure, and the risk of dementia: A systematic review and meta_analysis
SYNERGY	Cross_sectional relation of long_term glucocorticoids in hair with anthropometric measurements and their possible determinants: A systematic review and meta_analysis
SYNERGY	Efficacy of therapeutic plasma exchange for treatment of autoimmune hemolytic anemia: A systematic review and meta-analysis of randomized controlled trials
SYNERGY	Fragility Analysis of Statistically Significant Outcomes of Randomized Control Trials in Spine Surgery
SYNERGY	Human and animal evidence of potential transgenerational inheritance of health effects: An evidence map and state-of-the-science evaluation
SYNERGY	Long-term Outcomes of Cognitive Behavioral Therapy for Anxiety-Related Disorders
SYNERGY	Patients Retransitioning from Biosimilar TNF_ Inhibitor to the Corresponding Originator After Initial Transitioning to the Biosimilar: A Systematic Review
SYNERGY	Pharmacokinetics and Associated Efficacy of Emicizumab in Humans: A Systematic Review
SYNERGY	Poor nutritional condition promotes high_risk behaviours: a systematic review and meta_analysis
SYNERGY	Postmenopausal Hormone Replacement Therapy
SYNERGY	Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis
SYNERGY	Psychological theories of depressive relapse and recurrence: A systematic review and meta-analysis of prospective studies
SYNERGY	Sleep and Microdialysis: An Experiment and a Systematic Review of Histamine and Several Amino Acids
SYNERGY	Specialized psychotherapies for adults with borderline personality disorder: A systematic review and meta-analysis
SYNERGY	Systematic Review and Meta-Analysis of Early-Life Exposure to Bisphenol A and Obesity-Related Outcomes in Rodents
SYNERGY	The methodological rigour of systematic reviews in environmental health
SYNERGY	The rodent object-in-context task: A systematic review and meta-analysis of important variables
SYNERGY	Does the Source of Mesenchymal Stem Cell Have an Effect in the Management of Osteoarthritis of the Knee? Meta-Analysis of Randomized Controlled Trials
ES	Comparative efficacy of adjuvant non-opioid analgesia in adult cardiac surgical patients: A network meta-analysis
ES	The use of acupuncture in patients with Raynaud_ syndrome: A systematic review and meta-analysis of randomized controlled trials
ES	Promoting patient uptake and adherence in cardiac rehabilitation
ES	Oral anticoagulants for prevention of stroke in atrial fibrillation: systematic review, network meta-analysis, and cost effectiveness analysis

Dataset (continued)	Topic (continued)
ES	Oral anticoagulants for primary prevention, treatment and secondary prevention of venous thromboembolic disease
ES	Systematic review and meta-analysis of the effect of intraoperative α_2-adrenergic agonists on postoperative behaviour in children
ES	Systematic review of interventions for the prevention and treatment of postoperative urinary retention
ES	Psychological or pharmacological therapies for treatment-resistant depression in adults
COVID-19	Clinical efficacy and safety of SARS-CoV-2-neutralizing monoclonal antibody in patients with COVID-19: A living systematic review and meta-analysis
COVID-19	Efficacy and safety of corticosteroid regimens for the treatment of hospitalized COVID-19 patients: a meta-analysis
COVID-19	Efficacy and safety of ivermectin for the treatment of COVID-19: a systematic review and meta-analysis
COVID-19	Efficacy and safety of selective serotonin reuptake inhibitors in COVID-19 management: a systematic review and meta-analysis
COVID-19	Efficacy of chloroquine and hydroxychloroquine for the treatment of hospitalized COVID-19 patients: a meta-analysis
COVID-19	Efficacy of lopinavir_itonavir combination therapy for the treatment of hospitalized COVID-19 patients: a meta-analysis

Table 8: Criteria for COVID-19 topics.

Topic	Free Text Criteria	Structured Criteria	Inclusion/Exclusion Criteria
Clinical efficacy and safety of SARS-CoV-2-neutralizing monoclonal antibody in patients with COVID-19: A living systematic review and meta-analysis	Articles that met the following requirements were included in the study: 1) patients were adults with COVID-19; 2) interventions using neutralizing mAbs; 3) the comparator was a placebo or standard of care treatment; and 4) outcomes included primary or secondary outcomes (the primary outcomes included all-cause mortality at 28 days and hospitalization or hospital visit. Clinical recovery, ICU admission, mechanical ventilation, hospital discharge, and serious adverse events were classified as secondary outcomes.); 5) The study was a RCT. Only English and Korean studies were included in this meta-analysis. Two review authors independently and in duplicate evaluated publications for inclusion based on title and abstract and then reviewed relevant full-text articles. Disagreements during the review process were addressed by consensus, with the involvement of a third review author.	Randomised controlled trials Compared interventions using neutralizing monoclonal antibodies (Etesevimab, bamlanivimab, casirivimab, imdevimab, casirivimab, imdevimab, bamlanivimab, etesevimab, sotrovimab, bebtelovimab, tixagevimab, cilgavimab) with a placebo or standard of care treatment Included adults with COVID-19 Reported any of the following outcomes: all-cause mortality at 28 days and hospitalization or hospital visit. Clinical recovery, ICU admission, mechanical ventilation, hospital discharge, and serious adverse events Articles in English or Korean	Inclusion - Randomised controlled trials - Compared interventions using neutralizing monoclonal antibodies with a placebo or standard of care treatment - Included adults with COVID-19 - Reported any of the following outcomes: all-cause mortality at 28 days and hospitalization or hospital visit. Clinical recovery, ICU admission, mechanical ventilation, hospital discharge, and serious adverse events - Articles in English or Korean
Efficacy and safety of corticosteroid regimens for the treatment of hospitalized COVID-19 patients: a meta-analysis	We included both randomized and non-randomized comparative studies that met the following inclusion criteria: compared any corticosteroid therapy with standard of care, or compared any corticosteroid therapy with an adjuvant therapy to adjuvant therapy alone, and included hospitalized COVID-19 patients. We only included studies using adjuvant therapies if the adjuvant therapy between the control and experimental arms are the same in order to minimize the effect of the adjuvant therapies on treatment outcomes.	Randomised and non-randomised comparative studies Compared any corticosteroid therapy with standard of care, or compared any corticosteroid therapy with an adjuvant therapy to adjuvant therapy alone. We only included studies using adjuvant therapies if the adjuvant therapy between the control and experimental arms are the same in order to minimize the effect of the adjuvant therapies on treatment outcomes. Included hospitalized COVID-19 patients Reported any of the following outcomes: time to negative conversion of SARS-CoV-2 tests, overall length of hospitalization, length of ICU hospitalization, incidence of progression to ICU hospitalization, incidence of mortality, incidence of progression to mechanical ventilation, length of mechanical ventilation, incidence of all-cause adverse events, incidence of investigator-defined severe adverse events, incidence of hyperglycemia or corticosteroid-induced diabetes mellitus and incidence of nosocomial bacterial or fungal infections.	Inclusion - Randomised and non-randomised comparative studies - Compared any corticosteroid therapy with standard of care, or compared any corticosteroid therapy with an adjuvant therapy to adjuvant therapy alone. - Included hospitalized COVID-19 patients - Reported any of the following outcomes: time to negative conversion of SARS-CoV-2 tests, overall length of hospitalization, length of ICU hospitalization, incidence of progression to ICU hospitalization, incidence of mortality, incidence of progression to mechanical ventilation, length of mechanical ventilation, incidence of all-cause adverse events, incidence of investigator-defined severe adverse events, incidence of hyperglycemia or corticosteroid-induced diabetes mellitus and incidence of nosocomial bacterial or fungal infections. Exclusion - Studies that used different adjuvant therapies for its intervention and control arms
Efficacy and safety of ivermectin for the treatment of COVID-19: a systematic review and meta-analysis	We included both randomized and non-randomized comparative studies that met the following criteria: (i) compared ivermectin to standard of care or a control group receiving placebo; (ii) included adult COVID-19 inpatients and/or outpatients; and (iii) reported any of our outcomes of interest.	Randomized controlled trials and non-randomized comparative cohort studies Compared ivermectin against standard of care or control group/placebo Included adult (age >=18) COVID-19 inpatients and/or outpatients. Studies that used ivermectin for prophylaxis of COVID-19 were excluded. Reported any of the following outcomes: time to viral clearance, duration of hospitalization, mortality incidence, incidence of progression to mechanical ventilation, all-cause adverse events, incidence of investigator-defined serious adverse events Peer-reviewed articles only	Inclusion - Randomized controlled trials and non-randomized comparative cohort studies - Compared ivermectin against standard of care or control group/placebo - Included adult (age >=18) COVID-19 inpatients and/or outpatients - Reported any of the following outcomes: time to viral clearance, duration of hospitalization, mortality incidence, incidence of progression to mechanical ventilation, all-cause adverse events, incidence of investigator-defined serious adverse events Exclusion - Studies that used ivermectin for prophylaxis of COVID-19 - Non-peer-reviewed articles

Topic (continued)	Free Text Criteria (continued)	Structured Criteria (continued)	Inclusion/Exclusion Criteria (continued)
Efficacy and safety of selective serotonin reuptake inhibitors in COVID-19 management: a systematic review and meta-analysis	We included RCTs and prospective or retrospective observational studies that included inpatients and/or outpatients of any age with COVID-19 and compared the use of SSRIs to the use of the standard of care or placebo. Studies assessing patients with past or ongoing SSRI prescriptions at the time of diagnosis were specifically excluded. We did not impose any language restrictions.	Randomized controlled trials and non-randomized comparative cohort studies. Included COVID-19 inpatients and/or outpatients of any age. Compared the use of selective serotonin reuptake inhibitors (mainly fluvoxamine, fluoxetine, and citalopram/escitalopram) against standard of care or placebo. Studies assessing patients with past or ongoing prescriptions of selective serotonin reuptake inhibitors at the time of COVID-19 diagnosis were excluded. Reported any of the following outcomes: all-cause mortality, hospitalization, composite outcome of emergency room visits or hospitalization, hypoxemia, requirement for supplemental oxygen, requirement for ventilator support, serious adverse events	Inclusion - Randomized controlled trials and non-randomized comparative cohort studies - Included COVID-19 inpatients and/or outpatients of any age - Compared the use of selective serotonin reuptake inhibitors (mainly fluvoxamine, fluoxetine, and citalopram/escitalopram) against standard of care or placebo - Reported any of the following outcomes: all-cause mortality, hospitalization, composite outcome of emergency room visits or hospitalization, hypoxemia, requirement for supplemental oxygen, requirement for ventilator support, serious adverse events Exclusion - Studies assessing patients with past or ongoing prescriptions of selective serotonin reuptake inhibitors at the time of COVID-19 diagnosis
Efficacy of chloroquine and hydroxychloroquine for the treatment of hospitalized COVID-19 patients: a meta-analysis	We included both randomized and non-randomized comparative studies that met the following inclusion criteria in our review: compared hydroxychloroquine or chloroquine as a monotherapy or in combination with azithromycin with standard of care, or compared hydroxychloroquine or chloroquine with adjuvant therapies to adjuvant therapies alone, and included hospitalized COVID-19 patients. For studies involving hydroxychloroquine with adjuvant therapies, we only included studies that used the same adjunctive therapy for both its intervention and control arms to minimize the effect of adjuvant therapies on treatment outcomes, similar to the design of several other meta-analyses [4446].	Randomised and non-randomised comparative studies. Compared hydroxychloroquine or chloroquine as a monotherapy or in combination with azithromycin with standard of care, or compared hydroxychloroquine or chloroquine with adjuvant therapies to adjuvant therapies alone. For studies involving hydroxychloroquine with adjuvant therapies, we only included studies that used the same adjunctive therapy for both its intervention and control . Included hospitalized COVID-19 patients. Reported any of the following outcomes: time to negative conversion of SARS-CoV-2 tests, length of stay, mortality at the latest follow-up, incidence of mechanical ventilation, time to normalization of body temperature, rate of negative conversion of SARS-CoV-2 test at day 7 and day 14 and incidence of severe (or grade 3/4) adverse events.	Inclusion - Randomised and non-randomised comparative studies - Compared hydroxychloroquine or chloroquine as a monotherapy or in combination with azithromycin with standard of care, or compared hydroxychloroquine or chloroquine with adjuvant therapies to adjuvant therapies alone. - Included hospitalized COVID-19 patients - Reported any of the following outcomes: time to negative conversion of SARS-CoV-2 tests, length of stay, mortality at the latest follow-up, incidence of mechanical ventilation, time to normalization of body temperature, rate of negative conversion of SARS-CoV-2 test at day 7 and day 14 and incidence of severe (or grade 3/4) adverse events. Exclusion - Studies that used different adjuvant therapies for its intervention and control arms
Efficacy of lopinavir-Ritonavir combination therapy for the treatment of hospitalized COVID-19 patients: a meta-analysis	We included both RCTs and comparative non-randomized observational studies that satisfied the following inclusion criteria in our analysis: compared LPV/r with standard of care, or compared LPV/r with adjuvant therapies to adjuvant therapies alone, included laboratory-confirmed, hospitalized COVID-19 patients. While we included studies involving LPV/r with adjuvant therapies, we only included studies that used the same concurrent therapy for its intervention and control arms to minimize the effect of adjuvant therapies on treatment outcomes, similar to the design of several other meta-analyses [5658].	Randomized controlled trials and comparative non-randomized observational studies. Compared lopinavir-ritonavir with standard of care, or compared lopinavir-ritonavir with adjuvant therapies to adjuvant therapies alone. Studies involving lopinavir-ritonavir with adjuvant therapies were only included if they used the same concurrent therapy for their intervention and control arms. Included laboratory-confirmed, hospitalized COVID-19 patients	Inclusion - Randomized controlled trials and comparative non-randomized observational studies - Compared lopinavir-ritonavir with standard of care, or compared lopinavir-ritonavir with adjuvant therapies to adjuvant therapies alone - Included laboratory-confirmed, hospitalized COVID-19 patients Exclusion - Studies that used different adjuvant therapies for its intervention and control arms
Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis	Publications were included in the meta-analysis if they met the following criteria: (1) primary crosssectional or longitudinal studies conducted after January 2020 (or reported data from time points after January 2020); (2) included children and adolescents ≤ 18 years of age or students enrolled in primary or secondary educational institutions; and (3) reported the prevalence of depressive symptoms, anxiety symptoms, and/or sleep disturbances, as assessed using selfreported instruments or clinical interviews. Studies specifically targeting children with special needs or preexisting mental health disorders were excluded. No limitations on the publication's language or country of origin were placed.	Primary cross-sectional or longitudinal studies conducted after 2020 (or reported data from time points after January 2020). Included children and adolescents aged 18 years or older or students enrolled in primary or secondary educational institutions. Studies specifically targeting children with special needs or pre-existing mental health disorders were excluded. Reported any of the following outcomes: prevalence of depressive symptoms, anxiety symptoms or sleep disturbances	Inclusion - Primary cross-sectional or longitudinal studies conducted after 2020 (or reported data from time points after January 2020) - Included children and adolescents aged 18 years or older or students enrolled in primary or secondary educational institutions. - Reported any of the following outcomes: prevalence of depressive symptoms, anxiety symptoms or sleep disturbances Exclusion - Studies specifically targeting children with special needs or pre-existing mental health disorders were excluded.