A contextual genomic perspective on physical activity and its relationship to health, well-being and illness

Marco Galimberti[1,2], Daniel F. Levey[1,2], Joseph D. Deak[1,2], Keyrun Adhikari[1,2], Cassie Overstreet[1,2], Priya Gupta[1,2], Rachana Nitin[3], Hang Zhou[1,2], Nicole J. Lake[4], Kelly M. Harrington[5,6], Luc Djousse[6,7], Lea K. Davis[3], VA Million Veteran Program, J. Michael Gaziano[6-8], Murray B. Stein[9,10], Joel Gelernter[1,2,*]

1: Veterans Affairs Connecticut Healthcare System, West Haven, CT, USA
2: Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA
3: Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee, USA
4: Department of Genetics, Yale School of Medicine, New Haven, CT, USA
5: Department of Psychiatry, Boston University School of Medicine, Boston, MA, USA
6: Million Veteran Program (MVP) Coordinating Center, VA Boston Healthcare System, Boston, MA, USA
7: Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA
8: Department of Medicine, Harvard Medical School, Boston, MA, USA
9: VA San Diego Healthcare System, San Diego, CA, USA
10: Department of Psychiatry and School of Public Health, University of California San Diego, La Jolla, CA, USA
*: corresponding author

corresponding author email: joel.gelernter@yale.edu

52   Abstract
53
Physical activity (PA) is one of the most fundamental of all traits in the animal kingdom, has
pervasive health benefits, and is genetically influenced. Using data from the Million Veteran
Program (MVP), we conducted genetic analyses of leisure, work, and home-time PA. For leisure,
for individuals of European (EUR) ancestry, n=189,812 and SNP-based heritability
(h^2)=0.083±0.005; for African ancestry, n=27,044; h^2= 0.034±0.017; and for Latin-American
ancestry, n=10,263; h^2= 0.083±0.036. EUR and cross-ancestry meta-analyses with UK Biobank
identified 67 and 70 lead variants. Leisure-time PA was genetically distinct from PA at home or
work, with the latter two showing less health benefit on pathologies and lifespan. Mendelian
randomization analyses showed protective effects of leisure-time PA on cardiovascular and
respiratory system diseases, metabolic traits, aging, and other traits, and there was protective role
of leisure-time PA against COVID-19 hospitalization (β=-0.067±0.016; p-value=2.8×10^{-5}). These
findings provide new insights into the biology of PA, showing the causal health benefits of leisure-
time PA.

Introduction

Physical activity (PA) has pervasive and profound health benefits, including beneficial influences on chronic conditions like cancer, hypertension, and type 2 diabetes.[1] Very different kinds of PA – for example, weightlifting and aerobic exercise - can lead to a longer and healthier life[2].[2] An individual's level of PA is influenced by genetic factors, with estimated heritability (h^2) between 48% and 71% based on twin studies.[3] To identify genetic variants relevant to PA, genome-wide association studies (GWAS) have been conducted, initially focusing on PA during leisure time.[4-7] Identifying genetic associations with physical activity affords us a unique opportunity; because we are born with genetic variations prior to exposures to psychological, social, and environmental factors that may affect activity levels, we can begin to decompose some of the complex interactions which underlie and potentially confound the study of this important health trait. There is also a wide range of interindividual differences in requirements for physical activity in daily life. Some jobs require much more physical exertion than others; some individuals must exert more effort in their households than others. These kinds of physical activity tend to differ phenomenologically from exercise taken at leisure, occurring more in concentrated bursts in the first two cases, and with the potential to be more sustained in the latter case; yet level of activity in different contexts is often treated as if equivalent. There are, however, prior reports that comparable health benefits may not be assumed for activity at leisure vs. activity in other contexts[8,9], and genetic analyses may help shed light on any differences.

Two powerful previous studies of PA used UK Biobank (UKB) data.[10,11] The first study (N=91,105) identified three variants associated with PA, based on wrist-worn accelerometer data.[11] The second study ($N \leq 377,234$) identified ten significant loci across five phenotypes, three based on self-report data (moderate-to-vigorous PA, vigorous PA, and strenuous sport or other exercise (SSOE) during leisure time), and two on wrist-worn accelerometer data.[10] A multi-ancestry meta-analysis combining data from 703,901 individuals from 51 studies identified 11 associated loci for moderate-to-vigorous intensity PA.[12] Combining this phenotype with two traits phenotypically contrary to PA (leisure screen time and sedentary behaviour at work), 99 associated loci were found summing results from PA (11 variants), leisure screen time (89 variants), and sedentary behaviour at work (4 variants); that is, most loci were mapped for phenotypes correlated to a lack of physical activity.[12]

We sought to maximize power to investigate PA genetics by means of a quantitative trait definition, combining different intensities of PA by summing their frequencies, using data from the Million Veteran Program (MVP). We focused on PA *per se*, as opposed to sedentary behavior. Whereas previous studies considered multicontextual PA -- including data from leisure, at work, and home -- our primary quantitative trait, All-PA-Leisure, focused exclusively on PA during leisure time (see Supplementary Methods for PA context definitions). We studied this trait in two meta-analyses (designated "All-PA-Leisure+SSOE"), one of European (EUR) ancestry and one cross-ancestry (EUR, African (AFR), Latin-American (AMR)), with SSOE data from UKB, also based on information of PA during leisure time (Figure S1, Table S1).[10] This enabled us to investigate, from a genomic perspective, the relationship of PA to a range of health traits. We also investigated PA traits related to the different contexts (home and work, in addition to leisure). In so doing, we greatly increased the number of discovered loci while increasing the diversity of the sample beyond EUR. We investigated protective effects of PA against health traits including type 2 diabetes, gastroesophageal reflux disease, abdominal aortic aneurysm, heart failure, hospitalization caused by COVID-19, osteoarthritis, parental survival, and an aging phenotype. Moreover, we found substantial genetic differences considering PA during the three different contexts (leisure, work, home). These differences are supported by genetic correlation and structural equation modeling analyses.

124 **Results**
125 **GWAS of All-PA-Leisure (MVP) and All-PA-Leisure+SSOE (MVP+UKB) meta-analysis**
126
All-PA-Leisure and SSOE phenotypes
GWAS of All-PA-Leisure in MVP EUR data identified 14 lead SNPs at 13 loci, and GWAS of
SSOE in UKB identified 15 lead SNPs in 14 loci (Figure S2-S3, Table S2-S3). The two most
significant lead SNPs in the GWAS for All-PA-Leisure (EUR, rs761898, *p*-value=1.63×10^{-11};
rs7613360, *p*-value=3.15×10^{-11}) map to intergenic regions on chromosome 3 and 14 (Table S3).
Additional genome-wide significant loci included *SLC39A8**rs13107325 (*p*-value=1.57×10^{-10})
(solute carrier family 39 member 8). As previously described,[10] the most significant variant for
GWAS of SSOE in UKB is *CADM2**rs62253088 (*p*-value=1.0×10^{-19} in this study) (Cell Adhesion
Molecule 2). A sex-stratified analysis of the X chromosome in MVP (All-PA-Leisure) showed no
statistically significant associated variants (Figure S4).
137
SNP heritability and genetic correlation of PA phenotypes
In MVP data for All-PA-Leisure, SNP-based heritability (SNP-h^2) is 0.083±0.005 (standard error
(SE)) in EUR, 0.034±0.017 in AFR, and 0.083±0.036 in AMR (Table S4). SNP-h^2 of SSOE from
UKB is 0.061±0.003. The genetic correlation between All-PA-Leisure and SSOE (r_g=0.76±0.03; *p*-
value=4.7×10^{-118}) was higher than the genetic correlation between All-PA-Leisure and the other
two traits from the previous UKB report,[10] moderate-to-vigorous PA (r_g=0.25±0.04; *p*-
value=4.3×10^{-11}), and vigorous PA (r_g=0.55±0.04; *p*-value=3.9×10^{-53}). Accordingly, we selected the
SSOE phenotype for meta-analyses with MVP data, i.e. the All-PA-Leisure+SSOE meta-analysis
(summary statistics results are described in the Supplementary Text, Figure S2-S3, Table S5-S6).
147
All-PA-Leisure+SSOE meta-analyses
The EUR All-PA-Leisure+SSOE meta-analysis greatly increased discovery yielding 67 lead SNPs
in 55 loci (Figure 1A, Table S5). The cross-ancestry meta-analysis provided 70 lead SNPs in 56
loci (Figure 1B, Table S6). In both, the strongest association was with *CADM2**rs62253088 (EUR
meta-analysis *p*-value=2.0×10^{-21}; cross-ancestry meta-analysis *p*-value=6.1×10^{-20}). The second
strongest associated lead SNP was also on chromosome 3: *MST1R**rs3733134 (EUR meta-analysis
p-value=1.4×10^{-17}; cross-ancestry meta-analysis *p*-value=6.9×10^{-18}).
155
Gene based, gene-set, tissue expression, cell type enrichment analyses of All-PA-Leisure+SSOE
Focusing on the EUR meta-analysis, the gene-based test using MAGMA identified 119 significant
genes from 19,030 protein coding genes (genome-wide significance defined at *p*-value=2.6×10^{-6})
(Table S7). *CADM2* was the most significant gene (*p*-value=3.7×10^{-25}) followed by *NPC1* (*p*-
value=4.8×10^{-16}). Four significant gene ontology (GO) terms resulted from the gene-set analysis:
structural constituent of the presynapse, presynaptic active zone organization, maintenance of
presynaptic active zone structure, and structural constituent of the synapse (Table S8). Tissue
Expression Analysis showed 11 significant enrichments in brain tissues, with the two most
significant represented in the cerebellar hemisphere and in the cerebellum (Figure S5-S6). For gene-
based association analyses and gene-set analyses, similar results were obtained for the cross-
ancestry meta-analysis (Supplementary Text, Table S9-S10, Figure S7-S8).
Cell type enrichment analysis was performed with a collection of human cell types from brain,
blood, and pancreas tissue (all available tissues). In the EUR meta-analysis of All-PA-
Leisure+SSOE, the only significant cell type was *GABA*$^+$ *neurons* from the GSE76381 Linnarsson
Human Midbrain dataset (adjusted *p*-value per dataset after multiple testing correction=2.4×10^{-5}).
In the cross-ancestry meta-analysis, there were no significantly enriched cell types. The results
probably reflect the limited number of kinds of tissue included in this analysis.
173
174

Functional enrichment analysis of All-PA-Leisure+SSOE
Functional enrichment analysis with g:Profiler using significant genes from the gene-based test as
input, identified three significant gene ontology terms shared by the EUR meta-analysis and the
cross-ancestry meta-analysis (Table S11, S12): presynaptic active zone cytoplasmic component,
cell cortex region, and synapse. Also a human phenotype ontology term, polyclonal elevation of
IgM, was shared by the EUR meta-analysis and cross-ancestry meta-analysis.
181
PheWAS of All-PA-Leisure+SSOE
Phenome-wide association study (PheWAS) in BioVu revealed significant association with 159
traits (Figure 2, Table S13). The strongest PheWAS results included negative associations with
diabetes mellitus (odds ratio (OR)=0.868; p-value=5.89×10^{-36}), and chronic airway obstruction
(OR=0.839; p-value=4.22×10^{-32}). We also found significant negative associations for traits also
analyzed in the genetic correlation analyses: coronary artery disease (ischemic heart disease)
(OR=0.892; p-value=1.02×10^{-21}), gastroesophageal reflux disease (OR=0.929; p-value=5.07×10^{-13}),
cancer of bronchus or lung (OR=0.901; p-value=5.47×10^{-7}), and asthma (OR=0.937; p-
value=1.19×10^{-5}). We also found a negative correlation with obesity (OR=0.900; p-value=2.38×10^{-17}),
and several phenotypes related to heart failure: diastolic heart failure (OR=0.880; p-
value=1.76×10^{-9}), systolic or combined heart failure (OR=0.894; p-value=1.96×10^{-9}), and heart
failure (OR=0.902; p-value=1.88×10^{-5}).
194
Genetic correlations: All-PA-Leisure+SSOE and traits of interest
We estimated the genetic correlation of the All-PA-Leisure+SSOE phenotype (EUR meta-analysis)
with traits of interest selected to cover a broad range of health diseases and physiological traits,
using the Benjamini-Hochberg false discovery rate (FDR) correction (Figure 3, Table S14; more
details in *Methods* in Supplementary Text). Results showed significant negative genetic
correlations between All-PA-Leisure+SSOE with all disease traits studied, with the strongest values
for stroke (r_g=-0.62±0.11; p-value=3.54×10^{-9}) and gastroesophageal reflux disease (r_g=-
0.482±0023; p-value=9.9×10^{-97}) (Table S14 and Supplementary Text).
203
Local genetic correlation: All-PA-Leisure+SSOE
We next considered local correlations (LAVA[13]) as global genetic correlations may mask important
regional differences. There were a total of 91 significant local bivariate genetic correlations at 61
loci between the EUR All-PA-Leisure+SSOE meta-analysis and the 40 other traits used in the
genetic correlation analyses (Table S15). Of these 40 traits, there were 18 with at least one
significant local genetic correlation. Seventeen of these traits were previously found to have a
significant global genetic correlation with the EUR All-PA-Leisure+SSOE meta-analysis. The trait
with the highest number of significant local genetic correlations with All-PA-Leisure+SSOE was
educational attainment (31 loci), followed by BMI (15 loci) and income (9 loci). Only one trait- oral
cancer, had no significant global genetic correlation with EUR All-PA-Leisure+SSOE but still
showed a significant positive local genetic correlation (locus chr11: 118901214-120064528;
ρ=0.88; p-value=4.5×10^{-6}). For two traits, height and wellbeing, we observed local genetic
correlations with EUR All-PA-Leisure+SSOE in opposite directions (Table S15).
217
Mendelian Randomization analyses
We used MR analyses to investigate the causal relationship between All-PA-Leisure and traits with
which it had significant genetic correlation. Since several statistics for target traits included UKB
data, we used the GWAS of All-PA-Leisure in EUR MVP (and not the meta-analysis) to avoid
sample overlap. Four methods were tested: MR Egger, Weighted median, Inverse Variance
Weighted (IVW), and Simple Mode. We observed significant bidirectional negative (protective)
effects of All-PA-Leisure with respect to BMI, type 2 diabetes, leisure screen time, cigarettes per
day, and gastroesophageal reflux disease (Tables 1-2). There were unidirectional negative

(protective) effects of All-PA-Leisure on abdominal aortic aneurysm, heart failure, COVID-19
hospitalization, triglycerides, Phenotypic Age Acceleration (which is an age-adjusted biological age
measure, see Supplementary Methods), and osteoarthritis (Tables 1-2). We found significant
bidirectional positive effects between All-PA-Leisure and HDL cholesterol, parental survival, liking
of PA, household income, wellbeing, and educational attainment.
231
MVMR analyses testing BMI as confounder
To investigate the protective effect of All-PA-Leisure with respect to COVID-19 further, we
considered a possible role of BMI as confounder, as high BMI leads to worse COVID-19
outcomes.[14] We performed a multivariable MR (MVMR), where we used the All-PA-Leisure
phenotype and BMI as exposures, and COVID-19 hospitalization as the outcome. We observed a
significant protective effect of All-PA-Leisure on the COVID-19 trait which considers hospitalized
cases (effect=-0.067±0.016; p-value=2.8×10^{-5}; Table 3). Considering the role that BMI plays for
many other health outcomes, we also conducted MVMR including BMI for all other traits that had
significant MR results. Despite the reduced power of the analysis due to a decrease in number of
instrumental variables, which were reduced because of the required SNP overlap needed between
three GWAS compared to the two GWAS for the traditional MR analysis, we found significant
protective effects of All-PA-Leisure on HDL cholesterol, triglycerides, osteoarthritis, and
gastroesophageal reflux disease (Table 3).
245
All-PA-Leisure&Home and All-PA-Leisure&Home&Work phenotypes
To evaluate possible gain of information from combining data from PA during leisure time with PA
performed during home and work time, we ran two more GWAS in EUR. First, we summed All-
PA-Leisure with All-PA from physical activity at home (All-PA-Leisure&Home; Supplementary
Methods, Figure S9). We next added information from All-PA at work (All-PA-
Leisure&Home&Work; Supplementary Methods, Figure S10). In both the new GWASs we
observed a substantial reduction of the number of association peaks compared to All-PA-Leisure, as
would be expected given the relatively low genetic correlations among these traits.
254
255 For the EUR All-PA-Leisure+SSOE meta-analysis, we also conducted TWAS, fine-mapping, and
256 MTAG analysis with leisure screen time and liking of PA (see Supplementary Text, Table S16-
257 S20).
258
259 **GWAS of Vigorous PA phenotypes (Leisure, at Home, at Work)**
260
Kinds of vigorous physical activity and their genetic interrelationships
We ran independent MVP GWAS of all vigorous activity types depending by the context (Vig-PA-
Leisure, Vig-PA-Home, Vig-PA-Work) in EUR, and investigated correlations between each set of
two. For Vig-PA-Leisure and Vig-PA-Home, r_g=0.64±0.04 (p-value=5.0×10^{-59}); Vig-PA-Leisure
and Vig-PA-Work, r_g=0.46±0.07 (p-value=9.8×10^{-11}). Vig-PA-Home and Vig-PA-Work were more
strongly correlated with each other (r_g=0.97±0.05; p-value=1.2×10^{-73}). SNP-h^2 of vigorous PA
phenotypes are reported in Table S21.
268
269 For each one of the three vigorous PA traits, we ran GWAS, and for Vig-PA-Leisure we also
270 conducted XWAS, gene-based testing, TWAS, fine-mapping, functional enrichment, mitochondria-
271 SNV association, and gene-based association analysis on the mitochondrial genome (see
272 Supplementary Text, Table S22-S28, Figure S11-S15).
273
Genetic correlations between vigorous PA traits and traits of interest
To understand the differences between the vigorous activity traits better, we calculated the genetic
correlations between Vig-PA-Leisure, Vig-PA-Work, and Vig-PA-Home individually with the

previously analyzed traits of interest. We observed more statistically-significant differences
between Vig-PA-Leisure and Vig-PA-Work (for 15 traits), than when considering Vig-PA-Leisure
and Vig-PA-Home (for 12 traits) or Vig-PA-Home and Vig-PA-Work (for 3 traits) (Figure 4a).
Among these three pairs of vigorous activity traits, there were numerous significant differences
regarding the genetic correlations with income, leisure screen time, and educational attainment.
Analyses performed adjusting the vigorous PA traits with income using mtCOJO are described in
Supplementary Results, Table S29, Figure S16.
284
Genomic-SEM analysis
Considering the differences identified among the three vigorous PA traits (e.g. Vig-PA-Leisure,
Vig-PA-Work, and Vig-PA-Home), we used genomic-SEM analysis[15] to understand their
relationship better, inferring the overall genetic architecture among them and two other physical
activity traits in EUR, All-PA-Leisure+SSOE and liking of PA. The exploratory factor analysis
(EFA) indicated that a two-factor model fit the data best (Figure 4b). The two-factor model
accounted for 91% of cumulative variance explained (Factor 1: 50%; Factor 2: 41%). Both factors
had strong strength of sum of squares (SS) loadings (Factor 1 SS = 2.50; Factor 2 SS = 2.04).
Confirmatory factor analysis (CFA) confirmed that a two correlated-factors model fit the data well
with a comparative fit index (CFI) of 0.97, chi-square=109.03, Aikake information criterion
(AIC)=129.03, and a standardized root mean square residual (SRMR) of 0.069. Vig-PA-Leisure
(loading=1.00±0.03), All-PA-Leisure+SSOE (loading=0.89±0.03), and liking of PA
(loading=0.79±0.04) loaded most strongly on Factor 1, while Vig-PA-Home (loading=1.02±0.07)
and Vig-PA-Work (loading=0.82±0.08) loaded most strongly on Factor 2 (Figure 4b). Factor 1 and
Factor 2 were correlated at 0.50 (0.04).
300
Survival analyses
Survival analyses showed a significant reduction of risk of all-cause death for Vig-PA-Leisure
(beta=-0.1173±0.0037; *p*-value≤2×10^{-16}) and for Vig-PA-Work (beta=-0.0061±0.0029; *p*-
value=0.036). There was no statistically significant reduction of risk of death for Vig-PA-Home,
though the direction of effect and effect size were similar (beta=-0.0052±0.0030; *p*-value=0.082).
Discussion
Physical activity (PA) has long been appreciated to ameliorate risk for illness. Increasing physical
activity on a population level could greatly reduce morbidity and mortality – this is a major public
health undertaking and very difficult to accomplish. Understanding genetic factors that influence
level of activity for individuals could provide pathways for medical interventions to support efforts
to increase PA. We present here comprehensive information on the genetics of PA and investigate
the relationships between PA and health. This work provides tools to dissect causal relationships
between a range of other traits and PA, and to uncover previously unknown biology responsible in
part for inter-individual differences in activity. Those relationships were remarkably different
depending on the context of the physical activity – in particular, whether the activity was associated
with PA during leisure time (protective) rather than at work or around the house (often not
protective). Our main focus was accordingly on leisure-PA. All-PA-Leisure considered different
levels of PA intensity and frequency. We completed GWAS of All-PA-Leisure for EUR, AFR, and
AMR ancestry in MVP. We combined these data with UKB SSOE trait in two meta-analyses, one
EUR ancestry and the other cross-ancestry.
321
The polygenic architecture of leisure-time PA showed genetic liability to PA as a protective effect
with respect to numerous diseases. It was previously reported that PA has negative genetic
correlation with type 2 diabetes[10,12,16], triglyceride level[10], rheumatoid arthritis[12], coronary artery
disease[12], and lung cancer[12]. There is prior genetic evidence of the role of PA on preventing
colorectal cancer using MR methods[17], but we did not find a significant genetic correlation with

colorectal cancer. Our analyses confirmed the negative genetic correlations with type 2 diabetes,
coronary artery disease, and lung cancer. Moreover, we found significant negative genetic
correlation between PA and other medically relevant traits: thoracic and abdominal aortic
aneurysm, heart failure, stroke, asthma, type 1 diabetes, Alzheimer's disease, inflammatory bowel
disease, osteoarthritis, and gastroesophageal reflux disease. Notably, PA was also protective with
respect to COVID-19 and COVID-19-associated hospitalization, and positively correlated with
parental survival.

Whereas genetic correlations are an average of the shared association across the genome, local
genetic correlations can detect regions of shared genetic associations, including finding signals in
opposing directions that a genome-wide analysis would not be able to capture. Local genetic
correlation, like global correlation, supported a protective role of PA for multiple health traits,
showing negative local genetic correlations for diabetes types 1 and 2, gastroesophageal reflux
disease, and osteoarthritis. We also found negative local genetic correlations between All-PA-
Leisure+SSOE and BMI, triglycerides, Phenotypic Age Acceleration, leisure screen time,
neuroticism, and cigarettes per day. Positive local genetic correlations were found between All-PA-
Leisure+SSOE and bone mineral density, income, liking of PA, educational attainment, and,
unexpectedly, oral cancer: there was a single locus showing positive local genetic correlation
between PA and oral cancer – however, genome-wide genetic correlation between these traits
calculated with LDSC was not significant (*p*-value=0.418). We also found local genetic correlations
in opposite directions – at different genomic locations – with All-PA-Leisure+SSOE for height and
wellbeing.

Our MR results confirmed a protective effect of All-PA-Leisure genetic liability against type 2
diabetes, abdominal aortic aneurysm, heart failure, COVID-19 associated hospitalization,
triglyceride level, osteoarthritis, and gastroesophageal reflux disease. Most of these causal effects
were seen even when we tested for BMI as a confounder in the MR (MVMR) results (with the
exception of heart failure, which we could not test because of sample overlap for the BMI and heart
failure datasets). The low number of instrumental variables in our MVMR analyses could result in
false negatives. These findings nevertheless confirm the utility of All-PA-Leisure to decrease risk
for numerous diseases.[18,19] MR results also suggested a negative effect of All-PA-Leisure versus the
biological age measure Phenotypic Age Acceleration, and a bidirectional effect between All-PA-
Leisure and parental survival, uncovering the protective effect of genetic liability for this PA
phenotype versus aging and mortality.

PheWAS analysis in BioVU[20], which allowed a broad exploration of disease traits in an
independent sample, showed negative correlation with PA with several previously mentioned health
traits including diabetes mellitus, cardiovascular disease (e.g. coronary artery disease and heart
failure), gastroesophageal reflux disease, lung cancer, and asthma. The strongest association was
related to diabetes mellitus (*p*-value=5.89×10^{-36}).

The EUR meta-analysis of All-PA-Leisure+SSOE identified 163 independent significant SNPs, 91
of them novel compared to the independent significant SNPs of the two GWASs, All-PA-Leisure
and SSOE, in the meta-analysis. *CADM2**rs62253088 (encoding Cell Adhesion Molecule 2), on
chromosome 3, had the strongest association in both EUR and cross-ancestry All-PA-
Leisure+SSOE meta-analyses (EUR meta-analysis *p*-value=2.0×10^{-21}; cross-ancestry (EUR, AFR,
AMR) meta-analysis *p*-value=6.1×10^{-20}), but it was not significant in the GWAS of Vig-PA-
Leisure. However, considering Vig-PA-Leisure, we found another variant mapping to *CADM2*
which was a significant lead SNP in the GWAS (rs2326266; *p*-value=4.1×10^{-8}) and *CADM2* also
had the most significant *p*-value in the gene-based test (*p*-value=2.5×10^{-8}). *CADM2* variants,
including rs62253088, have previously been associated to PA, and also to decreased neuroticism

and decreased self-reported nervous and anxious feelings.[10] *CADM2* was also previously found to
be associated to obesity and metabolic traits,[21] cannabis use,[22] smoking and alcohol related traits,[23]
risk-taking behaviour,[24] educational attainment,[25] and autism spectrum disorder.[26] *CADM2* encodes
a member of the synaptic cell adhesion molecule 1 (SynCAM) family which belongs to the (Ig)
immunoglobulin superfamily.
383
The second most significant variant of the two meta-analyses, *MST1R**rs3733134 (EUR meta-
analysis p-value=1.4×10^{-17}; cross-ancestry meta-analysis p-value=6.9×10^{-18}), encoding macrophage
stimulating 1 receptor, was also significant in the GWAS of Vig-PA-Leisure (p-value=2.5×10^{-8}).
The third most significant variant was *EXD2**rs4899292 (EUR meta-analysis p-value=1.7×10^{-16};
cross-ancestry meta-analysis p-value=3.8×10^{-16}). *EXD2* (encoding exonuclease 3'-5' domain
containing 2) is involved in DNA replication to stabilize and restart stalled replication forks.[27]
Other strong significant associations included genes *LONRF2* (chromosome 2; encoding LON
peptidase N-terminal domain and ring finger 2) and *SLC39A8*, involved in the binding activity and
transportation of metal ions, and *NPC1* (chromosome 18; encoding NPC intracellular cholesterol
transporter 1), involved in cholesterol transport. *SLC39A8* was previously associated to several
psychiatric disorders, including problematic alcohol use,[28] schizophrenia,[29] and as a lead SNP in
opioid use disorder.[30] *C18orf8*, also known as *RMC1* (chromosome 18; encoding regulator of
MON1-CCZ1), mapped to the second most significant peak in the gene-based test of the cross-
ancestry meta-analysis, and is associated with type 2 diabetes and BMI.[31-33]
398
Post-GWAS analyses demonstrated that PA is related to brain structure and cellular architecture.
From MAGMA tissue expression analyses we determined that cerebellum and cerebellar
hemisphere showed significant enrichments. Cerebellar function is critical for skills relevant to PA,
such as coordination and movement. These results were similar to results of the previous GWAS of
SSOE.[10] Cell type specificity analyses showed a significant association of PA with the GABA$^+$
neurons.[34] EUR All-PA-Leisure+SSOE and cross-ancestry meta-analyses shared three significant
gene sets in their MAGMA gene-set analysis results: structural constituent of presynapse,
presynaptic active zone organization, and maintenance of presynaptic active zone structure. This
finding highlights the role of presynaptic cells in PA and the importance of brain in the biology of
PA level.
409
Vig-PA-Leisure was, as noted, genetically different from Vig-PA-Home and at Vig-PA-Work. We
observed moderate genetic correlation between Vig-PA-Leisure and Vig-PA-Home (r_g=0.64±0.04;
p-value=5.0×10^{-59}), but lower genetic correlation between Vig-PA-Leisure and Vig-PA-Work
(r_g=0.46±0.07; p-value=9.8×10^{-11}). While there was high genetic correlation between Vig-PA-
Home and Vig-PA-Work (r_g=0.97±0.05; p-value=1.2×10^{-73}), there were differences between these
two traits in genetic correlation analyses with other traits of interest, in particular for income,
educational attainment, and leisure screen time. Previous work on dementia also showed genetic
differences between PA during leisure vs. work time [35].
418
Consistent with what we observed with the divergent patterns of genetic correlation, the genomic-
SEM results supported that physical activity traits are genetically heterogeneous. Traits more
similar to voluntary physical activity and healthy lifestyle habits loaded on Factor 1 while Factor 2
captured traits more consistent with strenuous work and labor performed as part of work duties and
responsibilities at home. When we adjusted the vigorous physical activity traits for income using
mtCOJO, we still observed differences on genetic correlations with health traits. These findings
suggest that not all contexts of physical activity are equally beneficial, and some forms of vigorous
activity may be more strongly linked to improved mental and physical health.
427

A further confirmation of the different benefits by PA context was provided by the survival
analyses, which show a strong significant reduction of risk of death for Vig-PA-Leisure (beta=-
0.1173±0.0037; p-value≤$2×10^{-16}$) compared to a weak effect for Vig-PA-Work (beta=-
0.0061±0.0029; p-value=0.036), and none for for Vig-PA-Home (beta=-0.0052±0.0030; p-
value=0.082).The small protective effect for Vig-PA-Work contrasts with the opposing directions
for risk for several other important health traits (Figure 4a). The survival analyses may be
confounded by health status (i.e., healthier individuals are able to exercise more vigorously) and
competing risks (death). We conclude that PA at leisure is the most protective with respect to
negative medical outcomes.

This study has several limitations, including a relative lack of females in the MVP dataset
(approximately 9% for our data which nevertheless equates to >17,000 females for Vig-PA-
Leisure), which also made the potential interpretation of sex-specific associations infeasible due to
power imbalance. UKB has excellent female representation. Moreover, while there are many non-
EUR subjects included, their numbers are still insufficient, such that the cross-ancestry analysis
mostly reflects findings in EUR. It would also be important to verify the tissue expression analyses,
which are currently highlighting only significant associations with the brain, with more tissues and
larger samples. Indeed, our results may reflect a lack of power for other tissues or our sample size --
of the 53 tissue types contained in Genotype-Tissue Expression version 8 (GTEx v8), 13 tissues are
brain specific.

In conclusion, we have made progress towards revealing the genetic and biological nature of PA
during leisure time, which can differ greatly from PA undertaken in other contexts. In effect,
physical activity that one chooses to engage in in a leisure context has many positive effects with
respect to health traits, while activity that one must do, in a work or a household context, tends not
to have these beneficial effects. This difference was still observed after we adjusted for income.
Moreover, PA in the leisure context shows a much stronger decrease in the risk of death compared
to the other contexts. This may reflect the differences in intensity and duration of leisure compared
with work and home activity in the industrialized era. We detected a protective role of PA genetic
liability on several diseases, using global and local genetic correlation, PheWAS analyses and,
importantly, causal inference using MR. MR analyses showed a significant negative effect of PA on
several traits, with significant protective relationships on type 2 diabetes, abdominal aortic
aneurysm, heart failure, triglyceride level, osteoarthritis, gastroesophageal reflux disease, and
COVID-19 associated hospitalization. Finally, these results provide genetic insights into the
biological bases of PA, and add to strong experimental, epidemiological, and observational
literature showing the value to health of physical activity.

Data availability
The summary statistics produced in the two meta-analyses will be made available from PI lab
website upon publication: https://medicine.yale.edu/lab/gelernterlab/. The MVP GWAS summary
statistics will be available via the dbGaP study accession number phs001672.

Acknowledgements
This research is based in part on data from the Million Veteran Program, Office of Research and
Development, Veterans Health Administration.

This work was supported by funding from the Department of Veterans Affairs Office of Research
and Development, USVA, grants I01CX001849, and the VA Cooperative Studies Program study,
no. 575B; the VA National Center for PTSD Research, and the West Haven VA Mental Illness
Research, Education and Clinical Center; and by NIH grants R01DA054869 (JG), R01 AA026364

(JG). D.F.L. is supported by a Career Development Award CDA-2 from the Veterans Affairs Office of Research and Development (1IK2BX005058-01A2) and is Aimee Mann Fellow of Psychiatric Genetics.

This publication does not represent the views of the Department of Veteran Affairs or the United States Government.

Competing Interests

Dr. Gelernter is paid for editorial work for the journal "Complex Psychiatry". Dr. Murray Stein has in the past 3 years received consulting income from Acadia Pharmaceuticals, Aptinyx, atai Life Sciences, BigHealth, Bionomics, BioXcel Therapeutics, Boehringer Ingelheim, Clexio, Eisai, EmpowerPharm, Engrail Therapeutics, Janssen, Jazz Pharmaceuticals, NeuroTrauma Sciences, PureTech Health, Sumitomo Pharma, and Roche/Genentech. Dr. Stein has stock options in Oxeia Biopharmaceuticals and EpiVario. He has been paid for his editorial work on Depression and Anxiety (Editor-in-Chief), Biological Psychiatry (Deputy Editor), and UpToDate (Co-Editor-in-Chief for Psychiatry). He has also received research support from NIH, Department of Veterans Affairs, and the Department of Defense. He is on the scientific advisory board for the Brain and Behavior Research Foundation and the Anxiety and Depression Association of America. All other authors declare that they have no conflict of interest.

References

1. Bull, F.C.*, et al.* World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine* **54**, 1451-1462 (2020).
2. Gorzelitz, J.*, et al.* Independent and joint associations of weightlifting and aerobic activity with all-cause, cardiovascular disease and cancer mortality in the Prostate, Lung, Colorectal and Ovarian Cancer Screening Trial. *Br J Sports Med* **56**, 1277-1283 (2022).
3. Stubbe, J.H.*, et al.* Genetic influences on exercise participation in 37,051 twin pairs from seven countries. *PLoS One* **1**, e22 (2006).
4. De Moor, M.H.*, et al.* Genome-wide association study of exercise behavior in Dutch and American adults. *Med Sci Sports Exerc* **41**, 1887-1895 (2009).
5. Kim, J., Oh, S., Min, H., Kim, Y. & Park, T. Practical issues in genome-wide association studies for physical activity. *Annals of the New York Academy of Sciences* **1229**, 38-44 (2011).
6. Hara, M.*, et al.* Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. *Med Sci Sports Exerc* **50**, 2433-2441 (2018).
7. Lin, X.*, et al.* Genetic Determinants for Leisure-Time Physical Activity. *Med Sci Sports Exerc* **50**, 1620-1628 (2018).
8. Nabe-Nielsen, K.*, et al.* The effect of occupational physical activity on dementia: Results from the Copenhagen Male Study. *Scandinavian Journal of Medicine & Science in Sports* **31**, 446-455 (2021).
9. Zotcheva, E.*, et al.* Trajectories of occupational physical activity and risk of later-life mild cognitive impairment and dementia: the HUNT4 70+ study. *The Lancet Regional Health - Europe* **34**, 100721 (2023).
10. Klimentidis, Y.C.*, et al.* Genome-wide association study of habitual physical activity in over 377,000 UK Biobank participants identifies multiple variants including CADM2 and APOE. *International Journal of Obesity* **42**, 1161-1176 (2018).
11. Doherty, A.*, et al.* GWAS identifies 14 loci for device-measured physical activity and sleep duration. *Nature Communications* **9**, 5257 (2018).
12. Wang, Z.*, et al.* Genome-wide association analyses of physical activity and sedentary behavior provide insights into underlying mechanisms and roles in disease prevention. *Nature Genetics* **54**, 1332-1344 (2022).
13. Werme, J., van der Sluis, S., Posthuma, D. & de Leeuw, C.A. An integrated framework for local genetic correlation analysis. *Nature Genetics* **54**, 274-282 (2022).
14. Gao, M.*, et al.* Associations between body-mass index and COVID-19 severity in 6·9 million people in England: a prospective, community-based, cohort study. *Lancet Diabetes Endocrinol* **9**, 350-359 (2021).
15. Grotzinger, A.D.*, et al.* Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nature Human Behaviour* **3**, 513-525 (2019).
16. Carlsson, S., Ahlbom, A., Lichtenstein, P. & Andersson, T. Shared genetic influence of BMI, physical activity and type 2 diabetes: a twin study. *Diabetologia* **56**, 1031-1035 (2013).
17. Zhang, X.*, et al.* Genetically predicted physical activity levels are associated with lower colorectal cancer risk: a Mendelian randomisation study. *British Journal of Cancer* **124**, 1330-1338 (2021).
18. Guthold, R., Stevens, G.A., Riley, L.M. & Bull, F.C. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *Lancet Glob Health* **6**, e1077-e1086 (2018).

19. Guthold, R., Stevens, G.A., Riley, L.M. & Bull, F.C. Global trends in insufficient physical
activity among adolescents: a pooled analysis of 298 population-based surveys with 1·6
million participants. *Lancet Child Adolesc Health* **4**, 23-35 (2020).
20. Roden, D.M.*, et al.* Development of a large-scale de-identified DNA biobank to enable
personalized medicine. *Clin Pharmacol Ther* **84**, 362-369 (2008).
21. Morris, J.*, et al.* Genetic variation in CADM2 as a link between psychological traits and
obesity. *Scientific Reports* **9**, 7339 (2019).
22. Pasman, J.A.*, et al.* GWAS of lifetime cannabis use reveals new risk loci, genetic overlap
with psychiatric traits, and a causal effect of schizophrenia liability. *Nature Neuroscience*
21, 1161-1170 (2018).
23. Liu, M.*, et al.* Association studies of up to 1.2 million individuals yield new insights into the
genetic etiology of tobacco and alcohol use. *Nature genetics* **51**, 237-244 (2019).
24. Strawbridge, R.J.*, et al.* Genome-wide analysis of self-reported risk-taking behaviour and
cross-disorder genetic correlations in the UK Biobank cohort. *Translational Psychiatry* **8**, 39
(2018).
25. Lee, J.J.*, et al.* Gene discovery and polygenic prediction from a genome-wide association
study of educational attainment in 1.1 million individuals. *Nature Genetics* **50**, 1112-1121
(2018).
26. Casey, J.P.*, et al.* A novel approach of homozygous haplotype sharing identifies candidate
genes in autism spectrum disorder. *Hum Genet* **131**, 565-579 (2012).
27. Nieminuszczy, J.*, et al.* EXD2 Protects Stressed Replication Forks and Is Required for Cell
Viability in the Absence of BRCA1/2. *Molecular Cell* **75**, 605-619.e606 (2019).
28. Zhou, H.*, et al.* Genome-wide meta-analysis of problematic alcohol use in 435,563
individuals yields insights into biology and relationships with other traits. *Nature*
Neuroscience **23**, 809-818 (2020).
29. Pardiñas, A.F.*, et al.* Common schizophrenia alleles are enriched in mutation-intolerant
genes and in regions under strong background selection. *Nat Genet* **50**, 381-389 (2018).
30. Deak, J.D.*, et al.* Genome-wide association study in individuals of European and African
ancestry and multi-trait analysis of opioid use disorder identifies 19 independent genome-
wide significant risk loci. *Molecular Psychiatry* (2022).
31. Akiyama, M.*, et al.* Genome-wide association study identifies 112 new loci for body mass
index in the Japanese population. *Nat Genet* **49**, 1458-1467 (2017).
32. Pulit, S.L.*, et al.* Meta-analysis of genome-wide association studies for body fat distribution
in 694 649 individuals of European ancestry. *Hum Mol Genet* **28**, 166-174 (2019).
33. Vujkovic, M.*, et al.* Discovery of 318 new risk loci for type 2 diabetes and related vascular
outcomes among 1.4 million participants in a multi-ancestry meta-analysis. *Nat Genet* **52**,
680-691 (2020).
34. Maddock, R.J., Casazza, G.A., Fernandez, D.H. & Maddock, M.I. Acute Modulation of
Cortical Glutamate and GABA Content by Physical Activity. *The Journal of Neuroscience* **36**,
2449-2457 (2016).
35. Nabe-Nielsen, K.*, et al.* The effect of occupational physical activity on dementia: Results
from the Copenhagen Male Study. *Scand J Med Sci Sports* **31**, 446-455 (2021).

630
631 *Figure 1:* **(A) EUR All-PA-Leisure+SSOE meta-analysis and (B) cross-ancestry All-**
632 **PA-Leisure+SSOE meta-analysis.** Annotated genes are the closest to each significant
633 lead SNP. Different lead SNPs are highlighted in green. The red and blue horizontal lines
634 indicate the genome-wide significance (*p*-value=5×10^{-8}) and *p*-value=10^{-5} levels,
635 respectively.

636
637
638

640 *Figure 2:* **Traits genetically correlated with All-PA-Leisure+SSOE.** (A) PheWAS of
641 EUR All-PA-Leisure+SSOE meta-analysis. The red line represents the Bonferroni *p*-value
642 threshold=3.99x10^{-5}. (B) Genetic correlations of EUR meta-analysis for All-PA-
643 Leisure+SSOE phenotype with traits of interest. We display here the traits of interest that
644 were significant after the Benjamini-Hochberg false discovery procedure.

Figure 3: **Genetic differences among vigorous PA.** (A) Genetic correlation between each pair of vigorous PA (Vig-PA-Leisure, Vig-PA-Work, Vig-PA-Home) and other traits of interest. Significantly different correlations were defined having *p*-value < 4.76×10⁻⁴ after multiple testing correction: * represents significant difference between Vig-PA-Leisure and Vig-PA-Work, ^ between Vig-PA-Leisure and Vig-PA-Home, + between Vig-PA-Home and Vig-PA-Work. (B) genomic-SEM of the two-factor model, showing the loadings for the inferred traits.

Outcome	N	IVW		MR Egger		Weighted Median		Simple Mode	
		β (SE)	p	β (SE)	p	β (SE)	p	β (SE)	p
Aneurysm – thoracic aortic	70	-0.031 (0.020)	0.12	0.036 (0.071)	0.61	-0.025 (0.027)	0.35	-0.021 (0.064)	0.74
Coronary artery disease	53	-0.04 (0.02)	0.034	-0.01 (0.10)	0.89	-0.03 (0.02)	0.14	-0.03 (0.05)	0.51
Aneurysm – abdominal aortic	70	**-0.119 (0.028)**	**2.2×10^{-5}**	0.02 (0.10)	0.81	-0.089 (0.041)	0.029	-0.043 (0.096)	0.64
Heart failure	72	**-0.040 (0.009)**	**6×10^{-6}**	-0.04 (0.03)	0.23	-0.03 (0.01)	5×10^{-3}	-0.02 (0.03)	0.50
Stroke	72	-8.0×10^{-4} (3.4×10^{-4})	0.017	-1.3×10^{-3} (1.2×10^{-3})	0.30	-4.6×10^{-4} (4.2×10^{-4})	0.27	0.2×10^{-3} (1.3×10^{-3})	0.89
COVID-19	72	-0.015 (0.005)	5.3×10^{-3}	0.003 (0.021)	0.87	-0.007 (0.006)	0.27	-0.003 (0.017)	0.84
Asthma	72	-0.005 (0.002)	0.006	-0.008 (0.006)	0.23	-0.005 (0.002)	0.017	-0.007 (0.006)	0.23
COVID-19 hospitalization	71	**-0.047 (0.011)**	**3.3×10^{-5}**	-0.057 (0.043)	0.18	-0.043 (0.013)	1.4×10^{-3}	-0.045 (0.035)	0.21
Height	72	0.0079 (0.0062)	0.21	0.039 (0.022)	0.086	**0.0108 (0.0026)**	**4.1×10^{-5}**	0.0134 (0.0056)	0.019
Bone Mineral Density	73	0.0044 (0.0051)	0.39	0.003 (0.018)	0.86	0.0094 (0.0049)	0.055	0.012 (0.013)	0.36
BMI	53	**-0.028 (0.007)**	**2.9×10^{-5}**	-0.08 (0.03)	0.02	-0.020 (0.006)	1.9×10^{-3}	-0.018 (0.015)	0.23
HDL cholesterol	72	**0.0111 (0.0023)**	**2.5×10^{-6}**	0.0135 (0.0085)	0.12	**0.0069 (0.0014)**	**1.6×10^{-6}**	0.0070 (0.0034)	0.045
Type 1 diabetes	73	-0.026 (0.018)	0.14	-0.046 (0.065)	0.49	-0.043 (0.022)	0.048	-0.082 (0.056)	0.15
Triglycerides	72	**-0.0194 (0.0042)**	**3.5×10^{-6}**	-0.002 (0.015)	0.88	**-0.0150 (0.0037)**	**4.4×10^{-5}**	-0.0170 (0.0094)	0.073
Type 2 diabetes	53	**-0.068 (0.016)**	**1.6×10^{-5}**	**-0.061 (0.015)**	**4.0×10^{-5}**	-0.087 (0.084)	0.31	-0.078 (0.042)	0.072
Parental survival	67	**-0.0287 (0.0051)**	**2.4×10^{-8}**	0.009 (0.019)	0.63	**0.0208 (0.0061)**	**5.8×10^{-4}**	0.017 (0.015)	0.26
Biological Age Acceleration	72	-0.0121 (0.0094)	0.20	-0.021 (0.034)	0.54	-0.016 (0.012)	0.16	-0.014 (0.025)	0.57
Phenotypic Age Acceleration	72	**-0.112 (0.028)**	**6.5×10^{-5}**	0.02 (0.10)	0.88	-0.070 (0.030)	0.022	-0.046 (0.079)	0.56
Liking of PA	80	**0.0341 (0.0047)**	**2.9×10^{-13}**	0.003 (0.017)	0.86	**0.0302 (0.0052)**	**6.9×10^{-9}**	0.042 (0.013)	2.0×10^{-3}
Income	72	**0.038 (0.005)**	**7×10^{-16}**	0.002 (0.017)	0.90	**0.030 (0.004)**	**2×10^{-11}**	0.017 (0.013)	0.17
Wellbeing	62	**7.9×10^{-3} (2.2×10^{-3})**	**3.7×10^{-4}**	-7.6×10^{-3} (9.5×10^{-3})	0.43	3.3×10^{-3} (1.7×10^{-3})	0.056	1.7×10^{-3} (5.0×10^{-3})	0.74
Neuroticism	72	-0.014 (0.005)	0.004	0.022 (0.018)	0.21	-7.6×10^{-3} (4.4×10^{-3})	0.09	-0.015 (0.010)	0.15
Leisure screen time	73	**-0.0565 (0.0075)**	**7.0×10^{-14}**	-0.016 (0.027)	0.55	**-0.0366 (0.0066)**	**2.3×10^{-8}**	-0.034 (0.019)	0.082
Educational attainment	69	**0.0496 (0.0052)**	**2×10^{-21}**	0.003 (0.020)	0.86	**0.0272 (0.0035)**	**7.6×10^{-15}**	0.0248 (0.0084)	4.4×10^{-3}
Executive functioning	69	0.0015 (0.0018)	0.39	-0.0019 (0.0074)	0.79	0.0006 (0.0018)	0.75	-0.0083 (0.0055)	0.13
Alzheimer	73	-0.003 (0.002)	0.11	-0.004 (0.008)	0.65	-0.003 (0.003)	0.37	-0.001 (0.007)	0.84
Drinks per week	70	8.2×10^{-3} (3.3×10^{-3})	0.014	0.014 (0.013)	0.28	5.1×10^{-3} (3.0×10^{-3})	0.090	6.6×10^{-3} (8.6×10^{-3})	0.44
Cigarettes per day	70	**-4.07×10^{-2} (0.78×10^{-2})**	**1.77×10^{-5}**	0.010 (0.029)	0.73	**-4.17×10^{-2} (0.88×10^{-2})**	**1.98×10^{-6}**	-0.068 (0.027)	0.015
Fruit	72	**0.0118 (0.0024)**	**1.07×10^{-6}**	0.0040 (0.0087)	0.65	0.0089 (0.0034)	9.95×10^{-3}	0.0033 (0.0091)	0.72
Olive oil	72	0.0077 (0.0033)	0.019	0.031 (0.012)	8.80E-03	0.0092 (0.0042)	0.03	0.011 (0.012)	0.34
Vegetables	72	0.0010 (0.0022)	0.66	-0.0141 (0.0079)	0.078	-0.0006 (0.0031)	0.84	0.0011 (0.0080)	0.89
Cheese	72	0.0028	0.39	-0.004	0.74	0.0024	0.58	0.002	0.83

Exposure	N	IVW		MR Egger		Weighted Median		Simple Mode	
		(0.0032)		(0.012)		(0.0043)		(0.011)	
Inflammatory bowel disease	73	-0.0019 (0.0012)	0.10	-0.0002 (0.0029)	0.95	-0.0004 (0.0017)	0.80	-0.0017 (0.0033)	0.60
Cancer – lung	73	-0.044 (0.034)	0.19	-0.26 (0.12)	0.042	0.016 (0.043)	0.71	0.08 (0.11)	0.50
Osteoarthritis	72	**-0.0036 (0.0008)**	**1.15×10^{-5}**	-0.0032 (0.0030)	0.29	-0.0026 (0.0009)	5.3×10^{-3}	-0.0026 (0.0023)	0.27
Gastroesophageal reflux disease	70	**-0.063 (0.011)**	**5.8×10^{-9}**	0.009 (0.040)	0.83	**-0.058 (0.010)**	**1.5×10^{-9}**	-0.070 (0.031)	2.6×10^{-2}

668 *Table 1:* **MR analysis of causal effects of PA during leisure time (All-PA-Leisure) on**
669 **traits of interest.** Significant results after multiple testing correction (*p*-value < 6.9×10^{-4})
670 are highlighted in **bold.**

		β (SE)	p	β (SE)	p	β (SE)	p	β (SE)	p
Aneurysm – thoracic aortic	30	-0.005 (0.056)	0.93	0.11 (0.14)	0.43	0.117 (0.068)	0.086	0.11 (0.13)	0.41
Coronary artery disease	21	-0.01 (0.07)	0.85	-0.03 (0.13)	0.85	-0.07 (0.07)	0.31	-0.1 (0.1)	0.51
Aneurysm – abdominal aortic	23	0.029 (0.051)	0.57	-0.02 (0.13)	0.91	0.042 (0.062)	0.50	0.06 (0.13)	0.67
Heart failure	48	-0.2 (0.1)	0.07	0.8 (0.3)	7×10^{-3}	0.0 (0.1)	0.99	0.1 (0.2)	0.74
Stroke	22	-0.5 (4.4)	0.90	-8.4 (8.4)	0.33	5.1 (6.0)	0.39	7.6 (11.4)	0.51
COVID-19	52	-0.22 (0.15)	0.14	-0.09 (0.27)	0.73	-0.13 (0.20)	0.49	-0.74 (0.39)	0.07
Asthma	44	-0.3 (0.5)	0.55	-1.6 (1.5)	0.32	-1.1 (0.7)	0.10	-0.8 (1.2)	0.50
COVID-19 (hospitalized)	85	-0.04 (0.05)	0.48	0.01 (0.10)	0.96	-0.02 (0.08)	0.80	0.12 (0.17)	0.48
Height	722	0.208 (0.078)	8.0×10^{-3}	-0.12 (0.16)	0.44	0.14 (0.10)	0.17	0.36 (0.28)	0.19
Bone Mineral Density	366	-0.046 (0.063)	0.46	-0.10 (0.13)	0.44	-0.050 (0.091)	0.59	-0.02 (0.23)	0.93
BMI	141	**-1.02 (0.17)**	**3.3×10^{-9}**	-0.18 (0.48)	0.73	**-0.82 (0.19)**	**1.7×10^{-5}**	-1.63 (0.49)	1.1×10^{-3}
HDL cholesterol	360	**0.77 (0.22)**	**4.4×10^{-4}**	-0.32 (0.33)	0.33	0.18 (0.30)	0.55	-0.03 (0.72)	0.97
Type 1 diabetes	153	-0.004 (0.018)	0.81	-0.003 (0.025)	0.91	-0.018 (0.022)	0.41	0.011 (0.069)	0.87
Triglycerides	297	-0.262 (0.091)	4.1×10^{-3}	0.22 (0.13)	0.094	-0.09 (0.13)	0.45	0.13 (0.29)	0.66
Type 2 diabetes	210	**-0.156 (0.044)**	**3.7×10^{-4}**	-0.073 (0.066)	0.27	0.09 (0.10)	0.38	0.01 (0.15)	0.94
Parental survival	86	**0.67 (0.17)**	**9.6×10^{-5}**	-0.32 (0.39)	0.41	0.64 (0.22)	3.1×10^{-3}	1.42 (0.52)	7.8×10^{-3}
Biological Age Acceleration	76	-0.100 (0.071)	0.16	0.10 (0.16)	0.51	0.042 (0.096)	0.66	-0.06 (0.24)	0.79
Phenotypic Age Acceleration	116	-0.068 (0.025)	7.0×10^{-3}	-0.078 (0.070)	0.27	-0.037 (0.030)	0.21	0.029 (0.090)	0.75
Liking of PA	46	**1.42 (0.33)**	**2.2×10^{-5}**	-0.8 (1.2)	0.51	**1.51 (0.35)**	**1.5×10^{-5}**	1.88 (0.85)	0.031
Income	177	**2.12 (0.19)**	**4×10^{-28}**	0.85 (0.66)	0.20	**1.55 (0.22)**	**4×10^{-12}**	1.27 (0.79)	0.11
Wellbeing	276	**2.4 (0.4)**	**4×10^{-9}**	**6.6 (1.8)**	**3×10^{-4}**	**2.0 (0.5)**	**2×10^{-5}**	2.0 (1.7)	0.23
Neuroticism	186	**-0.97 (0.19)**	**2.5×10^{-7}**	-1.48 (0.77)	0.06	**-0.78 (0.22)**	**2.8×10^{-4}**	-0.86 (0.70)	0.22
Leisure screen time	274	**-1.45 (0.11)**	**1.3×10^{-37}**	-1.35 (0.44)	2.4×10^{-3}	**-1.09 (0.13)**	**6.6×10^{-17}**	-0.55 (0.44)	0.21
Educational attainment	544	**2.62 (0.12)**	**4.1×10^{-99}**	**2.46 (0.41)**	**5.2×10^{-9}**	**2.34 (0.16)**	**1.2×10^{-45}**	1.66 (0.70)	0.018
Executive functioning	235	0.93 (0.34)	6.6×10^{-3}	-0.62 (1.33)	0.64	1.09 (0.42)	9.8×10^{-3}	2.00 (1.46)	0.17
Alzheimer	50	-0.1 (0.3)	0.79	-0.1 (0.5)	0.80	0.2 (0.4)	0.60	1.0 (0.8)	0.21
Drinks per week	124	-0.1 (0.31)	0.59	-1.0 (0.70)	0.16	-0.86 (0.35)	0.014	0.24 (0.87)	0.78
Cigarettes per day	78	**-0.62 (0.10)**	**9.9×10^{-10}**	-0.07 (0.17)	0.68	-0.24 (0.13)	0.073	-1.01 (0.39)	0.012
Fruit	12	0.22 (0.88)	0.8	-4.05 (1.73)	0.042	-0.37 (0.95)	0.69	-1.42 (1.72)	0.43
Olive oil	12	0.08 (0.57)	0.89	-1.62 (1.44)	0.29	-0.35 (0.77)	0.65	-0.79 (1.35)	0.57
Vegetables	12	0.72 (1.10)	0.51	-1.16 (3.03)	0.71	1.71 (1.11)	0.12	2.00 (2.02)	0.34
Cheese	10	-1.22 (0.60)	0.042	-0.44 (1.50)	0.78	-1.53 (0.78)	0.050	-1.69 (1.24)	0.20
Inflammatory bowel disease	68	-1.08 (0.52)	0.035	-0.82 (2.79)	0.77	-0.95 (0.61)	0.12	-1.02 (1.20)	0.40
Cancer – lung	18	-0.006 (0.044)	0.90	-0.06 (0.13)	0.66	0.016 (0.059)	0.79	0.066 (0.099)	0.52

| Osteoarthritis | 34 | -4.5 (2.2) | 0.036 | -7.7 (5.8) | 0.20 | -0.5 (2.1) | 0.81 | 3.3 (4.4) | 0.46 |
| Gastroesophageal reflux disease | 110 | **-0.94 (0.13)** | **1.2×10^{-12}** | 0.10 (0.49) | 0.84 | **-0.73 (0.13)** | **2.2×10^{-8}** | -0.77 (0.43) | 0.07 |

711 **_Table 2:_ MR analysis of causal effects on PA during leisure time (All-PA-Leisure) of**
712 **traits of interest.** Significant results after multiple testing correction (p-value $< 6.9×10^{-4}$)
713 are highlighted in **bold.**

Exposure	Outcome	N	β (SE)	p
PA	Aneurysm – abdominal aortic	7	-0.087 (0.055)	0.11
BMI	Aneurysm – abdominal aortic	49	0.31 (0.18)	0.092
PA	**COVID-19 hospitalization**	**7**	**-0.067 (0.016)**	**2.8×10^{-5}**
BMI	COVID-19 hospitalization	50	0.37 (0.05)	8.2×10^{-13}
PA	**HDL cholesterol**	**7**	**0.0221 (0.0047)**	**2.3×10^{-6}**
BMI	HDL cholesterol	50	-0.042 (0.016)	7.0×10^{-3}
PA	**Triglycerides**	**7**	**-0.038 (0.011)**	**3.6×10^{-4}**
BMI	Triglycerides	50	0.040 (0.035)	0.26
PA	Type 2 diabetes	6	-0.058 (0.057)	0.30
BMI	Type 2 diabetes	43	0.42 (0.18)	0.19
PA	Parental survival	7	0.030 (0.010)	2.9×10^{-3}
BMI	Parental survival	48	-0.171 (0.034)	4.4×10^{-7}
PA	Phenotypic Age Acceleration	7	-0.091 (0.044)	3.7×10^{-2}
BMI	Phenotypic Age Acceleration	50	0.92 (0.15)	4.8×10^{-10}
PA	**Osteoarthritis**	**7**	**-0.0037 (0.0012)**	**1.4×10^{-3}**
BMI	Osteoarthritis	50	0.0253 (0.0039)	1.3×10^{-10}
PA	**Gastroesophageal reflux disease**	**7**	**-0.063 (0.015)**	**2.3×10^{-5}**
BMI	Gastroesophageal reflux disease	50	0.141 (0.050)	4.6×10^{-3}

759   *Table 3:* **MVMR analysis of All-PA-Leisure setting BMI as exposures.** Significant
760   results after multiple testing correction (*p*-value < 5.6x10^{-3} outcomes) are highlighted in
761   **bold.**
762
763