Using theater-based research to study views on contraception

Wieke Y. Beumer[a-c*], Merel Sprenger[d*], Ebru Eksi[e], Chris Bartels[e,f], Samira van Bohemen[g], Jenneke van Ditzhuijzen[b,c,h] Linda P. Dekker[f], Ingvil de Haan[i], Hanan El Marroun[e,f], Pauline W. Jansen[e,f], Hilmar H. Bijma[j,k], Clair A. Enthoven[e,f]

a. Amsterdam UMC location University of Amsterdam, Epidemiology and Data Science, Meibergdreef 9, Amsterdam, Netherlands.
b. Amsterdam UMC location University of Amsterdam, Obstetrics and Gynecology, Meibergdreef 9, Amsterdam, Netherlands.
c. Amsterdam Reproduction and Development, Amsterdam, The Netherlands.
d. Public Health and Primary Care/Health Campus The Hague, Leiden University Medical Center, Leiden, the Netherlands.
e. Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Center, Rotterdam, The Netherlands.
f. Department of Psychology, Education and Child Studies, Erasmus School of Social and Behavioral Sciences, Erasmus University, Rotterdam, The Netherlands.
g. Department of Public Administration and Sociology, Erasmus School of Social and Behavioral Sciences, Erasmus University, Rotterdam, The Netherlands.
h. Department of Interdisciplinary Social Science, Social Policy and Public Health, Utrecht University, Utrecht, The Netherlands.
i. Rutgers, Expert Centre on Sexuality, Utrecht, The Netherlands.
j. Department of Obstetrics and Gynaecology, Division of Obstetrics and Fetal Medicine, Erasmus University Medical Center MC Sophia, Rotterdam, The Netherlands.
k. Department of Care Ethics, University of Humanistic Studies, Utrecht, The Netherlands.

*Wieke Y. Beumer and Merel Sprenger contributed equally to this paper and share first authorship.

Correspondence to: Clair A. Enthoven, Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC, Rotterdam, Na Building, Floor 29 | E: c.enthoven@erasmusmc.nl

Wordcount abstract: 250; Wordcount text: 3060

Abstract

Objectives

Historically, contraceptive research is impeded by barriers to openly discuss sexual experiences. Theater-based research is a promising approach to foster deeper engagement and a richer exploration of lived experiences to gain insights in underlying implicit factors that shape contraceptive use. This study uses theater-based research to explore adults' views, responsibility, satisfaction, and knowledge about contraception.

Study Design

The study was conducted at a three-day music festival in the Netherlands with 1,024 participants (32.9% men, mean age 28.1 years). Theater was integrated by transforming the study setting into a dance club, where the research team performed characters. Participants completed an individual survey prior to the theater experience, a team-based contraceptive knowledge quiz, and a writings on the wall activity. Survey and quiz data were analyzed using linear regression, thematic analysis was applied for evaluating writings on the wall.

Results

On a scale from 1 to 10, the survey revealed general contraceptive satisfaction (women: 7.0, men: 7.7) and responsibility (women: 8.7, men: 7.7). Participant teams had a mean of 73.7% correct quiz answers. We found six themes that emerged in contraceptive views: side effects, fallibility, shared responsibility, information needs, dismissive counseling, and resistance to hormones.

Conclusion

This study revealed complementary findings between traditional survey findings, indicating overall satisfaction with contraception, and theater-based writings on the wall, eliciting more candid and critical reflections with a notably negative tone. Theater-based data collection may capture lived experiences more vividly and lower barriers for research participation, suggesting its value for studying sexual and reproductive health.

Keywords

Arts-based research; Interactive theater; Contraception; Satisfaction; Women; Men

Implications

Theater offers a promising complement to conventional contraception research. By fostering comfort, humor and participation, it reveals more candid perspectives and may therefore reduce social

62   desirability bias. Future research should further examine its added value in recruiting populations not

63   reached by traditional methods.

1   Introduction

Contraceptive methods play a central role in empowering people to make autonomous reproductive choices (Alspaugh et al., 2020). In settings where contraceptive care is widely available and generally accessible, discussions have focused on the reasons for a recent decline in (hormonal) contraceptive use (de Graaf et al., 2024; Le Guen et al., 2021; Schneider-Kamp & Takhar, 2023). A substantial body of qualitative research, primarily using semi-structured interviews and focus groups, shows that contraceptive use is not only shaped by the desire to avoid pregnancy, but also by negative views, satisfaction, and (sexual) partner dynamics (Alspaugh et al., 2020; Ayoola et al., 2007; Baxter et al., 2011; D'Souza et al., 2022; Hoga et al., 2014; Pratt et al., 2014; Williamson et al., 2009). These implicit factors may not be fully understood using traditional research methods, such as surveys, interviews and focus groups, as social desirability may limit gauging the lived experience of people, particularly when exploring intimate experiences and personal perspectives, such as contraception (Tourangeau & Yan, 2007). This complexity underscores the need for novel methodological approaches.

An approach to elucidate and evaluate implicit aspects of people's lived experiences is arts-based research. Arts-based research draws on diverse ways of knowing and experiencing the world (Knowles & Cole, 2008) and serves as an umbrella term encompassing a range of methodologies guided by constructivist, experiential, and empirical principles. Examples of arts-based methods include poetry, photography, painting, and theater: art forms that create interactions among participants extending beyond structured conversations, fostering deeper emotional and cognitive engagement and encouraging self-reflection (Myllyoja, 2024).

Theater-based research is an arts-based method in which theatrical elements are used during data collection (such as improvisation, theatrical costumes and role play), to lower participation barriers and create an open, playful environment that enables participants to articulate experiences and perspectives that might otherwise remain unspoken (Bleuer et al., 2018; Collins & Stockton, 2022; Myllyoja, 2024; Rossiter et al., 2008). Theater-based research may be particularly useful in research on contraception because it has the potential to engage a broader and typically less included sample (Knowles & Cole, 2008), addressing the underrepresentation of men commonly observed in contraception studies (Alspaugh et al., 2020). Furthermore, theater promotes knowledge exchange, allowing participants to gain new insights through active involvement in the research process (Bleuer et al., 2018; Myllyoja, 2024). As a result, theater has thus far been used mainly as a medium for disseminating knowledge (Bleuer et al., 2018; Fleckman et al., 2023; Marzi et al., 2025; Rossiter et al.,

95   2008), while its application as a method for data collection remains largely unexplored. Therefore, in this
96   study we aim to explore contraceptive views, responsibility, satisfaction and knowledge among adult
97   women and men using theater-based research.

2   Methods

2.1   Setting and participants

Participants aged 18 or older were recruited at Lowlands, a large yearly recurring music festival
attracting around 65,000 visitors in the Netherlands during three days in August 2023. Besides music,
the festival offers a wide variety of entertainment including theater, movies, comedy, literature, visual
arts and science. Lowlands Science is an initiative to foster science engagement by welcoming ten
different research projects at Lowlands each year (Docter-Loeb, 2024). The participants provided digital
informed consent, and individuals who were noticeably intoxicated could not participate. The protocol
received approval from the Ethics Review Committee of Erasmus School of Social and Behavioral
Sciences (ETH2223-0634). Our research project entitled: "Perception of Contraception".

2.2   Design

This study included an individual online survey, a team-based contraceptive knowledge quiz and
a writings on the wall activity. The online survey and quiz were co-created with stakeholders, including a
gynecologist and two program managers working at a societal organization to promote sexual and
reproductive health. The survey was conducted prior to the interactive theater experience, the quiz
during the theater experience and the writings on the wall at the end of this experience (See next
section 'Interactive theater').

The survey was conducted on participants' smartphones and included questions about age, sex,
gender, educational attainment, sexual orientation, presence of a steady sexual partner (hereafter
referred to as 'relationship'), and number of sexual partners in the past year. Participants then indicated
whether they did anything to prevent pregnancy (hereafter referred to as 'contraceptive use') and which
methods they used. Finally, they answered questions about satisfaction with their contraception and
their sense of responsibility on a scale from 0 (not at all) to 10 (very much). The survey questions and
answer options are described in Supplementary Table 1.

The contraceptive knowledge quiz was conducted among small teams (two to six participants
per team). The survey included eight questions with four answer options, based on existing questions

(Haynes et al., 2017; Sense). Topics included effectiveness and working mechanisms of contraceptive

methods, protection against sexually transmitted infections (STIs), sperm viability, and the fertile period.

A full description of the quiz is shown in Supplementary Table 2.

The writings on the wall consisted of wall panels featuring the invitation, "What is your opinion

about…" and contained large drawings of contraceptive methods, allowing participants to write about

these specific methods. The depicted contraceptive methods included: the contraceptive pill, injection,

ring, or patch, the condom, internal condom, diaphragm, (hormonal) intrauterine device (IUD), and

sterilization. Empty spaces around these drawings provided an opportunity to describe views toward

contraception in general.

2.3   Interactive theater

The theater design was developed during multiple one-on-one brainstorm sessions between

one of the researchers (CAE) and actors with extensive experience with development and performance

of interactive theater acts at festivals. Theater-based aspects of the study included the use of costumes,

role play, improvisation, attributes for participants to interact with, and a research area designed to

create a strong visual impact.

We created a dance club named 'Club Womb' as our project setting, visually presented in Figure

1. At the entrance, the 'bouncer' explained the purpose of the study, organized digital consent and

directed participants towards a brief online survey (Figure 1.1). Next, the 'host' welcomed the

participants, explained that they could "protect themselves from pregnancy" during their stay in 'Club

Womb' and invited them to choose one of ten enlarged contraceptive methods (e.g.., raincoat as

condom, hoop as contraceptive ring; Figure 1.2). Equipped with their method of choice, participants

entered the club by crawling through a large vulva after touching the 'clitoris' (a plasma ball light; Figure

1.3). Inside, the 'sexual health doctor' welcomed participants in the "most visited womb of the festival",

conducted a contraceptive knowledge quiz and explained the correct answers (Figure 1.4). Finally,

participants moved into the silent disco where the 'DJ/MC' (Figure 1.5) invited them to write their

opinions and experiences with contraceptive methods on the wall (Figure 1.6). The club 'mascot' (actor

in a penis suit) recruited participants and danced with them (Figure 1.1). Throughout the experience, all

actors helped participants feel at ease by using empathetic communication and humor, and by adapting

their approach to suit individual comfort levels.

Throughout data collection, we continuously reflected on how participants engaged with the
theatrical aspects of the study and the specific characters performed by the actors. Observations
focused on participants' movement through the three stages and on identifying possible improvements
to the design and interactions. Team reflection sessions were held four times—once midway through
the first day and at the end of each of the three days. Reflections on the benefits and drawbacks of the
use of theater-based research from three researchers (WB, MS, CAE) are described in the discussion
section.

[Figure 1, see end of doc]

2.4   Analyses

Regarding the quantitative analyses, for the online survey, means and standard deviations were
computed for contraceptive satisfaction and responsibility separately for women and men. For the
contraceptive knowledge quiz, the mean and standard deviation of the percentage of correct answers
per team were calculated separately for female, male, and mixed teams. We additionally conducted
multivariate linear regression analyses with sex as explanatory variable and contraceptive satisfaction,
responsibility, and knowledge as outcome variables, adjusted for age, educational attainment, and
relationship. We used IBM SPSS Statistics, Version 28.0.1.0.

Regarding the qualitative analyses, for the writings on the wall, we photographed the panels
and transcribed them verbatim into MS Excel for inductive data processing and analysis. Besides
verbatim contributions, participants interacted with other quotes through arrows or additional
punctuation marks. We used thematic analysis in our exploratory approach of identifying, analyzing, and
reporting of themes within the data (Braun & Clarke, 2006). Data extraction, open coding, axial coding,
and constructing the overarching themes was performed by CB, in consultation with CAE. Subsequent
analyses were performed by WB. Since this is an iterative process, the researchers went back and forth
between axial coding and constructing the themes. Decisions were made based on consensus within the
team (CB, CAE and WB).

3   Results

3.1   Sample characteristics

A total of 1,088 people started the study. We excluded 64 participants due to incomplete data,
leaving 1,024 participants. 744 Participants, divided over 292 teams, also participated in the
contraceptive knowledge quiz. Participants had a mean age of 28.1 years (range 18 to 62) and 33.6%
identified as male. The majority of the participants had a high educational attainment and were in a
relationship (Tables 1 and Supplementary Table 2). Most participants (91.3%) reported using
contraception. Of those who did not use contraception, most (73.0%) were either not engaging in penis-
in-vagina sex, were pregnant, or were trying to conceive. When contraception was used, the hormonal
IUD, condoms, and the contraceptive pill were most commonly reported (Supplementary Table 3).

Table 1. *Characteristics (n(%)) of the study sample (n=1,024) at Lowlands festival, The Netherlands, 2023.*

	Total sample n=1,024	Women n=683	Men n=341
Age			
18-24 years	288 (22.3)	161 (23.6)	67 (19.6)
25-29 years	497 (48.5)	354 (51.8)	143 (41.9)
≥30 years	299 (29.2)	168 (24.6)	131 (38.4)
Gender			
Female	677 (66.1)	677 (99.1)	0
Male	344 (33.6)	3 (0.4)	341 (100)
Other	3 (0.3)	3 (0.4)	0
Educational attainment[1]			
Low	5 (0.5)	1 (0.1)	4 (1.2)
Medium	158 (15.4)	95 (13.9)	63 (18.5)
High	857 (83.7)	585 (85.7)	272 (79.8)
Prefer not to say	4 (0.4)	2 (0.3)	2 (0.6)
Usually has sex[2] with			
Someone from the opposite sex	910 (88.9)	600 (87.8)	310 (90.9)
Someone from the same sex	82 (8.0)	56 (8.2)	26 (7.6)
Both	25 (2.4)	24 (3.5)	1 (0.3)
Never had sex[2]	6 (0.6)	3 (0.4)	3 (0.9)
Not listed	1 (0.1)	0	1 (0.3)
Relationship			
Yes	768 (75)	488 (71.4)	280 (82.1)
No/unsure/prefer not to say	256 (25.0)	195 (28.6)	61 (17.9)
Number of sexual[2] partners in past year			

≤ 1	678 (66.2)	435 (63.7)	244 (71.6)
2-10	325 (31.8)	236 (34.6)	89 (26.1)
>10	18 (1.8)	10 (1.5)	8 (2.3)
Using contraception			
No	89 (8.7)	50 (7.3)	39 (11.4)
Yes	935 (91.3)	633 (92.7)	302 (88.6)
If yes, which method[3]			
External condoms	350 (37.4)	222 (35.1)	128 (42.4)
Hormonal IUD	311 (33.3)	223 (35.2)	88 (29.1)
Contraceptive pill	253 (27.1)	175 (27.7)	78 (25.8)
I make sure my partner uses contraception	28 (3.0)	-	28 (9.3)
Cycle tracking	41 (4.4)	32 (5.1)	9 (3.0)
Copper IUD	73 (7.8)	49 (7.7)	24 (7.9)
Withdrawal	48 (5.1)	30 (4.7)	18 (6.0)
Morning after pill	27 (2.9)	20 (3.2)	7 (2.3)
Contraceptive implant	15 (1.6)	9 (1.4)	6 (2.0)
Sterilization	16 (1.7)	8 (1.3)	8 (2.6)
Contraceptive/vaginal ring	12 (1.3)	7 (1.1)	5 (1.7)
Internal condoms	7 (0.7)	5 (0.8)	2 (0.7)
Contraceptive injection	4 (0.4)	2 (0.3)	2 (0.7)
Contraceptive patch	3 (0.3)	2 (0.3)	1 (0.3)
Diaphragm	1 (0.1)	0 (0.0)	1 (0.3)

[1] Low: primary education or pre-vocational secondary education; Medium: lower/upper general secondary education, or secondary vocational education; High: higher professional education or university.

[2] Sex is broadly defined, such as manual sex, oral sex, penis-in-vagina sex, or anal sex.

[3] Multiple answers were allowed. Participants were asked about their and/or their partner's contraceptive methods, which is why men also reported contraceptives such as the contraceptive pill or IUDs. Given that some participants in the study were couples, there might be overlapping data in contraception methods. Cycle tracking includes periodic abstinence, calendar method, no penis-in-vagina sex mid-cycle, or using a fertility app.

3.2   Contraceptive satisfaction, responsibility, and knowledge

The mean satisfaction score of those using contraception (n=935) was 7.2 out of 10. Both women and men rated their contraception as satisfactory, with women being significantly less satisfied than men with their current method (Table 2). The mean sense of responsibility (n=1,024) was 8.4 out of 10. Women reported a significantly higher sense of responsibility towards contraceptive use compared to men (Table 2). The mean percentage of correct answers in the contraceptive knowledge quiz was 73.7%. Of all teams, 94.5% had 50% or more correct answers (Supplementary Table 3). The all-male teams had significantly lower knowledge (61.1%) than all-female (77.5%) or mixed (74.6%) teams (Table 2).

Table 2. *Mean scores of satisfaction, responsibility and knowledge for women and men separately at Lowlands festival, The Netherlands, 2023.*

Variable	Mean (SD)	Estimate (β)	95% Confidence Interval
Satisfaction (n=935)			
Women	7.0 (1.9)	-0.61	(-0.87; -0.35)
Men	7.7 (1.8)	Ref.	
Responsibility (n=1,024)			
Women	8.7 (1.9)	1.01	(0.74; 1.27)
Men	7.7 (2.3)	Ref.	
Knowledge (n=292 teams)			
All-male teams (n=35)	61.1 (17.9)	Ref.	
All-female teams (n=126)	77.5 (14.2)	16.79	(10.83; 22.75)
Mixed teams (n=131)	74.6 (15.9)	13.91	(7.94; 19.88)

Note: multivariate regression corrected for age, educational attainment, and relationship status. For educational attainment, low, medium and 'prefer not to say' were combined into one category due to the small number of participants in these categories.

3.3   Contraceptive views

We identified five themes (Figure 2). We defined the themes broadly, reflecting the exploratory nature of the study. The thematization is not intended to be rigid or mutually exclusive but aims to provide a framework for understanding the range of diverse views. Illustrative quotes per theme are displayed in Table 3.

Many participants expressed concerns about physical and mental **side effects** (Theme A) of contraception, often in a negative tone. These included general worries or personal experiences such as mood swings and heavier bleeding. At the same time, some noted perceived benefits, like menstrual regulation and reduced acne.

Various practical challenges related to the **fallibility of contraception** (Theme B) were also mentioned. Condoms were often described as unpleasant or unromantic interruptions during sex, and some participants reported issues with effectiveness, such as breakage or leakage. Other inconveniences included the painful insertion of an IUD, high costs of contraception, and forgetfulness—like missing a pill.

Further, there was a strong call for more **shared responsibility** (Theme C) in contraceptive use. Participants felt that the burden of side effects and costs disproportionately affects women. They

emphasized the importance of open communication and shared decision-making. Some advocated for
male contraceptive options, though others were skeptical about men's willingness to use them. Positive
examples of shared responsibility were also mentioned, such as using condom negotiation as a way to
signal consent.

Participants also expressed a clear **need for information** (Theme D). Questions were raised
about lesser-known methods like the internal condom and diaphragm, and many felt underinformed
about side effects. There was also a call for more research into non-hormonal alternatives and for
improved comprehensive sexuality education.

Participants also described **dismissive attitudes from healthcare providers during contraceptive**
counseling (Theme E). Others shared frustrations about not being taken seriously when seeking advice,
such as a participant whose concerns about an implant were dismissed with a referral to a psychologist.

Finally, participants voiced **resistance to hormonal methods** (Theme F), citing experienced side
effects or perceived long-term risks, such as potential impacts on fertility. These writings were often
related to other themes. For instance, some participants emphasized the importance of having their
concerns about hormonal contraception taken seriously by HCPs.

[Figure 2, see end of doc]

Table 3. *Illustrative quotes for each theme from the writings on the wall activity at Lowlands festival, The*
Netherlands, 2023.

Theme	Illustrative quotes
Side effects (A)	"Stupid side effects"
	"I feel heavier and more down with contraception"
	"Ups and downs, stable and unstable"
	"Being on your period for 3 months after getting the Implanon SUCKS"
	"What I like about the pill: Always taking it continuously"
	"I don't know if I experience any side effects, I use it since puberty"
Fallibility of contraception (B)	"Never had so much pain in my life EVER!" [about IUD insertion]
	"Contraception is too expensive for women with a low income"
	"Why does it [the condom] break so often? Then you need to do another STI test…"
	"I always forget taking the pill"
	"What a huge effort to think about it every single day"
Shared responsibility (C)	"Why is so much responsibility placed on women?"
	"Men, don't ask if she's on the pill after having sex, it's your responsibility too"
	"Would men take the pill if they read about all the side effects first?"

	"The costs of contraception too often rest with women" "I find it convenient how some men use 'Should I get a condom?' as consent for 'Do you feel like penetration now?'" "Stay in control yourself" [about condom as method to take responsibility]
Information needs (D)	"There is too little information about which contraception a woman/man can use" "Teach everyone about contraception! Not just 'girls'" "Make contraception a topic to openly talk about, for her <u>and</u> him" "I want more info about the internal condom!"
Dismissive contraceptive counselling of healthcare providers (E)	"A doctor prescribes you the pill at 16 but tells you 0 side effects" "Take women more seriously!!! Why do I have to go to the doctor 10 times before something is actually done?!" "Doctors are people too"
Resistance to hormones (F)	"Get rid of those hormones in your body!" "I don't want hormones that make me extra grumpy" "F*ck the normalized side effects of the pill!" "HormoNOOOOOO" "Anti-hormones, they are unnatural" "Hormones suck, but sometimes they're fun too, so just confusing"

4   Discussion

In this study, we applied theater-based data collection to study contraception. Our different

data collection methods complemented each other: while the quantitative methods showed sufficient

satisfaction, responsibility and knowledge, the writings on the wall revealed that participants also

perceived contraception as suboptimal due to (hormonal) side effects, the unequal burden on women,

and a lack of recognition in contraceptive concerns from healthcare providers.

Prior literature has suggested that drama methods may provide deeper insights into lived

experiences and embodied understanding (Myllyoja, 2024). While participants initially rated their

contraception as satisfactory in the survey, conducted outside the theater setting, more nuanced (and

primarily negative) views appeared on the writings on the wall towards the end of the study, after the

interaction with actors. The themes that emerged in the writings on the wall align with findings from

previous studies that used more traditional research methods, such as interviews (Alspaugh et al., 2020;

Ayoola et al., 2007; Baxter et al., 2011; Brown, 2015; D'Souza et al., 2022; Gollub et al., 2022; Hoga et al.,

2014; Kimport, 2018; Manzer et al., 2023; Pratt et al., 2014; Stewart et al., 2017; Wagner et al., 2024;

Williamson et al., 2009). However, compared to traditional data, participants' contributions to our

writings on the wall were noticeably more candid and nuanced – as if they were talking among friends –

thereby unveiling their lived experiences more vividly. For example, the language used in the writings on

the wall was informal and emotionally charged (for instance: "HormoNOOOO"), which may not be as

effectively captured in formal interview settings.

To the authors' knowledge, this is the first study to use a theater-based approach as a way to

evaluate views and lived experiences of contraception, yielding both quantitative and qualitative

insights in a group of over a thousand participants. Given its novelty, we will describe some reflections

on the use of theater as a data collection method to facilitate replication and support other researchers

who consider using this method. Firstly, we noticed that a theater-based approach offers unique ways to

increase recruitment and engagement. By making it fun, we lowered barriers for participation. This

resulted in some participants who were initially hesitant during recruitment– particularly men, who

sometimes felt they did not belong in research on contraception – to also take part. Ultimately, we

attracted a large number of participants, of whom one third were men. Through improvisation within

their character, the research team was able to tailor interactions to each individual, for example by

tapping into their curiosity about the theater setting of Club Womb, exploring why their participation

was relevant. Watching other festival visitors enter the experience, selecting enlarged contraceptive

methods and crawling through the door-sized vulva, in ponchos as condoms, increased the curiosity

about what was awaiting inside Club Womb, our theater research area. As the entrance also served as

the exit, festival visitors could see returning participants smiling and excitedly chatting about the study.

Meanwhile, the 'mascot', moving back and forth between the silent disco and the festival area, drew

festival visitors' attention and further increased participation. The setting was therefore very attractive

for participants to take part in the study.

Secondly, we notice an increased candor of participants to share their views and experiences

(Bleuer et al., 2018; Collins & Stockton, 2022; Myllyoja, 2024; Rossiter et al., 2008). The theatrical

playfulness visibly helped participants feel at ease, as we noticed them chatting to strangers already at

the start, when the host was welcoming them into the experience. As the study progressed, participants

became increasingly open, for example during the knowledge quiz, they laughed at the sexual health

doctor's jokes and started sharing personal experiences. When they then entered the club area,

participants danced together while reflecting on their contraceptive views and experiences. They

interacted with each other in their writings and drawings, and by talking to each other. Humorous one-

liners of the 'DJ/MC' amplified the sense that there were no limits on what could be shared. The text on

the writings on the wall showed that participants used their own language, as if among friends, which

they might less likely use in an in-depth interview. Therefore, theater-based data collection may help

reduce social desirability bias.

295   Our study shows the potential of theater-based research in evaluating contraceptive views,

evoking more open and authentic responses than is possible in traditional research. Nevertheless, this

study also has some limitations. As a result of our study setting, our sample consisted of relatively highly

educated festival participants and was therefore not representative of the Dutch population.

Additionally, results from the contraceptive knowledge quiz reflect shared rather than individual

knowledge and may therefore overestimate actual knowledge levels. Finally, as we did not collect data

on the participants' experiences of the theater-based research methods – wanting to first pilot this

method rather than setting up a formal evaluation in addition to what we were already asking of

participants – we were unable to provide an evaluation beyond our own perspectives as researchers and

actors in the project. More research is needed to further investigate the potential contribution of

theater-based data collection to research, especially regarding 1) the inclusion of populations that are

not reached by traditional research methods, and 2) the type of research questions for which this type

of data collection is most suitable.

308   5   Conclusion

This study applied theater-based data-collection to explore contraceptive views among adult women

and men. Regarding contraceptive satisfaction, the survey, conducted outside the theater-setting, and

the writings on the wall, towards the end of the theater-setting, complemented each other. While

participants appeared relatively satisfied with their contraception in the survey, the writings on the wall

evolved into a collection of candid, critical expressions, highlighting negative experiences with various

aspects of contraception, suggesting clear points for improvement. We believe theater may have further

encouraged participants to express themselves in their own words, as shown on the writings on the wall.

Theater-based data collection may therefore complement traditional research, fostering more inclusive,

engaging, and dynamic investigations of sexual and reproductive health, while capturing people's lived

experiences.

Acknowledgements

We gratefully acknowledge Lowlands Science 2023 for the organization. In addition, we would like to acknowledge all participants who took the time to participate in research while attending a music festival. Furthermore, we would like to thank Ineke van der Vlugt, program manager Rutgers, Expert Centre on Sexuality, for providing feedback on the questionnaire and knowledge quiz. Finally, we want to acknowledge the research team for their creativity in designing this project and for the energy they brought in the execution of the project during the festival weekend: Bart, Eveline, Jan, Nena, Rosa, Saskia, Teun, and Vincent.

Data availability statement

Anonymized survey, quiz data and syntaxes are openly available in the EUR Data Repository at (https://figshare.com/s/cc145e475ad1ba294d14).

Disclosure of interest

The authors report there are no competing interests to declare that are relevant to the content of this article.

Ethical statement

All participants provided digital informed consent, and the research protocol received approval from the Ethics Review Committee of Erasmus School of Social and Behavioral Sciences (ETH2223-0634).

Funding

This study was made possible by grants from the Netherlands Organization for Health Research and Development (ZonMw) Unintended Pregnancy Research Program, grant number 554002008 (Project UP), 554002012 (No Pink Clouds), 554002006 (RISE UP), 554001014 (Samen Groeien 010), the Vital Cities and Citizens initiative and the Erasmus School of Social and Behavioral Sciences Dragons' Den 2023. Hanan El Marroun was supported by the Stichting Volksbond Rotterdam, the Netherlands Organization for Scientific Research (NWO) Aspasia grant (No. 015.016.056), and the European Union's Horizon 2020 Research and Innovation Program (HappyMums, Grant agreement 101057390). Hilmar Bijma is supported by GGD-GHOR Netherlands (the Dutch national Association of GGD's (Regional Public Health Services) and GHOR (Regional Medical Emergency Preparedness and Planning)). The funders had no role in the design and conduct of the study or the writing of this article.

References

Alspaugh, A., Barroso, J., Reibel, M., & Phillips, S. (2020). Women's contraceptive perceptions, beliefs, and attitudes: An integrative review of qualitative research. *Journal of Midwifery & Women's Health*, *65*(1), 64–84. 10.1111/jmwh.12992

Ayoola, A. B., Nettleman, M., & Brewer, J. (2007). Reasons for unprotected intercourse in adult women. *Journal of Women's Health*, *16*(3), 302–310. 10.1089/jwh.2007.0210

Baxter, S., Blank, L., Guillaume, L., Squires, H., & Payne, N. (2011). Views regarding the use of contraception amongst young people in the UK: a systematic review and thematic synthesis. *The European Journal of Contraception & Reproductive Health Care*, *16*(3), 149–160. 10.3109/13625187.2011.556762

Bleuer, J., Chin, M., & Sakamoto, I. (2018). Why theater-based research works? Psychological theories from behind the curtain. *Qualitative Research in Psychology*, *15*(2-3), 395–411. 10.1080/14780887.2018.1430734

Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, *3*(2), 77–101. 10.1191/1478088706qp063oa

Brown, S. (2015). 'They think it's all up to the girls': Gender, risk and responsibility for contraception. *Culture, Health & Sexuality*, *17*(3), 312–325. 10.1080/13691058.2014.950983

Burke, H. M., Packer, C. A., Spector, H. L., & Hubacher, D. (2019). Opportunity, satisfaction, and regret: Trying long-acting reversible contraception in a unique scientific circumstance. *Women & Health*, *59*(3), 266–280. 10.1080/03630242.2018.1478363

Collins, C. S., & Stockton, C. (2022). The theater of qualitative research: The role of the researcher/actor. *International Journal of Qualitative Methods*, *21*, 160940692211031. 10.1177/16094069221103109

D'Souza, P., Bailey, J. V., Stephenson, J., & Oliver, S. (2022). Factors influencing contraception choice and use globally: A synthesis of systematic reviews. *The European Journal of Contraception & Reproductive Health Care*, *27*(5), 364–372. 10.1080/13625187.2022.2096215

de Graaf, H., Kraan, Y., Oldenhof, A., & Joemmanbaks, F. (2024). *Monitor seksuele gezondheid 2023* https://rutgers.nl/wp-content/uploads/2024/09/Rapport_monitor_seksuele_gezondheid_2023.pdf

Dixon, S. C., Herbert, D. L., Loxton, D., & Lucke, J. C. (2014). 'As many options as there are, there are just not enough for me': Contraceptive use and barriers to access among Australian women. *The European Journal of Contraception & Reproductive Health Care*, *19*(5), 340–351. 10.3109/13625187.2014.919380

Docter-Loeb, H. (2024). Lowlands: Where science meets music. *Nature*, *634*(8032), 251–253. 10.1038/d41586-024-03151-5

Fleckman, J., Brown, K., Lederer, A., Stoltman, S., & Craft, T. (2023). Tackling oppressive beliefs and sexual violence on college campuses: Evaluation of an innovative theater-based intervention. *Health Education & Behavior*, *50*(3), 318–327. 10.1177/10901981221120391

Gollub, E. L., Beauvais, S., & Roye, C. (2022). College-attending young men's sexual and reproductive health knowledge, attitudes and practices. *Journal of American College Health*, *70*(3), 706–716. 10.1080/07448481.2020.1762609

Haynes, M. C., Ryan, N., Saleh, M., Winkel, A. F., & Ades, V. (2017). Contraceptive knowledge assessment: Validity and reliability of a novel contraceptive research tool. *Contraception*, *95*(2), 190–197. 10.1016/j.contraception.2016.09.002

Hoga, L. A., Rodolpho, J. R., Sato, P. M., Nunes, M. C., & Borges, A. L. (2014). Adult men's beliefs, values, attitudes and experiences regarding contraceptives: a systematic review of qualitative studies. *Journal of Clinical Nursing*, *23*(7-8), 927–939. 10.1111/jocn.12262

Kimport, K. (2018). More than a physical burden: Women's mental and emotional work in preventing
pregnancy. *The Journal of Sex Research*, *55*(9), 1096–1105. 10.1080/00224499.2017.1311834
Knowles, J., & Cole, A. (2008). *Handbook of the arts in qualitative research: Perspectives, methodologies,*
examples, and issues. SAGE Publications, Inc. https://doi.org/10.4135/9781452226545
Kramer, R. D., Higgins, J. A., Everett, B., Turok, D. K., & Sanders, J. N. (2022). A prospective analysis of the
relationship between sexual acceptability and contraceptive satisfaction over time. *American*
Journal of Obstetrics and Gynecology, *226*(3), 396.e391–396.e311. 10.1016/j.ajog.2021.10.008
Le Guen, M., Schantz, C., Regnier-Loilier, A., & de La Rochebrochard, E. (2021). Reasons for rejecting
hormonal contraception in Western countries: A systematic review. *Social Science & Medicine*,
284, 114247. 10.1016/j.socscimed.2021.114247
Manzer, J. L., Berndt, V. K., & Bell, A. V. (2023). Intersections between culture, contraception, knowledge
and the body in the USA. *Culture, Health & Sexuality*, *25*(9), 1164–1179.
10.1080/13691058.2022.2141330
Marzi, T., Adembri, C., Vignozzi, L., Innocenti, B., Cruciata, M. A., & Lippi, D. (2025). Medicine at theater:
A tool for well-being and health-care education. *BMC Medical Education*, *25*(1).
10.1186/s12909-025-06793-9
Myllyoja, J. (2024). Applying drama methods in foresight: Supporting futures agency and creating in-
depth futures knowing. *Futures*, *156*, 103305. https://doi.org/10.1016/j.futures.2023.103305
Pratt, R., Stephenson, J., & Mann, S. (2014). What influences contraceptive behaviour in women who
experience unintended pregnancy? A systematic review of qualitative research. *Journal of*
Obstetrics and Gynaecology, *34*(8), 693–699. 10.3109/01443615.2014.920783
Rossiter, K., Kontos, P., Colantonio, A., Gilbert, J., Gray, J., & Keightley, M. (2008). Staging data: Theater
as a tool for analysis and knowledge transfer in health research. *Social Science & Medicine*, *66*(1),
130–146. 10.1016/j.socscimed.2007.07.021
Schneider-Kamp, A., & Takhar, J. (2023). Interrogating the pill: Rising distrust and the reshaping of health
risk perceptions in the social media age. *Social Science & Medicine*, *331*, 116081.
https://doi.org/10.1016/j.socscimed.2023.116081
Sense. *Anticonceptie-quiz*. https://sense.info/nl/condoom-pil-en-meer/anticonceptie-quiz
Steinberg, J. R., Marthey, D., Xie, L., & Boudreaux, M. (2021). Contraceptive method type and
satisfaction, confidence in use, and switching intentions. *Contraception*, *104*(2), 176–182.
10.1016/j.contraception.2021.02.010
Stewart, M., Ritter, T., Bateson, D., McGeechan, K., & Weisberg, E. (2017). Contraception - what about
the men? Experience, knowledge and attitudes: A survey of 2438 heterosexual men using an
online dating service. *Sexual Health*, *14*(6), 533–539. 10.1071/SH16235
Tourangeau, R., & Yan, T. (2007). Sensitive questions in surveys. *Psychological Bulletin*, *133*(5), 859–883.
10.1037/0033-2909.133.5.859
Wagner, B., Brogan, N., & Cleland, K. (2024). "A lot of my friends don't know how it works": Student
activists describe gaps in college students' emergency contraception knowledge. *Womens*
Health Issues, *34*(1), 66–71. 10.1016/j.whi.2023.07.002
Williamson, L. M., Parkes, A., Wight, D., Petticrew, M., & Hart, G. J. (2009). Limits to modern
contraceptive use among young women in developing countries: a systematic review of
qualitative research. *Reproductive Health*, *6*(1), 3. 10.1186/1742-4755-6-3

Figures

[Because the image contains photographs of people, readers can request access from the
corresponding author]

Figure 1. *The stages of the study at Lowlands festival, The Netherlands, 2023.*

Figure 2. *Schematic overview of overlapping themes in the writings on the wall at Lowlands festival, The*
Netherlands, 2023. Larger circles represent themes that were mentioned more frequently and indicate
greater overlap. Subthemes are in italics.