Title: Association between Discrimination in Healthcare and Doctor Visits Over Time
Short Title: Discrimination in Healthcare and Doctor Visits Over Time

Authors: Michael D. Green, Ph.D.;[1] Qing Yang, Ph.D.;[2] Hanzhang Xu, Ph.D.;[2,3] Radha Dhingra, M.D., Ph.D.;[4] Heather R. Farmer, Ph.D.;[5] M. Alan Brookhart;[4] Ann Marie Navar, M.D., Ph.D.;[6] Emily C. O'Brien, Ph.D.;[4] Roland J. Thorpe Jr. Ph. D;[1,7] Matthew E. Dupre, Ph.D.[4,8]

Affiliation:

1) Department of Health, Behavior, and Society, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
2) Duke University School of Nursing, Duke University, Durham, NC
3) Center for the Study of Aging and Human Development, Duke University, Durham, NC
4) Department of Population Health Sciences, Duke University School of Medicine, Durham, NC
5) Department of Human Development and Family Sciences, University of Delaware, Newark, DE
6) Department of Medicine, Division of Cardiology, University of Texas Southwestern, Dallas, TX
7) Johns Hopkins Alzheimer's Disease Resource Center for Minority Aging Research, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
8) Department of Sociology, Duke University, Durham, NC

Acknowledgements: This work was presented at the 2025 Gerontological Society of America Annual Meeting and won the "Health Science Section – Person in Training Award" for the best student paper presentation.

Funding: The research was supported by a National Institute on Aging (NIA) grant (PI: Dupre, R01AG069938), a NIA Diversity Supplement Award (PI: Dupre, R01AG069938-02S1), and a NIA pre-doctoral fellowship (PI: Green, F99AG088695). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Corresponding Author Contact Information:
Michael D. Green, Ph.D.
michaeldgreenphd@gmail.com
ORCID - 0000-0002-4982-8154

Word Count: 3,352

Credit Author Statement- Michael D. Green: Conceptualization, Funding acquisition, Data Curation, Software Writing - Original Draft, Writing - Review & Editing, Visualization, **Qing Yang:** Methodology, Supervision, Writing - Review & Editing. **Hanzhang Xu:** Writing - Review & Editing. **Radha Dhingra**: Writing - Review & Editing. **Heather R. Farmer:** Writing - Review & Editing. **M. Alan Brookhart:** Writing - Review & Editing. **Ann Marie Navar:** Writing - Review & Editing. **Emily C. O'Brien:** Writing - Review & Editing. **Roland J. Thorpe Jr.:** Writing - Review & Editing. **Matthew E. Dupre:** Funding acquisition, Investigation, Methodology, Writing - Original Draft, Writing - Review & Editing.

Abstract

Objective: Discrimination in healthcare can disrupt trust. However, it is unknown whether

discrimination is associated with the frequency and distribution of healthcare utilization. We

assessed how perceived discrimination in healthcare settings is associated with longitudinal

patterns of doctor visits among middle-aged and older adults in the United States.

Study Setting and Design: Prospective cohort study of US adults who were followed every 2

years for up to 12 years. Discrimination in healthcare was self-reported at baseline ("receiving

poorer treatment than others from doctors or hospitals") and dichotomized (any vs. never). The

primary outcome was the patterns of doctor visits over follow-up identified using group-based

trajectory models (GBTM).

Data source: Health and Retirement Study (HRS) from 2008-2020.

Principle findings: Among 13,422 participants (mean age 63.0 years [±8.0]), approximately

19.4%, reported experiencing discrimination in healthcare at baseline. Discrimination in

healthcare was more prevalent among participants with more visits, greater disease burden,

and greater social vulnerability. Five major patterns of doctor visits over time were identified:

"Low-Stable" (~2 visits/year, 35.5%), "Moderate-Stable" (~4.5 visits/year, 33.3%), "Increasing"

(~4.5 to 9 visits/year, 13.7%), "Decreasing" (~13.5 to 6 visits/year, 10.3%), and "Persistently-

High" (~11+ visits/year, 6.3%). Compared to participants with Low-Stable doctor visits, those

who reported discrimination were significantly more likely to exhibit Decreasing (relative risk

ratio [RRR]=1.38; 95% CI, 1.17–1.63) and Persistently-High (RRR=1.62; 95% CI, 1.30–2.03)

patterns of doctor visits.

Conclusions: Discrimination in healthcare settings is associated with greater numbers of doctor

visits, indicating potential opportunities to improve quality of care.

<u>Keywords</u> Healthcare Discrimination, Discrimination in Healthcare, Perceived Discrimination,

Group Based Trajectory Model (GBTM), Healthcare Utilization, Doctor Visits

What is known on this topic

• Perceived discrimination in healthcare is a potential barrier to quality care and is

frequently associated with poor health outcomes and patient mistrust in clinical care.

What this study found

• Middle-aged and older adults who perceived discrimination in healthcare settings were

significantly more likely to have high-frequency patterns of doctor visits.

• This association persisted after accounting for sociodemographic factors, behavioral

factors, and health status.

Introduction

Experiencing discrimination in healthcare is a barrier to equitable healthcare delivery,

and limits the possibility of optimal health outcomes.[1-3] From a patient standpoint, negative

experiences in healthcare can erode trust in providers and the healthcare system,[4] which in

turn, can influence how patients engage with health systems and the management of their

health conditions.[3,5,6] Despite this, the longitudinal association between experiencing

discrimination and healthcare utilization is unknown.

Discrimination in healthcare is when people are treated unfairly or overlooked because

of who they are or who others think they are.[7] Reasons that people attribute experiences of

discrimination can include sociodemographic characteristics such as age, sex, and race, as well

as social class, sexual orientation, or other factors.[8] Race, ethnicity, and disability are among

the factors most commonly associated with unfair treatment in healthcare settings.[9,10] Patients

who experience discrimination in healthcare settings may exhibit distinct patterns in their doctor

visits over time;[11] and these differences could stem from various reasons.[12-15] For example,

discrimination may signal poor quality of care leading to unmet medical needs, worsening

health-status, and, as a result, more frequent or fragmented healthcare utilization.[16] Additionally,

individual health needs significantly influence healthcare-seeking behaviors.[17,18] Middle-aged

and older adults have greater healthcare needs.[19,20] These adults also exhibit a high likelihood

of experiencing discrimination in healthcare settings.[21,22] Therefore, it is crucial to consider both

sociodemographic characteristics that are associated with perceived discrimination and

markers of health status that need different treatment intensities,[11,23-25] where treatment plans

and the number of visits will differ for conditions with varying etiologies and prognoses.[26]

Identifying heterogenous patterns of healthcare utilization will help us better understand the

potential implications of discrimination.

Using the largest ongoing prospective cohort study of middle-aged and older adults in

the United States, we examined whether individuals who reported discrimination in healthcare

had distinct patterns of doctor visits over time. To achieve this objective, we first identified the

major patterns of doctor visits occurring during follow-up. We then assessed how perceived

discrimination was associated with the likelihood of exhibiting specific patterns of doctor visits.

Finally, the associations were assessed while adjusting for a range of sociodemographic,

behavioral, and health-related factors.

117
118 **Methods**
119

Study Participants

Our analysis used data from the Health and Retirement Study (HRS), the largest

ongoing nationally representative longitudinal study of U.S. adults over age 50. The HRS is

sponsored by the National Institute on Aging (NIA U01AG009740) and is conducted by the

University of Michigan.[27] The HRS has accumulated over three decades of data on more than

40,000 individuals since its launch in 1992. The core survey for the HRS is conducted every 2

years, and comprehensive details on its methodology and response rates are documented

elsewhere.[28] Beginning in 2006, the HRS selected a random half-sample of respondents to

collect detailed self-reported psychosocial data.[29,30] Subsequent psychosocial data were

collected from the remaining half-sample in 2008, and follow-up data collection occurred every

four years after each half-sample initially provided data through 2020.

The study was limited to adults aged 50-80 at baseline who responded to the question

about their exposure to discrimination in healthcare settings from 2008 to 2020 (n=17,642). We

examined the patterns of doctor visits for individuals up to age 80 at baseline to limit the

influence of healthcare utilization related to end-of-life care at advanced ages. We excluded

individuals who did not report at least one doctor visit or hospitalization at baseline (n=1,619) to

capture those with potential exposure to discrimination in a healthcare setting. We also

excluded individuals who provided only baseline data and lacked longitudinal follow-up

(n=2,257). Participants with missing data on study covariates (n=344; ~2.5%) were also

excluded from the analyses. The final analytic sample included 13,422 individuals who provided

at least two waves of follow-up data with non-missing doctor visit information over the 14 years

of follow-up. All HRS participants provided written informed consent. Our study was approved

by the Duke University Health System Institutional Review Board (Pro00108869).

<u>Discrimination in Healthcare Settings</u>

Perceived discrimination in healthcare settings was measured using an item from the

Everyday Discrimination Scale (EDS), a validated instrument designed to capture the frequency

of discriminatory experiences across various daily contexts, including interactions in retail,

dining, and other social environments.[31] In 2008, the HRS incorporated an additional item into

the psychosocial survey specifically focused on discrimination in healthcare settings, which

asked respondents, "In your day-to-day life how often have any of the following things

happened to you: You receive poorer service or treatment than other people from doctors or

hospitals." Due to the limited variability in the reported frequency of experiencing healthcare

discrimination, we dichotomized the responses to distinguish those who had never encountered

discrimination (coded 0) from those who had (coded 1)—which included "less than once a year,"

"a few times a year," "a few times a month," "at least once a week," and "almost every day"—an

approach consistent with previous research.[32,33]

The primary analyses included self-reported exposure to discrimination in healthcare at

baseline. However, we recognize that it is difficult to disentangle the temporality of the

association between reported discrimination in healthcare settings and numbers of doctor visits.

To further account for this, we conducted additional analyses that included discrimination in

health as a time-varying exposure (as detailed below*).*

Outcome

Participants were asked at each wave if they had visited a doctor in the past two years

and, if so, how many times. Preliminary analyses assessed the full range of reported doctor

visits (0-900) and several winsorized thresholds were evaluated for upper limits (i.e., 20+, 30+,

40+, and 50+ visits) to account for extreme/uncommon values. Our final analyses included

counts of doctor visits from 0-50+ with approximately 2.92% (n=390) of participants reporting

more than 50 doctor visits in a reported wave.

170
Covariates

Analyses included the participants' baseline sociodemographic characteristics, health

behaviors, and clinical characteristics to account for factors that could influence exposure to

discrimination and/or patterns of healthcare utilization. Sociodemographic factors included age

(in years), gender (female or male), self-reported race and ethnicity (Non-Hispanic White, Non-

Hispanic Black, or Hispanic), educational attainment (in years), total household wealth based on

assets per household member (continuous, log-transformed), marital status (currently

married/partnered or not), and health insurance status (insured or uninsured). Health behaviors

included smoking status (never smoker, former smoker, or current smoker), daily alcohol

consumption (0 drinks, 1-2 drinks, or 3+ drinks), and physical inactivity (physically active or not).

Measures of health status included number of difficulties with activities of daily living ([ADL]s;

count), body mass index (BMI), and self-reported doctor diagnoses of high blood pressure,

diabetes, heart disease, stroke, cancer, and arthritis.

Statistical Analysis

We use group-based trajectory models (GBTM) to examine underlying heterogeneity in

the longitudinal patterns of doctor visits during follow-up. The GBTM were estimated with the *traj*

package in Stata 18.5.[34] This data-driven approach allowed for the identification of distinct

subgroups within the population that exhibited similar trajectories of doctor visits over time.[35,36,37]

The optimal number of groups was identified based on assessments of Bayesian information

criteria (BIC), average posterior probabilities (AvePP), odds of correct classification (OCC), 95%

confidence intervals (CI), and entropy.[38,39] In our preliminary analyses, we assessed the

distributions of doctor visits and determined that zero-inflated Poisson models were most

appropriate for modeling the counts of doctor visits over time.

Three sets of sensitivity analyses were also conducted. First, we examined mortality as

a potential source of non-random attrition using the "dropout" function in *traj*.[36,40] Mortality over

the 14-year follow-up period was generally low (16.6%) and did not change the results. Second,

we evaluated the results using baseline and time-varying indicators of discrimination in our

models. Approximately 82% of participants reported the same exposure to discrimination over

the entire follow-up period (with about 10% reporting no discrimination at baseline and

subsequently reporting discrimination in a later wave). The parameter estimates for the

trajectory models with the baseline indicator are available in (Supplementary Table 1). Because

there was substantially less follow-up data available for discrimination (measured quadrennially;

n=28,673 observations) that would omit nearly half of the doctor visits (measured biennially;

n=58,022 observations), our primary analyses included the baseline indicator of discrimination

in healthcare. Finally, we included time-varying adjustments for discrimination and major chronic

conditions at baseline (i.e. whether an individual had heart disease and/or cancer) to account for

potential changes in experiences of discrimination and health status.

Next, we used multinomial logistic regression models to examine the associations

between discrimination in healthcare settings reported at baseline and patterns of care visit over

time (trajectory group membership), in unadjusted (model 1) and covariate-adjusted (model 2)

models. For the multinomial logistic regression models, we used the most common trajectory of

doctor visits identified from GBTM as the reference group to compare with other trajectories. We

also examined differences among patients with similar starting (or ending) numbers of doctor

visits over time. As a final sensitivity analysis, we assessed the multinomial logistic regression

models in the trajectory model that included adjustments for time-varying discrimination and

comorbidities. *P* values < 0.05 were considered statistically significant. All analyses were

performed using Stata 18.5 (StataCorp LP, College Station, TX).

Results

The average age of our participants at baseline was 63.05 years (standard deviation ±

7.97), and 19.39% reported any discrimination in healthcare. Overall, reports of discrimination in

healthcare were significantly higher among adults who were relatively younger,

unmarried/unpartnered, uninsured, and belonged to a racial/ethnic minority group (Table 1).

Participants who reported discrimination were also more likely to have higher BMI, more

functional limitations, and higher prevalence of chronic conditions.

Patterns of Doctor Visits Over Time

We identified five distinct patterns of doctor visits over time (Figure 1). Baseline

characteristics of individuals across the five trajectory groups are presented in Table 2. The two

most common patterns of doctor visits were relatively stable over time, representing

approximately 2 yearly visits ("Low-Stable" group; 36.51%) and approximately 4.5 yearly doctor

visits ("Moderate-Stable" group; 33.26%). Two additional patterns showed significant changes in

doctor visits over time. The "Increasing" group (13.70%) indicated about 4.5 doctor visits each

year and gradually increased to more than 9 yearly doctor visits over the study period.

Conversely, the "Decreasing" group (10.28%) showed an inverse pattern, starting with over 13

yearly visits and gradually decreasing to a lower but still frequent level of about 6 yearly visits.

Lastly, there was a "Persistently-High" group (6.25%) which characterized participants with

consistently high numbers of doctor visits throughout the study period, with nearly monthly

visits.

Across all groups, the majority of individuals were Non-Hispanic White. The Low-Stable

group was the youngest (mean age 61.6), with the lowest rates of most chronic health

conditions and lowest levels of discrimination in healthcare (17.0%). The Moderate-Stable group

had relatively low experiences of discrimination (18.9%) but a greater burden of chronic

conditions. The Decreasing group reported greater discrimination in healthcare (24.0%) was the

oldest on average (64.4 years) and faced significant health challenges. The Increasing group

(13.7%) had a similar health profile to the Decreasing group, with a high prevalence of chronic

conditions but reported moderate discrimination (18.9%). The Persistently-High group (6.3%)

had the poorest health. This group had the lowest socioeconomic status and the highest

reported discrimination in healthcare (29.6%).

<u>Association between Discrimination and Trajectories</u>

<u>Table 3</u> presents results from the multinomial logistic regression models of the

association between discrimination in healthcare and doctor visit trajectory group membership.

Overall, in the unadjusted models, individuals who reported discrimination in healthcare settings

had significantly higher likelihood of following trajectories with more frequent doctor visits than

those who did not report discrimination. Compared with the Low-Stable group, results showed

that discrimination was associated with a 13% higher relative risk of being in the Moderate-

Stable group (relative risk ratio [RRR]=1.13, P=0.021), a 54% higher relative risk of being in the

Decreasing group (RRR=1.54, P<0.001), and more than double the relative risk of being in the

Persistently-High group (RRR=2.04, P<0.001). After adjusting for sociodemographic,

behavioral, and health-related covariates, the associations were attenuated but remained

significant for the Decreasing (RRR=1.38, P<0.001) and Persistently-High (RRR=1.62,

P<0.001) groups only. The association with the Moderate-Stable group was no longer

statistically significant (RRR=1.11, P=0.066).

Supplementary Figure 1 illustrates the results from the multinomial logistic regression

models using different reference groups that reflect similar levels of doctor visits at the start or

end of the trajectory (the estimates are provided in Supplementary Table 2). We highlight these

groups to better understand the association between discrimination and changing patterns of

doctor visits among individuals who begin or conclude care at comparable levels. When

comparing the Decreasing (~13.5 yearly visits decreasing to ~6) and the Moderate-Stable (~4

yearly visits), discrimination was associated with a 24% higher relative risk of membership in the

group which starts with high amounts of visits then decreasing. Discrimination was associated

with a 36% lower relative risk of starting with a high number of visits in the Increasing (~4.5

yearly visits increasing to ~9) compared to the Persistently-High Group (11+ Visits). Finally,

when considering time-varying changes in discrimination and major chronic conditions when

identifying trajectories of doctor visits (Supplementary Figure 2), the results for discrimination

were largely consistent and only partially attenuated (Supplementary Table 3 and

Supplementary Table 4).

280 Discussion
281
We identified 5 major longitudinal patterns of doctor visits in a large cohort of U.S.

middle-aged and older adults. Our findings suggest that discrimination in healthcare was

associated with exhibiting high-frequency patterns doctor visits over time. The associations

persisted after accounting for sociodemographic, behavioral, and health factors. Persistent

discrimination can hinder effective patient-provider communication, resulting in fragmented or

ineffective care that necessitates repeated visits.[41,42] Additionally, this pattern may reflect a

cycle where experiencing discrimination contributes to psychological stress and poor health

management, leading to more frequent doctor visits as health conditions progress. This

association indicates the potential for discrimination to deepen health inequities, as affected

individuals may experience both the direct harm of biased care and the indirect consequences

of unmanaged or poorly managed chronic conditions.

We found that middle-aged and older adults who had initially higher numbers of doctor

visits declined over long-term follow-up, with varying rates of decline. Discrimination in

healthcare was associated with distinct patterns of doctor visits over time. Those who reported

discrimination had a significantly higher probability of exhibiting patterns of doctor visits that

were initially high, but then declined int frequency (Decreasing) compared to those with

consistently low (Low-Stable) or moderate (Moderate-Stable) numbers of visits. Before

adjustment for sociodemographic and health factors, discrimination was also associated with a

greater probability of demonstrating high numbers of doctor visits (Persistently-High) compared

with those who started high and then largely declined (Decreasing). Because the Persistently-

High pattern of doctor visits was characterized by worse health status, it is plausible that

persistently high doctor visits may be more closely related to clinical needs than to

discrimination exposure. In comparison, the pattern of initially high but declining visits appears

more distinctively associated with experiencing discrimination in the healthcare setting.

However, because of the difficulty disentangling these factors, these conclusions warrant further

investigation.

One concern regarding discrimination in healthcare is that it may lead patients to avoid

care entirely. Our findings do not support this interpretation among middle-aged and older

adults. Individuals who reported discrimination were more likely to exhibit higher-frequency

patterns of visits rather than lower ones. However, a higher volume of visits does not

necessarily reflect effective care. Higher frequency of doctor visits among those who report

discrimination may reflect poorer quality care rather than higher quality treatment. It is also

important to acknowledge that the direction of this association may partly reflect exposure

opportunity. Individuals with more doctor visits have more encounters in which discrimination

can occur. Those with the greatest clinical need, and therefore the most in contact with the

healthcare system, may paradoxically be the most likely to encounter mistreatment.

Discrimination in healthcare, as measured in this study, represents a departure from the broader

framing of discrimination as a chronic psychological stressor experienced across various

settings.[43-45] By assessing reports of discrimination in healthcare specifically, these findings

illuminate potentially actionable targets for health systems seeking to improve care quality.

Assessing discrimination in healthcare can be effectively used as a tool to assess the

quality of care the services provided. Studies have shown that the cost of services provided to

individuals is higher for those who experience discrimination than those who do not.[46,47] Quality

of care is in the interest of both health systems focused on cost savings, alongside the doctors

who seek to improve their patients' health. Both patterns that exhibited the highest initial volume

of doctor visits (Decreasing and Persistently-High) demonstrated declining visits over the follow-

up period. Several interpretations of this general pattern are possible. Improvement in health

status could reduce the need for visits. For example, early frequent visits could reflect

successful management of a new diagnosis or recovery from an acute condition. The decline

could also reflect disengagement from care, potentially reacting to accumulated negative

healthcare experiences including discrimination, even as clinical need persists. Importantly,

even after such declines, these groups maintained frequent visits more so than other groups.

Further research is needed to help clarify whether the declines in visits reflect a change in their

health status, or whether discrimination is a driver of this change. Additionally, inclusion of more

racial/ethnic groups, consideration of broader age group categories, and time-varying covariates

would complement this analysis.

We believe our results are statistically rigorous and present a thoughtful analysis that

reflects real-world disease prognosis. GBTM has been critiqued as a method which can

generate spurious findings when Average Posterior Probability is the sole criterion for evaluating

model fit.[48] To address this concern we, assessed model fit using Average Posterior Probability,

entropy, Bayesian Information Criterion, Odds of Correct Classification, and 95% CIs. We

further evaluated our trajectories qualitatively by comparing our generated trajectories to

plausible care patterns. For example, if an individual had an onset of a condition such as kidney

failure, it would be plausible for them to start out with frequent visits and then require more

doctor visits over time. If an individual had a condition that was temporary, it would make sense

for them to start out with a high number of doctor visits, then decline.

Limitations

There are some limitations to consider when interpreting our results. We used a single

item to evaluate discrimation, instead of a full-scale assessment. While many scales have

been validated for population subgroups, with the intention of better understanding their

experiences, few have been constructed for broad use as a quality improvement tool.[49-54] A

psychometrically validated assessment, complemented by qualitative inquiry, could provide

additional context for impact of the perceived discrimination on subsequent doctor visits

patterns. Our sample is limited to those who responded to the HRS. While this study design and

sampling approach are robust, they may have excluded perspectives of marginalized individuals

or those who were unable to complete the survey. Furthermore, our approach may not have

adequately accounted for unobserved confounding, limiting our ability to draw causal

conclusions from our findings. We did not stratify by sociodemographic characteristics for our

analysis, which could cause different trends to emerge. For example, individuals who attribute

perceived discrimination to a disability are less likely to seek follow-up care.[55] Evaluating how

discrimination impacts the care trajectories of aging populations with disabilities and functional

limitations could be a crucial future consideration alongside other populations. The self-

perception of aging is both positively and negatively connected to perceived healthcare

discrimination.[56] We did not explore positive perception of aging in our analysis or ageism, but

future studies could consider this as a reason for divergence in discrimination experiences and

care-seeking behavior amongst middle-aged and older adults.

<u>Conclusion</u>

Our findings underscore the multifaceted impact of discrimination in healthcare on

patterns in healthcare utilization over time. This study highlights the persistent and potentially

costly implications of discriminatory experiences. It calls attention to the need for more nuanced,

inclusive research that captures the diversity of lived experiences across sociodemographic

lines. Addressing discrimination in healthcare is not only a moral imperative, but also a

pragmatic one. Discrimination in healthcare directly affects patient trust, care efficiency, and

system-wide costs. Interventions designed to reduce discriminatory practices could mitigate the

cycle of fragmented care and improve outcomes for those most affected.

References
380
1.   Tong JKC, Akpek E, Naik A, et al. Reporting of Discrimination by Health Care
Consumers Through Online Consumer Reviews. *JAMA Network Open*. 2022;5(2):e220715-
e220715. doi:10.1001/jamanetworkopen.2022.0715
2.   Borrell LN, Castor D. Benefits of Patient-Clinician Communication for Timely Health Care
Visits in Patients Perceiving Discrimination. *JAMA Network Open*. 2025;8(3):e250053-e250053.
doi:10.1001/jamanetworkopen.2025.0053
3.   Hausmann LRM, Hannon MJ, Kresevic DM, Hanusa BH, Kwoh CK, Ibrahim SA. Impact
of Perceived Discrimination in Healthcare on Patient-Provider Communication. *Medical Care*.
2011;49(7):626-633. doi:10.1097/MLR.0b013e318215d93c
4.   Williamson LD, A. SM, and Bigman CA. Does Discrimination Breed Mistrust? Examining
the Role of Mediated and Non-Mediated Discrimination Experiences in Medical Mistrust. *Journal*
of Health Communication. 2019/10/03 2019;24(10):791-799.
doi:10.1080/10810730.2019.1669742
5.   Rhoads K, Dohan D. Knowing it When You See it: The Need for Continuing Innovation in
Research on Healthcare Discrimination. *Journal of General Internal Medicine*. 2015/10/01
2015;30(10):1387-1388. doi:10.1007/s11606-015-3387-2
6.   Jafari Bidgoli M, Wang H, Macander C, Gregg A. Perceived Discrimination in Health
Care Settings and Care Delays in Patients With Diabetes and Hypertension. *JAMA Network*
Open. 2025;8(3):e250046-e250046. doi:10.1001/jamanetworkopen.2025.0046
7.   Togioka BM, Young E. Diversity and Discrimination in Health Care. *StatPearls*.
StatPearls Publishing
Copyright © 2025, StatPearls Publishing LLC.; 2025.
8.   Potter L, Zawadzki MJ, Eccleston CP, et al. The intersections of race, gender, age, and
socioeconomic status: Implications for reporting discrimination and attributions to discrimination.
Stigma and Health. 2019;4(3):264-281. doi:10.1037/sah0000099
9.   Abramson CM, Hashemi M, Sánchez-Jankowski M. Perceived discrimination in U.S.
healthcare: Charting the effects of key social characteristics within and across racial groups.
Preventive Medicine Reports. 2015/01/01/ 2015;2:615-621.
doi:https://doi.org/10.1016/j.pmedr.2015.07.006
10.   Rogers SE, Thrasher AD, Miao Y, Boscardin WJ, Smith AK. Discrimination in Healthcare
Settings is Associated with Disability in Older Adults: Health and Retirement Study, 2008–2012.
Journal of General Internal Medicine. 2015/10/01 2015;30(10):1413-1420. doi:10.1007/s11606-
015-3233-6
11.   Benjamin GC, DeVoe JE, Amankwah FK. Ending Unequal Treatment and Achieving
Optimal Health for All. *JAMA*. 2024;332(14):1143-1144. doi:10.1001/jama.2024.14268
12.   Aaron KJ, Colantonio LD, Deng L, et al. Cardiovascular Health and Healthcare
Utilization and Expenditures Among Medicare Beneficiaries: The REasons for Geographic And
Racial Differences in Stroke (REGARDS) Study. *Journal of the American Heart Association*.
2017;6(2):e005106. doi:doi:10.1161/JAHA.116.005106
13.   van der Vlegel M, Spronk I, Oude Groeniger J, et al. Health care utilization and health-
related quality of life of injury patients: comparison of educational groups. *BMC Health Services*
Research. 2021/09/19 2021;21(1):988. doi:10.1186/s12913-021-06913-3
14.   Jacobs MM, Ellis C. Heterogeneity among women with stroke: health, demographic and
healthcare utilization differentials. *BMC Women's Health*. 2021/04/17 2021;21(1):160.
doi:10.1186/s12905-021-01305-5
15.   Goel V, Rosella LC, Fu L, Alberga A. The Relationship Between Life Satisfaction and
Healthcare Utilization: A Longitudinal Study. *American Journal of Preventive Medicine*.
2018/08/01/ 2018;55(2):142-150. doi:https://doi.org/10.1016/j.amepre.2018.04.004

16.   Powell W, Jennifer R, Dinushika M, Irene Y, Allison J, and Corbie-Smith G. Medical
Mistrust, Racism, and Delays in Preventive Health Screening Among African-American Men.
Behavioral Medicine. 2019/04/03 2019;45(2):102-117. doi:10.1080/08964289.2019.1585327
17.   Bleser WK, Miranda PY, Jean-Jacques M. Racial/Ethnic Disparities in Influenza
Vaccination of Chronically Ill US Adults: The Mediating Role of Perceived Discrimination in
Health Care. *Medical Care.* 2016;54(6):570-577. doi:10.1097/mlr.0000000000000544
18.   Swift SL, Glymour MM, Elfassy T, et al. Racial discrimination in medical care settings
and opioid pain reliever misuse in a U.S. cohort: 1992 to 2015. *PLOS ONE.*
2019;14(12):e0226490. doi:10.1371/journal.pone.0226490
19.   Fisk AD, Czaja SJ, Rogers WA, Charness N, Sharit J. *Designing for older adults:*
Principles and creative human factors approaches. CRC press; 2020.
20.   Rowe JW, Fulmer T, Fried L. Preparing for Better Health and Health Care for an Aging
Population. *JAMA.* 2016;316(16):1643-1644. doi:10.1001/jama.2016.12335
21.   Doty M, Horstman C, Shah A, Ayo-Vaughan M, Zephyrin L. How Discrimination in health
care affects older Americans, and what health systems and providers can do: the
Commonwealth Fund; 2022. Issue Brief, April 21, 2022.
22.   Wyman MF, Shiovitz-Ezra S, Bengel J. *Ageism in the health care system: Providers,*
patients, and systems. Springer International Publishing; 2018.
23.   Ben J, Cormack D, Harris R, Paradies Y. Racism and health service utilisation: A
systematic review and meta-analysis. *PLOS ONE.* 2017;12(12):e0189900.
doi:10.1371/journal.pone.0189900
24.   MacIntosh T, Desai MM, Lewis TT, Jones BA, Nunez-Smith M. Socially-Assigned Race,
Healthcare Discrimination and Preventive Healthcare Services. *PLOS ONE.* 2013;8(5):e64522.
doi:10.1371/journal.pone.0064522
25.   Parker LJ, Hunte H, Ohmit A. Discrimination in Health Care: Correlates of Health Care
Discrimination Among Black Males. *American Journal of Men's Health.* 2017;11(4):999-1007.
doi:10.1177/1557988315585164
26.   Gonzales KL, Harding AK, Lambert WE, Fu R, Henderson WG. Perceived Experiences
of Discrimination in Health Care: A Barrier for Cancer Screening Among American Indian
Women with Type 2 Diabetes. *Women's Health Issues.* 2013/01/01/ 2013;23(1):e61-e67.
doi:https://doi.org/10.1016/j.whi.2012.10.004
27.   Health and Retirement Study. RAND HRS Longitudinal File (V1), RAND HRS 2020 Fat
File (E2A), RAND HRS 2018 Fat File (V2B), RAND HRS 2016 Fat File (V2C), RAND HRS 2014
Fat File (V2B), RAND HRS 2012 Fat File (V3A), RAND HRS 2010 Fat File (V6A), RAND HRS
2008 Fat File (V3A).
28.   Health and Retirement Study (HRS): A Longitudinal Study of Health, Retirement, and
Aging. Sponsored by the National Institute on Aging. Accessed 4/17/2022,
http://hrsonline.isr.umich.edu/
29.   Crimmins E, Faul J, Kim JK, et al. Documentation of biomarkers in the 2006 and 2008
Health and Retirement Study. *Ann Arbor, MI: Survey Research Center University of Michigan.*
2013;
30.   Crimmins E, Guyer H, Langa K, Ofstedal MB, Wallace R, Weir D. Documentation of
physical measures, anthropometrics and blood pressure in the Health and Retirement Study.
HRS Documentation Report DR-011. 2008;14(1-2):47-59.
31.   Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental
health: Socio-economic status, stress and discrimination. *Journal of health psychology.*
1997;2(3):335-351.
32.   Nguyen TT, Vable AM, Glymour MM, Nuru-Jeter A. Trends for Reported Discrimination
in Health Care in a National Sample of Older Adults with Chronic Conditions. *J Gen Intern Med.*
Mar 2018;33(3):291-297. doi:10.1007/s11606-017-4209-5

33.   Nguyen TT, Vable AM, Maria Glymour M, Allen AM. Discrimination in health care and
biomarkers of cardiometabolic risk in U.S. adults. *SSM - Population Health*. 2019/04/01/
2019;7:100306. doi:https://doi.org/10.1016/j.ssmph.2018.10.006
34.   L Jones B, Nagin D. A Stata Plugin for Estimating Group-Based Trajectory Models.
35.   Nagin DS. *Group-based modeling of development*. Harvard University Press; 2005:x,
201 p.
36.   Nagin DS, Odgers CL. Group-based trajectory modeling in clinical research. *Annu Rev*
Clin Psychol. 2010;6:109-38. doi:10.1146/annurev.clinpsy.121208.131413
37.   Hoang CT, Amin V, Behrman JR, Kohler H-P, Kohler IV. Heterogenous trajectories in
physical, mental and cognitive health among older Americans: Roles of genetics and life course
contextual factors. *SSM - Population Health*. 2023/09/01/ 2023;23:101448.
doi:https://doi.org/10.1016/j.ssmph.2023.101448
38.   Nagin DS. Group-based trajectory modeling: an overview. *Ann Nutr Metab*. 2014;65(2-
3):205-10. doi:10.1159/000360229
39.   Jones BL, Nagin DS. A Note on a Stata Plugin for Estimating Group-based Trajectory
Models. *Sociological Methods & Research*. 2013;42(4):608-613.
doi:10.1177/0049124113503141
40.   Haviland AM, Jones BL, Nagin DS. Group-based Trajectory Modeling Extended to
Account for Nonrandom Participant Attrition. *Sociological Methods & Research*. 2011;40(2):367-
390. doi:10.1177/0049124111400041
41.   Sorkin DH, Ngo-Metzger Q, De Alba I. Racial/Ethnic Discrimination in Health Care:
Impact on Perceived Quality of Care. *Journal of General Internal Medicine*. 2010/05/01
2010;25(5):390-396. doi:10.1007/s11606-010-1257-5
42.   Benjamins MR, Whitman S. Relationships between discrimination in health care and
health care outcomes among four race/ethnic groups. *Journal of Behavioral Medicine*.
2014/06/01 2014;37(3):402-413. doi:10.1007/s10865-013-9496-7
43.   Johnston DW, Lordan G. Discrimination makes me sick! An examination of the
discrimination–health relationship. *Journal of Health Economics*. 2012/01/01/ 2012;31(1):99-
111. doi:https://doi.org/10.1016/j.jhealeco.2011.12.002
44.   Gaston SA, Forde AT, Green M, Sandler DP, Jackson CL. Racial and Ethnic
Discrimination and Hypertension by Educational Attainment Among a Cohort of US Women.
JAMA Network Open. 2023;6(11):e2344707-e2344707.
doi:10.1001/jamanetworkopen.2023.44707
45.   Farmer HR, Ambroise AZ, Green MD, Dupre ME. Everyday discrimination and age-
related trajectories of blood pressure among Black and White middle-aged and older adults.
Stigma and Health. 2024;9(4):471-481. doi:10.1037/sah0000524
46.   Evans-Lacko S, Clement S, Corker E, et al. How much does mental health
discrimination cost: valuing experienced discrimination in relation to healthcare care costs and
community participation. *Epidemiology and Psychiatric Sciences*. 2015;24(5):423-434.
doi:10.1017/S2045796014000377
47.   Lippert-Rasmussen K. Cost-Effectiveness and the Avoidance of Discrimination in
Healthcare: Can We Have Both? *Cambridge Quarterly of Healthcare Ethics*. 2023;32(2):202-
215. doi:10.1017/S096318012200024X
48.   Mésidor M, Rousseau M-C, O'Loughlin J, Sylvestre M-P. Does group-based trajectory
modeling estimate spurious trajectories? *BMC Medical Research Methodology*. 2022/07/14
2022;22(1):194. doi:10.1186/s12874-022-01622-9
49.   Peek ME, Nunez-Smith M, Drum M, Lewis TT. Adapting the everyday discrimination
scale to medical settings: reliability and validity testing in a sample of African American patients.
Ethn Dis. Autumn 2011;21(4):502-9.

50.   López-Cevallos DF, Harvey SM. Psychometric Properties of a Healthcare Discrimination
Scale Among Young-Adult Latinos. *Journal of Racial and Ethnic Health Disparities*. 2019/06/01
2019;6(3):618-624. doi:10.1007/s40615-018-00560-x
51.   Sanford K, Pizzuto AE. The Healthcare Discrimination Experience Scale: Assessing a
Variable Crucial for Explaining Racial/Ethnic Inequities in Patient Activation and Health. *Journal*
of Racial and Ethnic Health Disparities. 2023/08/01 2023;10(4):1642-1652. doi:10.1007/s40615-
022-01350-2
52.   Almond AL. Measuring racial microaggression in medical practice. *Ethnicity & Health*.
2019/08/18 2019;24(6):589-606. doi:10.1080/13557858.2017.1359497
53.   Cruz D, Rodriguez Y, Mastropaolo C. Perceived microaggressions in health care: A
measurement study. *PLOS ONE*. 2019;14(2):e0211620. doi:10.1371/journal.pone.0211620
54.   Krieger N, Smith K, Naishadham D, Hartman C, Barbeau EM. Experiences of
discrimination: Validity and reliability of a self-report measure for population health research on
racism and health. *Social Science & Medicine*. 2005/10/01/ 2005;61(7):1576-1596.
doi:https://doi.org/10.1016/j.socscimed.2005.03.006
55.   Moscoso-Porras MG, Alvarado GF. Association between perceived discrimination and
healthcare–seeking behavior in people with a disability. *Disability and Health Journal*.
2018/01/01/ 2018;11(1):93-98. doi:https://doi.org/10.1016/j.dhjo.2017.04.002
56.   Sun N, Xu Z, Hua CL, et al. Self-perception of aging and perceived medical
discrimination. *Journal of the American Geriatrics Society*. 2023;71(10):3049-3058.
doi:https://doi.org/10.1111/jgs.18517
549
550

Figure 1. Estimated Trajectories of Doctor Visits Over Time in Middle-Aged and Older Adults, Health and Retirement Study 2008-2020 (N=13,422). *Note*: Estimates were derived using a group-based trajectory model that assumed a zero-inflated Poisson distribution for the number of doctor visits over time.

552 Table 1. Baseline Characteristics of by Discrimination Experience, Health and Retirement Study
553 2008-2020. (n=13,422)
554

Variable	No Discrimination, 10,836 (80.6%)	Discrimination, 2,606 (19.4%)	P-Value
Age	63.4 (8.0)	61.6 (7.7)	<0.001
Male	4,429 (40.9%)	1,121 (43.0%)	0.046
Race/Ethnicity			<0.001
Non-Hispanic White	8,027 (74.1%)	1,771 (68.0%)	
Non-Hispanic Black	2,007 (18.5%)	641 (24.6%)	
Hispanic	802 (7.4%)	194 (7.4%)	
Married/Partnered	7,015 (64.7%)	1,528 (58.6%)	<0.001
Education (Years)	13.2 (2.8)	13.2 (2.9)	0.265
Wealth (Log)	5.70 (3.44)	5.17 (3.85)	<0.001
Uninsured	654 (6.0%)	230 (8.8%)	<0.001
Smoking Status			<0.001
Never Smoker	4,779 (44.1%)	1,073 (41.2%)	
Former Smoker	4,450 (41.1%)	1,063 (40.8%)	
Current Smoker	1,607 (14.8%)	470 (18.0%)	
Drinking Status			<0.001
Moderate	2,380 (22.0%)	560 (21.5%)	
None	6,342 (58.5%)	1,618 (62.1%)	
Heavy	2,114 (19.5%)	428 (16.4%)	
Physically Inactive	5,564 (51.4%)	1,446 (55.5%)	<0.001
ADL Limitations	0.23 (0.78)	0.45 (1.09)	<0.001
BMI	29.1 (6.2)	30.2 (6.7)	<0.001
Hypertension	5,979 (55.2%)	1,503 (57.7%)	0.021
Heart Problem	1,986 (18.3%)	518 (19.9%)	0.068
Stroke	640 (5.9%)	187 (7.2%)	0.015
Diabetes	2,110 (19.5%)	611 (23.5%)	<0.001
Cancer	1,248 (11.5%)	333 (12.8%)	0.073
Arthritis	5,740 (53.0%)	1,492 (57.3%)	<0.001
Psychiatric Condition	1,652 (15.3%)	629 (24.1%)	<0.001
Note: Parenthesis next to continuous variables are standard deviations. P-values generated from Kruskal-Wallis tests and Chi-Square Tests. Abbreviations: BMI= Body Mass Index, ADL=Activities of Daily Living			

556 **Table 2.** Baseline Characteristics of Individuals in Identified Trajectory Groups of Doctor Visits,
557 Health and Retirement Study 2008-2020. (n=13,422)

Variables	Trajectory of Doctor Visits				
	Low-Stable	Moderate-Stable	Decreasing	Increasing	Persistently-High
n	4940 (36.51%)	4512 (33.26%)	1392 (10.28%)	1773 (13.70%)	825 (6.25%)
Discrimination in Healthcare	842 (17.0%)	851 (18.9%)	334 (24.0%)	335 (18.9%)	244 (29.6%)
Age	61.55 (7.88)	63.72 (7.83)	64.42 (7.93)	64.19 (8.00)	63.59 (7.97)
Male Gender	2250 (45.5%)	1776 (39.4%)	529 (38.0%)	705 (39.8%)	290 (35.2%)
Race and Ethnicity					
Non-Hispanic White	3480 (70.4%)	3377 (74.8%)	984 (70.7%)	1329 (75.0%)	628 (76.1%)
Non-Hispanic Black	1022 (20.7%)	852 (18.9%)	295 (21.2%)	321 (18.1%)	158 (19.2%)
Hispanic	438 (8.9%)	283 (6.3%)	113 (8.1%)	123 (6.9%)	39 (4.7%)
Married/Partnered	3218 (65.1%)	2933 (65.0%)	807 (58.0%)	1093 (61.6%)	492 (59.6%)
Education, Years	13.19 (2.83)	13.36 (2.72)	12.96 (2.83)	13.16 (2.84)	13.08 (2.74)
Wealth	5.64 (3.23)	5.80 (3.45)	5.28 (3.98)	5.43 (3.75)	5.09 (4.18)
Uninsured	434 (8.8%)	267 (5.9%)	54 (3.9%)	97 (5.5%)	32 (3.9%)
Smoking Status					
Never Smoker	2287 (46.3%)	1997 (44.3%)	547 (39.3%)	700 (39.5%)	321 (38.9%)
Former Smoker	1846 (37.4%)	1914 (42.4%)	620 (44.5%)	780 (44.0%)	353 (42.8%)
Current Smoker	807 (16.3%)	601 (13.3%)	225 (16.2%)	293 (16.5%)	151 (18.3%)
Drinking Status					
Moderate Consumption (1-2 drinks daily)	1266 (25.6%)	1019 (22.6%)	225 (16.2%)	324 (18.3%)	106 (12.8%)
No Consumption (0 drinks daily)	2641 (53.5%)	2633 (58.4%)	956 (68.7%)	1134 (64.0%)	596 (72.2%)
Heavy Consumption (3+ drinks daily)	1033 (20.9%)	860 (19.1%)	211 (15.2%)	315 (17.8%)	123 (14.9%)
Physically Inactive	2155 (43.6%)	2322 (51.5%)	921 (66.2%)	1040 (58.7%)	572 (69.3%)
ADL Limitations	0.07 (0.26)	0.12 (0.32)	0.25 (0.44)	0.18 (0.38)	0.34 (0.47)
Body Mass Index	28.42 (5.67)	29.56 (6.30)	29.85 (6.79)	29.84 (6.50)	30.97 (7.62)
Hypertension	2151 (43.5%)	2704 (59.9%)	941 (67.6%)	1130 (63.7%)	556 (67.4%)
Heart Problem	534 (10.8%)	849 (18.8%)	401 (28.8%)	428 (24.1%)	292 (35.4%)
Stroke	183 (3.7%)	252 (5.6%)	139 (10.0%)	137 (7.7%)	116 (14.1%)
Diabetes	551 (11.2%)	1013 (22.5%)	402 (28.9%)	482 (27.2%)	273 (33.1%)
Cancer	371 (7.5%)	527 (11.7%)	264 (19.0%)	256 (14.4%)	163 (19.8%)
Arthritis	1950 (39.5%)	2554 (56.6%)	942 (67.7%)	1155 (65.1%)	631 (76.5%)
Psychiatric Condition	468 (9.5%)	766 (17.0%)	324 (23.3%)	417 (23.5%)	306 (37.1%)

Continuous variables presented as means with standard deviations in parenthesis. Categorical and binary variables presented as counts with percentages in parenthesis. Wealth is log-adjusted. *Abbreviations:* ADL = Activities of Daily Living.

Table 2. Multinomial Estimates of Association Between Discrimination in Healthcare and Trajectory Group Membership of Middle Aged and Older Adults, Health and Retirement Study 2008-2020. (n=13,422)

Reference Group	Low-Stable											
Comparator Group	Moderate-Stable			Decreasing			Increasing			Persistently-High		
	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value
Unadjusted Model												
Discrimination in Healthcare	1.13	(1.02-1.26)	0.021	1.54	(1.33-1.77)	<0.001	1.13	(0.99-1.30)	0.079	2.04	(1.73-2.42)	<0.001
Adjusted Model												
Discrimination in Healthcare	1.11	(0.99-1.24)	0.066	1.38	(1.18-1.61)	<0.001	1.04	(0.89-1.20)	0.629	1.62	(1.35-1.95)	<0.001
Age	1.02	(1.02-1.03)	<0.001	1.02	(1.01-1.03)	<0.001	1.02	(1.02-1.03)	<0.001	1.01	(0.99-1.02)	0.379
Male	0.72	(0.66-0.79)	<0.001	0.74	(0.65-0.85)	<0.001	0.77	(0.68-0.87)	<0.001	0.68	(0.57-0.80)	<0.001
Non-Hispanic Black	0.81	(0.72-0.91)	<0.001	0.79	(0.66-0.94)	0.007	0.69	(0.59-0.81)	<0.001	0.62	(0.50-0.78)	<0.001
Hispanic	0.80	(0.67-0.95)	0.011	1.04	(0.81-1.33)	0.755	0.86	(0.68-1.08)	0.188	0.56	(0.39-0.82)	0.003
Married/Partnered	1.09	(0.99-1.20)	0.068	0.91	(0.79-1.04)	0.175	0.98	(0.87-1.11)	0.773	1.09	(0.92-1.30)	0.339
Education	1.07	(1.05-1.09)	<0.001	1.08	(1.05-1.11)	<0.001	1.08	(1.06-1.11)	<0.001	1.10	(1.07-1.14)	<0.001
Wealth	1.00	(0.99-1.01)	0.922	0.97	(0.96-0.99)	0.008	0.97	(0.96-0.99)	0.004	0.98	(0.96-1.00)	0.11
Uninsured	0.87	(0.73-1.03)	0.111	0.50	(0.37-0.68)	<0.001	0.77	(0.60-0.98)	0.034	0.50	(0.34-0.74)	0.001
Smoking Status												
Former Smoker	1.12	(1.02-1.23)	0.018	1.26	(1.10-1.45)	0.001	1.24	(1.09-1.41)	0.001	1.20	(1.00-1.43)	0.049
Current Smoker	0.97	(0.85-1.11)	0.651	1.11	(0.92-1.36)	0.280	1.23	(1.03-1.46)	0.024	1.17	(0.92-1.48)	0.212
Drinking Status												
No Consumption (0 drinks)	1.02	(0.91-1.13)	0.781	1.43	(1.20-1.69)	<0.001	1.25	(1.08-1.46)	0.003	1.73	(1.37-2.18)	<0.001
Heavy Consumption (3+ drinks)	1.01	(0.89-1.15)	0.878	1.14	(0.92-1.41)	0.237	1.15	(0.95-1.38)	0.144	1.46	(1.10-1.95)	0.009
Physically Inactive	1.08	(0.99-1.18)	0.085	1.62	(1.41-1.86)	<0.001	1.25	(1.11-1.42)	<0.001	1.61	(1.35-1.92)	<0.001
Activities of Daily Living	1.10	(1.03-1.18)	0.005	1.32	(1.23-1.43)	<0.001	1.22	(1.13-1.32)	<0.001	1.44	(1.33-1.56)	<0.001
BMI	1.01	(1.01-1.02)	0.001	1.00	(0.99-1.01)	0.608	1.01	(1.00-1.02)	0.090	1.01	(1.00-1.03)	0.058
Hypertension	1.57	(1.44-1.73)	<0.001	1.82	(1.58-2.10)	<0.001	1.65	(1.45-1.87)	<0.001	1.62	(1.35-1.94)	<0.001
Heart Problem	1.45	(1.28-1.65)	<0.001	2.13	(1.81-2.49)	<0.001	1.75	(1.51-2.04)	<0.001	2.65	(2.20-3.20)	<0.001
Stroke	1.09	(0.88-1.33)	0.435	1.38	(1.07-1.77)	0.013	1.21	(0.95-1.55)	0.121	1.74	(1.32-2.30)	<0.001
Diabetes	1.94	(1.72-2.20)	<0.001	2.21	(1.88-2.60)	<0.001	2.27	(1.95-2.64)	<0.001	2.61	(2.15-3.16)	<0.001
Cancer	1.40	(1.21-1.61)	<0.001	2.38	(1.99-2.86)	<0.001	1.72	(1.44-2.06)	<0.001	2.46	(1.98-3.06)	<0.001
Arthritis	1.53	(1.40-1.68)	<0.001	2.02	(1.76-2.32)	<0.001	1.97	(1.74-2.23)	<0.001	2.80	(2.33-3.38)	<0.001
Psychiatric Condition	1.78	(1.56-2.03)	<0.001	2.04	(1.72-2.42)	<0.001	2.337	(2.00-2.73)	<0.001	3.35	(2.77-4.06)	<0.001

Table 3. Multinomial Estimates of Association Between Discrimination in Healthcare and Trajectory Group Membership of Middle Aged and Older Adults, Health and Retirement Study 2008-2020. (n=13,422)

Reference Group	Low-Stable											
Comparator Group	Moderate-Stable			Decreasing			Increasing			Persistently-High		
	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value
Unadjusted Model												
Discrimination in Healthcare	0.99	(0.89-1.10)	0.889	1.29	(1.11-1.50)	0.001	1.06	(0.93-1.22)	0.401	1.57	(1.32-1.87)	<0.001
Adjusted Model												
Discrimination in Healthcare	1.02	(0.92-1.14)	0.661	1.23	(1.06-1.44)	0.008	1.03	(0.90-1.19)	0.642	1.38	(1.15-1.65)	0.001
Age	1.02	(1.01-1.02)	<0.001	1.02	(1.01-1.03)	0.001	1.02	(1.01-1.02)	<0.001	1.01	(1.00-1.02)	0.257
Male	0.73	(0.67-0.79)	<0.001	0.74	(0.64-0.84)	<0.001	0.74	(0.66-0.84)	<0.001	0.65	(0.54-0.77)	<0.001
Non-Hispanic Black	0.82	(0.73-0.92)	0.001	0.80	(0.67-0.95)	0.010	0.75	(0.64-0.87)	<0.001	0.67	(0.53-0.83)	<0.001
Hispanic	0.80	(0.67-0.95)	0.009	0.99	(0.77-1.27)	0.95	0.9	(0.72-1.12)	0.349	0.66	(0.46-0.93)	0.018
Married/Partnered	1.20	(1.09-1.32)	<0.001	0.93	(0.81-1.06)	0.287	0.96	(0.85-1.08)	0.522	1.04	(0.88-1.24)	0.622
Education	1.08	(1.07-1.10)	<0.001	1.07	(1.04-1.10)	<0.001	1.08	(1.06-1.11)	<0.001	1.10	(1.06-1.13)	<0.001
Wealth	1.01	(0.99-1.02)	0.385	0.98	(0.96-1.00)	0.019	0.99	(0.97-1.00)	0.154	0.98	(0.96-1.00)	0.097
Uninsured	0.85	(0.72-1.00)	0.055	0.5	(0.37-0.68)	<0.001	0.76	(0.61-0.96)	0.023	0.47	(0.31-0.70)	<0.001
Smoking Status												
Former Smoker	1.10	(1.01-1.21)	0.034	1.28	(1.11-1.47)	0.001	1.2	(1.06-1.36)	0.003	1.19	(1.00-1.42)	0.050
Current Smoker	0.93	(0.82-1.06)	0.258	1.15	(0.95-1.39)	0.159	1.09	(0.92-1.30)	0.296	1.19	(0.94-1.50)	0.153
Drinking Status												
No Consumption (0 drinks)	1.03	(0.93-1.15)	0.535	1.44	(1.22-1.71)	<0.001	1.2	(1.04-1.38)	0.014	1.65	(1.33-2.06)	<0.001
Heavy Consumption (3+ drinks)	0.95	(0.84-1.08)	0.437	1.26	(1.02-1.56)	0.029	1.2	(1.01-1.43)	0.037	1.29	(0.98-1.70)	0.065
Physically Inactive	1.05	(0.97-1.15)	0.235	1.57	(1.37-1.80)	<0.001	1.14	(1.01-1.28)	0.029	1.49	(1.26-1.77)	<0.001
Activities of Daily Living	1.09	(1.03-1.16)	0.006	1.34	(1.25-1.43)	<0.001	1.2	(1.12-1.29)	<0.001	1.38	(1.28-1.49)	<0.001
BMI	1.01	(1.00-1.02)	0.04	0.99	(0.98-1.00)	0.25	1.00	(0.99-1.01)	0.537	1.01	(0.99-1.02)	0.362
Hypertension	1.08	(0.99-1.18)	0.094	1.09	(0.95-1.25)	0.22	1.00	(0.89-1.13)	0.995	0.87	(0.73-1.03)	0.099
Heart Problem	0.95	(0.85-1.06)	0.341	1.14	(0.97-1.33)	0.121	1.05	(0.90-1.21)	0.544	1.29	(1.07-1.57)	0.009
Stroke	0.83	(0.70-1.00)	0.05	0.85	(0.66-1.10)	0.213	0.84	(0.67-1.06)	0.140	0.97	(0.72-1.29)	0.823
Diabetes	1.25	(1.12-1.39)	<0.001	1.17	(1.00-1.38)	0.055	1.28	(1.11-1.48)	0.001	1.19	(0.98-1.46)	0.080
Cancer	0.92	(0.81-1.05)	0.238	1.42	(1.19-1.70)	<0.001	1.01	(0.85-1.19)	0.934	1.30	(1.04-1.61)	0.021
Arthritis	1.02	(0.93-1.11)	0.654	1.18	(1.03-1.35)	0.017	1.26	(1.12-1.42)	<0.001	1.50	(1.26-1.79)	<0.001
Psychiatric Condition	1.08	(0.96-1.21)	0.212	1.02	(0.86-1.21)	0.815	1.29	(1.12-1.50)	0.001	1.58	(1.31-1.91)	<0.001