1 <u>Title</u>: Impact of Discrimination in Healthcare on Patterns of Doctor Visits Over Time
2
3 <u>Short Title</u>: Impact of Discrimination on Patterns of Doctor Visits
4

5 <u>Authors</u>: Michael D. Green, B.A.;[1] Qing Yang, Ph.D.;[2] Hanzhang Xu, Ph.D.;[2,3] Radha Dhingra,
6 M.D., Ph.D.;[1] Heather R. Farmer, Ph.D.;[4] Roland J. Thorpe Jr. Ph. D;[5] Matthew E. Dupre,
7 Ph.D.[1,6]
8
9 <u>Affiliation</u>:
10
1) Department of Population Health Sciences, Duke University School of Medicine,
Durham, NC
2) Duke University School of Nursing, Duke University, Durham, NC
3) Center for the Study of Aging and Human Development, Duke University, Durham, NC
4) Department of Human Development and Family Sciences, University of Delaware,
Newark, DE
5) Johns Hopkins Alzheimer's Disease Resource Center for Minority Aging Research,
Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
6) Department of Sociology, Duke University, Durham, NC
20
21

22 <u>Acknowledgements</u>: N/A
23

24 <u>Funding</u>: The research was supported by a National Institute on Aging (NIA) grant (PI: Dupre,
25 R01AG069938), a NIA Diversity Supplement Award (PI: Dupre, R01AG069938-02S1), and a
26 NIA pre-doctoral fellowship (PI: Green, F99AG088695). The content is solely the responsibility
27 of the authors and does not necessarily represent the official views of the National Institutes of
28 Health
29

<u>Corresponding Author Contact Information</u>:
Michael D. Green
Department of Population Health Sciences,
Duke University School of Medicine
215 Morris St,
Durham, NC, 27701
michael.d.greeen@duke.edu
ORCID - 0000-0002-4982-8154
38

39 <u>Word Count</u>: 2,994
40

Key Points

Question: Is perceived discrimination in healthcare settings associated with long-term patterns of doctor visits among middle-aged and older adults?

Findings: In this prospective cohort study of United States adults aged 50 to 80, perceived discrimination in healthcare was associated with increased likelihood of belonging to high-frequency doctor visit patterns over 14 years of follow-up, even after adjusting for sociodemographic and health-related factors.

Meaning: Discrimination in healthcare may contribute to fragmented or intensive healthcare utilization patterns, underscoring the need to address discrimination as part of efforts to improve care quality and equity.

Importance: Discrimination in healthcare disrupts trust and can negatively influence patients.

However, the impact of experiencing discrimination on long-term patterns of healthcare

utilization is unknown.

Objective: To assess how perceived discrimination in healthcare settings is associated with

longitudinal patterns of doctor visits among middle-aged and older adults in the United States.

Design: Prospective cohort study of US adults in the Health and Retirement Study (HRS) who

were followed every 2 years for up to 14 years (2008–2020).

Setting: US population-based cohort study.

Participants: Adults aged 50-80 years at baseline who completed the HRS psychosocial

questionnaire, answered the question on discrimination in healthcare settings, had complete

covariate information, and had at least two waves of follow-up data.

Exposure: Discrimination in healthcare settings was measured at baseline as the self-reported

frequency of "receiving poorer treatment than others from doctors or hospitals." Responses

were dichotomized to indicate any experience of discrimination.

Main Outcomes and Measures: Number of doctor visits in the prior two years were reported

by participants at each wave. Group-based trajectory models (GBTM) were used to identify

distinct longitudinal trajectories (patterns) of doctor visits over time. Multinomial logistic

regression models were used to examine associations between discrimination in healthcare and

trajectories of doctor visits while adjusting for sociodemographic, behavioral, and health-related

factors.

Results: Among 13,422 participants (mean age ~63.0 years [±8.0]), approximately 19.4%,

reported experiencing discrimination in healthcare at baseline. Results identified five major

patterns of doctor visits over time: "Low" (~2 visits/year, 35.5%), "Frequent" (~4.5 visits/year,

33.3%), "Frequent-to-High" (~4.5 to 9 visits/year, 13.7%), "High-to-Frequent" (~13.5 to 6

visits/year, 10.3%), and "High" (~11+ visits/year, 6.3%). Discrimination in healthcare was more

prevalent among participants with higher-utilization patterns and among those with greater

disease burden and social vulnerability. Compared to participants with Low doctor visits,

perceived discrimination was significantly greater in those with High-to-Frequent (relative risk

ratio [RRR]=1.38; 95% CI, 1.17–1.63; P<.001) and High (RRR=1.62; 95% CI, 1.30–2.03;

P<.001) patterns of doctor visits.

Conclusions and Relevance: Discrimination in healthcare settings is associated with greater

numbers of doctor visits, possibly indicating it as a marker of poor quality of care.

<u>Keywords (1-7)</u> Healthcare Discrimination, Discrimination in Healthcare, Perceived

Discrimination, Group Based Trajectory Model (GBTM), Healthcare Utilization, Doctor Visits

Introduction

Experiencing discrimination in healthcare is a barrier to equitable healthcare delivery,

and limits the possibility of optimal health outcomes.[1-3] From a patient-centered standpoint,

facing negative experiences in healthcare erodes patients' trust in their providers and the

healthcare system,[4] which in turn, potentially influences how patients engage with health

systems and the management of their health conditions.[3,5,6] Despite this, the long-term impact of

experiencing discrimination on healthcare utilization is unknown.

Discrimination in healthcare is when people are treated unfairly or overlooked because

of who they are or who others think they are.[7] Reasons that people attribute experiences of

discrimination can include sociodemographic characteristics such as age, sex, and race, as well

as social class, sexual orientation, or other factors.[8] Race, ethnicity, and disability are among

the factors most commonly associated with unfair treatment in healthcare settings.[15,16] Patients

who experience discrimination in healthcare settings may exhibit distinct patterns in their doctor

visits over time;[9] and these differences could stem from various reasons.[10-13] For example,

discrimination may signal poor quality of care leading to unmet medical needs, worsening

health-status, and, as a result, more frequent or fragmented healthcare utilization.[14] Additionally,

the health needs of individuals significantly influence their healthcare-seeking behaviors.[17,18]

Middle-aged and older adults have greater healthcare needs.[19,20] These adults also exhibit a

high likelihood of experiencing discrimination in healthcare settings.[21,22] Therefore, it is crucial

to consider both sociodemographic characteristics that are associated with varying reports of

discrimination and markers of health status that need different treatment intensities,[9,23-25] where

treatment plans and the number of visits will differ for conditions with varying etiologies and

prognoses.[26] Therefore, it is crucial to identify heterogenous patterns of healthcare utilization to

help understand the impact of discrimination.

Using the largest ongoing prospective cohort study of middle-aged and older adults in

the United States, we examined whether individuals who reported discrimination in healthcare

had distinct patterns of doctor visits over time. To achieve this objective, we first identified the

major patterns of doctor visits occurring during follow-up. We then assessed how perceived

discrimination was associated with the likelihood of exhibiting specific patterns of doctor visits.

Finally, the associations were assessed while adjusting for a range of sociodemographic,

behavioral, and health-related factors.

128
129 **Methods**
130

Study Participants

Our analysis used data from the Health and Retirement Study (HRS), the largest

ongoing nationally representative longitudinal study of U.S. adults over age 50. The HRS is

sponsored by the National Institute on Aging (NIA U01AG009740) and is conducted by the

University of Michigan.[29] The HRS has accumulated over three decades of data on more than

40,000 individuals since its launch in 1992. The core survey for the HRS is conducted every 2

years, and comprehensive details on its methodology and response rates are documented

elsewhere.[30] Beginning in 2006, the HRS selected a random half-sample of respondents to

collect detailed psychosocial data.[31,32] Subsequent psychosocial data were collected in the

remaining half-sample in 2008, and data collection occurred every four years after each half-

sample initially provided data through 2020.

The study was limited to adults aged 50-80 at baseline who responded to the question

about their exposure to discrimination in healthcare settings from 2008 to 2020 (n=17,642). We

examined the patterns of doctor visits for individuals up to age 80 at baseline to limit the

influence of healthcare utilization related to end-of-life care at advanced ages. We excluded

individuals who did not report at least one doctor visit or hospitalization at baseline (n=1,619) to

capture those with potential exposure to discrimination in a healthcare setting. We also

excluded individuals who provided only baseline data and lacked longitudinal follow-up

(n=2,257). Participants with missing data on study covariates (n=344; ~2.5%) were also

excluded from the analyses. The final analytic sample included 13,422 individuals who provided

at least two waves of follow-up data with non-missing doctor visits over the 14 years of follow-

up. All HRS participants provided written informed consent. Our study was approved by the

Duke University Health System Institutional Review Board (Pro00108869).

Discrimination in Healthcare Settings

Perceived discrimination in healthcare settings was measured using an item from the

Everyday Discrimination Scale (EDS), a validated instrument designed to capture the frequency

of discriminatory experiences across various daily contexts, including interactions in retail,

dining, and other social environments.[33] In 2008, the HRS incorporated an additional item into

the psychosocial survey specifically focused on discrimination in healthcare settings, which

asked respondents, "In your day-to-day life how often have any of the following things

happened to you: You receive poorer service or treatment than other people from doctors or

hospitals." Due to the limited variability in the reported frequency of experiencing healthcare

discrimination we dichotomized the responses to distinguish those who had never encountered

discrimination from those who had—categorizing "never" to represent "no experience of

healthcare discrimination" and combining all other response options ("less than once a year," "a

few times a year," "a few times a month," "at least once a week," "almost every day") into one to

represent "any experience of healthcare discrimination"—based on previously used

approaches.[34,35] This variable was employed as a time-constant exposure at baseline.

Outcome

Participants were asked at each wave if they had visited a doctor in the past two years

and, if so, how many times. Preliminary analyses assessed the full range of reported doctor

visits (0-900) and several winsorized thresholds were evaluated for upper limits (i.e., 20+, 30+,

40+, and 50+ visits) to account for extreme/uncommon values. Our final analyses included

counts of doctor visits from 0-50+ - with approximately 2.92% (n=390) of participants reporting

more than 50 doctor visits in a reported wave.

178
<u>Covariates</u>

Analyses included the participants' baseline sociodemographic characteristics, health

behaviors, and clinical characteristics to account for factors that could influence exposure to

discrimination and/or patterns of healthcare utilization. Sociodemographic factors included age

(in years), gender (female or male), race and ethnicity (Non-Hispanic White, Non-Hispanic

Black, or Hispanic), educational attainment (in years), total household wealth based on assets

per household member (log-transformed), marital status (currently married/partnered or not),

and health insurance status (insured or uninsured). Health behaviors included smoking status

(never smoker, former smoker, or current smoker), daily alcohol consumption (0 drinks, 1-2

drinks, or 3+ drinks), and physical inactivity (physically active or not). Measures of health status

included number of difficulties with activities of daily living ([ADL]s; count), body mass index

(BMI), and self-reported doctor diagnoses of high blood pressure, diabetes, heart disease,

stroke, cancer, and arthritis.

<u>Statistical Analysis</u>

We use group-based trajectory models (GBTM) to examine underlying heterogeneity in

the longitudinal patterns of doctor visits during follow-up. The GBTM were estimated with the *traj*

package in Stata 18.5.[36] This data-driven approach allowed for the identification of distinct

subgroups within the population that exhibited similar trajectories of doctor visits over time.[37,38,39]

The optimal number of groups was identified based on assessments of Bayesian information

criteria (BIC), average posterior probabilities (AvePP), odds of correct classification (OCC), 95%

confidence intervals (CI), and entropy.[40,41] In our preliminary analyses, we assessed the

distributions of doctor visits and determined that zero-inflated Poisson models were most

appropriate for modeling the counts of doctor visits over time. Preliminary analyses also

examined mortality as a source of non-random attrition using the "dropout" function in *traj*.[38,42]

Mortality over the 14-year follow-up period was generally low (16.6%) and did not change the

results. The parameter estimates for the final model are presented in Supplementary Table 1.

Next, we used multinomial logistic regression models to examine the associations

between discrimination in healthcare settings reported at baseline and patterns of care visit over

time (trajectory group membership), in both unadjusted (model 1) and covariate-adjusted (model

2) models. For the multinomial logistic regression models, we used the most common trajectory

of doctor visits identified from GBTM to compare with other trajectories. We then examined

additional comparisons between clinically informative trajectories of doctor visits. For instance, a

patient might have multiple comorbidities that require many doctor visits to develop a

comprehensive care management plan. Alternatively, a patient could have fewer initial visits, but

their needs may escalate over time. *P* values < 0.05 were considered statistically significant. All

analyses were performed using Stata 18.5 (StataCorp LP, College Station, TX).

216
Results

Patterns of Doctor Visits Over Time

We identified five distinct patterns of doctor visits over time (Figure 1). The two most

common patterns of doctor visits were relatively stable over time, representing approximately 2

yearly visits ("Low" group; 35.51%) and approximately 4.5 yearly doctor visits ("Frequent" group;

33.26%). Two additional patterns showed significant changes in doctor visits over time. The

"Frequent-to-High" group (13.70%) indicated about 4.5 doctor visits each year and gradually

increased to more than 9 yearly doctor visits over the study period. Conversely, the "High-to-

Frequent" group (10.28%) showed an inverse pattern, starting with over 13 yearly visits and

gradually decreasing to a lower but still frequent level of about 6 yearly visits. Lastly, there was

a "High" group (6.25%) which characterized participants with consistently high numbers of

doctor visits throughout the study period, with nearly monthly visits. In <u>Supplementary Figure 1</u>,

there was little variation for the shape of the distribution in responses for the discrimination

question.

The average age of our sample at is 63.05 (standard deviation ± 7.97), and 19.39%

reported any discrimination. Baseline characteristics of individuals across the five trajectory

groups are presented in <u>Table 1</u>. Across all groups, the majority of individuals were Non-

Hispanic White. The Low group was the youngest (mean age 61.6), with the lowest rates of

most comorbid health conditions and lowest levels of discrimination in healthcare (17.0%). The

Frequent group had relatively low experiences of discrimination (18.9%) but a greater burden of

chronic conditions. The High-to-Frequent group reported greater discrimination in healthcare

(24.0%) was the oldest on average (64.4 years) and faced significant health challenges. The

Frequent-to-High group (13.7%) had a similar health profile to the High-to-Frequent group, with

a high prevalence of chronic conditions but reported moderate discrimination (18.9%). The High

group (6.3%) had the poorest health. This group had the lowest socioeconomic status and the

highest reported discrimination in healthcare (29.6%).

<u>Association between Discrimination and Trajectories</u>

<u>Table 2</u> presents results from the multinomial logistic regression models of the

association between discrimination in healthcare and doctor visit trajectory group membership.

Overall, in the unadjusted models, individuals who reported discrimination in healthcare settings

had significantly higher likelihood of following trajectories with more frequent doctor visits than

those who did not report discrimination. Compared with the Low group, results showed that

discrimination was associated with a 13% higher relative risk of being in the Frequent group

(relative risk ratio [RRR]=1.13, P=0.021), a 54% higher relative risk of being in the High-to-

Frequent group (RRR=1.54, P<0.001), and more than double the relative risk of being in the

High group (RRR=2.04, P<0.001). After adjusting for sociodemographic, behavioral, and health-

related covariates, the associations were attenuated but remained significant for the High-to-

Frequent (RRR=1.38, *P*<0.001) and High (RRR=1.62, *P*<0.001) groups only. The association

with the Frequent group was no longer statistically significant (RRR=1.11, *P*=0.066).

Figure 2 illustrates the results from the multinomial logistic regression models using

different reference groups that reflect similar levels of doctor visits at the start or end of the

trajectory (the estimates are provided in Supplementary Table 2). We highlight these groups to

better understand the association between discrimination and changing patterns of doctor visits

among individuals who begin or conclude care at comparable levels. When comparing the High-

to-Frequent (~13.5 yearly visits decreasing to ~6) and the Frequent (~4 yearly visits),

discrimination was associated with a 24% higher relative risk of membership in the group which

starts with high amounts of visits then decreasing. Discrimination was associated with a 36%

lower relative risk of starting with a high number of visits in the Frequent-to-High (~4.5 yearly

visits increasing to ~9) compared to the High Group (11+ Visits).

267
268 **Discussion**
269
We identified 5 major longitudinal patterns of doctor visits in a large cohort of U.S.

middle-aged and older adults. Our findings suggest that discrimination in healthcare was

associated with exhibiting high-frequency patterns doctor visits over time. The associations

persisted after accounting for sociodemographic, behavioral, and health factors. Persistent

discrimination can hinder effective patient-provider communication, resulting in fragmented or

ineffective care that necessitates repeated visits.[27,43] Additionally, this pattern may reflect a

cycle where experiencing discrimination contributes to psychological stress and poor health

management, leading to more frequent doctor visits as health conditions progress. This

association indicates the potential for discrimination to deepen health inequities, as affected

individuals may experience both the direct harm of biased care and the indirect consequences

of unmanaged or poorly managed chronic conditions.

We identified that middle-aged and older adults who starts with higher doctor visits either

slightly decline to a still high level or sharply decline. Likelihood of exhibiting these patterns

varied for those who reported discrimination. Those who reported discrimination had a

significantly higher probability of exhibiting a High-to-Frequent pattern compared to the Low or

Frequent pattern. However, those who reported discrimination did not have a significant

association with the High pattern when compared to the High-to-Frequent pattern. Prior to

adjustment for sociodemographic and health factors, there was a higher likelihood of group

membership for the High pattern for those who report discrimination, but after considering these

factors, that association was attenuated. This is noteworthy because the High pattern was

associated with worse health-status, this might indicate that steepness of the decline is a

function of health status rather than experiencing discrimination.

Discrimination has a history of being evaluated as a chronic psychological stressor that

is a consequence of unequal social treatment.[44-46] Discrimination in healthcare as measured in

this study is a delineation from this framing as a general stressor experienced across various

settings, discrimination attributed to healthcare settings provide health systems an opportunity

to measure and modify their practices. Recently, healthcare dissatisfaction has been evaluated

alongside discrimination in healthcare as a marker of healthcare quality. These quality

measures are independently associated with an increased risk of dementia.[47] While

discrimination in healthcare can be stressful for patients, our findings suggest that individuals at

the highest risk of reported discrimination are also those who receive the most care. Notably,

the risk of discrimination is significantly higher than the risk associated with any of the

sociodemographic characteristics or the risk for most of the health-related factors.

Discrimination in healthcare can be effectively used as a tool to assess the quality-of-care

services provided. The cost of services provided to individuals is higher for those who

experience discrimination than those who do not.[48,49] Quality of care is in the interest of both

health systems focused on cost savings, alongside the doctors and patients who seek to

improve their patients' health. Further research into changes in their health status could clarify

whether the declines in visits reflect a change in their health status, or if the discrimination and

other factors they face are the cause of this change. Additionally, inclusion of more racial/ethnic

groups, consideration of broader age group categories, and time-varying covariates would

complement this analysis.

We believe our results are statistically rigorous and present a thoughtful analysis that

reflects real-world disease prognosis. GBTM has been critiqued as a method which can

generate spurious findings when Average Posterior Probability is the sole criterion for evaluating

model fit.[50] To address this concern we, assessed model fit using Average Posterior Probability,

entropy, Bayesian Information Criterion, Odds of Correct Classification, and 95% CIs. We

further evaluated our trajectories qualitatively by comparing our generated trajectories to

plausible care patterns. For example, if an individual had an onset of a condition such as kidney

failure, it would be plausible for them to start out with frequent visits and then require more

doctor visits over time. If an individual had a condition that was temporary, it would make sense

for them to start out with a high amount of doctor visits, then to decline to frequent.

<u>Limitations</u>

There are some limitations to consider when interpreting our results. We used a single

item to evaluate discrimination, instead of a full-scale assessment. To our knowledge, there are

no scales of discrimination in healthcare settings that have been psychometrically validated for

use in a diverse sample of middle-aged and older adults. While many scales have been

validated for population subgroups, with the intention of better understanding their experiences,

few have been constructed for broad use as a quality improvement tool.[51-56] A psychometrically

validated assessment, complemented by qualitative inquiry, could provide additional context for

impact of the perceived discrimination on subsequent doctor visits patterns. Our sample is

limited to those who responded to the HRS. While this study design and sampling approach are

robust, they may have excluded perspectives of marginalized individuals or those who were

unable to complete the survey. Furthermore, our approach may not have adequately accounted

for unobserved confounding, limiting our ability to draw causal conclusions from our findings.

We did not stratify by sociodemographic characteristics for our analysis, which could cause

different trends to emerge. For example, individuals who attribute perceived discrimination to a

disability are less likely to seek follow-up care.[57] Evaluating how discrimination impacts the care

trajectories of aging populations with disabilities and functional limitations, could be a crucial

future consideration alongside other populations. The self-perception of aging is both positively

and negatively connected to perceived healthcare discrimination.[58] We did not explore positive

perception of aging in our analysis or ageism, but future studies could consider this as a reason

for divergence in discrimination experiences and care-seeking behavior amongst middle-aged

and older adults.

Conclusion

Our findings underscore the multifaceted impact of discrimination in healthcare on

patterns in healthcare utilization over time. This study highlights the persistent and potentially

costly implications of discriminatory experiences. It calls attention to the need for more nuanced,

inclusive research that captures the diversity of lived experiences across sociodemographic

lines. Addressing discrimination in healthcare is not only a moral imperative, but also a

pragmatic one. Discrimination in healthcare directly affects patient trust, care efficiency, and

system-wide costs. Interventions designed to reduce discriminatory practices could mitigate the

cycle of fragmented care and improve outcomes for those most affected.

References

356
357
1.   Tong JKC, Akpek E, Naik A, et al. Reporting of Discrimination by Health Care
Consumers Through Online Consumer Reviews. *JAMA Network Open*. 2022;5(2):e220715-
e220715. doi:10.1001/jamanetworkopen.2022.0715
2.   Borrell LN, Castor D. Benefits of Patient-Clinician Communication for Timely Health Care
Visits in Patients Perceiving Discrimination. *JAMA Network Open*. 2025;8(3):e250053-e250053.
doi:10.1001/jamanetworkopen.2025.0053
3.   Hausmann LRM, Hannon MJ, Kresevic DM, Hanusa BH, Kwoh CK, Ibrahim SA. Impact
of Perceived Discrimination in Healthcare on Patient-Provider Communication. *Medical Care*.
2011;49(7):626-633. doi:10.1097/MLR.0b013e318215d93c
4.   Williamson LD, A. SM, and Bigman CA. Does Discrimination Breed Mistrust? Examining
the Role of Mediated and Non-Mediated Discrimination Experiences in Medical Mistrust. *Journal
of Health Communication*. 2019/10/03 2019;24(10):791-799.
doi:10.1080/10810730.2019.1669742
5.   Rhoads K, Dohan D. Knowing it When You See it: The Need for Continuing Innovation in
Research on Healthcare Discrimination. *Journal of General Internal Medicine*. 2015/10/01
2015;30(10):1387-1388. doi:10.1007/s11606-015-3387-2
6.   Jafari Bidgoli M, Wang H, Macander C, Gregg A. Perceived Discrimination in Health
Care Settings and Care Delays in Patients With Diabetes and Hypertension. *JAMA Network
Open*. 2025;8(3):e250046-e250046. doi:10.1001/jamanetworkopen.2025.0046
7.   Togioka BM, Young E. Diversity and Discrimination in Health Care. *StatPearls*.
StatPearls Publishing
Copyright © 2025, StatPearls Publishing LLC.; 2025.
8.   Potter L, Zawadzki MJ, Eccleston CP, et al. The intersections of race, gender, age, and
socioeconomic status: Implications for reporting discrimination and attributions to discrimination.
Stigma and Health. 2019;4(3):264-281. doi:10.1037/sah0000099
9.   Benjamin GC, DeVoe JE, Amankwah FK. Ending Unequal Treatment and Achieving
Optimal Health for All. *JAMA*. 2024;332(14):1143-1144. doi:10.1001/jama.2024.14268
10.   Aaron KJ, Colantonio LD, Deng L, et al. Cardiovascular Health and Healthcare
Utilization and Expenditures Among Medicare Beneficiaries: The REasons for Geographic And
Racial Differences in Stroke (REGARDS) Study. *Journal of the American Heart Association*.
2017;6(2):e005106. doi:doi:10.1161/JAHA.116.005106
11.   van der Vlegel M, Spronk I, Oude Groeniger J, et al. Health care utilization and health-
related quality of life of injury patients: comparison of educational groups. *BMC Health Services
Research*. 2021/09/19 2021;21(1):988. doi:10.1186/s12913-021-06913-3
12.   Jacobs MM, Ellis C. Heterogeneity among women with stroke: health, demographic and
healthcare utilization differentials. *BMC Women's Health*. 2021/04/17 2021;21(1):160.
doi:10.1186/s12905-021-01305-5
13.   Goel V, Rosella LC, Fu L, Alberga A. The Relationship Between Life Satisfaction and
Healthcare Utilization: A Longitudinal Study. *American Journal of Preventive Medicine*.
2018/08/01/ 2018;55(2):142-150. doi:https://doi.org/10.1016/j.amepre.2018.04.004
14.   Powell W, Jennifer R, Dinushika M, Irene Y, Allison J, and Corbie-Smith G. Medical
Mistrust, Racism, and Delays in Preventive Health Screening Among African-American Men.
Behavioral Medicine. 2019/04/03 2019;45(2):102-117. doi:10.1080/08964289.2019.1585327
15.   Abramson CM, Hashemi M, Sánchez-Jankowski M. Perceived discrimination in U.S.
healthcare: Charting the effects of key social characteristics within and across racial groups.
Preventive Medicine Reports. 2015/01/01/ 2015;2:615-621.
doi:https://doi.org/10.1016/j.pmedr.2015.07.006
16.   Rogers SE, Thrasher AD, Miao Y, Boscardin WJ, Smith AK. Discrimination in Healthcare
Settings is Associated with Disability in Older Adults: Health and Retirement Study, 2008–2012.

Journal of General Internal Medicine. 2015/10/01 2015;30(10):1413-1420. doi:10.1007/s11606-
015-3233-6
17.   Bleser WK, Miranda PY, Jean-Jacques M. Racial/Ethnic Disparities in Influenza
Vaccination of Chronically Ill US Adults: The Mediating Role of Perceived Discrimination in
Health Care. *Medical Care.* 2016;54(6):570-577. doi:10.1097/mlr.0000000000000544
18.   Swift SL, Glymour MM, Elfassy T, et al. Racial discrimination in medical care settings
and opioid pain reliever misuse in a U.S. cohort: 1992 to 2015. *PLOS ONE.*
2019;14(12):e0226490. doi:10.1371/journal.pone.0226490
19.   Fisk AD, Czaja SJ, Rogers WA, Charness N, Sharit J. *Designing for older adults:*
Principles and creative human factors approaches. CRC press; 2020.
20.   Rowe JW, Fulmer T, Fried L. Preparing for Better Health and Health Care for an Aging
Population. *JAMA.* 2016;316(16):1643-1644. doi:10.1001/jama.2016.12335
21.   Doty M, Horstman C, Shah A, Ayo-Vaughan M, Zephyrin L. How Discrimination in health
care affects older Americans, and what health systems and providers can do: the
Commonwealth Fund; 2022. Issue Brief, April 21, 2022.
22.   Wyman MF, Shiovitz-Ezra S, Bengel J. *Ageism in the health care system: Providers,*
patients, and systems. Springer International Publishing; 2018.
23.   Ben J, Cormack D, Harris R, Paradies Y. Racism and health service utilisation: A
systematic review and meta-analysis. *PLOS ONE.* 2017;12(12):e0189900.
doi:10.1371/journal.pone.0189900
24.   MacIntosh T, Desai MM, Lewis TT, Jones BA, Nunez-Smith M. Socially-Assigned Race,
Healthcare Discrimination and Preventive Healthcare Services. *PLOS ONE.* 2013;8(5):e64522.
doi:10.1371/journal.pone.0064522
25.   Parker LJ, Hunte H, Ohmit A. Discrimination in Health Care: Correlates of Health Care
Discrimination Among Black Males. *American Journal of Men's Health.* 2017;11(4):999-1007.
doi:10.1177/1557988315585164
26.   Gonzales KL, Harding AK, Lambert WE, Fu R, Henderson WG. Perceived Experiences
of Discrimination in Health Care: A Barrier for Cancer Screening Among American Indian
Women with Type 2 Diabetes. *Women's Health Issues.* 2013/01/01/ 2013;23(1):e61-e67.
doi:https://doi.org/10.1016/j.whi.2012.10.004
27.   Benjamins MR, Whitman S. Relationships between discrimination in health care and
health care outcomes among four race/ethnic groups. *Journal of Behavioral Medicine.*
2014/06/01 2014;37(3):402-413. doi:10.1007/s10865-013-9496-7
28.   Berke DS, Ruben MA, Liautaud MM, Meterko M, Kauth MR, Shipherd JC. Discrimination
exposure based on race, gender, and sexual orientation of veterans affairs health administration
patients. *Journal of Health Care for the Poor and Underserved.* 2022;33(2):714-725.
29.   Health and Retirement Study. RAND HRS Longitudinal File (V1), RAND HRS 2020 Fat
File (E2A), RAND HRS 2018 Fat File (V2B), RAND HRS 2016 Fat File (V2C), RAND HRS 2014
Fat File (V2B), RAND HRS 2012 Fat File (V3A), RAND HRS 2010 Fat File (V6A), RAND HRS
2008 Fat File (V3A).
30.   Health and Retirement Study (HRS): A Longitudinal Study of Health, Retirement, and
Aging. Sponsored by the National Institute on Aging. Accessed 4/17/2022,
http://hrsonline.isr.umich.edu/
31.   Crimmins E, Faul J, Kim JK, et al. Documentation of biomarkers in the 2006 and 2008
Health and Retirement Study. *Ann Arbor, MI: Survey Research Center University of Michigan.*
2013;
32.   Crimmins E, Guyer H, Langa K, Ofstedal MB, Wallace R, Weir D. Documentation of
physical measures, anthropometrics and blood pressure in the Health and Retirement Study.
HRS Documentation Report DR-011. 2008;14(1-2):47-59.

33.   Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental
health: Socio-economic status, stress and discrimination. *Journal of health psychology*.
1997;2(3):335-351.
34.   Nguyen TT, Vable AM, Glymour MM, Nuru-Jeter A. Trends for Reported Discrimination
in Health Care in a National Sample of Older Adults with Chronic Conditions. *J Gen Intern Med*.
Mar 2018;33(3):291-297. doi:10.1007/s11606-017-4209-5
35.   Nguyen TT, Vable AM, Maria Glymour M, Allen AM. Discrimination in health care and
biomarkers of cardiometabolic risk in U.S. adults. *SSM - Population Health*. 2019/04/01/
2019;7:100306. doi:https://doi.org/10.1016/j.ssmph.2018.10.006
36.   L Jones B, Nagin D. A Stata Plugin for Estimating Group-Based Trajectory Models.
37.   Nagin DS. *Group-based modeling of development*. Harvard University Press; 2005:x,
201 p.
38.   Nagin DS, Odgers CL. Group-based trajectory modeling in clinical research. *Annu Rev
Clin Psychol*. 2010;6:109-38. doi:10.1146/annurev.clinpsy.121208.131413
39.   Hoang CT, Amin V, Behrman JR, Kohler H-P, Kohler IV. Heterogenous trajectories in
physical, mental and cognitive health among older Americans: Roles of genetics and life course
contextual factors. *SSM - Population Health*. 2023/09/01/ 2023;23:101448.
doi:https://doi.org/10.1016/j.ssmph.2023.101448
40.   Nagin DS. Group-based trajectory modeling: an overview. *Ann Nutr Metab*. 2014;65(2-
3):205-10. doi:10.1159/000360229
41.   Jones BL, Nagin DS. A Note on a Stata Plugin for Estimating Group-based Trajectory
Models. *Sociological Methods & Research*. 2013;42(4):608-613.
doi:10.1177/0049124113503141
42.   Haviland AM, Jones BL, Nagin DS. Group-based Trajectory Modeling Extended to
Account for Nonrandom Participant Attrition. *Sociological Methods & Research*. 2011;40(2):367-
390. doi:10.1177/0049124111400041
43.   Sorkin DH, Ngo-Metzger Q, De Alba I. Racial/Ethnic Discrimination in Health Care:
Impact on Perceived Quality of Care. *Journal of General Internal Medicine*. 2010/05/01
2010;25(5):390-396. doi:10.1007/s11606-010-1257-5
44.   Johnston DW, Lordan G. Discrimination makes me sick! An examination of the
discrimination–health relationship. *Journal of Health Economics*. 2012/01/01/ 2012;31(1):99-
111. doi:https://doi.org/10.1016/j.jhealeco.2011.12.002
45.   Gaston SA, Forde AT, Green M, Sandler DP, Jackson CL. Racial and Ethnic
Discrimination and Hypertension by Educational Attainment Among a Cohort of US Women.
JAMA Network Open. 2023;6(11):e2344707-e2344707.
doi:10.1001/jamanetworkopen.2023.44707
46.   Farmer HR, Ambroise AZ, Green MD, Dupre ME. Everyday discrimination and age-
related trajectories of blood pressure among Black and White middle-aged and older adults.
Stigma and Health. 2024;9(4):471-481. doi:10.1037/sah0000524
47.   Aravena JM, Chen X, Levy BR. Association between experiencing low healthcare quality
and developing dementia. *Journal of the American Geriatrics Society*. 2024;72(7):2126-2132.
doi:https://doi.org/10.1111/jgs.18842
48.   Evans-Lacko S, Clement S, Corker E, et al. How much does mental health
discrimination cost: valuing experienced discrimination in relation to healthcare care costs and
community participation. *Epidemiology and Psychiatric Sciences*. 2015;24(5):423-434.
doi:10.1017/S2045796014000377
49.   Lippert-Rasmussen K. Cost-Effectiveness and the Avoidance of Discrimination in
Healthcare: Can We Have Both? *Cambridge Quarterly of Healthcare Ethics*. 2023;32(2):202-
215. doi:10.1017/S096318012200024X

50.   Mésidor M, Rousseau M-C, O'Loughlin J, Sylvestre M-P. Does group-based trajectory
modeling estimate spurious trajectories? *BMC Medical Research Methodology*. 2022/07/14
2022;22(1):194. doi:10.1186/s12874-022-01622-9
51.   Peek ME, Nunez-Smith M, Drum M, Lewis TT. Adapting the everyday discrimination
scale to medical settings: reliability and validity testing in a sample of African American patients.
Ethn Dis. Autumn 2011;21(4):502-9.
52.   López-Cevallos DF, Harvey SM. Psychometric Properties of a Healthcare Discrimination
Scale Among Young-Adult Latinos. *Journal of Racial and Ethnic Health Disparities*. 2019/06/01
2019;6(3):618-624. doi:10.1007/s40615-018-00560-x
53.   Sanford K, Pizzuto AE. The Healthcare Discrimination Experience Scale: Assessing a
Variable Crucial for Explaining Racial/Ethnic Inequities in Patient Activation and Health. *Journal
of Racial and Ethnic Health Disparities*. 2023/08/01 2023;10(4):1642-1652. doi:10.1007/s40615-
022-01350-2
54.   Almond AL. Measuring racial microaggression in medical practice. *Ethnicity & Health*.
2019/08/18 2019;24(6):589-606. doi:10.1080/13557858.2017.1359497
55.   Cruz D, Rodriguez Y, Mastropaolo C. Perceived microaggressions in health care: A
measurement study. *PLOS ONE*. 2019;14(2):e0211620. doi:10.1371/journal.pone.0211620
56.   Krieger N, Smith K, Naishadham D, Hartman C, Barbeau EM. Experiences of
discrimination: Validity and reliability of a self-report measure for population health research on
racism and health. *Social Science & Medicine*. 2005/10/01/ 2005;61(7):1576-1596.
doi:https://doi.org/10.1016/j.socscimed.2005.03.006
57.   Moscoso-Porras MG, Alvarado GF. Association between perceived discrimination and
healthcare–seeking behavior in people with a disability. *Disability and Health Journal*.
2018/01/01/ 2018;11(1):93-98. doi:https://doi.org/10.1016/j.dhjo.2017.04.002
58.   Sun N, Xu Z, Hua CL, et al. Self-perception of aging and perceived medical
discrimination. *Journal of the American Geriatrics Society*. 2023;71(10):3049-3058.
doi:https://doi.org/10.1111/jgs.18517
532
533

Figure 1. Estimated Trajectories of Doctor Visits Over Time in Middle-Aged and Older Adults, Health and Retirement Study 2008-2020 (N=13,422)

Note: The estimates were derived using a group-based trajectory model that assumes a zero-inflated Poisson distribution for the number of doctor visits over time.

 Table 1. Baseline Characteristics of Individuals in Identified Trajectory Groups of Doctor Visits,
536   Health and Retirement Study 2008-2020. (N=13,422)

Variables	Trajectory of Doctor Visits				
	Low	Frequent	High-to-Frequent	Frequent-to-High	High
n	4940 (35.51%)	4512 (33.26%)	1392 (10.28%)	1773 (13.70%)	825 (6.25%)
Discrimination in Healthcare	842 (17.0%)	851 (18.9%)	334 (24.0%)	335 (18.9%)	244 (29.6%)
Age	61.55 (7.88)	63.72 (7.83)	64.42 (7.93)	64.19 (8.00)	63.59 (7.97)
Male Gender	2250 (45.5%)	1776 (39.4%)	529 (38.0%)	705 (39.8%)	290 (35.2%)
Race and Ethnicity					
Non-Hispanic White	3480 (70.4%)	3377 (74.8%)	984 (70.7%)	1329 (75.0%)	628 (76.1%)
Non-Hispanic Black	1022 (20.7%)	852 (18.9%)	295 (21.2%)	321 (18.1%)	158 (19.2%)
Hispanic	438 (8.9%)	283 (6.3%)	113 (8.1%)	123 (6.9%)	39 (4.7%)
Married	3218 (65.1%)	2933 (65.0%)	807 (58.0%)	1093 (61.6%)	492 (59.6%)
Education, Years	13.19 (2.83)	13.36 (2.72)	12.96 (2.83)	13.16 (2.84)	13.08 (2.74)
Wealth	5.64 (3.23)	5.80 (3.45)	5.28 (3.98)	5.43 (3.75)	5.09 (4.18)
Uninsured	434 (8.8%)	267 (5.9%)	54 (3.9%)	97 (5.5%)	32 (3.9%)
Smoking Status					
Never Smoker	2287 (46.3%)	1997 (44.3%)	547 (39.3%)	700 (39.5%)	321 (38.9%)
Former Smoker	1846 (37.4%)	1914 (42.4%)	620 (44.5%)	780 (44.0%)	353 (42.8%)
Current Smoker	807 (16.3%)	601 (13.3%)	225 (16.2%)	293 (16.5%)	151 (18.3%)
Drinking Status					
Moderate Consumption (1-2 drinks daily)	1266 (25.6%)	1019 (22.6%)	225 (16.2%)	324 (18.3%)	106 (12.8%)
No Consumption (0 drinks daily)	2641 (53.5%)	2633 (58.4%)	956 (68.7%)	1134 (64.0%)	596 (72.2%)
Heavy Consumption (3+ drinks daily)	1033 (20.9%)	860 (19.1%)	211 (15.2%)	315 (17.8%)	123 (14.9%)
Physically Inactive	2155 (43.6%)	2322 (51.5%)	921 (66.2%)	1040 (58.7%)	572 (69.3%)
ADL Limitations	0.07 (0.26)	0.12 (0.32)	0.25 (0.44)	0.18 (0.38)	0.34 (0.47)
Body Mass Index	28.42 (5.67)	29.56 (6.30)	29.85 (6.79)	29.84 (6.50)	30.97 (7.62)
Hypertension	2151 (43.5%)	2704 (59.9%)	941 (67.6%)	1130 (63.7%)	556 (67.4%)
Heart Problem	534 (10.8%)	849 (18.8%)	401 (28.8%)	428 (24.1%)	292 (35.4%)
Stroke	183 (3.7%)	252 (5.6%)	139 (10.0%)	137 (7.7%)	116 (14.1%)
Diabetes	551 (11.2%)	1013 (22.5%)	402 (28.9%)	482 (27.2%)	273 (33.1%)
Cancer	371 (7.5%)	527 (11.7%)	264 (19.0%)	256 (14.4%)	163 (19.8%)
Arthritis	1950 (39.5%)	2554 (56.6%)	942 (67.7%)	1155 (65.1%)	631 (76.5%)
Psychiatric Condition	468 (9.5%)	766 (17.0%)	324 (23.3%)	417 (23.5%)	306 (37.1%)

Continuous variables presented as means with standard deviations in parenthesis. Categorical and binary variables presented as counts with percentages in parenthesis. Wealth is log-adjusted. *Abbreviations:* ADL = Activities of Daily Living.

Table 2. Multinomial Estimates of Association Between Discrimination in Healthcare and Trajectory Group Membership of Middle Aged and Older Adults, Health and Retirement Study 2008-2020. (N=13,422)

Reference Group	Low											
Comparator Group	Frequent			High-to-Frequent			Frequent-to-High			High		
	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value	RRR	95% CI	P-Value
Unadjusted Model												
Discrimination in Healthcare	1.13	(1.02-1.26)	0.021	1.54	(1.33-1.77)	<0.001	1.13	(0.99-1.30)	0.079	2.04	(1.73-2.42)	<0.001
Adjusted Model												
Discrimination in Healthcare	1.11	(0.99-1.24)	0.066	1.38	(1.18-1.61)	<0.001	1.04	(0.89-1.20)	0.629	1.62	(1.35-1.95)	<0.001
Age	1.02	(1.02-1.03)	<0.001	1.02	(1.01-1.03)	<0.001	1.02	(1.02-1.03)	<0.001	1.01	(0.99-1.02)	0.379
Male	0.72	(0.66-0.79)	<0.001	0.74	(0.65-0.85)	<0.001	0.77	(0.68-0.87)	<0.001	0.68	(0.57-0.80)	<0.001
Non-Hispanic Black	0.81	(0.72-0.91)	<0.001	0.79	(0.66-0.94)	0.007	0.69	(0.59-0.81)	<0.001	0.62	(0.50-0.78)	<0.001
Hispanic	0.80	(0.67-0.95)	0.011	1.04	(0.81-1.33)	0.755	0.86	(0.68-1.08)	0.188	0.56	(0.39-0.82)	0.003
Married	1.09	(0.99-1.20)	0.068	0.91	(0.79-1.04)	0.175	0.98	(0.87-1.11)	0.773	1.09	(0.92-1.30)	0.339
Education	1.07	(1.05-1.09)	<0.001	1.08	(1.05-1.11)	<0.001	1.08	(1.06-1.11)	<0.001	1.10	(1.07-1.14)	<0.001
Wealth	1.00	(0.99-1.01)	0.922	0.97	(0.96-0.99)	0.008	0.97	(0.96-0.99)	0.004	0.98	(0.96-1.00)	0.11
Uninsured	0.87	(0.73-1.03)	0.111	0.50	(0.37-0.68)	<0.001	0.77	(0.60-0.98)	0.034	0.50	(0.34-0.74)	0.001
Smoking Status												
Former Smoker	1.12	(1.02-1.23)	0.018	1.26	(1.10-1.45)	0.001	1.24	(1.09-1.41)	0.001	1.20	(1.00-1.43)	0.049
Current Smoker	0.97	(0.85-1.11)	0.651	1.11	(0.92-1.36)	0.280	1.23	(1.03-1.46)	0.024	1.17	(0.92-1.48)	0.212
Drinking Status												
No Consumption (0 drinks)	1.02	(0.91-1.13)	0.781	1.43	(1.20-1.69)	<0.001	1.25	(1.08-1.46)	0.003	1.73	(1.37-2.18)	<0.001
Heavy Consumption (3+ drinks)	1.01	(0.89-1.15)	0.878	1.14	(0.92-1.41)	0.237	1.15	(0.95-1.38)	0.144	1.46	(1.10-1.95)	0.009
Physically Inactive	1.08	(0.99-1.18)	0.085	1.62	(1.41-1.86)	<0.001	1.25	(1.11-1.42)	<0.001	1.61	(1.35-1.92)	<0.001
Activities of Daily Living	1.10	(1.03-1.18)	0.005	1.32	(1.23-1.43)	<0.001	1.22	(1.13-1.32)	<0.001	1.44	(1.33-1.56)	<0.001
BMI	1.01	(1.01-1.02)	0.001	1.00	(0.99-1.01)	0.608	1.01	(1.00-1.02)	0.090	1.01	(1.00-1.03)	0.058
Hypertension	1.57	(1.44-1.73)	<0.001	1.82	(1.58-2.10)	<0.001	1.65	(1.45-1.87)	<0.001	1.62	(1.35-1.94)	<0.001
Heart Problem	1.45	(1.28-1.65)	<0.001	2.13	(1.81-2.49)	<0.001	1.75	(1.51-2.04)	<0.001	2.65	(2.20-3.20)	<0.001
Stroke	1.09	(0.88-1.33)	0.435	1.38	(1.07-1.77)	0.013	1.21	(0.95-1.55)	0.121	1.74	(1.32-2.30)	<0.001
Diabetes	1.94	(1.72-2.20)	<0.001	2.21	(1.88-2.60)	<0.001	2.27	(1.95-2.64)	<0.001	2.61	(2.15-3.16)	<0.001
Cancer	1.40	(1.21-1.61)	<0.001	2.38	(1.99-2.86)	<0.001	1.72	(1.44-2.06)	<0.001	2.46	(1.98-3.06)	<0.001
Arthritis	1.53	(1.40-1.68)	<0.001	2.02	(1.76-2.32)	<0.001	1.97	(1.74-2.23)	<0.001	2.80	(2.33-3.38)	<0.001
Psychiatric Condition	1.78	(1.56-2.03)	<0.001	2.04	(1.72-2.42)	<0.001	2.337	(2.00-2.73)	<0.001	3.35	(2.77-4.06)	<0.001

Figure 2. Multinomial Estimates of Association between Discrimination in Healthcare Settings and Trajectory Group Membership with varying Comparator Groups based on the Similar Number of Doctor Visits, Health and Retirement Study (N=13,422)

Note: Models adjust for sociodemographic background, health behaviors, and general health status. Reference categories are indicated with parenthesis. **=p<0.01 ***=p<0.001. Abbreviations: RRR = Relative Risk Ratio.