**Doubling dolutegravir dosage reduces the viral reservoir in ART-treated people with HIV**

Céline Fombellida-Lopez[1,2], Aurelija Valaitienė[2], Lee Winchester[3], Nathalie Maes[4], Patricia Dellot[5], Céline Vanwinge[6], Aurélie Ladang[7], Etienne Cavalier[7], Fabrice Susin[8], Dolores Vaira[8], Marie-Pierre Hayette[8], Catherine Reenaers[9], Michel Moutschen[1,5], Courtney V. Fletcher[3], Alexander O. Pasternak[2]*, Gilles Darcis[1,5]*

[1]Laboratory of Immunology and Infectious Diseases, GIGA-Institute, University of Liège, Liège, Belgium; [2]Laboratory of Experimental Virology, Department of Medical Microbiology, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands; [3]Antiviral Pharmacology Laboratory, University of Nebraska Medical Center, Omaha, NE, USA; [4]Department of Biostatistics and Medico-Economic Information, University Hospital of Liège, Liège, Belgium; [5]Department of General Internal Medicine and Infectious Diseases, University Hospital of Liège, Liège, Belgium; [6]GIGA Flow Cytometry Platform, University of Liège, Liège, Belgium; [7]Department of Clinical Chemistry, University Hospital of Liège, Liège, Belgium; [8]Laboratory of Clinical Microbiology, University Hospital of Liège, Liège, Belgium; [9]Department of Gastroenterology, University Hospital of Liège, Liège, Belgium

*These authors contributed equally to the work and share the last authorship

Correspondence to Gilles Darcis (gdarcis@chuliege.be) and Alexander O. Pasternak (a.o.pasternak@amsterdamumc.nl)

**Abstract**

Whether antiretroviral therapy (ART) is always completely suppressive, or HIV might continue

to replicate at low levels despite ART in some people with HIV (PWH), is still debated. Here,

we intensified the ART regimen by doubling dolutegravir (DTG) dosage and investigated the

impact of this strategy on HIV blood and tissue reservoirs, immune activation, and

inflammation. Twenty HIV-infected adults, who had received a triple ART consisting of 50mg

DTG/600 mg abacavir/300 mg lamivudine pre-intensification and had been suppressed on ART

for at least two years, were enrolled in a phase 2 randomized clinical trial. Half of them received

an additional 50 mg of DTG for a period of 84 days. As expected, plasma and tissue DTG

concentrations significantly increased during the study period in the intensified group but not

in the control group. Accordingly, significant decreases in total HIV DNA, intact HIV DNA,

and cell-associated unspliced (US) HIV RNA in PBMCs, as well as in the US RNA/total DNA

ratio, were observed in the intensified group but not in the control group. Intensification also

reduced markers of immune activation and exhaustion but had no measurable impact on

systemic or tissue inflammation. Together with this, intensification resulted in a temporary

decrease in the CD4/CD8 ratio that returned to baseline by day 84. Our results strongly suggest

that the pre-intensification ART regimen was not completely suppressive. If confirmed in larger

clinical trials, these results could have an impact on the clinical management of PWH and HIV

curative strategies.

## Introduction

Human immunodeficiency virus (HIV) remains a major global public health issue, with an estimated 39.9 million people with HIV (PWH) at the end of 2023. Out of them, 30.7 million (77%) were accessing antiretroviral therapy (ART) (https://www.unaids.org/en/resources/fact-sheet). ART suppresses plasma viral load to undetectable levels, restores immune function, and eliminates the risk of developing AIDS (1). Current ART usually consists of one or two nucleotide or nucleoside reverse transcriptase inhibitors and the second or third drug of another class (e.g., a non-nucleoside reverse transcriptase inhibitor, a protease inhibitor (PI), or an integrase strand transfer inhibitor (INSTI)) (https://eacs.sanfordguide.com/). The mechanism of action of ART is interfering with different steps of the viral replication cycle, such as reverse transcription, integration, or viral particle maturation. Importantly, ART only blocks the infection of new cells and does not inhibit HIV gene expression in cells that were infected prior to ART initiation or in the progeny of such cells (2).

ART is not curative, and the main barrier to an HIV cure is the presence of latent viral reservoirs that arise due to the ability of HIV to integrate its genome into the host chromosome and persist in a latent form in some of the cells it infects, such as memory CD4+ T cells (3, 4). These reservoir cells harbour stably integrated and replication-competent provirus that can reactivate and, after cessation of ART, fuel a rapid viral rebound (5). HIV reservoirs persist lifelong, slowly decaying over time (6-8). Several mechanisms have been described to explain the long-term persistence of HIV reservoirs, including the infection of long-lived CD4+ T cells and clonal expansion by homeostatic or antigen-driven cellular proliferation or by HIV integration in or near genes that promote cellular proliferation (9, 10). However, if therapy does not prevent all new infections, then replenishment by *de novo* virus replication provides an additional potential mechanism of reservoir persistence (11).

Whether ART is fully suppressive is a topic of considerable and long-standing debate (12). Evidence of residual replication in ART-suppressed PWH has been previously presented (13-16), but other groups published reports that contest this notion (17-20). One strong argument against ongoing viral replication is the absence of detectable virus evolution on ART. However, if residual replication occurs intermittently and at a low level, with short infection chains continuously arising and terminating, then nucleotide substitutions are expected to be sporadic and not linked by temporal structure, unless the sampling is very deep and intensive (13, 21). In principle, failure to detect virus evolution does not form an unequivocal proof that residual

viral replication does not occur at all: absence of evidence is not evidence of absence. Indeed,

Paryad-Zanjani et al. recently developed a mathematical model to track the viral dynamics in

lymph nodes (LNs) after the initiation of ART (22), which suggested that the methods used by

the previous studies to detect genetic divergence were not sufficiently sensitive and that these

studies would have been unlikely to detect residual replication.

Over time, this persistent low-level residual replication may contribute to chronic immune

activation and inflammation in ART-treated PWH (23). Constant stimulation of the immune

system could lead to immune exhaustion and premature immune senescence, resulting, over

the years, in the development of non-AIDS co-morbidities, such as cardiovascular disease (24).

This replication may occur in anatomical sanctuaries/compartments, notably the LNs and the

gut-associated lymphoid tissue (GALT), two sites massively infected by HIV (25). Lower

penetration of antiretroviral drugs to these tissues reduces the local drug concentrations and,

consequently, their effectiveness in preventing HIV replication (26-28).

One way to investigate the residual replication is to intensify a standard ART regimen and

measure the impact of this intensification on latent reservoirs, inflammation, and immune

activation. If a reduction of a virological parameter is observed, it would strongly suggest that

the ART regimen prior to intensification did not completely prevent virus replication. Several

groups have conducted ART intensification trials, either by adding a CCR5 antagonist

(maraviroc) or an INSTI (raltegravir or dolutegravir (DTG)) to a standard three-drug regimen.

Some of these trials have indeed revealed residual virus replication at baseline, as treatment

intensification caused a decrease in immune activation, a decrease in cell-associated HIV RNA

in the ileum, and a transient increase in 2-long terminal repeat (2-LTR) circles (29-38).

However, other groups have failed to demonstrate any effects of ART intensification on

virological parameters (39-45).

In summary, is still unclear whether ongoing viral replication can occur in the presence of ART.

The consequences of residual replication for the HIV curative strategies are important since

approaches aimed at eliminating viral reservoirs such as latency reversal ("shock-and-kill")

(46) are not expected to be fully effective if the virus continues to replicate despite treatment.

In the worst-case scenario, the residual replication could lead to significant replenishment of

the HIV reservoir upon latency reversal, compensating for any reservoir depletion achieved by

this strategy. Therefore, understanding the phenomenon of residual replication and developing

strategies to eliminate it are key elements in the quest to cure PWH.

Here, we intensified ART in PWH who had been suppressed on ART for at least two years, by doubling the dosage of DTG, an antiretroviral drug that had already been part of the regimen pre-intensification. We investigated the impact of this intensification on HIV blood and tissue latent reservoirs, immune activation, and inflammation.

**Results**

**Study design and participants**

Between February and July 2019, twenty ART-suppressed participants receiving a triple antiretroviral regimen consisting of 50 mg DTG, 600 mg abacavir (ABC), and 300 mg lamivudine (3TC) daily were enrolled into two groups as follows: 10 participants continued to receive this regimen (control group) and 10 participants received an additional 50 mg of DTG daily for 84 days (intensified group) as shown in Figure S1. Study participants were mostly Caucasian males and the median age was 52 years. The median cumulative time of viral suppression on ART was 7.4 years and participants had maintained continuous viral suppression for a median of 3.9 years before the study. The clinical characteristics of the participants at baseline are shown in Table 1.

**ART intensification increases plasma and tissue DTG concentrations**

To assess if the desired experimental conditions were achieved, we monitored the concentrations of DTG and 3TC, two antiretroviral drugs that were part of the ART regimen (Figure 1, Figure S2). Baseline plasma and rectal tissue DTG and 3TC concentrations were not significantly different between the intensified and control groups (Table 1). At median, baseline DTG concentration in the rectal tissue was 16.7% (interquartile range (IQR), 15.1%-24.6%) of the corresponding DTG plasma concentration, strikingly close to the value (17%) reported previously (47). As expected, plasma and tissue DTG concentrations significantly increased between days 0 and 84 in the intensified group but not in the control group (Figures 1A, 1B). In fact, DTG concentrations almost exactly doubled in plasma (median fold increase, 1.97-fold) and more than doubled in tissue (median fold increase, 2.48-fold) on day 84, as compared to day 0, in the intensified group. However, at day 84, tissue DTG concentration in the intensified group was still much lower (median, 20.0%; IQR, 17.3%-22.3%) than the corresponding DTG plasma concentration, reflecting limited penetration of DTG into the

tissue. Plasma and tissue 3TC concentrations did not change on day 84 compared to day 0 in both groups, which was expected as the dosage of 3TC was not increased in any group (Figures 1C, 1D). However, we did observe a transient increase in the plasma DTG concentration in the control group and in the plasma 3TC concentration in both groups on days 1 and 56 of the study (Figure S2), with the concentrations returning to baseline levels by day 84. One possible explanation for this effect could be an increased adherence of the participants to their ART regimens during the study ("white coat compliance") (48).

## ART intensification reduces HIV reservoir

To investigate the effect of treatment intensification on the viral reservoir, we longitudinally measured total HIV DNA and cell-associated unspliced (US) HIV RNA in peripheral blood mononuclear cells (PBMCs) on days 0, 1, 28, 56, and 84 of the study. In addition, we measured intact HIV DNA in PBMCs and total HIV DNA in rectal biopsies on days 0 and 84 of the study. Strikingly, we observed significant longitudinal decreases in total HIV DNA, US RNA, and HIV transcription level per provirus (US RNA/total DNA ratio) in PBMCs in the intensified group (p=0.0022, p<0.001, and p<0.001, respectively) but not in the control group, with significant differences in the relative changes from baseline for two of these three markers between the groups (p<0.001, p=0.010, and p=0.090, respectively) (Figures 2A-2C, Figures S3-S5). While total HIV DNA in the intensified group transiently decreased in the first 28 days of the study (median (IQR) fold decrease from day 0 to day 28, 2.1 (1.1-4.7) fold, p=0.027) and increased afterwards (Figure 2A, Figure S3), US RNA and US RNA/total DNA ratio in the intensified group continued to decrease throughout the whole study period (Figures 2B-2C, Figures S4, S5). By day 84, US RNA and US RNA/total DNA ratio have decreased from day 0 by medians (IQRs) of 5.1 (3.3-6.4) and 4.6 (3.1-5.3) fold, respectively (p=0.016 for both markers). In comparison, in the control group these markers remained at the baseline level or even increased (Figures 2A-2C). These effects were confirmed by measurement of intact HIV DNA in PBMCs that was quantified by the intact proviral DNA assay (IPDA) (49). We measured a significant decrease in the intact HIV DNA between days 0 and 84 in the intensified group (median (IQR) fold decrease, 2.2 (1.3-3.1) fold, p=0.039) but not in the control group (Figure 2D). In contrast, total HIV DNA in rectal tissue did not change between days 0 and 84 in both groups (Figure 2E). Taken together, these results indicate that the intensification reduced the HIV reservoir in the peripheral blood of ART-treated PWH.

## ART intensification reduces markers of immune activation and exhaustion

To detect if intensification had an impact on immune activation and exhaustion, we performed longitudinal flow cytometry analysis to assess the expression of cell surface markers of activation (HLA-DR, CD38) and exhaustion (PD-1, TIGIT) on CD4+ and CD8+ T cells (Figure 3, Figure S6). A transient but significant increase in the percentage of CD4+ cells expressing PD-1 (p=0.022), and a significant decrease in the percentage of CD8+ cells expressing CD38 (p=0.034) were measured in the control group. A decrease in the percentage of CD4+ cells expressing TIGIT was measured in the intensified group (p=0.031), with a significant difference compared to the control group (p=0.048). We also measured a significant decrease in the percentage of CD8+ cells co-expressing CD38 and HLA-DR in the intensified group (p=0.049) (Figure 3). Overall, these results indicate that intensification reduced markers of immune activation and exhaustion.

## ART intensification does not impact systemic or tissue inflammation

To investigate the effect of intensification on systemic and tissue inflammation, we quantified several inflammatory cytokines in plasma and in rectal tissue. In plasma, we longitudinally measured the levels of IL-1β, IL-6, IFN-γ, IL-17 α, and TNF-α, and in rectal tissue, we measured expression of the same cytokines by quantifying their mRNA levels at days 0 and 84. We observed a significant increase in plasma IL-6 levels in the control group (p=0.049). Apart from that, no significant change from baseline for any cytokine was observed for both groups in both plasma and tissue (Figure 4, Figure S7). Overall, these results indicate that intensification did not measurably impact systemic or tissue inflammation.

## ART intensification leads to a transient decrease in the CD4/CD8 ratio

We longitudinally measured CD4 and CD8 T-cell counts, CD4/CD8 ratio, as well as plasma levels of soluble CD14 (sCD14), a marker of monocyte/macrophage activation, and of C-reactive protein (CRP), an inflammation marker (Figure 5, Figure S8). No significant changes were observed in the CD4 and CD8 counts in both groups, however we measured a transient but significant decrease in the CD4/CD8 ratio in the intensified group (p=0.0081) that returned

to the baseline level by day 84 (Figure 5). No change in the CD4/CD8 ratio was observed in the control group. We also observed a significant decrease in the sCD14 level in the control group ($p<0.001$) and a trend toward an increase in the intensified group ($p=0.079$), which resulted in a significant difference between the groups ($p<0.001$). Finally, we observed a small but significant increase in the CRP level in the control group ($p=0.045$) but not in the intensified group.

**Correlations between the measured parameters**

Finally, we assessed pairwise correlations between the measured parameters at baseline, as well as between their time-weighted changes from baseline during the study (time-weighted areas under curve) and between their relative changes between days 0 and 84 of the study (Figures S9-S11). At baseline, we found strong positive pairwise correlations between the levels of different plasma inflammatory cytokines, as well as between the percentages of peripheral CD8+ T cells expressing TIGIT and inflammatory cytokine levels in plasma and tissue. Of the latter, especially strong correlations were observed between CD8+TIGIT+ cell percentages and plasma or tissue levels of IL-6 (*rho*=0.74 and *rho*=0.71, respectively; $p<0.001$). We also found positive correlations at baseline between US RNA/total DNA ratio and percentages of CD4+ T cells expressing CD38 and co-expressing CD38 and HLA-DR, markers of immune activation (*rho*=0.53, $p=0.020$ and *rho*=0.50, $p=0.023$, respectively). Furthermore, US RNA changes between days 0 and 84 positively correlated with changes in levels of expression of immune activation/exhaustion markers CD38+ on CD4+ cells and PD-1+ on CD8+ cells (*rho*=0.53, $p=0.030$ and *rho*=0.61, $p=0.011$, respectively) and of tissue inflammatory cytokine IL-1β mRNA (*rho*=0.62, $p=0.0096$). These correlations were even stronger for the changes between days 0 and 84 of US RNA/total DNA ratio (CD4+CD38+: *rho*=0.55, $p=0.029$; CD8+PD-1+: *rho*=0.73, $p=0.0018$; IL-1β mRNA: *rho*=0.76, $p=0.0011$). This confirms the results of our earlier study, where US RNA/total DNA ratio was found to correlate with markers of immune exhaustion (50). Interestingly, intact, but not total, HIV DNA in PBMCs strongly negatively correlated with time of viral suppression (*rho*=-0.67, $p=0.0016$), confirming earlier observations that intact HIV DNA decays on ART much faster than the defective (or total) HIV DNA (51, 52).

**Discussion**

The aim of this study was to investigate whether ongoing viral replication contributes to HIV persistence under ART. To the best of our knowledge, this is the first study where ART intensification was achieved not by adding a new antiretroviral drug to the existing regimen but by increasing the dosage of a drug that was already part of the regimen pre-intensification. We evaluated the impact of doubling the dosage of DTG on HIV cellular and tissue reservoirs, immune activation, exhaustion, and inflammation in ART-suppressed PWH.

As expected, plasma and tissue DTG concentrations significantly increased between day 0 and day 84 in the intensified group but not in the control group. Accordingly, significant decreases in total HIV DNA, intact HIV DNA, US HIV RNA, and US RNA/total DNA ratio in PBMCs were observed during the study period in the intensified group but not in the control group. This strongly suggests that the pre-intensification ART regimen was not completely suppressive. The transient decrease in total DNA at day 28 may have been due to an initial decrease in newly infected cells upon ART intensification, however at the subsequent time points this effect was likely masked by proliferation (clonal expansion) of infected cells with defective proviruses that were obviously not affected by intensification. In line with this, the number of intact proviruses decreased, but that of total proviruses did not change, between days 0 and 84 (Figures 2A, 2D).

In addition, intensification significantly reduced percentages of CD4+ T cells expressing TIGIT, a T-cell exhaustion marker (53), and percentages of CD8+ T cells co-expressing CD38 and HLA-DR, T-cell activation markers. Interestingly, CD4+ cells expressing TIGIT have been shown to be enriched for HIV persistence in ART-treated PWH (54). Arguably, the reduction of HIV reservoir upon intensification has caused a corresponding reduction in chronic immune activation and exhaustion in this cohort. Indeed, changes in US HIV RNA levels and US RNA/total DNA ratio during the study positively correlated with changes in levels of markers immune activation, exhaustion, and inflammation, suggesting a link between changes in HIV reservoir activity and levels of immune activation and inflammation. These results are in line with previous studies reporting decreases in immune activation markers upon ART intensification (32-35).

In contrast, in the control group we measured significant longitudinal increases in total HIV DNA, percentages of CD4+ T cells expressing PD-1, another T-cell exhaustion marker, and in the plasma levels of IL-6, an inflammatory cytokine, and of CRP, another marker of

inflammation. Residual HIV replication, among other factors, could have contributed to these

effects in the control group. However, in the control group we also observed a reduction in the

percentages of CD8+ T cells co-expressing a T-cell activation marker CD38 and a strong

reduction in the plasma levels of sCD14. One possible explanation for these effects can be

increased adherence of the participants to their ART regimens during the study.

Two previous studies have measured a transient increase in HIV episomal DNA (2-LTR circles)

upon ART intensification with raltegravir, an INSTI (31, 36). As 2-LTR circles are a byproduct

of HIV integration and are expected to accumulate if integration is blocked, this was interpreted

as a strong evidence of new infections pre-intensification. However, another study did not

observe any change in 2-LTR circles upon ART intensification with DTG (45). In our study,

despite repeated attempts using different assays, we could only detect 2-LTR circles in a small

minority of participants (<5%) (data not shown). Notably, the effects of intensification on 2-

LTR circles in these previous studies were much more pronounced in participants who had

been treated with a PI-based ART regimen pre-intensification (31, 36). As PI-based regimens

have been shown to be less suppressive than other ART regimens (16, 55-57), they might have

allowed for a higher level of new infections and consequently of 2-LTR circles in these studies.

In contrast, in our study, participants had been treated with a DTG-based regimen for at least

two years and had had a suppressed plasma viral load (<50 copies/mL) for at least 2 years 11

months pre-intensification. Plausibly, the lack of detectable 2-LTR circles in our participants

reflects the high potency of their ART regimens and a long period of virological suppression

before the study. Remarkably, the intensification revealed that even this highly potent regimen

was not completely suppressive. Thus, our results emphasize that 2-LTR circles are not a very

sensitive marker of new infections due to their relative scarcity in infected cells. 2-LTR circles

represent only ~10% of total episomal DNA (58) and are not expected to be present in every

newly infected cell. In contrast, a newly infected CD4+ T lymphocyte can contain hundreds to

thousands of US HIV RNA copies on the peak of infection (59). Therefore, a change in the US

RNA level upon ART intensification is a much more sensitive indicator of new infections than

that of 2-LTR circles, as confirmed by this study.

Due to low penetration of antiretroviral drugs into tissue sites and anatomic sanctuaries (26-

28, 60, 61), ART pressure on HIV replication is generally expected to be more relaxed in tissues

than in peripheral blood. Surprisingly and in contrast to the effect of intensification on total

HIV DNA in peripheral blood, we did not observe any effect on total HIV DNA in the rectal

tissue. Although DTG concentration in the rectal tissue did increase upon intensification, it remained much lower than in the peripheral blood. This possibly contributed to the lack of detectable effect of intensification on the HIV DNA in tissue. Alternatively, intensification with DTG, an INSTI, had no effect in the tissue due to the preponderance of abortive infection events in lymphoid tissues that occur by cell-to-cell contact and are terminated prior to integration (62, 63).

The majority of previous studies that used INSTIs (raltegravir or DTG) as intensification drugs added it as a fourth drug using the standard dosage. For example, the DTG dosage in the study of Rasmussen et al. was 50 mg daily (45). In our study, we used a twice-higher dosage (100 mg daily). This DTG dosage is commonly used in clinical practice. Indeed, in PWH with confirmed or clinically suspected resistance to the integrase inhibitor class, the recommended dosage of DTG is 50 mg twice daily (https://www.cbip.be/fr/chapters/12?frag=10868). Based on our results (and pending the confirmation by larger studies), an increased dosage of DTG might be warranted in the future as a part of a standard regimen even in PWH without resistance. However, this has to be approached with caution as intensification has caused a transient decrease in the CD4/CD8 ratio in our study.

The main strength of our study is the measurement of a large number of virological and immunological parameters that allowed us to observe a significant impact of intensification on both virus and host. Our study is the first to observe significant reductions in as many as four HIV markers following ART intensification. Moreover, we measured plasma and tissue concentrations of DTG, allowing us to directly demonstrate their increase upon intensification. Our results are in line with some earlier studies that reported a decrease in cell-associated HIV RNA or a transient increase in 2-LTR circles upon intensification (31, 32, 36). However, they are in contrast with other studies, in which treatment intensification did not affect HIV markers (39, 40, 42). As this is the first ART intensification study to increase the dosage of an existing antiretroviral drug instead of adding a new drug, the results of this study and those of previous intensification studies cannot be directly compared.

Some limitations can be highlighted. First, we included twenty PWH (ten participants per arm). Although this small sample size does somewhat limit the robustness of the conclusions, we would like to emphasize that in this cohort, we performed a very detailed analysis, longitudinally measuring 30 biomarkers at five time points during ART. Secondly, despite the randomization, the groups were still not perfectly balanced as levels of total HIV DNA in

PBMCs were by chance higher at baseline in the intensified group. However, this is unlikely to have affected the conclusions of the study because (i) this study has a within-subject (repeated-measures) design, with the longitudinal change of a parameter within the same participant during the study being the main outcome, and (ii) this difference most probably reflects a higher number of infected cells, and not a higher level of residual replication, at baseline in the intensified group. No significant differences between the groups were found at baseline in the levels of the other four virological markers we measured (US RNA in PBMCs, US RNA/total DNA ratio, intact DNA in PBMCs, and total DNA in tissue), as well as between the antiretroviral drug concentrations in plasma or in tissue.

In conclusion, doubling the DTG dosage in PWH who had been suppressed on DTG-containing ART for a number of years resulted in a reduction in the levels of four HIV reservoir markers in peripheral blood, as well as in markers of T-cell activation and exhaustion. If confirmed in larger clinical trials, these results could have an impact on the clinical management of PWH. Moreover, if ongoing viral replication does indeed replenish HIV reservoirs over time, then improving ART regimens should be a necessary part of the curative strategies.

## Methods

### Study design and participants

This was a phase 2 open-label, interventional, monocentric, randomized, and controlled clinical trial performed at the Hospital University of Liège (Belgium) in which 20 HIV-infected adults were enrolled. Eligible participants had been treated with a combination of 50 mg DTG, 600 mg ABC, and 300 mg 3TC for more than two years and the purpose of this study was to assess the impact of DTG intensification (an additional daily dose of 50 mg) compared to a control group that continued with the above regimen. Participants were required to have a plasma viral load suppressed below 20 HIV RNA copies/mL during the 12 months before screening (allowing "blips") and an absolute CD4+ T lymphocyte count above 200 cells/mm$^3$. One participant out of 20 had an isolated "blip" of 115 copies/mL nine months before the study initiation. The detailed inclusion and exclusion criteria can be found in the supplementary data (Supplementary data 1). The study was approved by the Liège University Hospital-Faculty Ethics Committee (Comité d'Éthique Hospitalo-Facultaire Universitaire de Liège, 2018/228). Prior to inclusion in the study, each participant had dated and signed an informed consent form. ClinicalTrials.gov identifier: NCT05351684.

Historical plasma HIV RNA measurements, CD4+ T-cell counts, and treatment data were
retrieved from the outpatient medical records. The duration of continuous virological
suppression was calculated as the duration of the latest period with undetectable plasma HIV
RNA prior to the study initiation, allowing isolated 'blips' of 50–999 copies/ml. The duration
of cumulative suppression was calculated by adding together all such periods of continuous
suppression.

**Clinical procedures**

Each participant was randomly assigned to the intensified group or to the control group. This
study lasted 84 days in which 5 visits were programmed (days 0, 1, 28, 56, and 84). One visit
prior to inclusion was added to perform a blood test to ensure that the participant met the
inclusion/exclusion criteria for the study. The blood test included blood and platelet count,
electrolyte panel, creatinine, CPK, LDH counts, liver tests, coagulation tests, CD4 and CD8
lymphocyte counts and a plasma viral load. β-HCG levels were measured in women at each
visit to ensure that they were not pregnant. All participants completed the 12 week-long study
and no individuals were lost to follow-up.

Blood samples were collected in 4 EDTA tubes (~40mL) at days 0, 1, 28, 56, and 84 and directly
sent to the AIDS reference laboratory of Liège University Hospital for PBMC isolation. PBMC
samples were stored at -150°C for measurements of the HIV reservoir and immunological
parameters. Plasma samples were used to measure soluble inflammation markers. Ten flash-
frozen rectal biopsies were collected on days 0 and 84 during a routine gastroenterology visit.
The cryotubes containing the biopsies were directly placed in liquid nitrogen and stored at -
150°C.

**PBMC isolation**

PBMCs were isolated from whole blood by density gradient centrifugation using Ficoll-
Paque® Plus. The blood samples were collected in EDTA tubes and centrifuged at 1200 X g
for 10 min (acceleration 9, brake 9). The plasma was removed and stored at -150°C. The blood
was then well homogenized with RPMI 1640, gently transferred in a Falcon tube that contained
12,5 mL of Ficoll, and centrifuged at 700 X g for 20 minutes (acceleration 6, brake 0). The
PBMC layer was removed, diluted in 50 mL RPMI 1640, and centrifuged at 300 X g for 7

minutes (acceleration 9, brake 9). The supernatant was removed, and the process was repeated two more times, completing three washes. The pellet was resuspended, and the cells were counted. Foetal bovine serum (FBS) was added to the re-suspended pellet to obtain $10 \times 10^6$ cells/500 μL. For cryopreservation, 500 μL of freeze mixture (10% DMSO in FBS) was added to 500 μL of the cell suspension in the cryotubes. Cryotubes were slowly frozen in the Mr. Frosty™ freezing container at -80 °C and transferred to -150°C the next day.

**Quantification of total HIV DNA and cell-associated US HIV RNA in PBMCs**

Total nucleic acids were extracted from $2 \times 10^6$ PBMCs using Boom isolation method (64). Total HIV DNA was measured by qPCR using the previously described primer/probe combination that amplifies the HIV packaging signal (Ψ) region (49). Extracted cellular RNA was treated with DNase (DNA-free kit; ThermoFisher Scientific) to remove genomic DNA that could interfere with the quantitation and reverse transcribed into cDNA using random primers and SuperScript III reverse transcriptase (all from ThermoFisher Scientific). To quantify cell-associated US HIV RNA, this cDNA was pre-amplified using primer pair Ψ_F (49) and HIV-FOR (65). The product of this PCR was used as template for a seminested qPCR with the Ψ primer/probe combination (49). HIV DNA or RNA copy numbers were determined using a 7-point standard curve with a linear range of more than 5 orders of magnitude that was included in every qPCR run and normalized to the total cellular DNA (by measurement of β-actin DNA) or RNA (by measurement of 18S ribosomal RNA) inputs, respectively, as described previously (66). Non-template control wells were included in every qPCR run and were consistently negative. Total HIV DNA and US RNA were detectable in 90.7% and 66.7% of the samples, respectively. Undetectable measurements of US RNA or total DNA were assigned the values corresponding to 50% of the corresponding assay detection limits, with a maximum of 25 copies/μg total RNA or 50 copies/$10^6$ PBMCs, respectively. The detection limits depended on the amounts of the normalizer (input cellular DNA or RNA) and therefore differed among samples. Measurements with low input cellular DNA or RNA and undetectable total HIV DNA or US RNA (n=2 and n=3, respectively), were excluded from the analysis. HIV transcription levels per provirus (US RNA/total DNA ratios) were calculated taking into account that $10^6$ PBMCs contain 1 μg of total RNA (67).

**Quantification of intact HIV DNA in PBMCs**

Intact HIV DNA was quantified on days 0 and 84 of the study by the IPDA (49). In brief,

genomic DNA was isolated from $4 \times 10^6$ PBMCs using Puregene Cell Kit (QIAGEN Benelux

B.V.) according to the manufacturer's instructions and digested with *Bgl*I restriction enzyme

(ThermoFisher Scientific) as described previously (68). Notably, only a small minority (<8%)

of HIV clade B sequences contain *Bgl*I recognition sites between Ψ and *env* amplicons,

therefore *Bgl*I digestion is not expected to substantially influence the IPDA output, while

improving the assay sensitivity by increasing the genomic DNA input into a ddPCR reaction

(68). After desalting by ethanol precipitation, genomic DNA was subjected to two separate

multiplex droplet digital PCR (ddPCR) assays: one targeting HIV Ψ and *env* regions using

primers and probes described previously, including the unlabelled *env* competitor probe to

exclude hypermutated sequences (49), and one targeting the cellular *RPP30* gene, which was

measured to correct for DNA shearing and to normalize the intact HIV DNA to the cellular

input. The *RPP30* assay amplified two regions, with amplicons located at exactly the same

distance from each other as HIV Ψ and *env* amplicons. The first region was amplified using a

forward primer 5'-AGATTTGGACCTGCGAGCG-3', a reverse primer 5'-

GAGCGGCTGTCTCCACAAGT-3', and a fluorescent probe 5'-FAM-

TTCTGACCTGAAGGCTCTGCGCG-BHQ1-3' (69). The second region was amplified using

a forward primer 5'-AGAGAGCAACTTCTTCAAGGG-3', a reverse primer 5'-

TCATCTACAAAGTCAGAACATCAGA-3', and a fluorescent probe 5'-HEX-

CCCGGCTCTATGATGTTGTTGCAGT-BHQ1-3'. The ddPCR conditions were as described

previously (49) with some minor amendments: we used 46 cycles of

denaturation/annealing/extension and the annealing/extension temperature was 60°C. Intact

HIV DNA was detectable in 79.5% of the samples. Undetectable measurements of intact HIV

DNA were assigned the values corresponding to 50% of the assay detection limits. The

detection limits depended on the amounts of the normalizer (input cellular DNA) and therefore

differed among samples. For two participants, in whom intact HIV DNA on day 0 was

undetectable due to a technical issue, we included day 1 values in the analysis. Excluding these

two participants did not change the conclusions (data not shown).

**456 Quantification of total HIV DNA in rectal biopsies**

457 Prior to total HIV DNA isolation, rectal biopsies were homogenized with the TissueLyser

458 (QIAGEN Benelux B.V.). Total HIV DNA was isolated using ReliaPrep™ gDNA Tissue

Miniprep System (Promega) according to the manufacturer's instructions. Total HIV DNA in rectal biopsies was quantified using exactly the same method as total HIV DNA in PBMCs (see above). Total HIV DNA was detectable in 92.5% of the rectal tissue samples. Undetectable measurements of total HIV DNA were assigned the values corresponding to 50% of the corresponding assay detection limits. The detection limits depended on the amounts of the normalizer (input cellular DNA) and therefore differed among samples.

**Quantification of immune activation and exhaustion**

Flow cytometry was used to quantify surface markers of immune activation and exhaustion (CD38, HLA-DR, PD-1, TIGIT) on CD4+ and CD8+ T cells. Cryopreserved PBMCs ($10 \times 10^6$ cells) were thawed and stained with a fixable viability dye Zombie NIR (Sony), and the following antibodies: CD3-APC/Fire810, CD4-PE/Fire700, CD8a-PercP (all from Sony), PD-1-PE, TIGIT-BV421, HLA-DR-BB515, CD38-BV711 (all from BD Biosciences). Cells were then washed, fixed in 1% paraformaldehyde, and run on a FACS SONY ID7000. Analysis was performed using FlowJo software (v10.8.1). Cells were gated according to morphological parameters (forward and side scatter) and viability. The complete gating strategy can be found in the supplemental data (Fig. S12). To exclude the effect of inter-assay variation on the data, samples from each participant were always analysed together.

**Quantification of systemic and tissue inflammation**

To quantify systemic inflammation, the levels of five plasma inflammatory cytokines (IFN-γ, IL-1β, IL-6, IL-17α, TNF-α) were measured on MAGPIX (Luminex) using the ProcartaPlex™ Human Inflammation Panel, 20plex (ThermoFisher Scientific). sCD14 was quantified by Human CD14 Quantikine ELISA Kit (R&D Systems). CRP was measured by high-sensitivity C-reactive protein (hs-CRP) test.

To measure tissue inflammation, we determined rectal tissue levels of mRNAs encoding five inflammatory cytokines (IL-1β, IL-6, IFN-γ, IL-17 α, and TNF-α). Rectal tissue biopsies were manually homogenized and lysed in buffer L6 (64) and RNA was extracted using Boom isolation method (64). Extracted RNA was treated with DNase (DNA-free kit; ThermoFisher Scientific) and reverse transcribed using random primers and SuperScript III reverse transcriptase (all from ThermoFisher Scientific). Cytokine mRNAs were quantified using

TaqMan Gene Expression Assays (all from ThermoFisher Scientific): IL-1β -

Hs01555410_m1, IL-6 - Hs00174131_m1, IFN-γ - Hs00989291_m1, IL-17 α -

Hs00174383_m1, TNF-α - Hs00174128_m1. Levels of cytokine mRNAs were normalized to

that of *GAPDH* mRNA, measured using a TaqMan Gene Expression Assay Hs02758991_g1.

**Quantification of antiretroviral drug concentrations**

Briefly, DTG was extracted from plasma samples via methanolic protein precipitation and from

tissue samples via tissue homogenization, followed by protein precipitation. A DTG standard

calibration curve over the range 20.0-5,000 ng/mL and stable isotope labelled internal standard

were used to generate an analyte/internal standard response. Analyte/internal standard response

was detected by liquid chromatography tandem mass spectrometry (LCMS) based on

previously described methods (26, 60, 70). Tissue penetration ratios of DTG were calculated

as the ratios of DTG concentrations in tissue (as ng/mL after conversion from ng/g by dividing

the values in ng/g by 1.06 (71)) to the concentrations in plasma (ng/mL). Lamivudine (3TC)

concentrations were determined with an LCMS method similar to our previously described

procedures (72). Differences are as follows: 3TC stock and standard working solutions were

prepared in methanol and demonstrated similar stability. The MS used was a triple-quadrupole

AB Sciex 5500 series monitoring positive ion transitions 230.1 -> 112.1 (3TC) and 233.1 ->

115.1 (stable labeled internal standard). Mobile phase was 12.5% acetonitrile and 0.1% formic

acid in deionized water. No source rinsing mobile phase was needed. The limit of quantitation

was 30 ng/mL and the upper bound was 6,000 ng/mL. Total accuracy ranged from 94.3–96.9%

for. Within-day coefficients of variation were below 5%, between-day coefficients of variation

were below 5%. Stability of 3TC in human plasma was also similar.

**Statistical analysis**

Baseline parameters were compared using Mann-Whitney tests for continuous variables and

Fisher's exact tests for categorical variables. Virological parameters were $\log_{10}$-transformed

before analysis. Longitudinal dynamics of the measured parameters was modelled using linear

mixed-effects analysis. This analysis was used both to compare the longitudinal changes from

baseline between the intensified and the control groups and to compare the changes from

baseline with zero in each of these groups separately (intercept-only analysis). As a sensitivity

analysis, we performed a comparison between models that do and do not include time as a covariate. Including time in the models did not change the conclusions (Supplementary Tables 1 and 2). Paired Wilcoxon tests were used for within-group comparisons of parameters between days 0 and 84. For these comparisons, pairs where both values were undetectable were excluded from the analysis. Time-weighted changes from baseline were calculated by dividing the areas under curve (computed using the trapezoid rule) by the time period of the study. Time-weighted changes from baseline were not calculated if the baseline value and one or more of the subsequent values were undetectable. Time-weighted changes from baseline and ratios of the parameters between different study days were compared between the groups using Mann-Whitney tests. Spearman tests were used to produce correlation matrices of the parameters at baseline, as well as of time-weighted changes from baseline and of changes between day 0 and day 84 of the measured parameters. Changes of the parameters from baseline, as well as US RNA/total DNA ratios, were not calculated if both values in a pair were undetectable. Data were analysed using Prism 10.2.0 (GraphPad Software) or IBM SPSS Statistics 28.0.1.0. All statistical tests were two-sided, and p values of $<0.05$ were considered statistically significant.

**Acknowledgements**

We are grateful to Marine Hubert and Raquel Mateus for technical assistance in PBMC isolation. We acknowledge the participation and commitment of the trial participants who made the study possible. C.F.L. is a FRIA grantee of the Belgian Fund for Scientific Research (Fonds de la Recherche Scientifique –FNRS) and is supported by the Fondation Léon Fredericq. This work was supported in part by grant R01-AI124965 (to C.V.F.) from the US National Institutes of Health. A.O.P. is supported by the grant no. 09120011910035 from the Dutch Medical Research Council (ZonMw). G.D. is supported by the Belgian Fund for Scientific Research (Fonds de la Recherche Scientifique –FNRS).

**References**

1.      Deeks SG, Lewin SR, Havlir DV. The end of AIDS: HIV infection as a chronic disease. Lancet. 2013;382(9903):1525-33.
2.      Deeks SG, Overbaugh J, Phillips A, Buchbinder S. HIV infection. Nat Rev Dis Primers. 2015;1:15035.
3.      Pasternak AO, Berkhout B. HIV persistence: silence or resistance? Curr Opin Virol. 2023;59:101301.

4.      Darcis G, Berkhout B, Pasternak AO. The Quest for Cellular Markers of HIV Reservoirs: Any Color You Like. Front Immunol. 2019;10:2251.

5.      De Scheerder MA, Vrancken B, Dellicour S, Schlub T, Lee E, Shao W, et al. HIV Rebound Is Predominantly Fueled by Genetically Identical Viral Expansions from Diverse Reservoirs. Cell Host Microbe. 2019;26(3):347-58.e7.

6.      Siliciano JD, Kajdas J, Finzi D, Quinn TC, Chadwick K, Margolick JB, et al. Long-term follow-up studies confirm the stability of the latent reservoir for HIV-1 in resting CD4+ T cells. Nat Med. 2003;9(6):727-8.

7.      Crooks AM, Bateson R, Cope AB, Dahl NP, Griggs MK, Kuruc JD, et al. Precise Quantitation of the Latent HIV-1 Reservoir: Implications for Eradication Strategies. J Infect Dis. 2015;212(9):1361-5.

8.      Bachmann N, von Siebenthal C, Vongrad V, Turk T, Neumann K, Beerenwinkel N, et al. Determinants of HIV-1 reservoir size and long-term dynamics during suppressive ART. Nat Commun. 2019;10(1):3193.

9.      Cohn LB, Chomont N, Deeks SG. The Biology of the HIV-1 Latent Reservoir and Implications for Cure Strategies. Cell Host Microbe. 2020;27(4):519-30.

10.      Coffin JM, Hughes SH. Clonal Expansion of Infected CD4+ T Cells in People Living with HIV. Viruses. 2021;13(10).

11.      Chun TW, Nickle DC, Justement JS, Large D, Semerjian A, Curlin ME, et al. HIV-infected individuals receiving effective antiviral therapy for extended periods of time continually replenish their viral reservoir. J Clin Invest. 2005;115(11):3250-5.

12.      Martinez-Picado J, Deeks SG. Persistent HIV-1 replication during antiretroviral therapy. Curr Opin HIV AIDS. 2016;11(4):417-23.

13.      Sigal A, Kim JT, Balazs AB, Dekel E, Mayo A, Milo R, et al. Cell-to-cell spread of HIV permits ongoing replication despite antiretroviral therapy. Nature. 2011;477(7362):95-8.

14.      Lorenzo-Redondo R, Fryer HR, Bedford T, Kim EY, Archer J, Pond SLK, et al. Persistent HIV-1 replication maintains the tissue reservoir during therapy. Nature. 2016;530(7588):51-6.

15.      Pasternak AO, de Bruin M, Jurriaans S, Bakker M, Berkhout B, Prins JM, et al. Modest nonadherence to antiretroviral therapy promotes residual HIV-1 replication in the absence of virological rebound in plasma. J Infect Dis. 2012;206(9):1443-52.

16.      Pasternak AO, Vroom J, Kootstra NA, Wit FW, de Bruin M, De Francesco D, et al. Non-nucleoside reverse transcriptase inhibitor-based combination antiretroviral therapy is associated with lower cell-associated HIV RNA and DNA levels as compared with therapy based on protease inhibitors. Elife. 2021;10:e68174.

17.      Bozzi G, Simonetti FR, Watters SA, Anderson EM, Gouzoulis M, Kearney MF, et al. No evidence of ongoing HIV replication or compartmentalization in tissues during combination antiretroviral therapy: Implications for HIV eradication. Sci Adv. 2019;5(9):eaav2045.

18.      Kearney MF, Wiegand A, Shao W, McManus WR, Bale MJ, Luke B, et al. Ongoing HIV Replication During ART Reconsidered. Open Forum Infectious Diseases. 2017;4(3):ofx173.

19.      Van Zyl GU, Katusiime MG, Wiegand A, McManus WR, Bale MJ, Halvas EK, et al. No evidence of HIV replication in children on antiretroviral therapy. J Clin Invest. 2017;127(10):3827-34.

20.      Kearney MF, Spindler J, Shao W, Yu S, Anderson EM, O'Shea A, et al. Lack of detectable HIV-1 molecular evolution during suppressive antiretroviral therapy. PLoS Pathog. 2014;10(3):e1004010.

21.      Conway JM, Perelson AS. Residual Viremia in Treated HIV+ Individuals. PLoS Comput Biol. 2016;12(1):e1004677.

22.      Paryad-Zanjani S, Jagarapu A, Piovoso MJ, Zurakowski R. Ongoing HIV replication in lymph node sanctuary sites in treated individuals contributes to the total latent HIV at a very slow rate. J Theor Biol. 2023;575:111651.

23.      Zicari S, Sessa L, Cotugno N, Ruggiero A, Morrocchi E, Concato C, et al. Immune Activation, Inflammation, and Non-AIDS Co-Morbidities in HIV-Infected Patients under Long-Term ART. Viruses. 2019;11(3).

24.      Mazzuti L, Turriziani O, Mezzaroma I. The Many Faces of Immune Activation in HIV-1 Infection: A Multifactorial Interconnection. Biomedicines. 2023;11(1).

25. Chun TW, Nickle DC, Justement JS, Meyers JH, Roby G, Hallahan CW, et al. Persistence of HIV in gut-associated lymphoid tissue despite long-term antiretroviral therapy. J Infect Dis. 2008;197(5):714-20.

26. Fletcher CV, Staskus K, Wietgrefe SW, Rothenberger M, Reilly C, Chipman JG, et al. Persistent HIV-1 replication is associated with lower antiretroviral drug concentrations in lymphatic tissues. Proc Natl Acad Sci U S A. 2014;111(6):2307-12.

27. Fletcher CV, Kroon E, Schacker T, Pinyakorn S, Chomont N, Chottanapund S, et al. Persistent HIV transcription and variable antiretroviral drug penetration in lymph nodes during plasma viral suppression. AIDS. 2022;36(7):985-90.

28. Estes JD, Kityo C, Ssali F, Swainson L, Makamdop KN, Del Prete GQ, et al. Defining total-body AIDS-virus burden with implications for curative strategies. Nat Med. 2017;23(11):1271-1276.

29. Havlir DV, Strain MC, Clerici M, Ignacio C, Trabattoni D, Ferrante P, et al. Productive infection maintains a dynamic steady state of residual viremia in human immunodeficiency virus type 1-infected persons treated with suppressive antiretroviral therapy for five years. J Virol. 2003;77(20):11212-9.

30. Ramratnam B, Ribeiro R, He T, Chung C, Simon V, Vanderhoeven J, et al. Intensification of antiretroviral therapy accelerates the decay of the HIV-1 latent reservoir and decreases, but does not eliminate, ongoing virus replication. J Acquir Immune Defic Syndr. 2004;35(1):33-7.

31. Buzon MJ, Massanella M, Llibre JM, Esteve A, Dahl V, Puertas MC, et al. HIV-1 replication and immune dynamics are affected by raltegravir intensification of HAART-suppressed subjects. Nat Med. 2010;16(4):460-5.

32. Yukl SA, Shergill AK, McQuaid K, Gianella S, Lampiris H, Hare CB, et al. Effect of raltegravir-containing intensification on HIV burden and T-cell activation in multiple gut sites of HIV-positive adults on suppressive antiretroviral therapy. AIDS. 2010;24(16):2451-60.

33. Llibre JM, Buzon MJ, Massanella M, Esteve A, Dahl V, Puertas MC, et al. Treatment intensification with raltegravir in subjects with sustained HIV-1 viraemia suppression: a randomized 48-week study. Antivir Ther. 2012;17(2):355-64.

34. Vallejo A, Gutierrez C, Hernandez-Novoa B, Diaz L, Madrid N, Abad-Fernandez M, et al. The effect of intensification with raltegravir on the HIV-1 reservoir of latently infected memory CD4 T cells in suppressed patients. AIDS. 2012;26(15):1885-94.

35. Gutierrez C, Diaz L, Vallejo A, Hernandez-Novoa B, Abad M, Madrid N, et al. Intensification of antiretroviral therapy with a CCR5 antagonist in patients with chronic HIV-1 infection: effect on T cells latently infected. PLoS One. 2011;6(12):e27864.

36. Hatano H, Strain MC, Scherzer R, Bacchetti P, Wentworth D, Hoh R, et al. Increase in 2-long terminal repeat circles and decrease in D-dimer after raltegravir intensification in patients with treated HIV infection: a randomized, placebo-controlled trial. J Infect Dis. 2013;208(9):1436-42.

37. Puertas MC, Gómez-Mora E, Santos JR, Moltó J, Urrea V, Morón-López S, et al. Impact of intensification with raltegravir on HIV-1-infected individuals receiving monotherapy with boosted PIs. J Antimicrob Chemother. 2018;73(7):1940-8.

38. Puertas MC, Massanella M, Llibre JM, Ballestero M, Buzon MJ, Ouchi D, et al. Intensification of a raltegravir-based regimen with maraviroc in early HIV-1 infection. AIDS. 2014;28(3):325-34.

39. Dinoso JB, Kim SY, Wiegand AM, Palmer SE, Gange SJ, Cranmer L, et al. Treatment intensification does not reduce residual HIV-1 viremia in patients on highly active antiretroviral therapy. Proc Natl Acad Sci U S A. 2009;106(23):9403-8.

40. McMahon D, Jones J, Wiegand A, Gange SJ, Kearney M, Palmer S, et al. Short-course raltegravir intensification does not reduce persistent low-level viremia in patients with HIV-1 suppression during receipt of combination antiretroviral therapy. Clin Infect Dis. 2010;50(6):912-9.

41. Gandhi RT, Coombs RW, Chan ES, Bosch RJ, Zheng L, Margolis DM, et al. No effect of raltegravir intensification on viral replication markers in the blood of HIV-1-infected patients receiving antiretroviral therapy. J Acquir Immune Defic Syndr. 2012;59(3):229-35.

42. Gandhi RT, Zheng L, Bosch RJ, Chan ES, Margolis DM, Read S, et al. The effect of raltegravir intensification on low-level residual viremia in HIV-infected patients on antiretroviral therapy: a randomized controlled trial. PLoS Med. 2010;7(8).

43. Cillo AR, Hilldorfer BB, Lalama CM, McKinnon JE, Coombs RW, Tenorio AR, et al. Virologic and immunologic effects of adding maraviroc to suppressive antiretroviral therapy in individuals with suboptimal CD4+ T-cell recovery. AIDS. 2015;29(16):2121-9.

44.     Chaillon A, Gianella S, Lada SM, Perez-Santiago J, Jordan P, Ignacio C, et al. Size, Composition, and Evolution of HIV DNA Populations during Early Antiretroviral Therapy and Intensification with Maraviroc. J Virol. 2018;92(3).

45.     Rasmussen TA, McMahon JH, Chang JJ, Audsley J, Rhodes A, Tennakoon S, et al. The effect of antiretroviral intensification with dolutegravir on residual virus replication in HIV-infected individuals: a randomised, placebo-controlled, double-blind trial. Lancet HIV. 2018;5(5):e221-e30.

46.     Deeks SG. HIV: Shock and kill. Nature. 2012;487(7408):439-40.

47.     Greener BN, Patterson KB, Prince HM, Sykes CS, Adams JL, Dumond JB, et al. Dolutegravir pharmacokinetics in the genital tract and colorectum of HIV-negative men after single and multiple dosing. J Acquir Immune Defic Syndr. 2013;64(1):39-44.

48.     Podsadecki TJ, Vrijens BC, Tousset EP, Rode RA, Hanna GJ. "White coat compliance" limits the reliability of therapeutic drug monitoring in HIV-1-infected patients. HIV Clin Trials. 2008;9(4):238-46.

49.     Bruner KM, Wang Z, Simonetti FR, Bender AM, Kwon KJ, Sengupta S, et al. A quantitative approach for measuring the reservoir of latent HIV-1 proviruses. Nature. 2019;566(7742):120-5.

50.     Scherpenisse M, Kootstra NA, Bakker M, Berkhout B, Pasternak AO. Cell-Associated HIV-1 Unspliced-to-Multiply-Spliced RNA Ratio at 12 Weeks of ART Predicts Immune Reconstitution on Therapy. mBio. 2021;12(2).

51.     Peluso MJ, Bacchetti P, Ritter KD, Beg S, Lai J, Martin JN, et al. Differential decay of intact and defective proviral DNA in HIV-1-infected individuals on suppressive antiretroviral therapy. JCI Insight. 2020;5(4).

52.     Gandhi RT, Cyktor JC, Bosch RJ, Mar H, Laird GM, Martin A, et al. Selective Decay of Intact HIV-1 Proviral DNA on Antiretroviral Therapy. J Infect Dis. 2021;223(2):225-33.

53.     Chew GM, Fujita T, Webb GM, Burwitz BJ, Wu HL, Reed JS, et al. TIGIT Marks Exhausted T Cells, Correlates with Disease Progression, and Serves as a Target for Immune Restoration in HIV and SIV Infection. PLoS Pathog. 2016;12(1):e1005349.

54.     Fromentin R, Bakeman W, Lawani MB, Khoury G, Hartogensis W, DaFonseca S, et al. CD4+ T Cells Expressing PD-1, TIGIT and LAG-3 Contribute to HIV Persistence during ART. PLoS Pathog. 2016;12(7):e1005761.

55.     Darcis G, Maes N, Pasternak AO, Sauvage AS, Frippiat F, Meuris C, et al. Detectability of HIV Residual Viremia despite Therapy Is Highly Associated with Treatment with a Protease Inhibitor-Based Combination Antiretroviral Therapy. Antimicrob Agents Chemother. 2020;64(3).

56.     Darcis G, Berkhout B, Pasternak AO. Differences in HIV Markers between Infected Individuals Treated with Different ART Regimens: Implications for the Persistence of Viral Reservoirs. Viruses. 2020;12(5).

57.     Hatano H, Hayes TL, Dahl V, Sinclair E, Lee TH, Hoh R, et al. A randomized, controlled trial of raltegravir intensification in antiretroviral-treated, HIV-infected patients with a suboptimal CD4+ T cell response. J Infect Dis. 2011;203(7):960-8.

58.     Butler SL, Hansen MS, Bushman FD. A quantitative assay for HIV DNA integration in vivo. Nat Med. 2001;7(5):631-4.

59.     Hockett RD, Kilby JM, Derdeyn CA, Saag MS, Sillers M, Squires K, et al. Constant mean viral copy number per infected cell in tissues regardless of high, low, or undetectable plasma HIV RNA. J Exp Med. 1999;189(10):1545-54.

60.     Rothenberger M, Nganou-Makamdop K, Kityo C, Ssali F, Chipman JG, Beilman GJ, et al. Impact of Integrase Inhibition compared to non-nucleoside inhibition on HIV reservoirs in Lymphoid Tissues. J Acquir Immune Defic Syndr. 2019.

61.     Lee SA, Telwatte S, Hatano H, Kashuba ADM, Cottrell ML, Hoh R, et al. Antiretroviral Therapy Concentrations Differ in Gut vs. Lymph Node Tissues and Are Associated With HIV Viral Transcription by a Novel RT-ddPCR Assay. J Acquir Immune Defic Syndr. 2020;83(5):530-7.

62.     Galloway NL, Doitsh G, Monroe KM, Yang Z, Muñoz-Arias I, Levy DN, et al. Cell-to-Cell Transmission of HIV-1 Is Required to Trigger Pyroptotic Death of Lymphoid-Tissue-Derived CD4 T Cells. Cell Rep. 2015;12(10):1555-63.

63.     Doitsh G, Cavrois M, Lassen KG, Zepeda O, Yang Z, Santiago ML, et al. Abortive HIV infection mediates CD4 T cell depletion and inflammation in human lymphoid tissue. Cell. 2010;143(5):789-801.

64. Boom R, Sol CJ, Salimans MM, Jansen CL, Wertheim-van Dillen PM, van der Noordaa J.
Rapid and simple method for purification of nucleic acids. J Clin Microbiol. 1990;28(3):495-503.
65. Malnati MS, Scarlatti G, Gatto F, Salvatori F, Cassina G, Rutigliano T, et al. A universal real-
time PCR assay for the quantification of group-M HIV-1 proviral load. Nat Protoc. 2008;3(7):1240-8.
66. Pasternak AO, Jurriaans S, Bakker M, Prins JM, Berkhout B, Lukashov VV. Cellular levels of
HIV unspliced RNA from patients on combination antiretroviral therapy with undetectable plasma
viremia predict the therapy outcome. PLoS One. 2009;4(12):e8490.
67. Fischer M, Huber W, Kallivroussis A, Ott P, Opravil M, Luthy R, et al. Highly sensitive methods
for quantitation of human immunodeficiency virus type 1 RNA from plasma, cells, and tissues. J Clin
Microbiol. 1999;37(5):1260-4.
68. Levy CN, Hughes SM, Roychoudhury P, Reeves DB, Amstuz C, Zhu H, et al. A highly
multiplexed droplet digital PCR assay to measure the intact HIV-1 proviral reservoir. Cell Rep Med.
2021;2(4):100243.
69. Luo W, Yang H, Rathbun K, Pau CP, Ou CY. Detection of human immunodeficiency virus type
1 DNA in dried blood spots by a duplex real-time PCR assay. J Clin Microbiol. 2005;43(4):1851-7.
70. Fletcher CV, Podany AT, Thorkelson A, Winchester LC, Mykris T, Anderson J, et al. The
Lymphoid Tissue Pharmacokinetics of Tenofovir Disoproxil Fumarate and Tenofovir Alafenamide in
HIV-Infected Persons. Clin Pharmacol Ther. 2020;108(5):971-5.
71. Dyavar SR, Kumar S, Gautam N, Podany AT, Winchester LC, Weinhold JA, et al. Intramuscular
and subcutaneous administration of antiretroviral drugs, compared with oral, enhances delivery to
lymphoid tissues in BALB/c mice. J Antimicrob Chemother. 2021;76(10):2651-8.
72. Delahunty T, Bushman L, Fletcher CV. Sensitive assay for determining plasma tenofovir
concentrations by LC/MS/MS. J Chromatogr B Analyt Technol Biomed Life Sci. 2006;830(1):6-12.
741
742

 **Figure legends**

 **Figure 1**. Antiretroviral drug concentrations in plasma and rectal tissue. Plasma (A) and tissue

 (B) concentrations of DTG, as well as plasma (C) and tissue (D) concentrations of 3TC, were

 compared between days 0 (D0) and 84 (D84) in intensified and control groups. Wilcoxon tests

 were used to calculate statistical significance.

 **Figure 2**. Longitudinal dynamics of HIV reservoir markers. (A-C) Fold change of total HIV

 DNA in PBMCs (A), US HIV RNA in PBMCs (B), and US RNA/total DNA ratios (C) from

 baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups.

 Median values and IQRs are shown. Linear mixed-effects modelling on $\log_{10}$-transformed

 values was used to calculate statistical significance. P values at the bottom of the graphs show

 the significance of between-group comparisons and those on top of the graphs show the

 significance of comparisons of the changes from baseline with zero in each group separately

 (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a

 statistically significant increase or decrease from baseline, respectively. Participant numbers in

 both groups per time point are indicated below the graphs. (D, E) Comparisons of intact HIV

 DNA in PBMCs (D) and of total HIV DNA in rectal tissue (E) between days 0 (D0) and 84

 (D84) in the intensified and control groups. Open circles depict undetectable values, assigned

 the values corresponding to 50% of the assay detection limits. Wilcoxon tests were used to

 calculate statistical significance. All p values are marked red if significant ($<0.05$).

 **Figure 3**. Longitudinal dynamics of cellular markers of immune activation and exhaustion.

 Graphs show changes of (A) CD4+ cell markers and (B) CD8+ cell markers from baseline on

 days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median

 values and IQRs are shown. Linear mixed-effects modelling was used to calculate statistical

 significance. P values at the bottom of the graphs show the significance of between-group

 comparisons and those on top of the graphs show the significance of comparisons of the

 changes from baseline with zero in each group separately (intercept-only analysis). An upward

 or downward facing arrow next to a p value indicates a statistically significant increase or

 decrease from baseline, respectively. P values are marked red if significant ($<0.05$).

 **Figure 4**. Longitudinal dynamics of inflammation markers in plasma and tissue. (A) Plasma

 inflammation markers. Graphs show changes of markers from baseline on days 1, 28, 56, and

 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are

 shown. Linear mixed-effects modelling was used to calculate statistical significance. P values

at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. P values are marked red if significant ($<0.05$). (B) Tissue inflammation markers. Markers were compared between days 0 (D0) and 84 (D84) in the intensified and control groups. Open circles depict undetectable values, assigned the values corresponding to 50% of the assay detection limits. Wilcoxon tests were used to calculate statistical significance.

**Figure 5**. Longitudinal dynamics of clinical markers. Graphs show changes of markers from baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are shown. Linear mixed-effects modelling was used to calculate statistical significance. P values at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. P values are marked red if significant ($<0.05$).

792 Table 1. Clinical characteristics of participants at baseline.

| | All (n=20) | Control group (n=10) | DTG group (n=10) | p[a] |
|---|---|---|---|---|
| Sex, male | 19 (95.0)[b] | 9 (90.0) | 10 (100.0) | 1.00 |
| Age, years | 52 (43-60) | 52 (40-62) | 51 (46-59) | 0.94 |
| BMI, kg/m² | 24 (22-26) | 25 (22-27) | 23 (22-26) | 0.62 |
| Ethnicity | | | | |
| Caucasian | 17 (85.0) | 9 (90.0) | 8 (80.0) | 1.00 |
| African | 2 (10.0) | 1 (10.0) | 1 (10.0) | |
| Maghrebi | 1 (5.0) | 0 (0.0) | 1 (10.0) | |
| HLA typing B57/01, negative (n=19) | 19 (100.0) | 9 (100.0) | 10 (100.0) | - |
| Smoking, smoker or ex-smoker | 10 (50.0) | 4 (40.0) | 6 (60.0) | 0.66 |
| Time since first positive HIV serology, years | 12.6 (6.8-18.2) | 9.8 (7.3-13.3) | 15.6 (7.6-18.5) | 0.58 |
| Cumulative time of untreated HIV infection, years | 0.9 (0.1-3.5) | 1.8 (0.1-5.0) | 0.6 (0.1-2.1) | 0.47 |
| Cumulative time of viral suppression, years | 7.4 (4.8-9.2) | 5.6 (4.7-7.7) | 7.9 (5.8 – 12.4) | 0.32 |
| Continuous time of viral suppression before study, years | 3.9 (3.3-5.8) | 4.4 (3.7-6.9) | 3.4 (3.2-4.9) | 0.12 |
| Time on DTG-containing ART regimen before study, years | 3.5 (3.4-3.7) | 3.5 (3.4-3.8) | 3.5 (3.3-3.5) | 0.41 |
| Nadir CD4+ count, cells/mm³ | 277 (147-410) | 277 (205-340) | 292 (136-508) | 0.65 |
| First measured plasma viral load, $\log_{10}$ HIV RNA copies/mL | 4.52 (3.91-5.35) | 4.52 (4.36-4.96) | 4.55 (3.69-5.38) | 0.85 |
| CD4+ count, cells/mm³ | 775 (659-1144) | 732 (672-970) | 886 (668-1217) | 0.39 |
| CD8+ count, cells/mm³ | 745 (637-1288) | 723 (535-931) | 939 (690-1310) | 0.39 |
| CD4/CD8 ratio | 0.94 (0.66-1.47) | 1.00 (0.64-1.36) | 0.93 (0.69-1.56) | 0.82 |
| Total HIV DNA in PBMCs, copies/$10^6$ cells | 175.5 (70.0-467.3) | 99.4 (37.4-231.5) | 421.0 (139.5-764.3) | 0.015 |
| Intact HIV DNA in PBMCs, copies/$10^6$ cells | 55.2 (19.2-87.0) | 29.5 (16.2-97.1) | 58.6 (44.7-144.7) | 0.36 |
| US HIV RNA in PBMCs, copies/µg total RNA | 92.1 (13.3-290.3) | 72.2 (12.2-253.8) | 124.6 (17.0-591.0) | 0.58 |
| US RNA/total DNA ratio in PBMCs | 0.40 (0.07-1.47) | 1.21 (0.11-2.81) | 0.33 (0.06-0.70) | 0.21 |
| Total HIV DNA in rectal tissue, copies/$10^6$ cells | 477.0 (271.3-971.5) | 547.0 (291.0-1215) | 403.0 (194.6-952.5) | 0.44 |
| Plasma DTG concentration, ng/mL | 3287 (2602-5087) | 3560 (2953-4455) | 3266 (2245-5983) | 0.82 |
| Tissue DTG concentration, ng/g | 634 (533-830) | 737 (534-852) | 581 (535-714) | 0.44 |
| Plasma 3TC concentration, ng/mL | 316 (148-731) | 431 (214-941) | 246 (117-439) | 0.26 |
| Tissue 3TC concentration, ng/g | 2114 (1417-2345) | 2193 (1590-2784) | 2114 (1233-2317) | 0.50 |
| CDC classification | | | | - |
| A1 | 4 (20.0) | 1 (10.0) | 3 (30.0) | |
| A2 | 9 (45.0) | 6 (60.0) | 3 (30.0) | |
| A3 | 3 (15.0) | 1 (10.0) | 2 (30.0) | |
| B3 | 1 (5.0) | 0 (0.0) | 1 (10.0) | |
| C3 | 3 (15.0) | 2 (20.0) | 1 (10.0) | |
| HBV status | | | | - |
| Immune | 12 (60.0) | 8 (80.0) | 4 (40.0) | |
| Non-immune, not infected | 5 (25.0) | 2 (20.0) | 3 (30.0) | |
| Isolated HBc Ab | 2 (10.0) | 0 (0.0) | 2 (20.0) | |
| Cured Hepatitis B | 1 (5.0) | 0 (0.0) | 1 (10.0) | |
| HCV status | | | | 1.00 |
| Not Infected | 19 (95.0) | 10 (10.0) | 9 (90.0) | |
| Recovered | 1 (5.0) | 0 (0.0) | 1 (10.0) | |

**Figure 1**. Antiretroviral drug concentrations in plasma and rectal tissue. Plasma (A) and tissue (B) concentrations of DTG, as well as plasma (C) and tissue (D) concentrations of 3TC, were compared between days 0 (D0) and 84 (D84) in intensified and control groups. Wilcoxon tests were used to calculate statistical significance.

**Figure 2**. Longitudinal dynamics of HIV reservoir markers. (A-C) Fold change of total HIV DNA in PBMCs (A), US HIV RNA in PBMCs (B), and US RNA/total DNA ratios (C) from baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are shown. Linear mixed-effects modelling on log10-transformed values was used to calculate statistical significance. P values at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. Participant numbers in both groups per time point are indicated below the graphs. (D, E) Comparisons of intact HIV DNA in PBMCs (D) and of total HIV DNA in rectal tissue (E) between days 0 (D0) and 84 (D84) in the intensified and control groups. Open circles depict undetectable values, assigned the values corresponding to 50% of the assay detection limits. Wilcoxon tests were used to calculate statistical significance. All p values are marked red if significant (<0.05).

**Figure 3**. Longitudinal dynamics of cellular markers of immune activation and exhaustion. Graphs show changes of (A) CD4+ cell markers and (B) CD8+ cell markers from baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are shown. Linear mixed-effects modelling was used to calculate statistical significance. P values at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. P values are marked red if significant (<0.05).

**Figure 4**. Longitudinal dynamics of inflammation markers in plasma and tissue. (A) Plasma inflammation markers. Graphs show changes of markers from baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are shown. Linear mixed-effects modelling was used to calculate statistical significance. P values at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. P values are marked red if significant (<0.05). (B) Tissue inflammation markers. Markers were compared between days 0 (D0) and 84 (D84) in the intensified and control groups. Open circles depict undetectable values, assigned the values corresponding to 50% of the assay detection limits. Wilcoxon tests were used to calculate statistical significance.

**Figure 5**. Longitudinal dynamics of clinical markers. Graphs show changes of markers from baseline on days 1, 28, 56, and 84 of the study in the intensified (blue) and control (red) groups. Median values and IQRs are shown. Linear mixed-effects modelling was used to calculate statistical significance. P values at the bottom of the graphs show the significance of between-group comparisons and those on top of the graphs show the significance of comparisons of the changes from baseline with zero in each group separately (intercept-only analysis). An upward or downward facing arrow next to a p value indicates a statistically significant increase or decrease from baseline, respectively. P values are marked red if significant (<0.05).