Incidence, Persistence, and Steady-State Prevalence in Coding Intensity for Health Plan Payment

Thomas G. McGuire, PhD[1]; Oana M. Enache, MS[2*]; Michael Chernew, PhD[1]; J. Michael McWilliams, MD, PhD[1,3]; Tram Nham, BA[1]; Sherri Rose, PhD[4]

[1] Department of Healthcare Policy, Harvard Medical School, Boston, MA

[2] Department of Biomedical Data Science, Stanford School of Medicine, Stanford, CA

[3] Brigham and Women's Hospital, Boston, MA

[4] Department of Health Policy, Stanford University, Stanford, CA

*Correspondence to:

Oana M. Enache

Encina Commons, 615 Crothers Way

Stanford University

Stanford, CA 94305

oenache@stanford.edu

Structured Abstract

Objective. To define measures of Medicare diagnosis coding intensity that capture the dynamics of changes in coding prevalence.

Study setting and design. Retrospective analysis of coding practices for risk adjustment using observational claims data from Medicare beneficiaries assigned to Accountable Care Organizations (ACOs) in 2017 and 2018.

Data sources. Enrollment and claims data from 2017-2018 of a random 20 percent sample of Medicare beneficiaries were subset to include those assigned to an ACO in 2018. Beneficiaries were included in analyses if they were continuously enrolled in Medicare Parts A and B in 2017 and 2018 and aged 65 years or older during that time. Beneficiaries were excluded if they were originally enrolled in Medicare due to end-stage renal disease or were residing outside of the United States.

Principal findings. Borrowing terminology from epidemiology, prevalence of a diagnosis code can be decomposed into incidence (the proportion of beneficiaries that newly have the code) and persistence (the proportion of beneficiaries who previously had the code and continue to do so). Together, these two measures define what we call steady-state prevalence, the hypothetical long-run prevalence implied by no changes in current rates of incidence and persistence of coding. The concept of steady-state prevalence can help explain why observed prevalence tends to grow over time, apart from continued behavioral change. We demonstrate this and also illustrate the application of our measures for the most expensive diagnoses in Medicare's risk-adjustment payment model for ACOs.

Conclusions. Diagnostic coding is a highly heterogeneous behavior that can vary in response to changes in incentives. Researchers and policymakers can better monitor such behavior by

measuring the incidence and persistence of a given code separately. In addition, changes in coding practices can take years to be fully reflected in current coding prevalence even in the absence of further behavioral change.

<u>Keywords</u>

Clinical coding

Risk adjustment

Medicare

Accountable Care Organizations

Medicare Advantage

<u>Callout Box</u>

What is known on this topic:

- Medicare incentivizes providers and insurers to code for as many diagnoses per beneficiary as possible, contributing to billions of dollars of Medicare program spending.
- Empirical research indicates that the frequency of diagnostic coding in Medicare has increased steadily in recent years.
- Current measures of coding practices for a given diagnosis largely count the prevalence, or the proportion of beneficiaries with a specified code, in one time period.

What this study adds:

- This study proposes several new measures of coding practices decomposing diagnosis code prevalence into incident and persistent codes over two years, which policymakers can use to monitor coding behaviors.

- This study defines steady-state prevalence as the ultimate prevalence of a code implied by current coding patterns hypothetically remaining unchanged.

- Applying our measures to Medicare Accountable Care Organization coding finds that at current rates of incidence and persistence, observed prevalence of coding will increase even without further behavioral change.

<u>Introduction</u>

The Centers for Medicare and Medicaid Services (CMS) finances care for Medicare

beneficiaries using risk contracts in two major programs. The first is Medicare Advantage (MA),

where healthcare plans are administered by contracted private insurers[1]. MA enrollment has

grown rapidly in recent years and surpassed 51 percent of all eligible Medicare beneficiaries, or

31 million people[1]. Beneficiaries not enrolled in MA are said to be in Traditional Medicare (or

TM) where their care might be financed through a second risk contract with a group of providers

within the Accountable Care Organization (ACO) program[2]. Medicare assigns a beneficiary to an

ACO largely based on their use of primary care services[3].

Both MA and ACOs rely on the same baseline model (commonly referred to as the

CMS-HCC model) to determine payment per beneficiary. CMS-HCC is a linear regression

model with positive coefficients that predicts per-person spending as a function of each

beneficiary's diagnoses and demographic information[4]. For each enrollee, CMS obtains

diagnosis codes from physician, hospital inpatient, and hospital outpatient claims from the

previous year to predict costs for the performance year. These risk scores are then used to adjust

a benchmark for an ACO or MA plan's beneficiaries according to their expected expenditures, an

approach referred to as risk adjustment[4]. The MA and ACO programs risk adjust payments to

MA plans and ACOs with the aim to provide adequate resources to care for beneficiaries and to

incentivize insurers and provider groups to participate in the programs without avoiding

high-cost patients[4,5].

Although there are some differences in how payment is computed in ACO and MA

contexts specifically, both of these risk-adjusted systems are premised on accurate information

from providers and plans about patients' health conditions. If existing coding is irregular or

incomplete, then higher rates of coding by less intensive coders would improve the potential fit

of risk adjustment payment formulas and the overall functioning of health plan markets. On the

other hand, coding that does not accurately reflect a beneficiary's health conditions can interfere

with the efficiency and fairness of the risk adjustment payment formula. Both ACOs and MA

plans are incentivized to code more intensely by the current payment system, as more diagnoses

result in higher predicted spending by CMS-HCC[4,5].

Prior research has established that coding intensity is substantially higher in MA than in

TM[6] and highly variable across MA plans and ACOs[7-9]. Higher coding intensity in the risk

programs relative to TM for any reason increases federal spending on Medicare programs,

making it important to understand trends in the prevalence of codes. This paper proposes and

studies new measures of coding intensity meant to capture the built-in dynamics associated with

coding practices. In line with established terminology in epidemiology and other fields, we

decompose the components of coding into incidence, the rate at which a code appears among

those previously without the code corresponding to a given diagnosis, and persistence, or the rate

at which a code continues among those with that code at an earlier time point. Although

persistence in coding has been studied previously[10-12], the dynamic implications of incidence and

persistence have not been formally investigated to our knowledge.

To capture the long-run consequences of incidence and persistence, we also define

steady-state prevalence, the ultimate prevalence of a code implied by the current rate of

incidence and persistence if current coding patterns were to remain unchanged. Steady-state

prevalence treats coding as analogous to a one-period Markov process. A positive gap between

the steady-state prevalence implied by current patterns of coding and observed prevalence

signals that current prevalence will continue to increase even in the absence of any behavioral

change in coding practices. To illustrate these concepts, we study incidence, persistence, and steady-state prevalence in Medicare's ACO program with data from 2017 and 2018.

Methods

This project was approved by the CMS privacy board and the Harvard Medical School Institutional Review Committee, which also waived the requirement for obtaining informed consent because all administrative data were deidentified.

Data

We confined our study to the ACO sector in TM because the incentives for coding are stronger than in the non-ACO sector and the data were more readily available than MA data to researchers. We construct our measures from a random 20 percent sample of individuals continuously enrolled in Parts A and B of fee-for-service (FFS) Medicare in both 2017 and 2018. We exclude beneficiaries who are either aged younger than 65, originally enrolled at least in part due to end-stage renal disease (ESRD), or reside outside the United States.

Our study sample consists of 1,799,068 beneficiaries who were assigned to an ACO in 2018 as identified based on the CMS ACO Shared Savings Program Beneficiary-level file. These exclusion and inclusion criteria produce a sample of 1,798,319 individuals. We used the 2017-2018 version (version 22, or V22) of the CMS-HCC prospective risk adjustment formula to derive hierarchical condition category (HCC) variables from ICD-10 codes. This version of the formula uses demographic information as well as 79 HCC variables that represent presence or absence of various diagnoses. The outcome variable of the V22 formula is projected healthcare costs per beneficiary in 2018.

Statistical Analysis Procedure

One year of claims data contains the information needed to calculate *current prevalence,* p_t, the proportion of beneficiaries with a disease code in a population in year t. Two years of data are required to measure *incidence, i,* and *persistence, r,* the rate of codes occurring among those previously without and with the code, respectively. p_t depends on p_{t-1} and the rates of incidence and persistence: $p_t = (p_{t-1})r + (1 - p_{t-1})i$. The *steady-state prevalence, p,* is a hypothetical future value of prevalence that will be reached if the current rates of incidence and persistence do not change. In a steady state, $p = p_t = p_{t-1}$, so the steady-state prevalence satisfies

$p = pr + (1 - p)i$, which simplifies to $p = i/(1 - r + i)$. Therefore, steady-state prevalence depends only on incidence and persistence and can be computed directly from these two values independent of the current prevalence.

Results

Descriptive Data

Demographics of beneficiaries included as study participants are reported in Table 1, all of whom were enrolled in ACOs in 2017 and 2018. Geographically, 42.2 percent of participants reside in the South, 24.9 percent reside in the Midwest, 19.8 percent reside in the Northeast, and 13.0 percent reside in the Western United States. 9.2 percent of overall participants are eligible for Medicaid and 9.0 percent of overall participants have a disability. In terms of healthcare utilization, 17.0 percent of our sample had at least one inpatient stay and 73.7 percent of overall participants had at least one HCC code assigned in 2018, with each person having 2.1 HCCs

recorded on average. Average annual spending per beneficiary overall is $7,911.10, in part

because ACOs are not responsible for drug spending covered under Part D.

Incidence and Persistence

Figure 1 depicts incidence and persistence for the 10 HCCs with the highest contribution

to average risk scores among the ACOs in 2018, where contribution of an HCC to the risk score

is the product of the 2018 current prevalence among ACOs times the weight of that HCC in the

CMS-HCC community risk score formula. Together, these HCCs account for 39.8 percent of the

average risk score.

Incidence for diabetes with chronic complications, the HCC contributing the most to

ACO plan payment, is 5 percent, meaning that among the enrollees without diabetes in a year, 5

percent are newly diagnosed as having diabetes compared to the prior year. Vascular disease,

specified heart arrhythmias, and congestive heart failure all have incidence rates greater than 5

percent (Figure 1a), which is perhaps not unexpected since about half of the population in our

sample is 75 years or older (Table 1).

Figure 1b depicts persistence, showing that most enrollees with one of the top ten HCCs

continue to be coded with the HCC in a subsequent year. An exception is acute renal failure

(currently known as acute kidney injury), which can improve with management and without

dialysis for the majority of patients[13] and is therefore unlikely to persist as a code over several

years. However, rates of persistence, even for highly chronic conditions like diabetes and chronic

obstructive pulmonary disease fall well below 100 percent. Some unexpectedly low rates of

persistence may be due to miscoding or undercoding. For example, an analysis of data from

private health insurance in 2013 and 2014 found that persistence of the HCC indicating the very

permanent diagnosis of paraplegia was only 57 percent[4] and a more recent analysis of 2017-2019 data found that TM undercoding accounted for 22.3 percent of the difference in risk scores between TM and MA[12.]. This suggests that such coding-related issues may be present across the insurance sector more broadly.

The empirical distinction between current and steady-state prevalence can be illustrated using HCC58 (Major Depression, Bipolar and Paranoid Disorders) in Table 2. These conditions are often, but not always, chronic and roughly half of patients with major depression experience just one episode in their lifetime[14]. Clinically, bipolar and paranoid disorders persist at higher rates, but even for these conditions, disease remission and recovery take place.

The first two columns in Table 2 show current prevalence for HCC58 rising from 7.1 percent in 2017 to 8.8 percent in 2018. The next two columns introduce the decomposition of current prevalence. Incidence of the code between 2017 and 2018 is 4.1 percent, and persistence is 70.1 percent. In other words, among the 92.9 percent of beneficiaries without an indication for HCC58 in 2017 (one minus the current prevalence in 2017), 4.1 percent gain the code in 2018. Among the 7.1 percent with the code in 2017, 70.1 percent remain with the code in 2018. Steady-state prevalence calculated according to the formula above is shown in the final column. If present practice of coding in terms of incidence and persistence do not change, the prevalence of HCC58 is expected to increase above the current prevalence of 8.8 percent to closer to 12.0 percent.

Current and Steady-State Prevalence

Figure 2 compares current and steady-state prevalence for the top 10 HCCs in 2018. In all cases, steady-state prevalence exceeds current prevalence, indicating that current prevalence is

likely to increase over time for all ten HCCs. In some cases, the gap between steady-state and current prevalence is particularly large, as in the 28.0 percent steady-state and 18.7 percent current prevalence for specified heart arrhythmias, signaling that particularly rapid increases in current prevalence can be expected for this condition.

We further illustrate this in Figure 3, which shows that current prevalence can continue to increase even if coding behaviors do not change for the same 10 HCCs. Incidence and persistence are held fixed at their 2017-2018 values. These are then used to calculate hypothetical projected prevalence values from 2018 current prevalence (time=0), which are plotted in subsequent years. Although the slope of each HCC's projected prevalence varies over time, all of them increase for several years beyond 2018 and some (for instance, specified heart arrhythmias) increase for all ten years shown. This means that a one-time change in coding behavior alone could account for continued growth of ACOs or MA coding intensity (relative to non-ACO TM or TM coding respectively) for a significant amount of time.

Discussion

In most of the research and policy literature, the current prevalence of diagnostic codes is reported as the frequency of a code within a year's data. Given disease chronicity, it is also useful to describe such coding as a pair of behaviors. Analogous to epidemiology, current prevalence in diagnostic coding depends on two underlying parameters: incidence and persistence of codes. Each of these behaviors are useful to monitor on their own and together define the steady-state prevalence implied by current coding practices.

Notably, viewing coding prevalence as a simple dynamic process has a number of limitations that can be improved upon in future research. First, our analysis is descriptive, not

normative; our proposed measures do not by themselves answer the question whether changes in prevalence are due to better coding or opportunistic coding. Further, although we only consider two years of data for simplicity of computation, including more years in future analyses may be more informative. For example, a multiperiod Markov model or analyses involving more than two years of data would allow a richer characterization of the disease coding process but is not pursued here to keep the analysis straightforward. Doing so might be especially relevant in evaluating coding intensity changes as CMS-HCC model updates are phased in, which typically happens over the course of several years. Relatedly, we note that we illustrate our approach using an older version of the CMS-HCC model (V22 instead of the more current V28) as it best aligns with our data; however, future work could extend to other model versions as well.

In addition, we do not explicitly define comparison populations to use in computing rates of coding; we do so intentionally to emphasize the flexibility of our proposed measures. However, we note that additional comparison populations (for example non-ACO TM beneficiaries in comparison to ACO beneficiaries) may be informative to examine in assessing coding patterns. Although we only apply our dynamic characterization of coding in ACO populations, the measures we propose can readily be extended to other groups of individuals as well, including MA beneficiaries.

In any case, the novel measures we propose are a simple but informative tool for both clinicians and policymakers in a number of ways. Besides applications mentioned above, clinicians could evaluate the incidence and persistence of disease codes against the expected incidence and persistence of disease itself in a population, ideally using epidemiologic data free of coding incentives. Similarly, policymakers could compare incidence in codes from ACOs or

MA plans to incidence from community-based surveys where indications of disease are not used for payment to inform further development in Medicare payment policies[9].

<u>Conclusion</u>

Concern about coding intensity plays a large role in payer policy and in the evaluation of the functioning of diagnostic-based risk adjustment systems, including but not limited to the CMS-HCC model. The decomposition of coding patterns into incidence and persistence, and the use of these to identify steady-state prevalence provide an informative tool to characterize coding and offer a partial explanation of why the observed prevalence of codes increases even in the absence of continued behavioral change.

Acknowledgments

We thank Tim Layton, Richard Kronick, and members of the Risk Adjustment Network for comments on an earlier version. TGM, MC, JMM, and TN were funded by the National Institute on Aging of the National Institutes of Health under award number P01-AG032952. OME was funded by the National Library of Medicine grant 5T15LM007033- 40 and the Stanford Interdisciplinary Graduate Fellowship. SR was funded by the NIH Director's New Innovator Award DP2-MD012722.

Conflicts of Interest

TGM, OME, TN, and SR report no conflicts of interest. MC reports being the Chair of MedPAC and sits on the board of Massachusetts Health Connector and previously the HCCI, serving on advisory boards and panels for NIHCM, CBO, BCBSA, BHI, Aledade Inc., and equity in Waymark Inc. JMM reports research support from the National Institute on Aging, Agency for Healthcare Research and Quality, Arnold Ventures, and the Commonwealth Fund, and from an institutional gift for research and education from Humana to Harvard Medical School; personal income from Oak Ridge Associated Universities for services as a senior adviser to the Center for Medicare and Medicaid Innovation; consulting fees from RTI International, MITRE, Analysis Group, Phillips and Cohen, Action Now Initiative, and the West Health Policy Center; speaker honorarium from America's Physician Groups; and service as a member of the board of directors for the Institute for Accountable Care. The content of this article is solely the responsibility of the authors and does not necessarily reflect the views of any organization with which they are affiliated.

Data availability statement

No data are available. Under the requisite data use agreements, data for the analyses are not publicly available.

References

1. Biniek JF, Freed M, Damico A, Neuman T. Half of All Eligible Medicare Beneficiaries Are Now Enrolled in Private Medicare Advantage Plans. Kaiser Family Foundation. Accessed June 15, 2023. https://www.kff.org/policy-watch/half-of-all-eligible-medicare-beneficiaries-are-now-enr olled-in-private-medicare-advantage-plans/

2. *CMS Innovation Center Strategy Refresh*. Centers for Medicare and Medicaid Services; 2021. Accessed April 10, 2024. https://innovation.cms.gov/strategic-direction-whitepaper

3. *Assignment of Beneficiaries to Accountable Care Organizations Participating in the Medicare Shared Savings Program*. Centers for Medicare and Medicaid Services; 2011. Accessed October 10, 2024. https://www.cms.gov/newsroom/fact-sheets/assignment-beneficiaries-accountable-care-or ganizations-participating-medicare-shared-savings

4. Ellis RP, Martins B, Rose S. Chapter 3 - Risk Adjustment for Health Plan Payment. In: McGuire TG, van Kleef RC, eds. *Risk Adjustment, Risk Sharing and Premium Regulation in Health Insurance Markets*. Academic Press; 2018:55-104. doi:10.1016/B978-0-12-811325-7.00003-8

5. Carlin CS, Feldman R, Jung J. The mechanics of risk adjustment and incentives for coding intensity in Medicare. *Health Serv Res*. 2024;59(3):e14272. doi:10.1111/1475-6773.14272

6. Medicare Payment Advisory Commission. *Report to Congress: Medicare Payment Policy*; 2024. Accessed April 30, 2024.

https://www.medpac.gov/wp-content/uploads/2024/03/Mar24_MedPAC_Report_To_Con
gress_SEC.pdf

7. Rose S, Zaslavsky AM, McWilliams JM. Variation In Accountable Care Organization Spending And Sensitivity To Risk Adjustment: Implications For Benchmarking. *Health Aff* . 2016;35(3):440-448. doi:10.1377/hlthaff.2015.1026

8. Chernew ME, Carichner J, Impreso J, et al. Coding-Driven Changes In Measured Risk In Accountable Care Organizations. *Health Aff* . 2021;40(12):1909-1917. doi:10.1377/hlthaff.2021.00361

9. McWilliams JM, Weinreb G, Landrum MB, Chernew ME. Use Of Patient Health Survey Data For Risk Adjustment To Limit Distortionary Coding Incentives In Medicare: Article examines use of patient health survey data for risk adjustment to limit distortionary coding incentives in Medicare. *Health Aff (Millwood)*. 2025;44(1):48-57. doi:10.1377/hlthaff.2023.01351

10. *Announcement of Calendar Year (CY) 2008 Medicare Advantage Capitation Rates and Payment Policies*. Centers for Medicare and Medicaid Services; 2007. Accessed October 10, 2024. https://www.cms.gov/Medicare/Health-Plans/MedicareAdvtgSpecRateStats/Downloads/a nnouncement2008.pdf

11. Copeland L. CMS Finalizes 2024 Medicare Advantage Payment Policies. Medicare Rights Center. April 6, 2023. Accessed October 10, 2024. https://www.medicarerights.org/medicare-watch/2023/04/06/cms-finalizes-2024-medicar e-advantage-payment-policies

12. Datta NG, Chernew ME, McWilliams JM. Lack of Persistent Coding in Traditional Medicare May Widen The Risk-Score Gap With Medicare Advantage. *Health Aff (Millwood)*. doi:10.1377/hlthaff.2024.00169

13. Vijayan A, Abdel-Rahman EM, Liu KD, et al. Recovery after Critical Illness and Acute Kidney Injury. *Clin J Am Soc Nephrol*. 2021;16(10):1601-1609. doi:10.2215/CJN.19601220

14. *Manic-Depressive Illness: Bipolar Disorders and Recurrent Depression*, Second edition. By Frederick K. Goodwin and Kay Redfield Jamison. New York, Oxford University Press, 2007. 978-0-19-513579-4

Tables

Table 1. Demographic information for the analysis cohort, which corresponds to a random 20 percent sample of Medicare beneficiaries assigned to an Accountable Care Organization in 2018.

Total individuals			1,798,319
Age group and sex (% of total individuals)	Age 65-74	Female	26.9%
	Age 65-74	Male	21.1%
	Age 75-84	Female	20.8%
	Age 75-84	Male	15.6%
	Age 85 and above	Female	10.2%
	Age 85 and above	Male	5.4%
Geographic region in the U.S. (% of total individuals)	South		42.2%
	Midwest		24.9 %
	Northeast		19.8%
	West		13.0%
With Medicaid eligibility (% of total individuals)			9.2%
With Disability (%			9.0%

of total individuals)		
With any inpatient stays (% of total individuals)		17.0%
With at least one HCC (% of total individuals)		73.7%
Average number of HCCs (% of total individuals)		2.1
Spending ($)	Average total spending per beneficiary	$7911.10
	25th percentile spending per beneficiary	$478.40
	50th percentile spending per beneficiary	$1415.90

Table 2. Current prevalence, incidence, persistence, and steady state prevalence for HCC58:

Major Depressive, Bipolar, and Paranoid Disorders

Current prevalence (2017)	Current prevalence (2018)	Incidence	Persistence	Steady-State Prevalence
7.1%	8.8%	4.1%	70.1%	12.0%

Figures

Figure 1. Incidence and Persistence for the 10 HCCs with highest risk score contributions in version 22 of CMS-HCC for ACOs in 2018. HCCs are ordered by ACO risk score contribution, and the percentage following each HCC label corresponds to that HCC's contribution to the ACO risk score.

Figure 2. Current vs. Steady-State Prevalence for the 10 HCCs with highest risk score contribution in the ACOs for version 22 of CMS-HCC. HCCs are ordered by ACO risk score contribution, and the percentage following each HCC label corresponds to that HCC's contribution to the ACO risk score.

Figure 3. Current coding prevalence can continue to increase over time even if incidence and persistence do not change. Beginning at 2018 current prevalence values for coding in ACOs (shown on the y-axis at number of years = 0), each point within a given HCC represents the hypothetical current prevalence value if incidence and persistence for that HCC were to remain fixed as they were in 2017-2018. The percentage in parentheses following each HCC label corresponds to that HCC's contribution to the ACO risk score. Even absent behavioral change (e.g. changes in incidence or prevalence), current prevalence can continue to increase for several years. The number of years can vary considerably; for instance, the HCC corresponding to acute kidney injury (labeled and formerly known as acute renal failure) has the same current prevalence after year 2, while the HCC corresponding to specified heart arrhythmias increases for all 10 years shown.