Sibling Psychiatric Disorders as Risk Markers for Postpartum Psychosis

Authors: Shelby Smout[1,2], Floriana Milazzo[2], Thalia K. Robakis (MD, PhD)[2],
Veerle Bergink (MD, PhD)[2,3*], Behrang Mahjani (PhD)[1-2,4-8*]

*Shared authors

Affiliations:

[1]Seaver Autism Center for Research and Treatment, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[2]Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[3]Department of Psychiatry, Erasmus Medical Center, Rotterdam, The Netherlands.

[4]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[5]Department of Artificial Intelligence and Human Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[6]The Mindich Child Health and Development Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[7]Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm.

[8]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

Location of work and address for reprints:

Behrang Mahjani, PhD, behrang.mahjani@mssm.edu, One Gustave L. Levy Place, Box 1230, New York, NY 10029

Abstract:

Background: Postpartum psychosis affects 1–2 per 1,000 women following childbirth and constitutes a psychiatric emergency with significant risks for maternal and infant outcomes. While a personal history of bipolar disorder is the most established risk factor, nearly half of affected women have no prior psychiatric diagnosis, underscoring the need for improved risk stratification strategies.

Methods: In this nationwide Swedish register-based cohort study, we analyzed 676,293 women with at least one sister who delivered their first liveborn child between 1980 and 2017. A secondary analysis included 100,652 women with at least one brother. Postpartum psychosis was defined as psychiatric hospitalization for mania, psychosis, or depression with psychotic features within 90 days postpartum. Associations between sibling psychiatric diagnoses and postpartum psychosis were estimated using logistic regression, adjusting for maternal age and birth year. Hierarchical clustering was applied to identify familial subtypes based on sibling psychiatric profiles.

Results: Among 885 women (3.7 per 1,000) diagnosed with postpartum psychosis, we observed strong associations with sister diagnoses of schizoaffective disorder (OR, 14.50; 95% CI, 4.44–34.30) and postpartum psychosis (OR, 10.69; 95% CI, 6.60–16.26). Significant associations were also found for sister diagnoses of schizophrenia (OR, 7.10; 95% CI, 1.76–18.60) and bipolar disorder (OR, 6.26; 95% CI, 4.00–9.29). In brother-sister pairs, the strongest association was with bipolar disorder (OR, 6.23; 95% CI, 2.21-13.67). Cluster analysis revealed four familial subtypes, with 6.3% of cases showing distinct patterns of familial psychiatric burden: predominantly internalizing (anxiety/bipolar/personality disorders), predominantly externalizing (substance use disorders), and a broad affective-psychotic pattern with multiple comorbidities.

Conclusions: Familial psychiatric disorders, particularly those with psychotic or affective features, are strong risk markers for postpartum psychosis. These findings support the conceptualization of postpartum psychosis as a disorder with heterogeneous familial liability and highlight the potential utility of incorporating sibling psychiatric history into perinatal mental health screening, especially for women without prior psychiatric diagnoses.

1. Introduction

Postpartum psychosis is one of the most acute psychiatric emergencies associated with childbirth, affecting approximately 1–2 per 1,000 women (Meltzer-Brody et al. 2018; Munk-Olsen et al. 2016; VanderKruik et al. 2017). Onset typically occurs within the first two to four weeks postpartum and is characterized by rapid-onset mania, depression with psychotic features, or mixed affective states (Munk-Olsen et al. 2016; Bergink, Rasgon, and Wisner 2016). Despite its rarity, postpartum psychosis carries substantial risks for severe outcomes such as suicide, infanticide, and long-term psychiatric morbidity, making early detection and intervention critical. While a personal history of bipolar disorder remains the strongest known predictor of postpartum psychosis, with postpartum periods triggering bipolar relapses in approximately 20% to 50% of women with pre-existing bipolar disorder (Gilden et al. 2021), nearly half of all women affected with postpartum psychosis have no prior psychiatric diagnosis (Bergink, Rasgon, and Wisner 2016; Albers et al. 2023), underscoring the urgent need for improved risk stratification tools.

Recent large-scale epidemiological studies have revealed that postpartum psychosis exhibits one of the strongest familial aggregation patterns among psychiatric disorders (Kępińska, Robakis, and Humphreys 2025). In a nationwide Swedish cohort of 1.65 million women, having a full sister – meaning they have the same biological parents – with postpartum psychosis conferred a 10.7-fold increased risk (95% CI, 6.6–16.3) of developing the condition (Kępińska, Robakis, and Humphreys 2025). This relative recurrence risk surpasses that reported for many chronic psychiatric conditions, including bipolar disorder and schizophrenia (Lichtenstein et al. 2009). Notably, risk of bipolar and schizophrenia appears to aggregate similarly across male and female relatives, suggesting a sex-independent familial liability (Florio, Jones, and Forty 2021). Yet, to what extent this familial risk reflects transmission of postpartum psychosis specifically rather than severe psychiatric disorders broadly, remains unknown.

The clinical and diagnostic trajectory of postpartum psychosis further complicates our understanding of its etiology. Longitudinal studies suggest that postpartum psychosis frequently represents an initial manifestation of bipolar disorder: 72% to 88% of women with postpartum psychosis are subsequently diagnosed with bipolar or schizoaffective disorder, compared to only 12% diagnosed with schizophrenia (Bergink, Rasgon, and Wisner 2016; I. Jones and Craddock 2001). This has led to speculation that postpartum psychosis lies on the bipolar spectrum. However, recent genetic studies have begun to clarify these diagnostic relationships, suggesting that postpartum psychosis may have both overlapping and distinct genetic underpinnings with other psychiatric disorders (Jung et al. 2024; Di Florio et al. 2021).

Indeed, emerging genetic evidence supports this interpretation. Polygenic risk score (PRS) analyses indicate that women with postpartum psychosis share risk alleles with those who have bipolar disorder, but not major depression, suggesting a partially distinct genetic architecture (Di Florio et al. 2021). We expanded on these findings by combining family-based epidemiology and whole genome sequencing (WGS), and demonstrated that postpartum psychosis is highly heritable, with estimates of 55% from familial data and 37% from genomic data, including evidence for disproportionate heritability from the X chromosome (Jung et al. 2024). Rare variant analysis further identified enrichment of ultra-rare deleterious variants in constrained genes, with convergent associations across psychiatric and autoimmune disorders (Jung et al. 2024). These findings point to a combination of polygenic, rare variant, and possibly immune-metabolic pathways contributing to postpartum psychosis risk.

Despite this growing body of evidence, no study to date has systematically assessed whether familial risk of postpartum psychosis is conferred by other psychiatric diagnoses in relatives beyond postpartum psychosis itself. Prior registry-based research from Denmark suggests that a family history of bipolar disorder increases the risk of any postpartum psychiatric illness (HR = 2.86; 95% CI, 1.88–4.35), while family histories of schizophrenia and major depression confer more modest risks (Florio, Jones, and Forty 2021). However, these studies have not isolated postpartum psychosis specifically, nor have they focused on sibling diagnoses, a clinically accessible source of familial psychiatric information. Shared genetic liability, pleiotropic effects, and common environmental exposures are likely contributors to cross-disorder familial transmission, but the relative importance of these mechanisms in postpartum psychosis remains unclear.

In this population-based cohort study, we use comprehensive Swedish national register data to evaluate whether having a full sibling with any psychiatric disorder increases a woman's risk of developing postpartum psychosis. We assess cross-disorder familial risk based on sibling diagnoses of bipolar disorder, schizophrenia, schizoaffective disorder, major depressive disorder, anxiety disorders, obsessive-compulsive disorder, post-traumatic stress disorder, personality disorders, substance use disorders, sleep disorders, and dissociative or conversion disorders. By systematically quantifying these associations, we aim to determine whether familial transmission of postpartum psychosis reflects shared risk with other psychiatric conditions, and whether this information can inform perinatal risk screening in women without personal psychiatric history.

2. Methods

2.1 Study Population

This nationwide cohort study used data from multiple Swedish population registers. Sweden assigns a unique personal identification number to all residents, enabling linkage across multiple national registers (Ludvigsson et al. 2016).

From the Swedish Medical Birth Register, we identified all women who delivered their first liveborn child between January 1, 1980, and October 31, 2017. This register includes more than 98% of births in Sweden (Cnattingius et al. 2023). We assessed postpartum psychiatric outcomes through December 31, 2017, using linked register data.

Full siblings—defined as individuals sharing both biological parents—were identified using the Multi-Generation Register, which documents parent-child relationships with high completeness (97% for mothers and 95% for fathers) (Ekbom 2011). Because the Medical Birth Register includes only individuals who have had a child born in Sweden, our sibling dataset includes only sisters who gave birth (by design) and brothers who fathered at least one child. As a result, psychiatric diagnoses in siblings were only observable for those who had become parents and were captured in national registers.

The Swedish Ethical Review Authority approved the study and waived the requirement for informed consent owing to the use of de-identified registry data.

2.2 Exposure, Outcome, and Covariates

Diagnostic data were obtained from the Swedish National Patient Register (NPR), which includes all inpatient psychiatric care (nationwide since 1987) and outpatient visits (available from 2001 onward). Diagnoses are made by clinicians and registered using the International Classification of Diseases (ICD): ICD-8 (1969–1986), ICD-9 (1987–1996), and ICD-10 (1997–2017).

The primary outcome, postpartum psychosis, was defined as any psychiatric episode occurring within 90 days following a woman's first liveborn delivery. Postpartum psychosis lacks official diagnostic criteria in major classification systems like the Diagnostic and Statistical Manual of Mental Disorders (DSM) and is often considered an umbrella term (Di Florio, Munk-Olsen, and Bergink 2016). Therefore, to identify postpartum psychosis cases, we applied a diagnostic classification widely used in the literature: patients with mania and/or psychosis with onset in the postpartum period (Meltzer-Brody et al. 2018).

Similar to our previous study (Kępińska, Robakis, and Humphreys 2025), our case definition included postpartum-onset mania, psychosis, or depression with psychotic features (Kępińska, Robakis, and Humphreys 2025), using the following diagnostic codes: ICD-10 (F23, F28, F29, F30, F31, F323, F333, F531); ICD-9 (296A, 296G, 298A, 298B, 298X); and ICD-8 (2961, 2963, 2968, 2969, 2981, 2944). To maximize case ascertainment, we included both primary and secondary diagnoses with symptom onset within 0-3 months following a woman's first liveborn childbirth. We selected this three-month period to ensure comprehensive coverage of the typical onset window, consistent with previous epidemiological studies (Harlow et al. 2007; Langan Martin et al. 2016; Munk-Olsen et al. 2016; Valdimarsdóttir et al. 2009; VanderKruik et al. 2017).

The NPR has been extensively validated for psychiatric diagnoses. Positive predictive values for major psychiatric disorders, including bipolar disorder and schizophrenia-spectrum illnesses, exceed 85% to 95%, with kappa statistics ranging from 0.74 to 0.87 for diagnostic agreement in validation studies (Sellgren et al. 2011; Dalman et al. 2002). While specialist outpatient coverage is limited in earlier years and milder conditions may be underreported, postpartum psychosis generally necessitates hospitalization, thereby minimizing outcome misclassification.

For exposure assessment, the primary exposure was having a full sibling with a psychiatric diagnosis recorded in the NPR. Excluding postpartum psychosis, all sibling diagnoses needed to have been recorded prior to their own respective pregnancies if any occurred. Instances of postpartum psychosis among siblings were identified using the same criteria already described. We examined diagnostic categories using codes from ICD-8, ICD-9, and ICD-10 (see Supplementary Table S1). Each psychiatric disorder category was analyzed separately.

Maternal age was included as a potential confounder based on its established association with psychiatric disorders. Year of childbirth was included to adjust for temporal changes in diagnostic practices and registration patterns in the Swedish healthcare system over the study period.

2.3 Statistical Analysis

Our statistical analysis followed two complementary approaches: logistic regression to estimate familial risk, and unsupervised clustering to explore potential subtypes of familial psychiatric liability among women with postpartum psychosis.

All full sibling pairs (sibling1, sibling2) were identified, including both same-sex and opposite-sex combinations. Although postpartum psychosis occurs exclusively in women, both male and female siblings were included to assess whether familial risk differed by sibling sex. Within each sibling pair, one sibling (sibling1) was defined as the index individual whose psychiatric diagnosis

history constituted the exposure. The outcome was postpartum psychosis in sibling2, conditional on sibling2 being female and having experienced childbirth.

We first conducted logistic regression analyses restricted to sister–sister pairs to estimate the association between specific sibling psychiatric diagnoses and risk of postpartum psychosis. We then extended these models to include brother–sister pairs to evaluate potential sex-specific familial risk. All models were adjusted for sibling1's year of childbirth at first delivery. Results are presented as odds ratios (ORs) with 95% confidence intervals (CIs).

To investigate heterogeneity in familial psychiatric liability among women diagnosed with postpartum psychosis, we performed unsupervised hierarchical clustering based on their siblings' psychiatric diagnoses. Binary indicators for 22 diagnoses (11 psychiatric conditions, separately for brothers and sisters) were included in the clustering input. Women were clustered using Ward's method with Euclidean distance, and the number of clusters (k = 4) was selected based on silhouette width (average = 0.87) and interpretability. Cluster characteristics were summarized using diagnosis burden and frequencies of key sibling diagnoses.

3. Results

3.1 Study Population and Descriptive Statistics

A total of 676,293 women with at least one sister were included in the analysis. Among these, 885 were diagnosed with postpartum psychosis within 90 days of their first liveborn delivery.

The median maternal age at first delivery was 29 years (IQR: 26–33), and the median year of childbirth was 1996 (IQR: 1986–2007). The overall prevalence of postpartum psychosis in the cohort was 3.7 per 1,000. The distribution of pre-pregnancy psychiatric diagnoses among women with postpartum psychosis is summarized in Table 1, with the most common comorbidities being bipolar disorder (48.4%), major depressive disorder (33.1%), and anxiety disorders (23.5%), followed by substance use disorders (13.7%) and personality disorders (13.2%).

Table 1: Distribution of Psychiatric Diagnoses before Childbirth Among Women with Postpartum Psychosis (N = 885)

Diagnosis	Count	Percent
Bipolar Disorder	428	48.4%
Major Depressive Disorder	293	33.1%
Anxiety Disorders	208	23.5%

Substance Use Disorders	121	13.7%
Personality Disorders	117	13.2%
Post-Traumatic Stress Disorder	102	11.5%
Obsessive-Compulsive Disorder	38	4.3%
Schizophrenia	36	4.1%
Sleep Disorders	29	3.3%
Schizoaffective Disorder	15	1.7%
Dissociative Disorders	6	0.7%

Note: Percentages sum to more than 100% due to comorbidity among diagnoses

A total of 100,652 women with at least one brother were included in the analysis. Among these, 203 were diagnosed with postpartum psychosis within 90 days of their first liveborn delivery.

3.2 Familial Recurrence Risk

In sister-sister pairs, the strongest associations were observed for sibling diagnoses of schizoaffective disorder (OR 14.50, 95% CI 4.44-34.30) and postpartum psychosis itself (OR 10.69, 95% CI 6.60-16.26) (Table 1, Figure 3). Substantial familial risk was also evident for schizophrenia (OR 7.10, 95% CI 1.76-18.60) and bipolar disorder (OR 6.26, 95% CI 4.00-9.29). Moderate but statistically significant associations were found for several other conditions, including personality disorders (OR 3.18, 95% CI 1.97-4.83), major depressive disorder (OR 2.86, 95% CI 2.24-3.61), and sleep disorders (OR 2.74, 95% CI 0.98-5.94).

Table 2: Risk of Postpartum Psychosis Associated with Psychiatric Disorders in Siblings (Sister-Brother Relationships)

Disorder	Odds Ratio	95% Confidence Interval	p-value
Schizoaffective disorder	14.50	[4.44–34.30]	<0.001
Postpartum psychosis (sibling)	10.69	[6.60–16.26]	<0.001
Schizophrenia	7.10	[1.76–18.60]	0.001
Bipolar disorder	6.26	[4.00–9.29]	<0.001
Personality disorders	3.18	[1.97–4.83]	<0.001
Major depressive disorder	2.86	[2.24–3.61]	<0.001
Sleep disorders	2.74	[0.98–5.94]	0.025
Anxiety disorders	2.54	[1.95–3.25]	<0.001
Obsessive-compulsive disorder	2.00	[0.86–3.90]	0.068

Dissociative disorders	1.96	[0.32–6.08]	0.345
Substance use disorders	1.88	[1.34–2.55]	<0.001
Post-traumatic stress disorder	1.58	[1.05–2.27]	0.020

In brother-sister pairs, the strongest association was observed for bipolar disorder (OR 6.23, 95% CI 2.21-13.67), followed by personality disorders (OR 2.88, 95% CI 0.71-7.58) and obsessive-compulsive disorder (OR 2.65, 95% CI 0.44-8.31), though the latter two did not reach statistical significance (Table 3, Figure 3). Significant associations were also found for anxiety disorders (OR 2.45, 95% CI 1.46-3.86), major depressive disorder (OR 2.31, 95% CI 1.31-3.77), and substance use disorders (OR 1.87, 95% CI 1.15-2.86). We did not have cases for schizophrenia, schizoaffective disorder, or dissociative disorders to conduct this analysis.

Figure 1. Cross–Sibling Odds Ratios for Postpartum

Table 3: Risk of Postpartum Psychosis Associated with Psychiatric Disorders in Siblings (Sister-Sister Relationships)

Disorder	Odds Ratio	95% Confidence Interval	p-value
Bipolar disorder	6.23	[2.21-13.67]	<0.001
Anxiety disorders	2.45	[1.46-3.86]	<0.001
Personality disorders	2.88	[0.71-7.58]	0.070
Obsessive-compulsive disorder	2.65	[0.44-8.31]	0.172
Sleep disorders	2.23	[0.37-7.02]	0.259
Major depressive disorder	2.31	[1.31-3.77]	0.002
Substance use disorders	1.87	[1.15-2.86]	0.007
Post-traumatic stress disorder	1.72	[0.42-4.51]	0.354

We do not have enough cases for Schizophrenia, Schizoaffective disorder, and Dissociative disorders

3.3 Familial Subtypes

To explore heterogeneity in familial psychiatric liability among women with postpartum psychosis, we applied hierarchical clustering to binary indicators of sibling psychiatric diagnoses. In other words, these clusters describe the distribution of psychiatric disorders among the family members of women with postpartum psychosis, thereby creating distinct subtypes of familial psychiatric liability. Four distinct clusters emerged, supported by a high average silhouette width (0.87), indicating good cluster separation. The largest group (Cluster 1; 93.7%) exhibited minimal familial burden, with virtually no sibling psychiatric diagnoses recorded. Cluster 2 (4.2%) was characterized by anxiety disorders in sisters (45.7%), along with moderate rates of bipolar disorder (17.1%) and personality disorders (9.5%), suggesting a familial internalizing subtype. Cluster 3 (1.4%) showed high prevalence of substance use disorders in brothers (57.1%) with minimal other diagnoses, representing a narrow externalizing subtype. The smallest cluster (Cluster 4; 0.7%) exhibited the highest familial burden, with an average of 4.67 sibling diagnoses per woman. This group included high rates of anxiety (77.8%), personality disorders (50.0%), and bipolar disorder (27.8%) in sisters, indicating a broad affective-psychotic familial subtype. These findings support the existence of distinct familial subtypes of postpartum psychosis and reinforce the hypothesis that the disorder arises from diverse genetic and familial pathways, consistent with its polygenic architecture.

4. Discussion

This nationwide cohort study provides evidence that postpartum psychosis is strongly associated with a familial history of psychiatric disorders, particularly those with psychotic and affective components. We observed that the presence of a full sibling with postpartum psychosis conferred an exceptionally high relative risk (OR = 10.69, 95% CI 6.60–16.26), with a comparable risk from sisters with schizoaffective disorder (OR = 14.50, 95% CI 4.44–34.30). These findings are consistent with prior evidence demonstrating high familial aggregation of postpartum psychosis (Kępińska, Robakis, and Humphreys 2025). Notably, we extend these findings by quantifying cross-disorder familial risk using systematically coded sibling psychiatric diagnoses, revealing that sibling histories of schizophrenia (OR = 7.10, 95% CI 1.76–18.60) and bipolar disorder (OR = 6.26, 95% CI 4.00–9.29) are also significantly associated with postpartum psychosis. These results support the hypothesis that postpartum psychosis shares etiological features with broader psychotic and mood disorders.

To our knowledge, this is the first population-based study to systematically assess cross-disorder familial risk for postpartum psychosis based on sibling psychiatric diagnoses. Prior work using Swedish registers reported strong familial recurrence of postpartum psychosis but did not examine the contribution of other psychiatric disorders in siblings (Kępińska, Robakis, and Humphreys 2025). The present findings extend the literature by demonstrating that familial risk is not limited to recurrence of postpartum psychosis, but includes liability shared with bipolar and schizophrenia-spectrum disorders. These results are congruent with longitudinal studies showing that the majority of affected women are eventually diagnosed with bipolar disorder or schizoaffective disorder (Ian Jones and Craddock 2002; Bergink, Rasgon, and Wisner 2016), further suggesting overlapping neurobiological and genetic architectures.

Of particular note is the observed familial association with schizoaffective disorder in sister-sister analyses, which yielded an odds ratio exceeding that of postpartum psychosis itself. This observation reinforces the conceptualization of postpartum psychosis as a pleiotropic manifestation of affective-psychotic liability, potentially modulated by hormonal and immunological shifts in the postpartum period (Bergink, Rasgon, and Wisner 2016; Meltzer-Brody et al. 2018). This interpretation is consistent with polygenic risk score analyses demonstrating elevated bipolar, but not major depressive, genetic risk among women with postpartum psychosis (Di Florio et al. 2021) and with whole-genome sequencing studies identifying rare deleterious variants in constrained genes implicated in both psychiatric and autoimmune disorders (Jung et

al. 2024). The observed cross-disorder associations may therefore reflect both common polygenic and rare variant liability converging on a shared etiological axis for postpartum psychosis.

The identification of four distinct familial subtypes via hierarchical clustering underscores the heterogeneity in the familial transmission of postpartum psychosis. While most affected women (93.7%) had no recorded sibling psychiatric diagnoses, a meaningful subset exhibited clustered familial patterns characterized by anxiety disorders (Cluster 2, 4.2%), substance use disorders (Cluster 3, 1.4%), and a broad affective-psychotic pattern with multiple comorbidities (Cluster 4, 0.7%). These subtypes likely represent divergent etiological pathways and may reflect variable contributions of polygenic risk, rare variant burden, and environmental exposures. The existence of such subgroups may explain the inconsistent penetrance and clinical variability observed across postpartum psychosis cases and may inform future stratified risk models or targeted interventions.

Our findings carry important clinical implications. Existing perinatal mental health screening tools emphasize personal psychiatric history, particularly bipolar disorder. However, given that approximately 50% of women with postpartum psychosis have no prior psychiatric diagnosis (Bergink, Rasgon, and Wisner 2016), our results suggest that inclusion of sibling psychiatric history, especially of psychotic or mood disorders, could enhance early risk stratification. Implementation could involve integrating targeted questions about sibling psychiatric history into prenatal screening protocols, with specific focus on bipolar disorder, schizophrenia, and schizoaffective disorder. These adjustments may improve clinical preparedness, especially given the acute onset and potential severity of postpartum psychosis.

This study has several strengths. The use of a population-based national cohort minimizes selection bias and enables detection of rare cross-disorder associations with sufficient statistical power. By leveraging Sweden's comprehensive and validated national registers, we were able to examine familial psychiatric transmission at a resolution not achievable in clinical samples.

However, several limitations merit consideration. Although Swedish registers provide high diagnostic validity, under-ascertainment of milder or primary care-managed conditions remains possible Despite adjustment for maternal age and birth year, residual confounding remains possible, including unmeasured shared environmental factors such as familial adversity or socioeconomic status. Nonetheless, the magnitude and specificity of observed risk, especially for schizoaffective and psychotic disorders, strongly support a role for shared genetic liability. Third, our sibling dataset only includes individuals who became parents and were thus captured in the Medical Birth Register. While all index women were mothers by design, brothers were only

included if they fathered at least one child born in Sweden. As a result, our analysis may underestimate familial psychiatric burden in male siblings, particularly if severe mental illness reduces the likelihood of becoming a parent. This selection bias could attenuate risk estimates in brother–sister comparisons and may underrepresent certain familial risk profiles.

In conclusion, this study provides compelling evidence that sibling psychiatric disorders, particularly those involving psychotic or affective features, serve as strong familial risk markers for postpartum psychosis. The exceptionally high risk associated with sibling schizoaffective disorder (OR = 14.50, 95% CI 4.44–34.30) and postpartum psychosis itself (OR = 10.69, 95% CI 6.60–16.26), along with significant associations for schizophrenia (OR = 7.10, 95% CI 1.76–18.60) and bipolar disorder (OR = 6.26, 95% CI 4.00–9.29), advances our understanding of the disorder's familial architecture. These findings reinforce the genetic heterogeneity of postpartum psychosis and highlight the potential clinical value of incorporating sibling psychiatric history into perinatal risk assessment.

Data Sharing

Data may be obtained from a third party and are not publicly available. Data cannot be shared publicly owing to restrictions by law. Data are available from the National Medical Registries in Sweden after approval by the Swedish Ethical Review Authority.

Acknowledgments:

This study was supported by a grant from the Beatrice and Samuel A. Seaver Foundation (BM, SS); the National Institute of Mental Health (NIMH), R21MH131933 (BM), R01 HD111117 (TKR), and R01MH122869 (VB).

Author Contributions:

- Study concept and design: VB, BM, SS
- Acquisition, analysis, or interpretation of data: VB, BM, FM, TMO, SS
- Drafting of the manuscript: All authors
- Critical revision of the manuscript for important intellectual content: All authors
- Statistical analysis: BM, FM, SS
- Obtained funding: VB, BM
- Study supervision: VB, BM

References

Albers, Saundra M., Timothy Wen, Catherine Monk, Teresa C. Logue, Mary E. D'Alton, Whitney A. Booker, and Alexander M. Friedman. 2023. "Postpartum Psychosis during Delivery Hospitalizations and Postpartum Readmissions, 2016-2019." *American Journal of Obstetrics & Gynecology MFM* 5 (5): 100905.

Bergink, Veerle, Natalie Rasgon, and Katherine L. Wisner. 2016. "Postpartum Psychosis: Madness, Mania, and Melancholia in Motherhood." *The American Journal of Psychiatry* 173 (12): 1179–88.

Cnattingius, Sven, Karin Källén, Anna Sandström, Henny Rydberg, Helena Månsson, Olof Stephansson, Thomas Frisell, and Jonas F. Ludvigsson. 2023. "The Swedish Medical Birth Register during Five Decades: Documentation of the Content and Quality of the Register." *European Journal of Epidemiology* 38 (1): 109–20.

Dalman, C., J. Broms, J. Cullberg, and P. Allebeck. 2002. "Young Cases of Schizophrenia Identified in a National Inpatient Register: Are the Diagnoses Valid?" *Soc Psychiatry Psychiatr Epidemiol* 37 (11): 527–31.

Di Florio, Arianna, Jessica Mei Kay Yang, Karen Crawford, Veerle Bergink, Ganna Leonenko, Antonio F. Pardiñas, Valentina Escott-Price, et al. 2021. "Post-Partum Psychosis and Its Association with Bipolar Disorder in the UK: A Case-Control Study Using Polygenic Risk Scores." *The Lancet. Psychiatry* 8 (12): 1045–52.

Di Florio, Arianna, Trine Munk-Olsen, and Veerle Bergink. 2016. "The Birth of a Psychiatric Orphan Disorder: Postpartum Psychosis." *The Lancet Psychiatry* 3 (6): 502.

Ekbom, Anders. 2011. "The Swedish Multi-Generation Register." *Methods in Molecular Biology (Clifton, N.J.)* 675: 215–20.

Florio, Di, A. Jones, and I. R. Forty. 2021. "Bipolar Disorder, Affective Psychosis, and Schizophrenia in Women with Postpartum Psychosis." *Lancet Psychiatry* 8 (12): 1045–52.

Gilden, Janneke, Eline M. P. Poels, Simon Lambrichts, Annabel Vreeker, Marco P. M. Boks, Roel A. Ophoff, René S. Kahn, Astrid M. Kamperman, and Veerle Bergink. 2021. "Bipolar Episodes after Reproductive Events in Women with Bipolar I Disorder, A Study of 919 Pregnancies." *Journal of Affective Disorders* 295 (December): 72–79.

Harlow, Bernard L., Allison F. Vitonis, Par Sparen, Sven Cnattingius, Hadine Joffe, and Christina M. Hultman. 2007. "Incidence of Hospitalization for Postpartum Psychotic and Bipolar Episodes in Women With and Without Prior Prepregnancy or Prenatal Psychiatric Hospitalizations." *Archives of General Psychiatry* 64 (1): 42.

Jones, I., and N. Craddock. 2001. "Familiality of the Puerperal Trigger in Bipolar Disorder: Results of a Family Study." *The American Journal of Psychiatry* 158 (6): 913–17.

Jones, Ian, and Nick Craddock. 2002. "Do Puerperal Psychotic Episodes Identify a More Familial Subtype of Bipolar Disorder? Results of a Family History Study." *Psychiatric Genetics* 12 (3): 177–80.

Jung, Seulgi, Madison Caballero, Adrianna Kępińska, Shelby Smout, Trine Munk-Olsen, Thalia K. Robakis, Veerle Bergink, and Behrang Mahjani. 2024. "Genetic Architecture of Postpartum Psychosis: From Common to Rare Genetic Variation." *MedRxiv*. https://doi.org/10.1101/2024.12.09.24318732.

Kępińska, A. P., T. K. Robakis, and K. Humphreys. 2025. "Familial Risk of Postpartum Psychosis. MedRxiv." *MedRXiv*. https://doi.org/10.1101/2023.07.20.23292910.

Langan Martin, Julie, Gary McLean, Roch Cantwell, and Daniel J. Smith. 2016. "Admission to Psychiatric Hospital in the Early and Late Postpartum Periods: Scottish National Linkage Study." *BMJ Open* 6 (1): e008758.

Lichtenstein, Paul, Benjamin H. Yip, Camilla Björk, Yudi Pawitan, Tyrone D. Cannon, Patrick F. Sullivan, and Christina M. Hultman. 2009. "Common Genetic Determinants of Schizophrenia and Bipolar Disorder in Swedish Families: A Population-Based Study." *Lancet* 373 (9659): 234–39.

Ludvigsson, Jonas F., Catarina Almqvist, Anna-Karin Edstedt Bonamy, Rickard Ljung, Karl Michaëlsson, Martin Neovius, Olof Stephansson, and Weimin Ye. 2016. "Registers of the Swedish Total Population and Their Use in Medical Research." *European Journal of Epidemiology* 31 (2): 125–36.

Meltzer-Brody, Samantha, Louise M. Howard, Veerle Bergink, Simone Vigod, Ian Jones, Trine Munk-Olsen, Simone Honikman, and Jeannette Milgrom. 2018. "Postpartum Psychiatric Disorders." *Nature Reviews. Disease Primers* 4 (1): 18023.

Munk-Olsen, T., M. L. Maegbaek, B. M. Johannsen, X. Liu, L. M. Howard, A. di Florio, V. Bergink, and S. Meltzer-Brody. 2016. "Perinatal Psychiatric Episodes: A Population-Based Study on Treatment Incidence and Prevalence." *Translational Psychiatry* 6 (10): e919.

Sellgren, C., M. Landén, P. Lichtenstein, C. M. Hultman, and N. Långström. 2011. "Validity of Bipolar Disorder Hospital Discharge Diagnoses: File Review and Multiple Register Linkage in Sweden." *Acta Psychiatrica Scandinavica* 124 (6): 447–53.

Valdimarsdóttir, Unnur, Christina M. Hultman, Bernard Harlow, Sven Cnattingius, and Pär Sparén. 2009. "Psychotic Illness in First-Time Mothers with No Previous Psychiatric Hospitalizations: A Population-Based Study." Edited by Phillipa Hay. *PLoS Medicine* 6 (2): e1000013.

VanderKruik, Rachel, Maria Barreix, Doris Chou, Tomas Allen, Lale Say, Lee S. Cohen, and Maternal Morbidity Working Group. 2017. "The Global Prevalence of Postpartum Psychosis: A Systematic Review." *BMC Psychiatry* 17 (1): 272.