Biosecurity uptake and perceived risk of avian influenza among people in contact with birds in the UK

Harry Whitlow[1]*, Suzanne Gokool[1,2], Genevieve Clapp[1], Irene Bueno[3], Mariam Logunleko[3], Peter Moore[3], Sarah Masterton[4], Jo Taylor-Egbeyemi[4], Ian Brown[5], Riinu Pae[4], Louise E Smith[4], Ellen Brooks-Pollock[1,6], Amy C Thomas[1]*

1: Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

2: Gastrointestinal Infections, Food Safety and One Health Team (GIFSOH), UK Health Security Agency, London, UK

3: Bristol Veterinary School, University of Bristol, Langford, UK

4: Behavioural Science and Insights Unit, UK Health Security Agency, London, UK

5: The Pirbright Institute, Pirbright, England, UK

6: UK NIHR Health Protection Research Unit in Behavioural Science and Evaluation

* Corresponding authors: ja23401@bristol.ac.uk and amyc.thomas@bristol.ac.uk

Abstract

Recent intercontinental spread of highly pathogenic avian influenza (HPAI) A(H5N1) among kept and wild birds and transmission to mammalian hosts, including cattle and humans has heightened the need to review public health risk assessments. Biosecurity is essential for limiting disease spread, but how widely practices are implemented is not fully known. Here, we report on the uptake of biosecurity measures using questionnaire data collected in 2024 from people in contact with birds in the UK (Avian Contact Study). Almost half of the sample were poultry farmers (104/225). We found hand washing after contact with birds was the most common biosecurity measure implemented (89%, 196/218), followed by using disinfecting footwear dips (78%, 170/218). Individuals in contact with a higher number of birds were more likely to use at least one PPE measure for the face or body ($\chi 2$ (1, n=217) = 32.452, p<0.001) or at least one footwear-related PPE measure (Df=1, n=217, p<0.001), indicating the increasing number of birds as a likely proxy for farming practice, given the skewed poultry farmer sample. The perceived risk of avian influenza to the health of birds was generally high for individuals in contact with large flocks (≥1001 birds) and was associated with uptake of at least one footwear-related PPE measure ($\chi 2$ (1, n=185)= 9.171, p=0.002). The risk to human health was not associated with any biosecurity measure. These results suggest that routinely used biosecurity measures are implemented to limit disease spread among birds, but not with the primary view to limit zoonotic transmission from birds to humans. Future work should investigate attitudes towards avian influenza and biosecurity in larger sample sizes across varying populations to guide zoonotic influenza policy and inform targeted interventions.

Introduction

Between 2021 and 2022, 48 million birds were culled across Europe following outbreaks of highly pathogenic avian influenza (HPAI) A(H5N1) virus, belonging to clade 2.3.4.4b, in commercial poultry establishments and wild bird populations [1]. Rapid and widespread A(H5N1) transmission in Europe [1], Africa [2], North America [3], South America and the Antarctic [4, 5], constituted a panzootic in wild birds [3]. In the UK, a total of 9.38 million birds were estimated to have been culled in the six months to December 2022 [6]. To minimise avian transmission, in October 2022 the English Department for Environment, Food and Rural Affairs (Defra) and the Animal and Plant Health Agency (APHA) introduced a mandatory avian influenza prevention zone (AIPZ) and on-farm biosecurity measures across England [7]. Transmission among domestic poultry (kept birds) subsided during 2024 [8], but resurgence as of January

2025 has led to the declaration of another AIPZ in England, demonstrating the virus' persistent

threat to birds and mammalian wildlife [9].

Risk of avian influenza virus (AIV) infection for the general population in the UK and Europe

remains low [10], but concern and risk assessments are heightened given A(H5N1) infection of

wild and marine mammals [4], US dairy cattle [11], and higher mortality rates among wild birds

[12]. In humans, infection with A(H5N1) is rare following exposure to infected animals and no

evidence of human-to-human transmission has been recorded [13]. In the UK, asymptomatic

surveillance detected four human A(H5N1) cases in 2023, although two were suspected to be

due to contamination [10]. One further case was detected in January 2025 [14]. In the US,

A(H5N1) has infected 1,021 dairy herds across 17 states with known spillover to 70 people [11];

human infection has been asymptomatic or mild, with one fatality in an individual ≥65 years

with underlying health conditions (last reported 21 April 2025) [15]. Despite the current low

public health risk, genetic evolution and/or co-infection with human seasonal influenza virus

increases the risk of virus adaption to humans [16]. Limiting AIV transmission among birds and

mammalian wildlife is essential for the prevention and control of panzootics and pandemics.

Biosecurity measures (BMs) for individuals in contact with birds are recommended to prevent

AIV transmission in the UK [17] and in developing countries [18, 19]. A recent qualitative study

in the UK found high uptake of footwear dips among commercial poultry producers [20],

contrasting with backyard flock keepers in the UK who largely report non-compliance with Defra

enforced biosecurity measures, including footwear dips [21]. This demonstrates diverse uptake

trends among different demographics in contact with birds, requiring assessment of all possible

groups to inform a more nuanced understanding of biosecurity. Currently, BM uptake is largely

focused on the prevention of on-farm avian transmission [22, 23], resulting in a limited

understanding of BM uptake to prevent avian-to-human transmission.

BM uptake among people in contact with birds can be influenced by the perceived risk of

zoonotic disease transmission and AIV in birds [24]. Protection motivation theory [25] suggests

that uptake of BMs is driven by perceived susceptibility and severity of AIV infection in birds,

alongside the effectiveness of actions intended to prevent transmission (response efficacy).

Studies in Nigeria and China have shown that biosecurity uptake to prevent avian-to-avian

transmission varies by education status, occupation duration and farming operation size [22,

26]. In Taiwan and Europe, poultry farmers have documented personal attitudes, knowledge and

concern for business viability to influence uptake of BMs [23, 27, 28]. Factors such as age,

gender and education, have also been shown to either influence or have no effect upon AIV

awareness and knowledge for both avian- and human-specific BMs in Europe, Vietnam and

China [23, 29, 30]. Moreover, existing studies on motivation are concentrated in South East Asia

[30, 31], limiting generalisability to the UK. This suggests that the uptake and source of

motivation for BMs is heterogenous and requires investigation regarding avian-to-human

transmission within a UK context.

Here we use questionnaire data from the Avian Contact Study [32] to investigate trends in BM

uptake and risk perception of AIV (to oneself and to one's birds) in the context of preventing

zoonotic influenza infection among individuals in contact with kept and wild birds in the UK.

Building upon previous research regarding farming operation size [30] and the influence of risk

perception upon BM uptake [24], we hypothesised that the number of birds a person had

contact with on a typical day ('bird contact') and the perceived risk of AIV may influence uptake

of different types of measures to prevent human infection with AIV.

Materials and methods

Brief description of the Avian Contact Study

The Avian Contact Study (www.bristol.ac.uk/avian-contact) [32] aimed to inform current

zoonotic influenza public health measures in light of increased transmission in birds. A cross-

sectional, self-reported questionnaire was co-developed with avian influenza public health

teams (UK Health Security Agency; UKHSA), veterinary government scientists (Animal and Plant

Health Agency; APHA) and poultry farmers. This questionnaire was designed to capture the

knowledge and practices of people who have contact with kept and/or wild birds in the UK.

Eligible participants were aged 18 years or over, a UK resident (England, Northern Ireland,

Scotland, Wales) and had contact with domestic and/or wild birds. Full details of the

questionnaire methodology can be found in the Data Note [32], and the questionnaire is

available as extended data [33].

Ethics and consent

105   Ethical approval for the study was obtained from the University of Bristol, Faculty Research

106   Ethics Committee, approval number 17048 on 16 January 2024.

107   Informed written consent (using e-consent hosted on Research Electronic Data CAPture tools,

108   REDCap [34, 35]) for the use of data collected via the questionnaire was obtained from

109   respondents.

Data collection

We used a convenience sampling method, where the questionnaire was initially delivered in-person at the British Pig and Poultry Fair (15 and 16 May 2024) - an agricultural conference aimed at pig and poultry keepers in the UK [36]. The questionnaire was also distributed online via individuals met at the conference, animal health protection teams from APHA and poultry networks on social media. All questions were optional. Questionnaire data were collected using the Research Electronic Data Capture (REDCap) platform, hosted at the University of Bristol [34, 35]. Data captured between 15 May 2024 to 31 July 2024 were included in these analyses.

The questionnaire captured demographic data including age, gender, occupation and health. We analysed the following questions relating to bird contact, biosecurity measures and risk perception:

- How many birds of any type were you in direct contact with on a typical day?
  - 1-10
  - 11-100
  - 101-1000
  - 1001-10,000
  - 10,001-100,000
  - More than 100,000
- If you are a bird owner, how many birds do you own?
  - 1-10
  - 11-100
  - 101-1000
  - 1001-10,000
  - 10,001-100,000
  - More than 100,000+
  - I am not a bird owner
- Do you use any biosecurity measures to limit the risk of catching avian influenza (bird flu)? Please select all that apply.
  - I use face masks
  - I use goggles
  - I use gloves
  - I change clothing when in contact with birds
  - I use outer garments when in contact with birds

o   I change boots

o   I use boot covers

o   I use a disinfecting footwear dip

o   I wash/clean my hands after touching birds

o   I wash/clean my hands before and after handling raw poultry meat

o   I make sure raw poultry meat is fully cooked before eating

o   I do not eat undercooked or raw poultry meat

o   I make sure eggs are thoroughly cooked

o   I do not eat raw eggs

o   I do not remember

o   None

o   Other

• How much risk do you think avian influenza (bird flu) currently poses to your physical

health?

o   Very high risk

o   High risk

o   Medium risk

o   Low risk

o   No risk at all

o   I do not know

• How much risk do you think avian influenza (bird flu) currently poses to the health of

your birds?

o   Very high risk

o   High risk

o   Medium risk

o   Low risk

o   No risk at all

o   I do not know

Data analysis

RStudio (R version 4.4.1) was used to clean and analyse the dataset. The underlying code can

be found on GitHub (https://github.com/amythomas/aviancontactstudy). Completeness of data

varied by question. Use of biosecurity measures were recorded using radio buttons that

respondents "checked" for yes. We assumed responses with all measures "unchecked"

indicated a skipped question and were therefore coded as missing.

Descriptive statistics, including frequency counts, proportions and medians were calculated for

questions relating to respondent characteristics, bird contact, biosecurity measures and risk

perception. Occupational titles were derived from the National Statistics Extended Standard

Occupational Classification (SOC) 2020 Framework [37]. Where an appropriate job title was not

listed, respondents could input their own job title into a free-text box which was later combined

into appropriate categories. Differences in uptake of BMs and risk perception ranks between

bird contact groups were tested using the two proportions z-test, and confidence intervals

(95%) for a single proportion were also calculated. Chi-squared tests of independence or

Fisher's exact tests (where cell counts were below five) were used to test for an association

between the uptake of BMs and the number of birds a person was in contact with ('bird

contact') alongside perceived risk of AIV (to oneself and birds). Missing responses for BM

uptake, bird contact and risk perception were removed from these analyses.

Due to low cell counts among categories of bird contact, BM uptake and risk perception of AIV,

responses were categorised according to the groupings shown in Table 1. Respondents who

selected "I do not know" for the questions relating to risk perception of AIV to physical health

and bird health were considered as missing and excluded from subsequent analyses. The

frequency distribution of BM uptake and risk perception variables before grouping are shown in

the supplementary information (Figures S1 and S2). Distribution of bird contact is shown in

Table 2.

Table 1. Groupings for biosecurity measures, risk perception and bird contact. *Aggregate groups were created for*
biosecurity measures, risk perception scores and bird contact quantities for inferential analyses.

Group	Biosecurity measure
Handwashing	I wash/clean my hands after touching birds
Footwear-related PPE	I change boots
	I use boot covers
	I use a disinfecting footwear dip
Food-safety measures	I wash/clean my hands before and after handling raw poultry meat
	I make sure raw poultry meat is fully cooked before eating
	I do not eat undercooked or raw poultry meat
	I make sure eggs are thoroughly cooked
	I do not eat raw eggs
PPE for the face or body	I use face masks
	I use goggles
	I use gloves
	I change clothing when in contact with birds

	I use outer garments when in contact with birds
None	None
Other	Other
I do not remember	I do not remember
Group	**Risk perception**
No/low/medium risk	No risk at all
	Low risk
	Medium Risk
High/very high risk	High risk
	Very high risk
I do not know	I do not know
Group	**Quantity of birds a person was in contact with on a 'typical day'**
Low	1 – 10
	11 – 100
	101 – 1000
High	1001 – 10,000
	10,001 – 100,000
	More than 100,000

Results

Respondent characteristics

A total of 225 respondents completed the questionnaire between 15 May 2024 and 31 July 2024. Demographic characteristics are shown in Table 2. In-person respondents accounted for approximately one third of the sample (28%, 63 of 225) compared to online completion of the questionnaire (72%, 162 of 225). The median age was 48 years, ranging from 19 to 89 years. The largest age category were respondents aged between 50 and 59 (27%, 60 of 225).

Respondents provided information on their occupation, bird contact and bird ownership. Almost half were poultry farmers (46%, 104 of 225). Fewer respondents were veterinarians (7%, 16 of 225), zookeepers (7%, 15 of 225), retired (4%, 10 of 225), farm managers (3%, 7 of 225) or another type of farmer (3%, 6 of 225). Other occupations which could not be grouped into these categories accounted for 67 of 225 (30%) respondents.

Over half of respondents reported high contact levels with birds (≥1001 birds/day; 57%; 129 of 225). Low bird contact (≤1000 birds/day) was reported by 39% of all respondents (88 of 225). Respondents were most likely to have daily contact with large flocks of 10,001–100,000 birds (35%; 79 of 225), or presumed hobbyist/back-yard keepers in contact with 1–10 birds (16%;35 of 225) or 11–100 birds (19%; 42 of 225). Most respondents did not own birds (49%; 110 of 225). Of

215   those who did, ownership of birds was evenly distributed across the categories: 11% (24 of 225)

216   owned 1–10 birds, 11% (24 of 225) owned 11–100 birds and 12% (28 of 225) owned 10,001–

217   100,000 birds.

218   ***Table 2. Avian Contact Study respondent characteristics.*** *Note, proportions may not sum to 100% due to rounding.*

	Frequency (N=225)
Demographics	n (%) [min, max]
Questionnaire response type	
Online	162 (72%)
In-person	63 (28%)
Median age	48 [19 – 89]
Age categories	
19-29	21 (9%)
30-39	48 (21%)
40-49	46 (20%)
50-59	60 (27%)
60-64	22 (10%)
≥65	28 (12%)
Gender	
Male	142 (63%)
Female	81 (36%)
Other	<5 (<2%)
Occupation	
Poultry farmers	104 (46%)
Veterinarian	16 (7%)
Zookeeper	15 (7%)
Retired	10 (4%)
Farm manager	7 (3%)
Other type of farmer (mixed/livestock/arable)	6 (3%)
Other*	67 (30%)
Bird contact on a 'typical day'	
Low contact	**88 (39%)**
1–10 birds	35 (16%)
11–100 birds	42 (19%)
101–1000 birds	11 (5%)
High contact	**129 (57%)**
1001–10,000 birds	20 (9%)
10,001–100,000 birds	79 (35%)
>100,000 birds	30 (13%)
Missing	8 (4%)

Bird ownership	
1–10 birds	24 (11%)
11–100 birds	24 (11%)
101–1000 birds	<5 (<2%)
1001–10,000 birds	<5 (<2%)
10,001–100,000 birds	28 (12%)
>100,000 birds	11 (5%)
Non-bird owners	110 (49%)
Missing	22 (10%)

* Only groups with ≥5 observations are shown. All other groups are combined as 'Other'. Includes missing responses.

Biosecurity measures to limit avian-to-human transmission

Respondents were asked what biosecurity measures they used to prevent infection with AIV,

and 7 of 225 responses were excluded due to missing BM uptake data. In total, 98% (213 of 218)

reported using at least one BM, and 2% (5 of 218) reported using none (Figure 1). The single most

reported BM was washing/cleaning hands after touching birds, used by 89% (196 of 218) of

respondents, followed by using disinfecting footwear dips, reported by 78% (170 of 218). Nine

out of fourteen specific BMs were reportedly used by at least 50% of respondents, including

hand washing, using a disinfecting footwear dip, changing boots, washing hands after handling

raw poultry meat, ensuring raw poultry meat is fully cooked before eating, not eating

undercooked or raw poultry, ensuring eggs are fully cooked before eating, not eating raw eggs

and using face masks.

Figure 1. Proportion of participants reporting the use of biosecurity measures to prevent personal infection with avian influenza virus. *Colours represent biosecurity measures grouped into similar aggregate categories. Error bars represent 95% confidence intervals for a single proportion.*

We hypothesised that the number of birds an individual was in contact with would affect uptake of biosecurity measures. A total of 217 respondents provided information on both bird contact and uptake of BMs. There was substantial evidence that a larger proportion of high bird contact respondents (≥1001 birds/day) reported using at least one footwear related PPE measure (z=6.49, p<0.001) or at least one PPE measure for the face or body (z=5.88, p<0.001), compared to low bird contact respondents (Figure 2). There was limited evidence for a difference in the proportion between high and low bird contact respondents reporting the use of at least one food

242 safety measure (z=0.94, p=0.447) and washing their hands after touching birds (z=1.41,

243 p=0.238). Only low bird contact respondents reported using no measures (6%, 5 of 88) (Figure

244 2).

Figure 2. Biosecurity measure uptake by bird contact quantity. *Proportion of participants using at least one biosecurity measure as per the aggregate biosecurity categories by low and high bird contact group. Handwashing after touching birds was considered as its own category. Low contact: 1–1000 birds; high contact: ≥1001 birds. Error bars represent 95% confidence intervals for a single proportion.*

Uptake of BMs was higher among the high bird contact group than the low bird contact group

(Table 3). There was evidence for an association between bird contact levels and the use of at

least one PPE measure for the face or body (χ2 (1, n=217) = 32.452, p<0.001), and use of at

least one footwear related PPE measure (Df=1, n=217, p<0.001) (Table 3). There was no

evidence of association for use of at least one food safety measure (χ2 (1, n=217) = 0.888,

p=0.346) or hand washing after touching birds (χ2 (1, n=217) = 1.988, p=0.159). There was

moderate evidence for the use of no measures (Df=1, n=217, p=0.010). However, frequency

counts were small.

Table 3. Relationship between biosecurity measures and bird contact. *Chi-square test of independence between*
biosecurity measures and bird contact level (Df = Degrees of freedom). P-values are rounded to 3 decimal places.
Due to rounding, some percentages do not add to 100%.

Biosecurity measure		Low bird contact (n=88, %)	High bird contact (n=129, %)
Use of at least one food safety measure*	Reported	70 (80%)	109 (85%)
	Not reported	18 (20%)	20 (15%)
	Df[§], $\chi2$, p-value	1, 0.888, 0.346	
Use of at least one PPE measure for the face or body[†]	Reported	57 (65%)	123 (95%)
	Not reported	31 (35%)	6 (5%)
	Df, $\chi2$, p-value	1, 32.452, <0.001	
Use of at least one footwear-related PPE measure[‡]	Reported	59 (67%)	127 (98%)
	Not reported	29 (33%)	2 (2%)
	Df, $\chi2$, p-value	1, <0.001[‖]	
Handwashing after touching birds	Reported	76 (86%)	119 (92%)
	Not reported	12 (14%)	10 (8%)
	Df, $\chi2$, p-value	1, 1.988, 0.159	
Use of no measures	Reported	5 (6%)	0 (0%)
	Not reported	83 (94%)	129 (100%)
	Df, $\chi2$, p-value	1, 0.010[‖]	

[*] I wash/clean my hands before and after handling raw poultry meat; I make sure raw poultry meat is fully cooked before eating; I do not eat undercooked or raw poultry meat
[†] I use face masks; I use goggles; I use gloves; I use change clothing when in contact with birds; I use outer garments when in contact with birds
[‡] I change boots; I use boot covers; I use a disinfecting footwear dip
[§] Degrees of freedom
[‖] Fisher's Exact test

Risk perception of avian influenza to own health and birds' health

263 We collected respondents' perceptions of AIV risk to their own health and the health of their

264 birds. After removing missing data and those with unknown bird contact (34 of 225), 116

265 respondents reported high bird contact, and 75 respondents reported low bird contact (n=191)

266 (Figure 3). Most respondents perceived the risk of AIV to their physical health to be low or

267 medium (80%; 152 of 191). There was moderate evidence that a greater proportion of low bird

[13]

contact than high bird contact respondents perceived the risk to be low (z=2.38, p=0.017; n-Low=52/75, n-High=59/116). There was no evidence for a difference between bird contact groups in all other risk categories for this question. Risk to the health of respondents' birds was largely perceived to be medium, high or very high (147 of 191). There was substantial evidence for a greater proportion of low bird contact respondents perceiving the risk to be low compared to high bird contact respondents (z=3.74, p<0.001; n-Low=24/75, n-High=11/116), and for increased high bird contact respondents perceiving the risk to be very high compared to low bird contact respondents (z=3.53, p<0.001; n-Low=8/75, n-High=40/116).

Figure 3. Risk perception of avian influenza by bird contact. *Proportion self-reporting agreement with risk statements by bird contact group (low and high) for a) risk to human health and b) risk to the health of their birds. Low contact: 1 to 1000 birds; high contact: ≥1001 birds. Error bars represent 95% confidence intervals for a single proportion.*

To test the association between perceived risk and uptake of BMs, risk categories were grouped

into similar themes. Respondents who selected "I do not know" for physical health (n=5) and

bird health (n=6) were coded as missing and were excluded from analyses, resulting in 186 and

185 respondents in the personal health and bird health categories respectively.

Respondents who considered AIV risk as high to bird health were more likely to use at least one

footwear related PPE measure than those who perceived the risk as low ($\chi2$ (1, n=185)= 9.171,

p=0.002). For all other biosecurity groups, there was no evidence of an association with

perceived risk to bird health or perceived risk to physical health (Table 4).

Table 4: Relationship between biosecurity measure and perceived risk of avian influenza virus to bird and
personal health. *Chi-squared or Fisher's exact tests of association between the use of biosecurity measures and*
perceived risk of avian influenza virus to personal and bird health. Biosecurity measures were grouped into themes
where appropriate. Risk groups were aggregated into low (no, low and medium) and high (high and very high) risk. P-
values are rounded to 3 decimal places. Due to rounding, some percentages do not add to 100%.

Biosecurity measure		Risk to personal health (n=186)		Risk to bird health (n=185)	
		Low (n, %)	High (n, %)	Low (n, %)	High (n, %)
Use of at least one food safety measure*	Reported	144 (77%)	11 (6%)	74 (40%)	83 (45%)
	Not reported	27 (15%)	4 (2%)	10 (5%)	18 (10%)
	Df[§], $\chi2$, p-value	1, 0.283[‖]		1, 0.832, 0.362	
Use of at least one PPE measure for the face or body[†]	Reported	139 (75%)	15 (8%)	64 (35%)	89 (48%)
	Not reported	32 (17%)	0 (0%)	20 (11%)	12 (6%)
	Df, $\chi2$, p-value	1, 0.078[‖]		1, 3.766, 0.052	
Use of at least one footwear-related PPE measure[‡]	Reported	144 (77%)	15 (8%)	64 (35%)	94 (51%)
	Not reported	27 (15%)	0 (0%)	20 (11%)	7 (4%)
	Df, $\chi2$, p-value	1, 0.133[‖]		1, 9.171, 0.002[¶]	
Handwashing after touching birds	Reported	155 (83%)	14 (8%)	77 (42%)	90 (49%)
	Not reported	16 (9%)	1 (<1%)	7 (4%)	11 (6%)
	Df, $\chi2$, p-value	1, 1.000[‖]		1, 0.112, 0.737[¶]	
Use of no measures	Reported	3 (2%)	0 (0%)	3 (2%)	0 (0%)
	Not reported	168 (90%)	15 (8%)	81 (44%)	101 (55%)
	Df, $\chi2$, p-value	1, 1.000[‖]		1, 0.092[‖]	

[*] I wash/clean my hands before and after handling raw poultry meat; I make sure raw poultry meat is fully cooked before eating; I do not eat undercooked or raw poultry meat

[†] I use face masks; I use goggles; I use gloves; I use change clothing when in contact with birds; I use outer garments when in contact with birds

[‡] I change boots; I use boot covers; I use a disinfecting footwear dip

Discussion

This study used a cross-sectional self-reported questionnaire to explore biosecurity practices and risk perceptions of zoonotic avian influenza virus (AIV) among individuals in contact with kept and wild birds in the UK. Our analyses identified that handwashing was the most common biosecurity measure (BM) reported by respondents to prevent infection with AIV, followed by use of disinfecting footwear dips and changing boots. Five participants reported that they used no measures. Uptake of at least one footwear-related PPE measure or at least one PPE measure for the face or body was associated with bird contact; uptake was higher for respondents in contact with a high number of birds. The perceived risk of AIV to respondents own health was perceived as low, regardless of the number of birds a respondent had daily contact with. Risk of AIV to bird health was perceived as very high for the majority of respondents reporting 'high' daily bird contact (≥1001 birds/day), compared to a mixed perception of risk for those reporting 'low' daily bird contact (<1000 birds/day). Perceived risk of AIV to physical health had no association with the uptake of any biosecurity measures, but the perceived risk to the health of birds was associated with use of at least one footwear related PPE measure.

We found that hand washing after touching birds was the most frequently reported BM, contrasting with a European study on poultry farms where it was among the least reported measures for the prevention of avian transmission [23]. This could be explained by the framing of our questionnaire being around avian-to-human transmission risk, leading respondents to consider the personal activities they undertake to prevent the spread of infection in general, and the improved response efficacy of this specific action as opposed to avian-centred BMs [25]. Another explanation could be social desirability, as evidenced by bias in reporting trends during the COVID-19 pandemic [38]. When completing the questionnaire face-to-face with the research team, respondents may have been more inclined to report an action commonly expected of them. The impact of this bias is likely reduced as most respondents completed our questionnaire online. The high uptake of PPE, footwear dips and boots are consistent with a recent qualitative study of UK poultry farmers [20], strengthening our understanding of these being the most common among the poultry industry.

A study on French duck farms reported outdoor disinfection practices, such as disinfecting footwear dips, as one of the lowest scoring BMs for preventing AIV transmission [27]. Although

we cannot directly compare uptake, we found footwear dips are frequently used and has been commonly cited in other studies [20]. This could be a distinctive finding among the psycho-social cohort singled out for poor attitudes to biosecurity but may also relate to the farming operation sizes, species on the premises and the practices of the farmers recruited. We also found that the number of birds a person is in daily contact with was associated with biosecurity uptake to prevent zoonotic transmission. As a proxy for farming operation size, this finding is consistent with other studies where preventive measure uptake is more common among larger farm sizes and practices [22, 27]. However, this evidence is centred on the prevention of avian-to-avian transmission and is likely influenced by unmeasured covariates, such as farm assurance schemes. This association likely does not capture all mechanisms and requires a scoping assessment of other influences.

Perceived risk to human health was not associated with biosecurity uptake, and instead perceived risk to bird health was associated with uptake of at least one footwear related PPE measure. Using protection motivation theory [25], this may be explained by participants' relative perceived susceptibility to, and severity of AIV to human health as low or medium; a tendency which has been demonstrated in other demographics [30]. This is in line with other findings that perceived human susceptibility to AIV is low and that this may affect uptake of certain protective behaviours [39]. In addition, the lack of sustained human-to-human transmission is also likely to influence this perception [13], resulting in very little motivation to use BMs to protect their personal health.

Strengths and limitations

This study compliments existing knowledge regarding biosecurity usage to prevent avian infection among bird keepers in Europe and South East Asia [21, 30], and the changing transmission dynamics witnessed in North America and Europe in the last four years [3, 12, 13]. Our sample included those both in contact with domestic and wild birds, enabling a range of individuals to complete the questionnaire who have low or high contact with birds, such as hobbyist bird keepers. This goes beyond the existing efforts to understand biosecurity behaviour among members of the poultry industry alone. A limitation is the overrepresentation of commercial poultry farmers in the sample, as shown in the study Data Note, likely owing to recruitment at a national agricultural show [32]. This limits the generalisability of our findings to small flock-holders, like those studied in the UK Flockdown survey [21], and the wild bird sector. As a first data release of this pilot study, our sample is small and some covariates, such as farm assurance schemes and previous AIV outbreaks, were not measured. Future iterations of the

[17]

study will aim to record these covariates and reach a larger sample to perform these tests.

Questions recording BM uptake were not time-bound and were framed around the

interpretation of a 'typical day' that a respondent was in contact with birds. As a result,

consistent uptake could not be assessed. Finally, collapsing biosecurity and risk perception

groups to reduce small frequency counts may have led to less variation in our data - constituent

categories should be considered when interpreting findings.

Conclusions

Our findings suggest that current use of biosecurity measures by people who have contact with

birds in the UK are influenced by the perceived risk to bird health (high) and the number of birds

a person was in contact with (as a proxy for farming operation size), but not perceived risk to

human health (low). Further work should be undertaken to explore attitudes towards AIV and

biosecurity use regarding avian-to-human transmission among a larger sample, including

barriers and facilitators, to guide zoonotic influenza public health policy. Improving our

understanding of human behaviour in the context of zoonotic influenza will improve public

health responses through targeted interventions.

Data availability

Repository data.bris: The Avian Contact Study: questionnaire data 15 May – 31 July 2024. Data

are openly available at the University of Bristol Research Data Repository data.bris,

at https://doi.org/10.5523/bris.3nmqsrbv5ruom2abn0ql6e8yh2.

This project contains the following underlying data:

- • - Data file 1. (Raw underlying questionnaire data – csv file)

- • - Data file 2. (Raw underlying questionnaire data – .RDS file)

- • - Data file 2. (Associated data dictionary – csv file)

- • - Data file 3. (Code for importing underlying data in csv format into R for setting up
labelled data – .r file)

- • - Data file 4. (Blank consent form and participant information sheet – pdf file)

Data are available under the terms of National Archives' Non-Commercial Government Licence

for public sector information.

Software availability

Source code available from: https://github.com/amythomas/aviancontactstudy.

Conflicts of interest

LES, SG and RP are employees of the UK Health Security Agency. LES receives consultancy fees from the Sanofi group of companies and other life sciences companies. PM is an employee of the Animal Plant and Health Agency. The views expressed are those of the authors and not necessarily those of the UKHSA or the Department of Health and Social Care.

Funding statement

Funding for the Avian Contact Study was awarded by PolicyBristol from the Research England QR Policy Support Fund (QR PSF) 2022-24 for investigating 'Zoonotic spillover of avian influenza'. AT is funded by the Wellcome Trust, Early Career Award [227041/Z/23/Z].

Acknowledgements

We thank all participants who completed the questionnaire and those who contributed to the development of the study. EBP acknowledges supported from the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Evaluation of Interventions at the University of Bristol.

Supplementary materials

1. aiv_biosecurity_supplementary.docx - document containing Figure S1 (frequency counts of different biosecurity measures in each biosecurity group) and Figure S2 (frequency counts of risk perception categories for risk of avian influenza to respondents' health and the health of their birds)

References

1. European Centre for Disease Prevention and Control (ECDC). 2021-2022 data show largest avian flu epidemic in Europe ever: ECDC (European Union) [Internet]. 2022 [cited 2024 Nov 18]. Available from: https://www.ecdc.europa.eu/en/news-events/2021-2022-data-show-largest-avian-flu-epidemic-europe-ever.
2. Lo FT, Zecchin B, Diallo AA, Racky O, Tassoni L, Diop A, et al. Intercontinental Spread of Eurasian Highly Pathogenic Avian Influenza A(H5N1) to Senegal. Emerg Infect Dis. 2022;28(1):234-7. Doi: www.doi.org/10.3201/eid2801.211401.
3. Peacock T, Moncla L, Dudas G, VanInsberghe D, Sukhova K, Lloyd-Smith JO, et al. The global H5N1 influenza panzootic in mammals. Nature. 2024;637:304-313. Doi: https://doi.org/10.1038/s41586-024-08054-z
4. Uhart MM, Vanstreels RET, Nelson MI, Olivera V, Campagna J, Zavattieri V, et al. Epidemiological data of an influenza A/H5N1 outbreak in elephant seals in Argentina

419   indicates mammal-to-mammal transmission. Nat Commun. 2024;15(1):9516. Doi:
420   https://doi.org/10.1038/s41467-024-53766-5

421 5. Aguado B, Begeman L, Günther A, Iervolino M, Soto F, Vanstreels RET, et al. Searching
422   for high pathogenicity avian influenza virus in Antarctica. Nat Microbiol.
423   2024;9(12):3081-3. Doi: https://doi.org/10.1038/s41564-024-01868-7

424 6. World Organisation for Animal Health (WOAH). SMR_161191 - United Kingdom - Second
425   Semester 2022 – Terrestrial [Internet]. Paris: World Organisation for Animal Health
426   (WOAH); 2022 [cited 2024 Dec 19]. 30p. Available from: https://wahis.woah.org/#/smr-
427   review/161193?reportInfoId=98997&fromPage=smr-management-url

428 7. Department for Environment Food & Rural Affairs (DEFRA), Animal and Plant Health
429   Agency (APHA). Avian Influenza Prevention Zone (AIPZ): the whole of England excluding
430   Norfolk, Suffolk and parts of Essex (2022) [Internet]. 2022 [cited 2024 Jul 25]. [Available
431   from: https://www.gov.uk/animal-disease-cases-england/avian-influenza-prevention-
432   zone-aipz-the-whole-of-england-excluding-norfolk-suffolk-and-parts-of-essex-2022.

433 8. UK Health Security Agency. HAIRS risk statement: Avian influenza A(H5N1) in livestock
434   [Internet]. 2024 [cited 2024 Nov 25]. [Available from:
435   https://www.gov.uk/government/publications/hairs-risk-statement-avian-influenza-
436   ah5n1-in-livestock/hairs-risk-statement-avian-influenza-ah5n1-in-livestock.

437 9. Department for Environment Food & Rural Affairs (DEFRA), Animal and Plant Health
438   Agency (APHA). Avian Influenza Prevention Zone declared for whole of England
439   [Internet]. 2025 [cited 2025 Feb 01]. Available from:
440   https://www.gov.uk/government/news/avian-influenza-prevention-zone-declared-for-
441   whole-of-england

442 10. UK Health Security Agency. Investigation into the risk to human health of avian influenza
443   (influenza A H5N1) in England: technical briefing 5 [Internet]. 2023 [cited 2025
444   27/01/25]. Available from: https://www.gov.uk/government/publications/avian-
445   influenza-influenza-a-h5n1-technical-briefings/investigation-into-the-risk-to-human-
446   health-of-avian-influenza-influenza-a-h5n1-in-england-technical-briefing-5.

447 11. US Centre for Disease Control. H5 Bird Flu: Current Situation [Internet]. 2025 [cited
448   2025 Feb 16]. Available from: https://www.cdc.gov/bird-flu/situation-
449   summary/?CDC_AAref_Val=https://www.cdc.gov/flu/avianflu/avian-flu-summary.htm

450 12. European Food Safety Authority (EFSA), European Centre for Disease Prevention
451   (EcfDP), European Union Reference Laboratory for Avian Influenza (EURLfAI), Adlhoch C,
452   Fusaro A, Gonzales JL, Kuiken T, et al. Avian influenza overview March – April 2023. EFSA
453   J. 2023;21(6):e08039. Doi: https://doi.org/10.2903/j.efsa.2023.8039

454 13. Alexakis L, Buczkowski H, Ducatez M, Fusaro A, Gonzales JL, Kuiken T, et al. Avian
455   influenza overview June–September 2024. EFSA J. 2024;22(10). Doi: https://doi.org/
456   10.2903/j.efsa.2024.9057

457 14. UK Health Security Agency. Human case of avian flu detected in England [Internet].
458   2025 [cited 2025 Feb 01]. Available from: https://www.gov.uk/government/news/human-
459   case-of-avian-flu-detected-in-england.

460 15. US Centre for Disease Control. First H5 Bird Flu Death Reported in United States
461   [Internet]. 2025 [cited 2025 Jan 15]. Available from:
462   https://www.cdc.gov/media/releases/2025/m0106-h5-birdflu-death.html

16. Thomas A, Gokool S, Whitlow H, Clapp G, Moore P, Puleston R, et al. Seasonal Influenza Vaccination in People who Have Contact With Birds. Influenza Other Respir Viruses. 2025;19(4):e70101. Doi: https://doi.org/10.1111/irv.70101

17. Department for Environment Food & Rural Affairs (Defra), Animal and Plant Health Agency (APHA). Bird flu (avian influenza): how to prevent it and stop it spreading - Biosecurity and hygiene standards to keep birds safe from bird flu (avian influenza) [Internet]. 2022 [cited 2025 Jan 28]. Available from: https://www.gov.uk/guidance/bird-flu-avian-influenza-how-to-prevent-it-and-stop-it-spreading.

18. Gupta SD, Fournié G, Hoque MA, Henning J. Factors influencing chicken farmers' decisions to implement prevention and control measures to reduce avian influenza virus spread under endemic conditions. Transbound Emerg Dis. 2021;68(1):194-207. Doi: https://doi.org/10.1111/tbed.13757

19. Conan A, Goutard FL, Sorn S, Vong S. Biosecurity measures for backyard poultry in developing countries: a systematic review. BMC Vet Res. 2012;8(1):240. Doi: https://dx.doi.org/10.1186/1746-6148-8-240

20. Hosseini P, Syndicus I, Houghton E, Alarcon P, Fournie G, Guitian J, et al. Qualitative studies on farm managers perception of risk from HPAIV and implementation attitudes of biosecurity measures on UK poultry farms. Prev Vet Med. 2025;239:106445. Doi: https://doi.org/10.1016/j.prevetmed.2025.106445

21. McClaughlin E, Elliott S, Jewitt S, Smallman-Raynor M, Dunham S, Parnell T, et al. UK flockdown: A survey of smallscale poultry keepers and their understanding of governmental guidance on highly pathogenic avian influenza (HPAI). Prev Vet Med. 2024;224:106117. Doi: https://doi.org/10.1016/j.prevetmed.2024.106117.

22. Cui B, Liu ZP. Determinants of Knowledge and Biosecurity Preventive Behaviors for Highly Pathogenic Avian Influenza Risk Among Chinese Poultry Farmers. Avian Dis. 2016;60(2):480-6. Available from: https://www-jstor-org.bris.idm.oclc.org/stable/26431788

23. Souillard R, Allain V, Dufay-Lefort CA, Rousset N, Amalraj A, Spaans A, et al. Biosecurity implementation on large-scale poultry farms in Europe: A qualitative interview study with farmers. Prev Vet Med. 2024;224:106119. Doi: https://doi.org/10.1016/j.prevetmed.2024.106119

24. Dowd K, Taylor M, Toribio J-ALML, Hooker C, Dhand NK. Zoonotic disease risk perceptions and infection control practices of Australian veterinarians: Call for change in work culture. Prev Vet Med. 2013;111(1-2):17-24. Doi: https://dx.doi.org/10.1016/j.prevetmed.2013.04.002

25. Maddux JE, Rogers RW. Protection motivation and self-efficacy: A revised theory of fear appeals and attitude change. J Exp Soc Psychol. 1983;19(5):469-79. Doi: https://doi.org/10.1016/0022-1031(83)90023-9

26. Ameji ON, Abdu PA, Sa'idu L, Kabir J, Assam A. Awareness, Knowledge, Readiness to Report Outbreak and Biosecurity Practices Towards Highly Pathogenic Avian Influenza in Kogi State, Nigeria. Int J Poult Sci. 2012;11(1):11-5. Doi: http://dx.doi.org/10.3923/ijps.2012.11.15

27. Delpont M, Racicot M, Durivage A, Fornili L, Guerin JL, Vaillancourt JP, et al. Determinants of biosecurity practices in French duck farms after a H5N8 Highly Pathogenic Avian Influenza epidemic: The effect of farmer knowledge, attitudes and

personality traits. Transbound Emerg Dis. 2021;68(1):51-61. Doi:
https://doi.org/10.1111/tbed.13462

28.   Pao H-N, Jackson E, Yang T-S, Tsai J-S, Sung WHT, Pfeiffer DU. Determinants of farmers'
biosecurity mindset: A social-ecological model using systems thinking. Front Vet Sci.
2022;9. Doi: https://doi.org/10.3389/fvets.2022.959934

29.   Manabe T, Thuy PTP, Can VV, Takasaki J, Huyen DTT, Chau NTM, et al. Impact of
Educational Intervention Concerning Awareness and Behaviors Relating to Avian
Influenza (H5N1) in a High-Risk Population in Vietnam. PLoS ONE. 2011;6(8):e23711.
Doi: https://doi.org/10.1371/journal.pone.0023711

30.   Cui B, Wang F, Wang LD-L, Pan C, Ke J, Tian Y. A Comparative Analysis of Risk Perception
and Coping Behaviors among Chinese Poultry Farmers Regarding Human and Poultry
Infection with Avian Influenza. Int J Environ Res Public Health. 2019;16(20):3832. Doi:
https://doi.org/10.3390/ijerph16203832

31.   Pao H-N, Jackson E, Yang T-S, Tsai J-S, Hwang Y-T, Sung WHT, et al. The attitude-
behaviour gap in biosecurity: Applying social theories to understand the relationships
between commercial chicken farmers' attitudes and behaviours. Front Vet Sci. 2023;10.
Doi: https://doi.org/10.3389/fvets.2023.1070482

32.   Thomas A, Gokool S, Clapp G, Whitlow H, McGrath C, Moore P, et al. Data note on the
Avian Contact Study: a questionnaire resource for avian influenza public health planning
[version 2; peer review: 2 approved]. Wellcome Open Res. 2025;9(604). Doi:
https://doi.org/10.12688/wellcomeopenres.23064.2

33.   Thomas A, Gokool S, Clapp G, Whitlow H, Carmel M, Moore P et al. Avian Contact Study
Questionnaire and Data Dictionary [Internet]: Zenodo; 2024. Available from:
https://zenodo.org/records/13617061

34.   PA Harris, R Taylor, R Thielke, J Payne, N Gonzalez, JG. Conde. Research electronic data
capture (REDCap) – A metadata-driven methodology and workflow process for providing
translational research informatics support. J Biomed Inform. 2009;42(2):377-81.

35.   PA Harris, R Taylor, BL Minor, V Elliott, M Fernandez, L O'Neal, et al. The REDCap
consortium: Building an international community of software partners. J Biomed Inform.
2019. Doi: https://doi.org/ 10.1016/j.jbi.2019.103208

36.   Royal Agricultural Society of England. The British Pig and Poultry Fair [Internet]. 2024
[cited 2024 Jul 15]. Available from: https://pigandpoultry.org.uk/

37.   Office for National Statistics. Extended SOC 2020 [Internet]. 2023 [cited 2024 Aug 9].
Available from:
https://www.ons.gov.uk/methodology/classificationsandstandards/standardoccupation
alclassificationsoc/standardoccupationalclassificationsocextensionproject.

38.   Mieth L, Mayer MM, Hoffmann A, Buchner A, Bell R. Do they really wash their hands?
Prevalence estimates for personal hygiene behaviour during the COVID-19 pandemic
based on indirect questions. BMC Public Health. 2021;21(1). Doi:
https://dx.doi.org/10.1186/s12889-020-10109-5

39.   Pae R, Findlater L, Amlôt R, Capelastegui F, Dabrera G, Humphreys C, et al. "I definitely
cannot afford to be feeling poorly if there's no need to be": a qualitative evaluation of
antiviral uptake following suspected occupational exposure to avian influenza. BMC
Public Health. 2025;25(1). Doi: https://dx.doi.org/10.1186/s12889-025-21459-3

Supplementary information: Biosecurity uptake and perceived risk of avian influenza among people in contact with birds in the UK

Harry Whitlow[1]*, Suzanne Gokool[1,2], Genevieve Clapp[1], Irene Bueno[3], Mariam Logunleko[3], Peter Moore[3], Sarah Masterton[4], Jo Taylor-Egbeyemi[4], Ian Brown[5], Riinu Pae[4], Louise E Smith[4], Ellen Brooks-Pollock[1,6], Amy C Thomas[1]*

1: Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

2: Gastrointestinal Infections, Food Safety and One Health Team (GIFSOH), UK Health Security Agency, London, UK

3: Bristol Veterinary School, University of Bristol, Langford, UK

4: Behavioural Science and Insights Unit, UK Health Security Agency, London, UK

5: The Pirbright Institute, Pirbright, England, UK

6: UK NIHR Health Protection Research Unit in Behavioural Science and Evaluation

* Corresponding authors: ja23401@bristol.ac.uk and amyc.thomas@bristol.ac.uk

Figure S1. Frequency counts of different biosecurity measures in each biosecurity group. (a) Use of at least one PPE measure for the face or body, (b) use of at least one footwear related PPE measure and (c) use of at least one food safety measure. Use of biosecurity measures were recorded using radio buttons that respondents "checked" for yes. We assumed responses with all measures "unchecked" indicated a skipped question and were therefore coded as missing.

Figure S2. Frequency counts of risk perception categories for risk of avian influenza to (a)
respondents' health and (b) risk to the health of their birds.