# Biosecurity uptake and perceived risk of avian influenza among people in contact with birds

Harry Whitlow[1]*, Suzanne Gokool[1,2], Genevieve Clapp[1], Irene Bueno[3], Mariam Logunleko[3], Peter Moore[3], Sarah Masterton[4], Jo Taylor-Egbeyemi[4], Ian Brown[5], Riinu Pae[4], Louise E Smith[4], Ellen Brooks-Pollock[1,6], Amy C Thomas[1]**

1: Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

2: Gastrointestinal Infections, Food Safety and One Health Team (GIFSOH), UK Health Security Agency, London, UK

3: Bristol Veterinary School, University of Bristol, Langford, UK

4: Behavioural Science and Insights Unit, UK Health Security Agency, London, UK

5: The Pirbright Institute, Pirbright, England, UK

6: UK NIHR Health Protection Research Unit in Behavioural Science and Evaluation

*Corresponding author 1: ja23401@bristol.ac.uk

**Corresponding author 2: amyc.thomas@bristol.ac.uk

**Keywords:** Avian Influenza, Biosecurity, Public Health, Risk Perception, Spillover, Zoonoses

**Impacts:**

- Biosecurity uptake among individuals in contact with birds is likely influenced by the perceived risk to avian health, but not human health.
- Individuals working in contact with a greater number of birds are more likely to use biosecurity measures, reflective of farm operation size. Public health strategies may need to specifically target small flock holders to improve biosecurity.
- Attitudes towards avian influenza virus infections in humans need further characterisation to determine barriers for increasing uptake and effectiveness of avian influenza public health measures.

# Data availability

Repository data.bris: The Avian Contact Study: questionnaire data 15 May – 31 July 2024. Data are openly available at the University of Bristol Research Data Repository data.bris, at https://doi.org/10.5523/bris.3nmqsrbv5ruom2abn0ql6e8yh2.

[1]

This project contains the following underlying data:

- - Data file 1. (Raw underlying questionnaire data – csv file)

- - Data file 2. (Raw underlying questionnaire data – .RDS file)

- - Data file 2. (Associated data dictionary – csv file)

- - Data file 3. (Code for importing underlying data in csv format into R for setting up
labelled data – .r file)

- - Data file 4. (Blank consent form and participant information sheet – pdf file)

Data are available under the terms of National Archives' Non-Commercial Government Licence
for public sector information.

# Software availability

41    Source code available from: https://github.com/amythomas/aviancontactstudy.

# Conflicts of interest

43    LES, SG, SM, JT and RP are employees of the UK Health Security Agency. LES receives
44    consultancy fees from the Sanofi group of companies and other life sciences companies. PM is
45    an employee of the Animal Plant and Health Agency. The views expressed are those of the
46    authors and not necessarily those of the UKHSA or the Department of Health and Social Care.

# Funding statement

Funding for the Avian Contact Study was awarded by PolicyBristol from the Research England
QR Policy Support Fund (QR PSF) 2022-24 for investigating 'Zoonotic spillover of avian
influenza'. AT is funded by the Wellcome Trust, Early Career Award [227041/Z/23/Z]. EBP
acknowledges support from the National Institute for Health Research Health Protection
Research Unit (NIHR HPRU) in Evaluation and Behavioural Science at the University of Bristol
(NIHR207385).

# Abstract

**Introduction:** Recent intercontinental spread of highly pathogenic avian influenza (HPAI) A(H5N1) among kept and wild birds, and transmission to mammalian hosts, including cattle and humans, has heightened the need to review public health risk assessments. Biosecurity is essential for limiting disease spread, but how widely practices are implemented is not fully known.

**Methods:** Here, we report on uptake of biosecurity measures and risk perception of avian influenza virus (AIV) in the context of preventing zoonotic transmission to persons potentially at high risk of exposure. Questionnaire data from 225 people in contact with birds in the UK (Avian Contact Study, May to July 2024) was analysed.

**Results:** We found hand washing after contact with birds was the most common biosecurity measure implemented (89%, 196/218), followed by using disinfecting footwear dips (78%, 170/218). Individuals in contact with a higher number of birds were more likely to use at least one PPE measure for the face or body ($\chi 2$ (1, n=217) = 32.452, p<0.001) or at least one footwear-related PPE measure (Df=1, n=217, p<0.001). The perceived risk of AIV to the health of birds was high for individuals in contact with large flocks (≥1001 birds) and associated with uptake of at least one footwear-related PPE measure ($\chi 2$ (1, n=185)= 9.171, p=0.002). Perceived risk of AIV to respondents' own health was low, regardless of the number of birds a respondent had daily contact with.

**Conclusions:** Routinely used biosecurity measures are implemented to limit AIV spread among birds, but not with the purpose of limiting zoonotic transmission from birds to humans. Identifying cohort characteristics which could lead to low BM uptake, alongside barriers and facilitators to BM uptake is important for informing zoonotic AIV public health campaigns.

# Introduction

Between 2021 and 2022, 48 million birds were culled across Europe following outbreaks of highly pathogenic avian influenza (HPAI) A(H5N1) virus, belonging to clade 2.3.4.4b, in commercial poultry establishments and wild bird populations (1). Rapid and widespread A(H5N1) transmission in Europe (1), Africa (2), North America (3), South America and the Antarctic (4,5), constituted a panzootic in wild birds (3). In the UK, a total of 9.38 million birds were estimated to have been culled in the six months to December 2022 (6). To minimise avian transmission, in October 2022 the English Department for Environment, Food and Rural Affairs (Defra) and the Animal and Plant Health Agency (APHA) introduced a mandatory avian influenza prevention zone (AIPZ) and on-farm biosecurity measures across England (7). Transmission among domestic poultry (kept birds) subsided during 2024 (8), but resurgence as of January 2025 has led to the declaration of another AIPZ in England, demonstrating the virus' persistent threat to birds and mammalian wildlife (9).

Risk of avian influenza virus (AIV) infection for the general population in the UK and Europe remains low (10), but concern and risk assessments are heightened given A(H5N1) infection of wild and marine mammals (4), US dairy cattle (11), and higher mortality rates among wild birds (12). In humans, infection with A(H5N1) is rare following exposure to infected animals and no evidence of human-to-human transmission has been recorded (13). In the UK, asymptomatic surveillance detected four human A(H5N1) cases in 2023, although two were suspected to be due to contamination (10). One further case was detected in January 2025 (14). In the US, A(H5N1) has infected 1,077 dairy herds across 17 states with known spillover to 70 people (11); human infection has been asymptomatic or mild, with one fatality in an individual ≥65 years with underlying health conditions (last updated 1 July 2025) (15). Despite the current low public health risk, genetic evolution and/or co-infection with human seasonal influenza virus increases the risk of virus adaption to humans (16). Limiting AIV transmission among birds and mammalian wildlife is essential for the prevention and control of panzootics and pandemics.

Biosecurity measures (BMs) for individuals in contact with birds are recommended to prevent AIV transmission in the UK (17) and in developing countries (18,19).  A recent qualitative study in the UK found high uptake of footwear dips among commercial poultry producers (20), contrasting with backyard flock keepers in the UK who largely report non-compliance with Defra enforced biosecurity measures, including footwear dips (21). This demonstrates diverse uptake trends among different demographics in contact with birds, requiring assessment of all possible groups to inform a more nuanced understanding of biosecurity. Currently, BM uptake is largely

focused on the prevention of on-farm avian transmission (22,23), resulting in a limited

understanding of BM uptake to prevent avian-to-human transmission.

BM uptake among people in contact with birds can be influenced by the perceived risk of

zoonotic disease transmission and AIV in birds (24). Protection motivation theory (25) suggests

that uptake of BMs is driven by perceived susceptibility and severity of AIV infection in birds,

alongside the effectiveness of actions intended to prevent transmission (response efficacy).

Studies in Nigeria and China have shown that biosecurity uptake to prevent avian-to-avian

transmission varies by education status, occupation duration and farming operation size

(22,26). In Taiwan and Europe, poultry farmers have documented personal attitudes, knowledge

and concern for business viability to influence uptake of BMs (23,27,28). Factors such as age,

gender and education, have also been shown to either influence or have no effect upon AIV

awareness and knowledge for both avian- and human-specific BMs in Europe, Vietnam and

China (23,29,30). Moreover, existing studies on motivation are concentrated in South East Asia

(30,31), limiting generalisability to the UK. This suggests that the uptake and source of

motivation for BMs is heterogenous and requires investigation regarding avian-to-human

transmission within a UK context.

Using data from the Avian Contact Study, a cross-sectional questionnaire of individuals in

contact with kept and wild birds in the UK (32), we investigate trends in BM uptake and risk

perception of AIV in the context of preventing zoonotic influenza infection. Building upon

previous research regarding farming operation size (30) and the influence of risk perception

upon BM uptake (24), we hypothesised that the number of birds a person had contact with on a

typical day ('bird contact') and the perceived risk of AIV may influence uptake of different types

of measures to prevent human infection with AIV.

# 133 Materials and methods

## 134 Brief description of the Avian Contact Study

The Avian Contact Study ([www.bristol.ac.uk/avian-contact](www.bristol.ac.uk/avian-contact)) (32) aimed to inform current

zoonotic influenza public health measures given increased transmission in birds. A cross-

sectional, self-reported questionnaire was co-developed with avian influenza public health

teams (UK Health Security Agency; UKHSA), veterinary government scientists (Animal and Plant

Health Agency; APHA) and poultry farmers. This questionnaire was designed to capture the

knowledge and practices of people who have contact with kept and/or wild birds in the UK.

Eligible participants were aged 18 years or over, a UK resident (England, Northern Ireland,
Scotland, Wales) and had contact with domestic and/or wild birds. Full details of the
questionnaire methodology can be found in the Data Note (32), and the questionnaire is
available as extended data (33).

## Ethics and consent

146 Ethical approval for the study was obtained from the University of Bristol, Faculty Research
147 Ethics Committee, approval number 17048 on 16 January 2024.

148 Informed written consent (using e-consent hosted on Research Electronic Data CAPture tools,
149 REDCap (34,35)) for the use of data collected via the questionnaire was obtained from
150 respondents.

## Data collection

We used a convenience sampling method, where the questionnaire was initially delivered in-
person at the British Pig and Poultry Fair (15 and 16 May 2024) - an agricultural conference
aimed at pig and poultry keepers in the UK (36). The questionnaire was also distributed online
via individuals met at the fair, animal health protection teams from APHA and poultry networks
on social media. All questions were optional. Questionnaire data were collected using the
Research Electronic Data Capture (REDCap) platform, hosted at the University of Bristol
(34,35). Data captured between 15 May 2024 to 31 July 2024 were included in these analyses.

The questionnaire captured demographic data including age, gender, occupation and health.
We analysed the following questions relating to bird contact, biosecurity measures and risk
perception:

• How many birds of any type were you in direct contact with on a typical day?
o 1-10
o 11-100
o 101-1000
o 1001-10,000
o 10,001-100,000
o More than 100,000
• If you are a bird owner, how many birds do you own?
o 1-10
o 11-100

o   101-1000

o   1001-10,000

o   10,001-100,000

o   More than 100,000+

o   I am not a bird owner

•   Do you use any biosecurity measures to limit the risk of catching avian influenza (bird

flu)? Please select all that apply.

o   I use face masks

o   I use goggles

o   I use gloves

o   I change clothing when in contact with birds

o   I use outer garments when in contact with birds

o   I change boots

o   I use boot covers

o   I use a disinfecting footwear dip

o   I wash/clean my hands after touching birds

o   I wash/clean my hands before and after handling raw poultry meat

o   I make sure raw poultry meat is fully cooked before eating

o   I do not eat undercooked or raw poultry meat

o   I make sure eggs are thoroughly cooked

o   I do not eat raw eggs

o   I do not remember

o   None

o   Other

•   How much risk do you think avian influenza (bird flu) currently poses to your physical

health?

o   Very high risk

o   High risk

o   Medium risk

o   Low risk

o   No risk at all

o   I do not know

•   How much risk do you think avian influenza (bird flu) currently poses to the health of

your birds?

| 206 | o | Very high risk |
| 207 | o | High risk |
| 208 | o | Medium risk |
| 209 | o | Low risk |
| 210 | o | No risk at all |
| 211 | o | I do not know |

## Data analysis

RStudio (R version 4.4.1) was used to clean and analyse the dataset. The underlying code can

be found on GitHub (https://github.com/amythomas/aviancontactstudy). Completeness of data

varied by question. Use of biosecurity measures were recorded using radio buttons that

respondents 'checked' for yes. We assumed responses with all measures 'unchecked' indicated

a skipped question and were therefore coded as missing.

Descriptive statistics, including frequency counts, proportions and medians were calculated for

questions relating to respondent characteristics, bird contact, biosecurity measures and risk

perception. Occupational titles were derived from the Office for National Statistics' Extended

Standard Occupational Classification (SOC) 2020 Framework (37). Where an appropriate job

title was not listed, respondents could input their own job title into a free-text box which was

later combined into appropriate categories. Differences in uptake of BMs and risk perception

ranks between bird contact groups were tested using the two proportions z-test, and

confidence intervals (95%) for a single proportion were also calculated. Chi-squared tests of

independence or Fisher's exact tests (where cell counts were below five) were used to test for

an association between the uptake of BMs and the number of birds a person was in contact with

('bird contact') alongside perceived risk of AIV (to oneself and birds). Missing responses for BM

uptake, bird contact and risk perception were removed from these analyses.

Due to low cell counts among categories of bird contact, BM uptake and risk perception of AIV,

responses were categorised according to the groupings shown in Table 1. Respondents who

selected "I do not know" for the questions relating to risk perception of AIV to physical health

and bird health were considered as missing and excluded from subsequent analyses. The

frequency distribution of BM uptake and risk perception variables before grouping are shown in

the supplementary information (Figures S1 and S2).

**Table 1. Groupings for biosecurity measures, risk perception and bird contact.** Aggregate
groups were created for biosecurity measures, risk perception scores and bird contact
quantities for inferential analyses.

| Group | Biosecurity measure |
|---|---|
| Handwashing | I wash/clean my hands after touching birds |
| Footwear-related PPE | I change boots |
| | I use boot covers |
| | I use a disinfecting footwear dip |
| Food-safety measures | I wash/clean my hands before and after handling raw poultry meat |
| | I make sure raw poultry meat is fully cooked before eating |
| | I do not eat undercooked or raw poultry meat |
| | I make sure eggs are thoroughly cooked |
| | I do not eat raw eggs |
| PPE for the face or body | I use face masks |
| | I use goggles |
| | I use gloves |
| | I change clothing when in contact with birds |
| | I use outer garments when in contact with birds |
| None | None |
| Other | Other |
| I do not remember | I do not remember |

| Group | Risk perception |
|---|---|
| No/low/medium risk | No risk at all |
| | Low risk |
| | Medium Risk |
| High/very high risk | High risk |
| | Very high risk |
| I do not know | I do not know |

| Group | Quantity of birds a person was in contact with on a 'typical day' |
|---|---|
| Low | 1 – 10 |
| | 11 – 100 |
| | 101 – 1000 |
| High | 1001 – 10,000 |
| | 10,001 – 100,000 |
| | More than 100,000 |

# **Results**

## Respondent characteristics

A total of 225 respondents completed the questionnaire between 15 May 2024 and 31 July 2024.

Demographic characteristics are shown in Table 2. In-person respondents accounted for

approximately one third of the sample (28%, 63 of 225) compared to online completion of the

[9]

questionnaire (72%, 162 of 225). The median age was 48 years, ranging from 19 to 89 years. The
largest age category were respondents aged between 50 and 59 (27%, 60 of 225).

Respondents provided information on their occupation, bird contact and bird ownership.
Almost half were poultry farmers (46%, 104 of 225). Fewer respondents were veterinarians (7%,
16 of 225), zookeepers (7%, 15 of 225), retired (4%, 10 of 225), farm managers (3%, 7 of 225) or
another type of farmer (3%, 6 of 225). Other occupations which could not be grouped into these
categories accounted for 67 of 225 (30%) respondents.

Over half of respondents reported high contact levels with birds (≥1001 birds/day; 57%; 129 of
225). Low bird contact (≤1000 birds/day) was reported by 39% of all respondents (88 of 225).
Respondents were most likely to have daily contact with large flocks of 10,001–100,000 birds
(35%; 79 of 225), or presumed hobbyist/back-yard keepers in contact with 1–10 birds (16%;35 of
225) or 11–100 birds (19%; 42 of 225). Most respondents did not own birds (49%; 110 of 225). Of
those who did, ownership of birds was evenly distributed across the categories: 11% (24 of 225)
owned 1–10 birds, 11% (24 of 225) owned 11–100 birds and 12% (28 of 225) owned 10,001–
100,000 birds.

**Table 2. Avian Contact Study respondent characteristics.** Note, proportions may not sum to
100% due to rounding.

|  | Frequency (N=225) |
| --- | --- |
| **Demographics** | n (%) [min, max] |
| Questionnaire response type |  |
|     Online | 162 (72%) |
|     In-person | 63 (28%) |
| Median age | 48 [19 – 89] |
| Age categories |  |
|     19-29 | 21 (9%) |
|     30-39 | 48 (21%) |
|     40-49 | 46 (20%) |
|     50-59 | 60 (27%) |
|     60-64 | 22 (10%) |
|     ≥65 | 28 (12%) |
| Gender |  |
|     Male | 142 (63%) |
|     Female | 81 (36%) |
|     Other | <5 (<2%) |
| **Occupation** |  |
| Poultry farmers | 104 (46%) |

[10]

| | |
|---|---|
| Veterinarian | 16 (7%) |
| Zookeeper | 15 (7%) |
| Retired | 10 (4%) |
| Farm manager | 7 (3%) |
| Other type of farmer (mixed/livestock/arable) | 6 (3%) |
| Other[†] | 67 (30%) |
| **Bird contact on a 'typical day'** | |
| Low contact | **88 (39%)** |
|     1–10 birds | 35 (16%) |
|     11–100 birds | 42 (19%) |
|     101–1000 birds | 11 (5%) |
| High contact | **129 (57%)** |
|     1001–10,000 birds | 20 (9%) |
|     10,001–100,000 birds | 79 (35%) |
|     >100,000 birds | 30 (13%) |
| Missing | 8 (4%) |
| **Bird ownership** | |
|     1–10 birds | 24 (11%) |
|     11–100 birds | 24 (11%) |
|     101–1000 birds | <5 (<2%) |
|     1001–10,000 birds | <5 (<2%) |
|     10,001–100,000 birds | 28 (12%) |
|     >100,000 birds | 11 (5%) |
|     Non-bird owners | 110 (49%) |
|     Missing | 22 (10%) |

[†]Only groups with ≥5 observations are shown. All other groups are combined as 'Other'. Includes missing responses.

## BM measures to limit avian-to-human transmission are influenced by bird contact

Respondents were asked what biosecurity measures they used to prevent infection with AIV, and 7 of 225 responses were excluded due to missing BM uptake data. In total, 98% (213 of 218) reported using at least one BM, and 2% (5 of 218) reported using none (Figure 1). The single most reported BM was handwashing after touching birds, used by 89% (196 of 218) of respondents, followed by using disinfecting footwear dips, reported by 78% (170 of 218) and changing boots, reported by 70% (152 of 218).

**Figure 1. Proportion of participants reporting the use of biosecurity measures to prevent personal infection with avian influenza virus**. Colours represent biosecurity measures grouped into similar aggregate categories. Error bars represent 95% confidence intervals for a single proportion.

We hypothesised that the number of birds an individual was in contact with would affect uptake of biosecurity measures. A total of 217 respondents provided information on both bird contact and uptake of BMs. There was substantial evidence that a larger proportion of high bird contact respondents (≥1001 birds/day) reported using at least one footwear related PPE measure (z=6.49, p<0.001) or at least one PPE measure for the face or body (z=5.88, p<0.001), compared to low bird contact respondents (Figure 2). There was limited evidence for a difference in the proportion between high and low bird contact respondents reporting the use of at least one food

[12]

283    safety measure (z=0.94, p=0.447) and washing their hands after touching birds (z=1.41,

284    p=0.238). Only low bird contact respondents reported using no measures (6%, 5 of 88) (Figure

285    2).

**Figure 2. Biosecurity measure uptake by bird contact quantity.** Proportion of participants
using at least one biosecurity measure as per the aggregate biosecurity categories by low and
high bird contact group. Handwashing after touching birds was considered as its own category.
Low contact: 1–1000 birds; high contact: ≥1001 birds. Error bars represent 95% confidence
intervals for a single proportion.

292    Uptake of BMs was higher among the high bird contact group than the low bird contact group

293    (Table 3). There was evidence for an association between bird contact levels and the use of at

294    least one PPE measure for the face or body ($\chi 2$ (1, n=217) = 32.452, p<0.001), and use of at

295    least one footwear related PPE measure (Df=1, n=217, p<0.001) (Table 3). There was no

[13]

evidence of association for use of at least one food safety measure ($\chi$2 (1, n=217) = 0.888, p=0.346) or hand washing after touching birds ($\chi$2 (1, n=217) = 1.988, p=0.159). There was moderate evidence for the use of no measures (Df=1, n=217, p=0.010). However, frequency counts were small.

**Table 3. Relationship between biosecurity measures and bird contact.** Chi-square test of independence between biosecurity measures and bird contact level (Df = Degrees of freedom). P-values are rounded to 3 decimal places. Due to rounding, some percentages do not add to 100%.

| Biosecurity measure | | Low bird contact (n=88, %) | High bird contact (n=129, %) |
|---|---|---|---|
| Use of at least one food safety measure[†] | Reported | 70 (80%) | 109 (85%) |
| | Not reported | 18 (20%) | 20 (15%) |
| | Df[¶], $\chi$2, p-value | 1, 0.888, 0.346 | |
| Use of at least one PPE measure for the face or body[‡] | Reported | 57 (65%) | 123 (95%) |
| | Not reported | 31 (35%) | 6 (5%) |
| | Df[¶], $\chi$2, p-value | 1, 32.452, <0.001 | |
| Use of at least one footwear-related PPE measure[§] | Reported | 59 (67%) | 127 (98%) |
| | Not reported | 29 (33%) | 2 (2%) |
| | Df[¶], $\chi$2, p-value | 1, <0.001[††] | |
| Handwashing after touching birds | Reported | 76 (86%) | 119 (92%) |
| | Not reported | 12 (14%) | 10 (8%) |
| | Df[¶], $\chi$2, p-value | 1, 1.988, 0.159 | |
| Use of no measures | Reported | 5 (6%) | 0 (0%) |
| | Not reported | 83 (94%) | 129 (100%) |
| | Df[¶], $\chi$2, p-value | 1, 0.010[††] | |

[†] I wash/clean my hands before and after handling raw poultry meat; I make sure raw poultry meat is fully cooked before eating; I do not eat undercooked or raw poultry meat

[‡] I use face masks; I use goggles; I use gloves; I use change clothing when in contact with birds; I use outer garments when in contact with birds

[§] I change boots; I use boot covers; I use a disinfecting footwear dip

[¶] Degrees of freedom

[††] Fisher's Exact test

## Perceptions of avian influenza risk for bird health are higher than human health

We collected respondents' perceptions of AIV risk to their own health and the health of their birds. After removing missing data and those with unknown bird contact (34 of 225), 116 respondents reported high bird contact, and 75 respondents reported low bird contact (n=191) (Figure 3). Most respondents perceived the risk of AIV to their physical health to be low or medium (80%; 152 of 191). There was moderate evidence that a greater proportion of low bird contact than high bird contact respondents perceived the risk to be low (z=2.38, p=0.017; n-Low=52/75, n-High=59/116). There was no evidence for a difference between bird contact groups in all other risk categories for this question. Risk to the health of respondents' birds was largely perceived to be medium, high or very high (147 of 191). There was substantial evidence for a greater proportion of low bird contact respondents perceiving the risk to be low compared to high bird contact respondents (z=3.74, p<0.001; n-Low=24/75, n-High=11/116), and for increased high bird contact respondents perceiving the risk to be very high compared to low bird contact respondents (z=3.53, p<0.001; n-Low=8/75, n-High=40/116).

**Figure 3. Risk perception of avian influenza by bird contact.** Proportion self-reporting agreement with risk statements by bird contact group (low and high) for a) risk to human health and b) risk to the health of their birds. Low contact: 1 to 1000 birds; high contact: ≥1001 birds. Error bars represent 95% confidence intervals for a single proportion.

To test the association between perceived risk and uptake of BMs, risk categories were grouped into similar themes. Respondents who selected "I do not know" for physical health (n=5) and bird health (n=6) were coded as missing and were excluded from analyses, resulting in 186 and 185 respondents in the personal health and bird health categories respectively.

Respondents who considered AIV risk as high to bird health were more likely to use at least one footwear related PPE measure than those who perceived the risk as low ($\chi 2$ (1, n=185)= 9.171,

[16]

p=0.002). For all other biosecurity groups, there was no evidence of an association with

perceived risk to bird health or perceived risk to physical health (Table 4).

**Table 4: Relationship between biosecurity measure and perceived risk of avian influenza**
**virus to bird and personal health.** Chi-squared or Fisher's exact tests of association between
the use of biosecurity measures and perceived risk of avian influenza virus to personal and bird
health. Biosecurity measures were grouped into themes where appropriate. Risk groups were
aggregated into low (no, low and medium) and high (high and very high) risk. P-values are
rounded to 3 decimal places. Due to rounding, some percentages do not add to 100%.

| Biosecurity measure | | Risk to personal health (n=186) | | Risk to bird health (n=185) | |
|---|---|---|---|---|---|
| | | Low (n, %) | High (n, %) | Low (n, %) | High (n, %) |
| Use of at least one food safety measure[†] | Reported | 144 (77%) | 11 (6%) | 74 (40%) | 83 (45%) |
| | Not reported | 27 (15%) | 4 (2%) | 10 (5%) | 18 (10%) |
| | Df[¶], $\chi 2$, p-value | 1, 0.283[††] | | 1, 0.832, 0.362 | |
| Use of at least one PPE measure for the face or body[‡] | Reported | 139 (75%) | 15 (8%) | 64 (35%) | 89 (48%) |
| | Not reported | 32 (17%) | 0 (0%) | 20 (11%) | 12 (6%) |
| | Df[¶], $\chi 2$, p-value | 1, 0.078[††] | | 1, 3.766, 0.052 | |
| Use of at least one footwear-related PPE measure[§] | Reported | 144 (77%) | 15 (8%) | 64 (35%) | 94 (51%) |
| | Not reported | 27 (15%) | 0 (0%) | 20 (11%) | 7 (4%) |
| | Df[¶], $\chi 2$, p-value | 1, 0.133[††] | | 1, 9.171, 0.002[‡‡] | |
| Handwashing after touching birds | Reported | 155 (83%) | 14 (8%) | 77 (42%) | 90 (49%) |
| | Not reported | 16 (9%) | 1 (<1%) | 7 (4%) | 11 (6%) |
| | Df[¶], $\chi 2$, p-value | 1, 1.000[††] | | 1, 0.112, 0.737[‡‡] | |
| Use of no measures | Reported | 3 (2%) | 0 (0%) | 3 (2%) | 0 (0%) |
| | Not reported | 168 (90%) | 15 (8%) | 81 (44%) | 101 (55%) |
| | Df[¶], $\chi 2$, p-value | 1, 1.000[††] | | 1, 0.092[††] | |

[†] I wash/clean my hands before and after handling raw poultry meat; I make sure raw poultry meat is fully cooked before eating; I do not eat undercooked or raw poultry meat

[‡] I use face masks; I use goggles; I use gloves; I use change clothing when in contact with birds; I use outer garments when in contact with birds

[§] I change boots; I use boot covers; I use a disinfecting footwear dip

[¶] Degrees of freedom

[††] Fisher's Exact test

[‡‡] Yate's continuity correction

# Discussion

We determined biosecurity practices and risk perceptions of zoonotic avian influenza virus (AIV) among individuals in contact with kept and wild birds in the UK. Our analyses identified that handwashing was the most common biosecurity measure (BM) reported by respondents to prevent infection with AIV, followed by use of disinfecting footwear dips and changing boots. BM uptake was higher for respondents in daily contact with high numbers of birds. The perceived risk of AIV to respondents' own health was perceived as low, regardless of the number of birds a respondent had daily contact with. The risk of AIV to bird health was perceived as very high for those reporting 'high' daily bird contact (≥1001 birds/day). There were mixed perceptions for those reporting 'low' daily bird contact (<1000 birds/day). Our findings suggest that BM uptake in people in contact with birds is likely influenced by the perceived risk to avian health, but not human health and is further impacted by bird contact/farming scale.

The reported high uptake of footwear-related PPE measures is consistent with a recent qualitative study of UK poultry farmers (20), strengthening our understanding of these being common practices among the poultry industry. In contrast to a European study on poultry farms, we found hand washing after touching birds was also frequently reported to limit AIV transmission (23). This could reflect differences in questionnaire framing, where questions intended to elicit answers for preventing avian-to-human transmission led to respondents considering personal hygiene activities with improved response efficacy, as opposed to avian-centred BMs (25). This suggests that public health campaigns framed around avian-to-avian transmission risk may encourage the uptake of different BMs compared to those centred on avian-to-human transmission risk. In turn, this could affect the ability to control onward transmission of AIV following spillover to people and highlights the importance of appropriate risk framing.

Our results show that daily contact with a high number of birds was associated with greater BM uptake to prevent zoonotic AIV transmission. As a proxy for farming operation size, this is similar to other studies where larger farm sizes report higher BM uptake for the prevention of avian-to-avian transmission (22,27). Although small flock systems present a lower risk of AIV outbreaks (21), lower BM uptake may put individuals in contact with birds at a greater risk of exposure. Identifying cohorts with low BM uptake, like those in contact with small flocks, is useful to optimise targeted preventive public health programmes for zoonotic AIV.

The perceived risk of AIV to human health was not associated with the use of any BM measure, whereas risk to bird health was associated with the uptake of at least one footwear-related PPE

measure. Using protection motivation theory (25), this may be explained by participants' perceived susceptibility and severity of AIV to human health, demonstrated by risk rankings being low and medium; a tendency which has been observed in other demographics (30). This is in line with other findings that perceived human susceptibility to AIV is low, possibly affecting uptake of certain protective behaviours (38). This reinforces the importance of how the public health narrative of zoonotic AIV is framed when increased BM uptake is a desired outcome. Moreover, limited evidence for sustained human-to-human transmission may result in little motivation to use BMs (13).

## Strengths and limitations

This study compliments existing knowledge regarding biosecurity usage to prevent avian and human infection among bird keepers in Europe and South East Asia (21,30), and the changing transmission dynamics in North America and Europe over the last four years (3,12,13). Our questionnaire was co-developed with poultry farmers, veterinary public health agencies and stakeholders (32). This approach ensured relevancy of questions, accessibility of language and provided timely insight to the behaviours of groups potentially at risk of AIV exposure. Inclusion of individuals in contact with any bird (wild and/or domestic) provided nuanced information from different risk groups - going beyond the often-commercial poultry-centred approach to recruitment.

Despite this, there was an overrepresentation of commercial poultry farmers in the sample, as shown in the study Data Note (32), likely owing to the launch (and recruitment) at a national agricultural fair. This limits the generalisability of our findings to small-flock holders, like those studied in the UK Flockdown survey (21). Some covariates, such as farm assurance schemes and previous AIV outbreaks were not measured, preventing some statistical tests from being robustly performed. Future iterations of the study will aim to record these and ensure that questions are time-bound to enable the assessment of consistent biosecurity uptake. This will improve our understanding of how live outbreaks and wider public health contexts influence behaviour.

## Conclusions

Our findings suggest that use of BMs by people who have contact with birds in the UK are influenced by the number of birds a person is in contact with (as a proxy for farming operation size) and the perceived risk to bird health (high), but not perceived risk to human health (low).

[19]

404 Further work should be undertaken to identify the characteristics which lead to low BM uptake

405 among different cohorts in contact with birds to help inform zoonotic AIV public health

406 messaging and campaigns. Moreover, a more extensive investigation into the barriers and

407 facilitators of BM uptake is vital for improving our understanding of nuanced behaviour in the

408 context of zoonotic influenza spillover.

# Acknowledgements

410 We thank all participants who completed the questionnaire and those who contributed to the
411 development of the study.

# Supplementary materials

413 1. aiv_biosecurity_supplementary.docx - document containing Figure S1 (frequency counts of
414    different biosecurity measures in each biosecurity group) and Figure S2 (frequency counts
415    of risk perception categories for risk of avian influenza to respondents' health and the health
416    of their birds)

# References

1. European Centre for Disease Prevention and Control. 2021-2022 data show largest avian flu
epidemic in Europe ever [Internet]. 2022 [cited 2024 Nov 18]. Available from:
https://www.ecdc.europa.eu/en/news-events/2021-2022-data-show-largest-avian-flu-
epidemic-europe-ever

2. Lo FT, Zecchin B, Diallo AA, Racky O, Tassoni L, Diop A, et al. Intercontinental Spread of
Eurasian Highly Pathogenic Avian Influenza A(H5N1) to Senegal. Emerg Infect Dis [Internet].
2022 Jan;28(1):234–7. Available from: doi.org/10.3201/eid2801.211401

3. Peacock T, Moncla L, Dudas G, VanInsberghe D, Sukhova K, Lloyd-Smith JO, et al. The global
H5N1 influenza panzootic in mammals. Nature [Internet]. 2024 Sept 24; Available from:
doi.org/10.1038/s41586-024-08054-z

4. Uhart MM, Vanstreels RET, Nelson MI, Olivera V, Campagna J, Zavattieri V, et al.
Epidemiological data of an influenza A/H5N1 outbreak in elephant seals in Argentina
indicates mammal-to-mammal transmission. Nat Commun [Internet]. 2024 Nov
11;15(1):9516. Available from: doi.org/10.1038/s41467-024-53766-5

5. Aguado B, Begeman L, Günther A, Iervolino M, Soto F, Vanstreels RET, et al. Searching for
high pathogenicity avian influenza virus in Antarctica. Nat Microbiol [Internet]. 2024 Dec
1;9(12):3081–3. Available from: doi.org/10.1038/s41564-024-01868-7

6. World Organisation for Animal Health (WOAH). SMR_161191 - United Kingdom - Second
Semester 2022 – Terrestrial [Internet]. 2022 [cited 2024 Dec 19]. Available from:
https://wahis.woah.org/#/smr-review/161193?reportInfoId=98997&fromPage=smr-
management-url

7.  Department for Environment Food & Rural Affairs (DEFRA), Animal and Plant Health Agency
(APHA). Notifiable animal disease cases and control zones. 2022 [cited 2024 July 25]. Avian
Influenza Prevention Zone (AIPZ): the whole of England excluding Norfolk, Suffolk and parts
of Essex (2022). Available from: https://www.gov.uk/animal-disease-cases-england/avian-
influenza-prevention-zone-aipz-the-whole-of-england-excluding-norfolk-suffolk-and-parts-
of-essex-2022

8.  UK Health Security Agency. Research and analysis. 2024 [cited 2024 Nov 25]. HAIRS risk
statement: Avian influenza A(H5N1) in livestock. Available from:
https://www.gov.uk/government/publications/hairs-risk-statement-avian-influenza-ah5n1-
in-livestock/hairs-risk-statement-avian-influenza-ah5n1-in-livestock

9.  Department for Environment Food & Rural Affairs (DEFRA), Animal and Plant Health Agency
(APHA). Avian Influenza Prevention Zone declared for whole of England. 2025 Jan 23 [cited
2025 Feb 1]; Available from: https://www.gov.uk/government/news/avian-influenza-
prevention-zone-declared-for-whole-of-england

10. UK Health Security Agency. Research and Analysis. 2023 [cited 2025 July 1]. Investigation
into the risk to human health of avian influenza (influenza A H5N1) in England: technical
briefing 5. Available from: https://www.gov.uk/government/publications/avian-influenza-
influenza-a-h5n1-technical-briefings/investigation-into-the-risk-to-human-health-of-avian-
influenza-influenza-a-h5n1-in-england-technical-briefing-5#sources-and-
acknowledgments

11. US Centre for Disease Control. H5 Bird Flu: Current Situation [Internet]. 2025 [cited 2025
July 1]. Available from: https://www.cdc.gov/bird-flu/situation-
summary/?CDC_AAref_Val=https://www.cdc.gov/flu/avianflu/avian-flu-summary.htm

12. European Food Safety Authority EC for DP, Control EURL for AI, Adlhoch C, Fusaro A,
Gonzales JL, Kuiken T, et al. Avian influenza overview March – April 2023. EFSA J [Internet].
2023 [cited 2024 Nov 1];21(6):e08039. Available from:
https://doi.org/10.2903/j.efsa.2023.8039

13. Alexakis L, Buczkowski H, Ducatez M, Fusaro A, Gonzales JL, Kuiken T, et al. Avian influenza
overview June–September 2024. EFSA J [Internet]. 2024;22(10). Available from:
doi.org/10.2903/j.efsa.2024.9057

14. UK Health Security Agency. GOV.UK. 2025 [cited 2025 Feb 1]. Human case of avian flu
detected in England. Available from: https://www.gov.uk/government/news/human-case-
of-avian-flu-detected-in-england

15. US Centre for Disease Control. First H5 Bird Flu Death Reported in United States. 2025 Jan 6
[cited 2025 July 1]; Available from: https://www.cdc.gov/media/releases/2025/m0106-h5-
birdflu-death.html

16. Thomas A, Gokool S, Whitlow H, Clapp G, Moore P, Puleston R, et al. Seasonal Influenza
Vaccination in People who Have Contact With Birds. Influenza Other Respir Viruses
[Internet]. 2025;19(4):e70101. Available from: https://doi.org/10.1111/irv.70101

17. Department for Environment Food & Rural Affairs (Defra), Animal and Plant Health Agency
(APHA). Animal and Plant Health: guidance. 2022 [cited 2025 Jan 28]. Bird flu (avian
influenza): how to prevent it and stop it spreading - Biosecurity and hygiene standards to
keep birds safe from bird flu (avian influenza). Available from:

https://www.gov.uk/guidance/bird-flu-avian-influenza-how-to-prevent-it-and-stop-it-
spreading

18. Gupta SD, Fournié G, Hoque MA, Henning J. Factors influencing chicken farmers' decisions
    to implement prevention and control measures to reduce avian influenza virus spread
    under endemic conditions. Transbound Emerg Dis [Internet]. 2021 Jan 1;68(1):194–207.
    Available from: https://doi.org/10.1111/tbed.13757

19. Conan A, Goutard FL, Sorn S, Vong S. Biosecurity measures for backyard poultry in
    developing countries: a systematic review. BMC Vet Res [Internet]. 2012;8(1):240. Available
    from: doi.org/10.1186/1746-6148-8-240

20. Hosseini P, Syndicus I, Houghton E, Alarcon P, Fournie G, Guitian J, et al. Qualitative studies
    on farm managers perception of risk from HPAIV and implementation of biosecurity
    measures on UK poultry farms. Prev Vet Med [Internet]. 2025 June 1;239:106445. Available
    from: https://doi.org/10.1016/j.prevetmed.2025.106445

21. McClaughlin E, Elliott S, Jewitt S, Smallman-Raynor M, Dunham S, Parnell T, et al. UK
    flockdown: A survey of smallscale poultry keepers and their understanding of governmental
    guidance on highly pathogenic avian influenza (HPAI). Prev Vet Med [Internet]. 2024 Mar
    1;224:106117. Available from: https://doi.org/10.1016/j.prevetmed.2024.106117

22. Cui B, Liu ZP. Determinants of Knowledge and Biosecurity Preventive Behaviors for Highly
    Pathogenic Avian Influenza Risk Among Chinese Poultry Farmers. Avian Dis [Internet]. 2016
    [cited 2024 Aug 1];60(2):480–6. Available from: https://www-jstor-
    org.bris.idm.oclc.org/stable/26431788

23. Souillard R, Allain V, Dufay-Lefort CA, Rousset N, Amalraj A, Spaans A, et al. Biosecurity
    implementation on large-scale poultry farms in Europe: A qualitative interview study with
    farmers. Prev Vet Med [Internet]. 2024;224:106119. Available from:
    doi.org/10.1016/j.prevetmed.2024.106119

24. Dowd K, Taylor M, Toribio JALML, Hooker C, Dhand NK. Zoonotic disease risk perceptions
    and infection control practices of Australian veterinarians: Call for change in work culture.
    Prev Vet Med [Internet]. 2013;111(1–2):17–24. Available from:
    doi.org/10.1016/j.prevetmed.2013.04.002

25. Maddux JE, Rogers RW. Protection motivation and self-efficacy: A revised theory of fear
    appeals and attitude change. J Exp Soc Psychol [Internet]. 1983 Sept 1;19(5):469–79.
    Available from: https://doi.org/10.1016/0022-1031(83)90023-9

26. Ameji ON, Abdu PA, Sa'idu L, Kabir J, Assam A. Awareness, Knowledge, Readiness to Report
    Outbreak and Biosecurity Practices Towards Highly Pathogenic Avian Influenza in Kogi State,
    Nigeria. Int J Poult Sci [Internet]. 2012;11(1):11–5. Available from:
    http://dx.doi.org/10.3923/ijps.2012.11.15

27. Delpont M, Racicot M, Durivage A, Fornili L, Guerin JL, Vaillancourt JP et al. Determinants of
    biosecurity practices in French duck farms after a H5N8 Highly Pathogenic Avian Influenza
    epidemic: The effect of farmer knowledge, attitudes and personality traits. Transbound
    Emerg Dis [Internet]. 2021 Jan 1 [cited 2024 June 19];68(1):51–61. Available from:
    doi.org/10.1111/tbed.13462

523     28.   Pao HN, Jackson E, Yang TS, Tsai JS, Sung WHT, Pfeiffer DU. Determinants of farmers'
524           biosecurity mindset: A social-ecological model using systems thinking. Front Vet Sci
525           [Internet]. 2022 Aug 15 [cited 2024 June 19];9. Available from:
526           doi.org/10.3389/fvets.2022.959934

527     29.   Manabe T, Thuy PTP, Can VV, Takasaki J, Huyen DTT, Chau NTM, et al. Impact of Educational
528           Intervention Concerning Awareness and Behaviors Relating to Avian Influenza (H5N1) in a
529           High-Risk Population in Vietnam. PLoS ONE [Internet]. 2011 Aug 22 [cited 2024 June
530           19];6(8):e23711. Available from: doi.org/10.1371/journal.pone.0023711

531     30.   Cui B, Wang F, Wang LDL, Pan C, Ke J, Tian Y. A Comparative Analysis of Risk Perception and
532           Coping Behaviors among Chinese Poultry Farmers Regarding Human and Poultry Infection
533           with Avian Influenza. Int J Environ Res Public Health [Internet]. 2019 Oct 11 [cited 2024 June
534           19];16(20):3832. Available from: doi.org/10.3390/ijerph16203832

535     31.   Pao HN, Jackson E, Yang TS, Tsai JS, Hwang YT, Sung WHT, et al. The attitude-behaviour gap
536           in biosecurity: Applying social theories to understand the relationships between
537           commercial chicken farmers' attitudes and behaviours. Front Vet Sci [Internet]. 2023 Feb 9
538           [cited 2024 June 19];10. Available from: doi.org/10.3389/fvets.2023.1070482

539     32.   Amy C. Thomas, Suzanne Gokool, Genevieve Clapp, Harry Whitlow, Carmel McGrath, Peter
540           Moore, et al. Data note on the Avian Contact Study: a questionnaire resource for avian
541           influenza public health planning [version 2; peer review: 2 approved]. Wellcome Open Res
542           [Internet]. 2025;9(604). Available from: https://doi.org/10.12688/wellcomeopenres.23064.1

543     33.   Thomas A, Gokool S, Clapp G, Whitlow H, McGrath C, Moore P, et al. Avian Contact Study
544           Questionnaire and Data Dictionary [Internet]. Zenodo; 2024 [cited 2025 July 28]. Available
545           from: https://zenodo.org/records/13617061

546     34.   PA Harris, R Taylor, R Thielke, J Payne, N Gonzalez, JG. Conde. Research electronic data
547           capture (REDCap) – A metadata-driven methodology and workflow process for providing
548           translational research informatics support. J Biomed Inform. 2009;42(2):377–81.

549     35.   PA Harris, R Taylor, BL Minor, V Elliott, M Fernandez, L O'Neal, et al. The REDCap
550           consortium: Building an international community of software partners. J Biomed Inform
551           [Internet]. 2019; Available from: doi.org/10.1016/j.jbi.2019.103208

552     36.   The British Pig & Poultry Fair [Internet]. [cited 2024 July 15]. Available from:
553           https://pigandpoultry.org.uk/

554     37.   Office for National Statistics. Classifications and Standards. 2023 [cited 2024 Sept 9].
555           Extended SOC 2020. Available from:
556           https://www.ons.gov.uk/methodology/classificationsandstandards/standardoccupationalcl
557           assificationsoc/standardoccupationalclassificationsocextensionproject

558     38.   Pae R, Findlater L, Amlôt R, Capelastegui F, Dabrera G, Humphreys C, et al. "I definitely
559           cannot afford to be feeling poorly if there's no need to be": a qualitative evaluation of
560           antiviral uptake following suspected occupational exposure to avian influenza. BMC Public
561           Health [Internet]. 2025;25(1). Available from: doi.org/10.1186/s12889-025-21459-3