Decoding subphenotypes in electronic medical records within late-onset Alzheimer's disease reveals heterogeneity and sex-specific differences

Yukari Katsuhara[1,2], Umair Khan[1,3], Zachary A. Miller[4,5], Isabel E. Allen[6,7,8], Tomiko T. Oskotsky[1,9], Marina Sirota[1,11,12,*], Alice S. Tang[1,10,11,*]

[1]Bakar Computational Health Sciences Institute, UCSF, San Francisco, CA, USA

[2]Health Data Science Graduate Program, UCSF, San Francisco, CA, USA

[3]Biological and Medical Informatics Graduate Program, UCSF, San Francisco, CA, USA

[4]Memory and Aging Center, Department of Neurology, UCSF Weill Institute for Neurosciences, UCSF, San Francisco, CA, USA

[5]Dyslexia Center, Department of Neurology and Psychiatry, UCSF Weill Institute for Neurosciences, UCSF, San Francisco, CA, USA

[6]Department of Epidemiology and Biostatistics, UCSF, San Francisco, CA, USA

[7]Global Brain Health Institute, UCSF, San Francisco, CA, USA

[8]Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

[9]Division of Clinical Informatics and Digital Transformation, Department of Medicine, UCSF, San Francisco, CA, USA

[10]School of Medicine, UCSF, San Francisco, CA, USA

[11]Senior author

[12]Lead contact

[*]Correspondence: marina.sirota@ucsf.edu (M.S.) and alice.tang@ucsf.edu (A.S.T.) share co-corresponding authorship.

Summary

We applied unsupervised learning techniques to electronic medical records from UCSF to identify distinct Alzheimer's disease subgroups based on comorbidity profiles. Given the well-known female sex predominance in Alzheimer's disease prevalence, we performed sex-stratified analyses to evaluate differences in disease manifestations based on sex. Findings were validated using an independent UC-Wide dataset. Among 8,363 patients, we identified five Alzheimer's disease subphenotypes, characterized by comorbidities related to cardiovascular conditions, gastrointestinal disorders, and frailty-related conditions such as pneumonia and pressure ulcers. We further refined significant comorbidity variations across clusters through sex-stratified analyses, observing a higher prevalence of circulatory diseases in males in Cluster 2 and bladder stones in females in Cluster 3. Key results were consistent across the UCSF and UC-Wide datasets. Our study identifies clinically meaningful Alzheimer's disease subgroups, along with sex-specific variations, suggesting underlying biological factors, and indicates the potential utility of these findings in informing individualized therapeutic regimens.

Keywords

Introduction

Alzheimer's disease (AD) is the most common cause of dementia. With rising prevalence and no curative treatments, AD represents a major public health crisis worldwide. Clinically, AD has traditionally been conceptualized as an amnestic disorder predominantly affecting memory in those 65 and older, but it can also manifest at an earlier age and with non-amnestic behavioral, executive, language, visuospatial, and asymmetric motor coordination phenotypes. Pathologically, AD is defined by the accumulation of β-amyloid (Aβ) plaques, hyperphosphorylated tau neurofibrillary tangles, glial changes (microglia and astrocytes), and subsequent neurodegeneration [1,2]. Despite decades of research, its biological mechanism remains incompletely understood, making early diagnosis and effective management challenging.

The greatest known risk factor for developing AD is age followed by Apolipoprotein E (*APOE*) genetic status [3,4]. Cardiovascular and metabolic risk factors have also been shown to affect risk in typical amnestic late-onset AD (LOAD) presentations, while novel risk factors may apply in earlier onset and non-amnestic presentations [5], highlighting our incomplete understanding of AD vulnerability. Increasingly, sex has emerged as a critical factor in understanding AD risk, progression, and clinical manifestations, as women account for approximately two-thirds of AD cases in the United States, with a higher lifetime risk of developing AD than men [6]. Recent studies indicate that sex may modify multiple aspects of AD, including susceptibility of AD, progression, and molecular pathology including *APOE* [7-12]. Men with AD tend to have a shorter survival time, while women experience greater memory impairment [13-15]. Yet, here too, the mechanisms underlying these sex differences remain poorly understood, underscoring the need for more research into sex-specific risk factors associated with AD.

Recent advances in electronic medical records (EMRs) and machine learning have enabled large-scale, data-driven approaches to studying AD heterogeneity. EMRs capture a wide range of real-world health data, including multimorbidity patterns, medication histories, and disease trajectories, making them valuable resources for identifying novel AD subtypes and potential therapeutic targets [16]. Data-driven phenotyping approaches have revealed distinct AD subtypes influenced by factors including sex and race

[17-20], but comprehensive and unbiased analyses for sex-specific patterns among subphenotypes of AD remain largely unexplored.

In this study, we apply unsupervised clustering to a large EMR dataset of AD patients, consisting largely of LOAD presentations, to identify if subphenotypes within this group exist. By leveraging real-world clinical data, we aim to uncover novel insights into AD heterogeneity, which may inform precision medicine approaches and improve comprehensive, sex-specific care for patients suffering from neurodegenerative disease. Deeper characterization of AD-related comorbidities and their subphenotypes may uncover novel risks for the disease and facilitate the development of individualized therapeutic regimens that could serve as adjuvants to the current anti-amyloid- and upcoming anti-tau-based treatments, maximizing therapeutic benefits for patients through a truly personalized therapeutic approach.

Results

Patient characteristics

AD cohorts were identified from University of California (UC) San Francisco (UCSF) data and UC-Wide data, with UCSF patients excluded from the latter, respectively. We identified 8,363 AD patients from the UCSF EMRs (5,315 female (64%); median age: 90.0 [IQR: 84.0–91.0]) (Figure 1). From the UC-Wide validation dataset, we identified 25,896 AD patients (16,036 female (61.9%); median age: 89.0 [IQR: 82.0–90.0]). In both datasets, AD patients were predominantly "White" (62.0% at UCSF, 65.7% in the UC-Wide dataset), followed by those classified as "Other" and "Asian". At UCSF, most AD patients (75.0%) were recorded as "Alive," whereas in the UC-Wide dataset, only 50.9% were alive. Additionally, the median number of comorbidities recorded at UCSF was lower compared to the UC-Wide dataset (UCSF: median = 23.0 [IQR: 9.0–63.0]; UC-Wide: median = 40.0 [IQR: 16.0–88.0]). Other demographic characteristics of AD patients are presented in Table 1 and 2.

Low-dimensional embeddings derived from principal components of diagnostic data reveal five subphenotypes of AD

The analysis data consisted of patient-level diagnostic records, where each row represented an individual patient and each column corresponded to one of 33,031 unique diagnosis names recorded by International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM) codes across 8,363 AD patients. Diagnosis names were one-hot encoded, with a binary value indicating the presence or absence of each diagnosis for a given patient. To mitigate the impact of redundant or noisy features, Principal Component Analysis (PCA) was applied to diagnosis names (33,031 features) prior to K-means clustering. For the UCSF dataset, we selected 1,000 principal components, capturing approximately 80% of the cumulative explained variance, as input features for clustering.

Cluster determination was based on the analysis of within-cluster sum of squares (WSS) and silhouette scores. The WSS analysis indicated that after four clusters, the rate of WSS reduction diminished substantially, suggesting that increasing the number of clusters beyond four did not provide meaningful improvements in partitioning. Conversely, a sharp decline in the silhouette score was observed between five and six clusters (88.1%), indicating a considerable loss in clustering quality. These trends suggest that either a four- or five-cluster (Cluster 1–Cluster 5) solution could be appropriate. To balance cluster compactness with the representation of heterogeneity, the five-cluster solution was selected, as the four-cluster solution may not fully capture the diversity of AD subphenotypes (Supplemental Table 1).

Within the UCSF clusters, Cluster 5 had significantly fewer comorbidities (median = 12.0 [IQR: 5.0–21.0], Bonferroni-corrected p-values < 0.05) compared to the other clusters, whereas Cluster 1 had the highest number of comorbidities (median = 313.5 [IQR: 258.0–379.0], Bonferroni-corrected p-values < 0.05). In terms of age distribution, Cluster 4 (median = 91.0 [IQR: 90.0–91.0], Bonferroni-corrected p-values < 0.05) was significantly older than Cluster 1, 3, and 5, followed by Cluster 2. Conversely, Cluster 3 (median = 88.0 [IQR: 82.0–90.0], Bonferroni-corrected p-values < 0.05) was significantly younger than all other clusters (Supplemental Table 2). Black or African American individuals comprised 13% of Cluster 2 and 12% of Cluster 4, representing a higher proportion compared to other clusters (Bonferroni-corrected p-values < 0.05, Supplemental Table 2). Regarding sex, Cluster 5 included a significantly

higher proportion of males than Cluster 4, with these males displaying a younger age distribution. Other demographic characteristics and sex-stratified distributions for each cluster are presented in Table 1 and Supplemental Table 3.

For the UC-Wide dataset, 19,835 unique diagnosis names from 25,896 AD patients were used as features for PCA. 1,100 principal components covering approximately 80% of the cumulative explained variance were selected as input features for K-means clustering. Consistent with the UCSF dataset, five clusters (Cluster A–Cluster E) were identified based on silhouette scores and WSS (Supplemental Table 1). The WSS reduction analysis indicated that the decrease from five to six clusters was relatively small, suggesting that increasing the number of clusters beyond five does not substantially improve clustering compactness. In contrast, the silhouette score analysis revealed a significant decline in clustering structure when increasing from four to five clusters (50.0%), indicating a reduction in cluster separation. Based on these findings, a four- or five-cluster solution appeared to provide the most stable clustering structure. Given that this study aims to capture the heterogeneity of comorbidity patterns in AD, the five-cluster solution was selected.

Significant differences in the number of comorbidities were observed between Cluster A (median = 201.0 [IQR: 166.0–253.0], Bonferroni-corrected p-values < 0.05) and Cluster B (median = 15.0 [IQR: 7.0–25.0], Bonferroni-corrected p-values < 0.05) (Supplemental Table 2). Other demographic characteristics of each cluster are presented in Table 2. A low-dimensional Uniform Manifold Approximation and Projection (UMAP) visualization of the PCA-transformed components illustrates the distribution of AD patient clusters, sex differences, the number of comorbidities, and locations (Figure 2).

Comorbidity enrichment analysis shows significant cluster-related comorbidities at UCSF

Our analysis identified significant diagnoses in each cluster within the UCSF dataset (two-sided Fisher's exact or Chi-squared test, Bonferroni-corrected p-value < 0.05) (Figure 3). Cluster 1 had 1,911 significant comorbidities, most of which were positively associated. In contrast, 1,650 significant comorbidities were

identified in Cluster 5, predominantly showing negative associations. However, a few conditions exhibited positive associations, including Pick's disease. Cluster 2 – 4 displayed unique comorbidity enrichment patterns. For example, in Cluster 2 (985 significant comorbidities), the top positively associated diagnoses included laceration, phlebitis and thrombophlebitis of femoral vein, and benign neoplasm of the conjunctiva. Interestingly, Cluster 3 (682 significant comorbidities) showed relatively lower prevalence of mental disorders and personality changes. Finally, Cluster 4 (245 significant comorbidities) exhibited positive associations with pneumonia and disseminated intravascular coagulation (Supporting Data). Manhattan plots illustrated the distribution of statistical significance (Supplemental Figure 1).

Significant cluster-specific comorbidities at UCSF are subsequently validated in the UC-Wide validation cohort

We identified cluster-specific comorbidities using UpSet plots (Figure 4). Cluster 1 exhibited the highest number of positively associated comorbidities (812 comorbidities), followed by Cluster 2 (225 comorbidities). Cluster 3 had 96 positively associated comorbidities and one negatively associated comorbidity (personality change), while Cluster 4 had 13 positively and 6 negatively associated comorbidities. In contrast, Cluster 5 had only two positively associated comorbidities, whereas 1,581 comorbidities were negatively associated. Examining the most significantly associated comorbidities specific to each cluster, we found that Cluster 3 was characterized by acute systolic heart failure and ileus, whereas pneumonia and brief psychotic disorder were observed in Cluster 4. In Cluster 5, we observed essential hypertension, anemia, and urinary tract infection as the most significantly associated comorbidities (Figure 5).

Additionally, cluster-specific comorbidities with high odds ratios were identified (Figure 6A). Cluster 1 showed numerous significant associations with a variety of conditions, including diabetes mellitus with other diabetic kidney complication, contusion of the right upper arm, displaced fracture of shaft of right clavicle, complication of kidney transplant, encounter for aftercare following kidney transplant, and

mononeuropathy of right lower (all ORs = 227, Bonferroni-corrected p-values < 0.05). In Cluster 2, phlebitis and thrombophlebitis of femoral vein had the highest odds ratio (OR = 44.1, Bonferroni-corrected p-value < 0.05). Cluster 3 was characterized by infection and inflammatory reaction due to a urinary catheter (OR = 103, Bonferroni-corrected p-value < 0.05), followed by intestinal obstruction (OR = 84.3, Bonferroni-corrected p-value < 0.05). Additionally, dysphagia was significantly associated with Cluster 3 (OR = 5.3, Bonferroni-corrected p-value < 0.05, Supporting Data). Cluster 4 was significantly associated with poisoning by cardiac-stimulant glycosides (OR = 23.6, Bonferroni-corrected p-value < 0.05), pneumonia (OR = 10.6, Bonferroni-corrected p-value < 0.05), and pressure ulcer (OR = 7.5, Bonferroni-corrected p-value < 0.05). Lastly, in Cluster 5, Pick's disease was significantly enriched (OR = 13.4, Bonferroni-corrected p-value < 0.05). However, ORs in our study should be interpreted with caution, as the frequency of certain diagnoses in each cluster was relatively small, which may have led to inflated odds ratios (Supporting Data).

In the UC-Wide validation dataset, we identified corresponding UC-Wide clusters for each UCSF cluster based on the overlap in significant comorbidities, as visualized by Sankey plots (Figure 7A). The proportion of significant comorbidities in each UCSF cluster that were also captured in the corresponding UC-Wide clusters ranged from 20% to 55%. For instance, 239 of the 812 (29.4%) significant comorbidities identified in UCSF Cluster 1 were also observed in the matched UC-Wide Cluster A, which itself contained 1,976 cluster-specific significant comorbidities.

To further evaluate the consistency of comorbidity enrichment across datasets, we examined the correlation of shared comorbidities using Log-Log plots (Figure 7B). UCSF cluster-specific comorbidities with high odds ratios including phlebitis and thrombophlebitis in Cluster 2, intestinal obstruction and infection and inflammatory reaction due to a urinary catheter in Cluster 3, and pneumonia and pressure ulcer in Cluster 4, were largely replicated in their corresponding UC-Wide clusters. However, Cluster 2 did not show a significant correlation with its matched UC-Wide Cluster A (Pearson r = 0.07, p = 0.68).

Sex-stratified analysis identifies significantly associated cluster-sex-specific comorbidities at UCSF, which are subsequently validated in the UC-Wide validation cohort

After illustrating the distribution of statistical significance in Miami plots, we identified cluster-sex-specific comorbidities using UpSet plots (Figure 4, Supplemental Figure 2). When we focus on the significant comorbidities exclusive to each cluster, in Cluster 1, females showed a strong association with pain-related conditions, whereas males were primarily associated with symptoms of the genitourinary system. In Cluster 2, females exhibit a strong association with tension-type headaches, whereas males are more frequently associated with circulatory diseases, including unstable angina and hypertensive heart disease. In Cluster 3, females show significant associations with pressure ulcers and osteoarthritis, while males are predominantly linked to respiratory diseases, such as acute pulmonary edema, and circulatory conditions, including acute systolic heart failure. In Cluster 4, females are significantly associated with pneumonia. In Cluster 5, we observed essential hypertension, anemia, and urinary tract infection across both sexes (Figure 8).

Considering odds ratios, diabetes-related conditions were significantly associated with Cluster 1 in both sexes, with diabetes mellitus due to an underlying condition with diabetic chronic kidney disease being predominant in females (OR = 146) and diabetes mellitus with hyperosmolarity in males (OR = 169). Similarly, Pick's disease remained highly associated with Cluster 5 across both sexes (Female OR = 8.6, Male OR = 4.1). In Cluster 2, females exhibit stronger associations with eye diseases, such as pingueculitis (OR=50.4, Bonferroni-corrected p-value < 0.05), whereas males are more associated with circulatory diseases, including hypertension (OR=43.5, Bonferroni-corrected p-value < 0.05) and elevated erythrocyte sedimentation rate (ESR) (OR=32.7, Bonferroni-corrected p-value < 0.05). In Cluster 3, females demonstrate high odds ratios for tinnitus (OR=57.7, Bonferroni-corrected p-value < 0.05), intestinal obstruction (OR=57.7, Bonferroni-corrected p-value < 0.05), and bladder calculus (OR=25.3, Bonferroni-corrected p-value < 0.05), while males are significantly associated with infection and inflammatory reaction (OR=75.5, Bonferroni-corrected p-value < 0.05) and enterocolitis (OR=55.5,

Bonferroni-corrected p-value < 0.05). Lastly, in Cluster 4, females have strong associations with poisoning by cardiac-stimulant glycosides (OR=20.6, Bonferroni-corrected p-value < 0.05) and pneumonia (OR=15.2, Bonferroni-corrected p-value < 0.05), whereas males are associated with pressure ulcers (OR=12.1, Bonferroni-corrected p-value < 0.05) (Figure 6B and 6C, Supporting Data). However, sex-stratified odds ratios should be interpreted with caution due to the limited number of cases for certain diagnoses, which may lead to variability in estimates.

In the UC-Wide dataset, we identified corresponding UC-Wide clusters based on the coverage rate of UCSF cluster-specific comorbidities in each sex (Supplemental Figure 3). The matched UC-Wide clusters remained consistent with those identified in the overall population, except for UCSF Cluster 3 among females. For UCSF Cluster 4 among males, two UC-Wide clusters had the same coverage rate. In this case, the cluster that was consistently matched in the overall analysis and had one of the highest coverage rates was selected as the matched cluster. Between 15% and 60% of significant comorbidities in each UCSF cluster were recaptured by the matched UC-Wide clusters. Overall, UCSF cluster-specific comorbidities with high odds ratios —such as elevated ESR in Cluster 2 among males, intestinal obstruction in Cluster 3 among females, and infection and inflammation reaction due to urinary catheter, and enterocolitis in Cluster 3 among males—were largely replicated in the UC-Wide clusters. However, some Pearson correlation coefficients were neither strong nor statistically significant (Figure 9).

Discussion

In this study of 8,363 AD patients from UCSF and 25,896 from an independent UC-Wide cohort, we analyzed multi-site EMR by applying dimensionality reduction, followed by clustering analysis, and in the process, we identified five subphenotypes of AD based on comorbidities. Enrichment analysis comparing a specific cluster with the other clusters revealed clinical heterogeneity. Sex-stratified analysis further highlighted sex-specific differences within the identified clusters. In the following sections, we present our findings for each cluster, followed by the identification of notable sex-specific differences, and discuss their implications.

Cluster 1: high comorbidity burden with complex clinical profiles

We performed comorbidity enrichment analysis for overall populations. Cluster 1 was characterized by a high comorbidity burden, which may indicate the presence of patients with complex clinical profiles. In particular, diseases with the highest odds ratios and lowest p-values included several kidney-related conditions—diabetes mellitus with other diabetic kidney complications, complications of kidney transplant, and encounter for aftercare following kidney transplant. These findings highlight the prominence of renal comorbidities in this cluster. However, additional enriched conditions such as contusion of the right upper arm, displaced fracture of shaft of right clavicle, and mononeuropathy of right lower were also identified, reflecting a broader range of systemic complications beyond kidney disease alone. Kidney transplantation is a recognized risk factor for AD and is often associated with multiple comorbidities due to long-term immunosuppressive therapy and underlying chronic conditions[21,22], therefore, kidney-related conditions are suggested as a key component of this cluster, although they do not solely define it. Further stratification of this cluster is warranted to identify subgroups with specific comorbidity patterns, which may facilitate the development of more targeted therapeutic treatment.

Cluster 2: African American individuals, phlebitis and cardiovascular conditions

Cluster 2 presented with a relatively higher proportion of Black or African American individuals. AD prevalence is about twice as high in this population compared to non-Hispanic Whites. Studies have suggested that *APOE* ε4-based mechanisms of AD may be less relevant to Black or African American than non-Hispanic Whites, and thus this cluster may be highlighting *APOE* ε4-independent risks, such as social determinants of health [6,23-25]. Patients in Cluster 2 also exhibited enriched comorbidities such as phlebitis and thrombophlebitis of femoral vein. Superficial thrombophlebitis, the most common form of thrombophlebitis, has been increasingly recognized as a condition associated with deep vein thrombosis and pulmonary embolism, both of which can lead to serious clinical outcomes [26-28]. Established risk factors for these thrombotic conditions include a hypercoagulable state and the presence of abnormal

fibrin clot structures [29-31]. Previous literature suggests that increased fibrinogen aggregation and Aβ

fibrillization occur in AD brain parenchyma and vessels and induce the formation of abnormal fibrin clots

that are more resistant to degradation [32,33]. Additionally, recent studies have shown that blood-derived

fibrin deposits in the central nervous system are a common feature of autoimmune and neurodegenerative

diseases, including AD [34,35]. These findings highlight fibrin as a potential therapeutic target for

immunotherapy designed to reduce fibrin-induced neurotoxicity [36]. There is no strong evidence for the

impact on peripheral blood vessels caused by the abnormal fibrin clots in the brain; however, previous

evidence shows that peripheral treatment with a low molecular weight heparin reduces plaques and Aβ

accumulation in a mouse model of Alzheimer's disease [37]. This suggests a possibility that increased

fibrinogen aggregation and Aβ in AD patients may contribute to vascular abnormalities such as phlebitis.

Cluster 3: lower rates of mental illness and increased infections and gastrointestinal dysfunction

This group exhibited significant negative associations with mental disorders. Given that mental disorders

are commonly reported comorbidities in AD patients [38], this finding is particularly noteworthy. One

possible explanation for the lower prevalence of mental disorders in Cluster 3 is the significantly younger

age of patients in this cluster compared to others. This aligns with prior research indicating that AD is a

progressive disease in which the risk of comorbidities, including mental disorders, increases over time [39].

On the other hand, previous evidence has demonstrated that early-onset AD (EOAD) is associated with a

more aggressive clinical course and distinct biological mechanisms compared to LOAD and in some

reports actually possesses a higher amount of psychiatric symptoms burden [5,40-43]. However, since this

study focused on LOAD population and the ICD-10-CM's broad disease classification, such as "Other

specified mental disorders," we cannot ascertain whether this relatively younger population reflects an

early onset presentation of AD or is instead at the younger age range of LOAD.

Additionally, we identified infection and inflammatory reactions due to a urinary catheter, as well as

intestinal obstruction, as positively associated comorbidities in Cluster 3. Increasing evidence shows

reciprocal associations between AD, infection, and inflammatory reactions [44-46,]. Therefore, the

combination of inflammation and reduced mobility associated with AD may contribute to the comorbidity profile observed in Cluster 3.

Regarding intestinal obstruction, several possible mechanisms can be considered. First, reduced mobility, including oropharyngeal dysphagia, and altered nutritional intake associated with AD progression may contribute to gastrointestinal dysfunction such as constipation [47-49]. Notably, dysphagia was identified as a Cluster 3-specific comorbidity (OR = 5.3, Bonferroni-corrected p-value < 0.05, Supporting Data), suggesting that individuals in this cluster may be at higher risk of malnutrition and associated gastrointestinal deficiency. Secondly, recent studies have explored the associations between AD and gut microbiota [50,51]. Research has shown that AD patients exhibit reduced microbial diversity compared to cognitively healthy individuals [52]. It is hypothesized that AD-related alterations in gut microbiota may contribute to gastrointestinal dysfunction, potentially increasing the risk of intestinal obstruction. However, further research is required to elucidate the interplay between AD progression, gut microbiota alterations, and gastrointestinal complications, including intestinal obstruction.

Cluster 4: oldest age and strong association with pneumonia

Patients in Cluster 4 constituted the oldest subgroup and exhibited a higher prevalence of pneumonia, a condition commonly associated with aging and characterized by functional decline, including impaired swallowing function [53]. This observation is particularly noteworthy given the longstanding hypothesis that infectious respiratory diseases may precipitate the onset of AD. This hypothesis was originally proposed over a century ago and has since gained support from a growing body of scientific evidence [54-56], including recent studies that demonstrate lower risks of developing AD in those who were vaccinated against respiratory pathogens, including pneumococcal and influenza[57-60]. Our findings underscore the clinical relevance of implementing preventive strategies, such as vaccination, to reduce modifiable risk factors for neurodegeneration in older populations.

Cluster 5: male sex, low comorbidity burden, potential overlap with frontotemporal dementia

Cluster 5 includes a significantly higher proportion of males with a relatively younger age distribution. Given that AD possesses a female predominance, the enrichment of younger males is particularly intriguing. While this showed the fewest positively associated comorbidities, there was one notable exception, Pick's disease. Pick's disease, also referred to as behavioral variant frontotemporal dementia (bvFTD), is second to AD in being the most common neurodegenerative disorder in individuals with first symptoms younger than 65 [61,62]. Pick's disease is also notable for its higher prevalence of male sex [63,64]. Considering the demographics of Cluster 5, we wonder if this cluster might be identifying groups where conditions like bvFTD were on the differential diagnosis.

Sex specific analysis:

While sex related results only came out in our first pass analysis in cluster 5, given the significant female bias to AD, we performed follow up sex specific analyses. Here, our findings underscored the presence of sex-specific disease manifestations in Clusters 2 and 3. In Cluster 2, hypertension and elevated ESR were strongly associated with males. Alongside hypertension, diseases known to increase ESR, such as diabetes [65] and cardiovascular disease (CVD) [66], are well-established risk factors for AD. Previous studies have demonstrated strong associations between these conditions and cerebrovascular pathology, as well as cognitive decline [38,67].

Our findings suggest that Cluster 2 represents a specific subgroup of AD males with pronounced vascular and metabolic dysfunction, which contrasts with previous reports indicating that these conditions are more common in AD females [10,68,69]. A potential explanation for these sex differences may involve *APOE* genotype, which is known as a risk factor for hypertension, CVD, and diabetes [70-72]. Furthermore, existing evidence suggests that the impact of *APOE* varies by sex and genotypes [12,71-73]. Given this, it is plausible that male sex and *APOE* genotype could contribute to the observed vascular-metabolic phenotype in Cluster 2. However, our dataset lacks *APOE* information, and we were not able to directly assess the potential combined effects of male sex and *APOE* genotype. Further studies incorporating genetic data are needed to test this hypothesis.

Another potential explanation for the observed comorbidity patterns in Cluster 2 involves the interaction between racial composition and susceptibility to APOE ε4-related mechanisms. Although Cluster 2 includes a relatively higher proportion of Black or African American individuals overall, sex-stratified demographics revealed that the male subgroup had a higher proportion of White individuals and a lower proportion of Black individuals compared to the female subgroup.

Previous studies suggest that White populations are more susceptible to *APOE* ε4-associated AD than Black populations, and *APOE* ε4 has been linked to an increased risk of vascular dysfunction [23,24,71,72,74]. Therefore, this racial stratification may partially explain the higher burden of *APOE* ε4-related cardiovascular conditions observed in males within Cluster 2, who predominantly belong to the White population.

Our findings hypothesized that male AD patients in Cluster 2 may represent a distinct subset characterized by more pronounced vascular and metabolic dysfunction, potentially influenced by the complex interplay of *APOE* genotypes and ethnic background. However, due to the absence of *APOE* genomic information in our dataset, this hypothesis remains speculative and requires further investigation to confirm its validity. Future research should integrate genetic data to further explore the role of *APOE* and other genetic factors in shaping comorbidity patterns. Additionally, understanding the interplay between sex, race, *APOE* genotype, and vascular risk factors could provide valuable insights for personalized risk assessment and targeted interventions in AD patients.

In Cluster 3, our study found that calculus in the bladder was more strongly associated with females. Generally, urolithiasis, including bladder calculi, is more prevalent in males, and the most common kidney stone is calcium oxalate [75-77]. One possible explanation for this finding is the influence of sex-specific differences in microbiome alterations. Dysbiosis may lead to increased oxalate accumulation, which plays a central role in bladder stone formation [78]. Recent studies have highlighted the relationship between gut microbiome alterations, AD, and sex differences. These findings suggest that microbial diversity changes observed in AD patients may be more relevant to females, potentially contributing to

sex-specific differences in disease susceptibility and progression [52,79]. While the relationship between microbiome alteration, AD progression, and oxalate metabolism remains unclear, future research integrating microbiome data may provide new therapeutic opportunities targeting the gut microbiome to improve gastrointestinal and neurological health in AD patients [80,81].

If AD and bladder calculus are linked through microbiome dysregulation, Cluster 3 may overrepresent females in a later stage post-diagnosis. Since urinary retention is rare in females [82], symptoms may go unnoticed, leading to delayed diagnosis. Consequently, bladder calculus may only be detected once both urological and AD symptoms have advanced. However, as our study lacks data on diagnosis timing, further research incorporating disease onset and progression timelines is needed to validate this hypothesis. Such investigations may also facilitate the development of personalized therapeutic regimens tailored to disease stage and progression.

Final thoughts:

Through unsupervised learning, we identified five patient subgroups and revealed that AD-related comorbidities can be categorized into clinically meaningful subgroups: Cluster 2 was primarily associated with cardiovascular conditions, Cluster 3 with gastrointestinal disorders, and Cluster 4 with frailty-related conditions. Our findings suggest the presence of subphenotypes related to *APOE* genotypes and microbiome alterations related to AD, which may contribute to the observed differences in comorbidity patterns across sexes. Future studies incorporating genomic and microbiome data could provide deeper insights into the complex interplay between genetic susceptibility, microbiome composition, and disease progression in AD.

Currently, anti-beta amyloid therapies are nearly universal and FDA-approved, with the next step being the addition of anti-tau therapies. Our findings suggest that, following these approaches, individuals may benefit from targeted therapeutics tailored to their specific cluster profiles, such as interventions to reduce systemic inflammation, protect against urinary tract infections, or prevent gastrointestinal diseases.

Limitations of the study

There are several limitations in this study. First, the presence of imbalanced data resulted in some odds ratios being disproportionately inflated. While key findings were consistent in validation analysis, suggesting a certain level of reliability, potential biases due to sample size variation cannot be ruled out.

Second, some Pearson correlation coefficients in validation analysis were not particularly high. This finding suggests the presence of unexplored heterogeneity within both datasets. This discrepancy may partly result from demographic differences between the UCSF and UC-Wide cohorts, which could have influenced the distribution of AD severity and comorbidity patterns.

Third, the potential for misdiagnosis of AD remains a concern, as our study did not incorporate biomarkers for confirmation, which may have introduced diagnostic inaccuracies.

Fourth, data quality poses another limitation, as EMRs are primarily maintained for clinical rather than research purposes, leading to missing values and potential inconsistencies. In addition, age data was not consistently recorded across datasets, with UCSF capping reported ages at 91 and UC-Wide data capping at 90, which may have impacted age-related analyses.

Fifth, the case-control study design precludes establishing causal relationships or determining the temporal sequence of disease progression. Regarding generalizability, our analysis was limited to medical institutions in California, necessitating caution when extrapolating these findings to other regions or populations.

Finally, while our study primarily focused on LOAD, the lack of EOAD analysis constrains the understanding of AD heterogeneity. Future research should extend the investigation to include EOAD populations to provide a more comprehensive understanding of AD subphenotypes.

Despite these limitations, our research provides a comprehensive, robust, analysis of AD comorbidities, identifying distinct subphenotypes that highlight the heterogeneity in disease manifestation.

Resource availability

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Marina Sirota (marina.sirota@ucsf.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Ethical review and approval were waived for this study because it did not meet the definition of human participants research, and obtaining informed consent would not be possible or necessary. The datasets from UCHDW and UCDDP were fully de-identified prior to analysis, and the requirement for written informed consent was waived by the respective institutions in compliance with applicable ethical guidelines. The data supporting the findings of this study are not publicly available due to their sensitive nature. Access is restricted to UCSF-affiliated individuals and approved collaborators. Individuals not affiliated with UCSF may establish an official collaboration with a UCSF-affiliated investigator by contacting the principal investigator, Marina Sirota (marina.sirota@ucsf.edu). Requests for collaboration are typically processed within a few weeks. UCSF-affiliated individuals seeking access to the UCSF EMR database may contact UCSF's Clinical and Translational Science Institute (ctsi@ucsf.edu) or the UCSF Information Commons team (Info.Commons@ucsf.edu) for more details. The UCDDP database is accessible only to UC researchers who have completed analyses within their respective UC institutions and have provided a justified rationale for scaling their analyses across multiple UC health centers. Censored source data for phenotype comorbidity enrichment analysis used to create Figure 3 and 6 are provided in Supporting Data.

The code corresponding to this research is available at https://github.com/yukari-katsuhara/SexSubphenotypes_AD.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request

Acknowledgements

This research was primarily supported by grants from the National Institute on Aging (NIA) (R01AG060393 and R01AG057683). The authors sincerely thank John Kornak, Thomas Hoffmann, Christopher Seaman, and all members of the Sirota Lab for their insightful discussions and valuable guidance throughout the preparation of this manuscript. The authors also thank the Center for Data-driven Insights and Innovation at UC Health (CDI2; https://www.ucop.edu/uc-health/functions/center-for-data-driven-insights-and-innovations-cdi2.html), for its analytical and technical support related to use of the UC Health Data Warehouse and related data assets, including the UC Data Discovery Platform (UCDDP).

Author contributions

Y.K., U.K., T.T.O., A.S.T., and M.S. designed research studies, conducted experiments, and acquired data. Y.K. analyzed data and wrote the manuscript. A.S.T. mentored Y.K. All authors interpreted results, edited and reviewed the manuscript.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Document S1. Figures S1–S3 and Tables S1 and S2

Table S3. Excel file containing additional data too large to fit in a PDF, related to Figures 3 and 6.

STAR Methods

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Deposited data		
Original Code	This paper	https://github.com/yukari-katsuhara/SexSubphenotypes_AD/tree/main
Software and algorithms		
Python version 3.7.9	Python Software Foundation	https://www.python.org/
sklearn.decomposition.PCA	scikit-learn	https://scikit-learn.org/stable/modules/generated/sklearn.decomposition.PCA.html
sklearn.cluster.KMeans	scikit-learn	https://scikit-learn.org/stable/modules/generated/sklearn.cluster.KMeans.html
sklearn.preprocessing.OneHotEncoder	scikit-learn	https://scikit-learn.org/stable/modules/generated/sklearn.preprocessing.OneHotEncoder.html
sklearn.preprocessing.StandardScaler	scikit-learn	https://scikit-learn.org/stable/modules/generated/sklearn.preprocessing.StandardScaler.html
umap	umap	https://umap-learn.readthedocs.io/en/latest/index.html
scipy	Python Package Index (PyPI)	https://scipy.org/
statsmodels.api	Python Package Index (PyPI)	https://www.statsmodels.org/stable/index.html
upsetplot.UpSet	Python Package Index (PyPI)	https://upsetplot.readthedocs.io/
R version 4.3.1	R Development Core Team	https://www.r-project.org/
tidyverse	R Development Core Team	https://www.tidyverse.org/
gt	R Development Core Team	https://gt.rstudio.com/
gtsummary	R Development Core Team	https://www.danieldsjoberg.com/gtsummary/

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Cohort identification

All clinical data used in this study were obtained from the University of California's Health Data Warehouse (UCHDW) and the University of California Data Discovery Platform (UCDDP). This study identified subgroups of AD patients from UCSF EMR and validated the results using independent UC-Wide data from UCDDP to ensure the robustness of the findings.

AD patients were identified from over five million records in the UCSF EMR dataset, which contains data from 1982 to 2020. Due to the de-identification process, all dates were shifted by up to one year while preserving relative dates. Additionally, birth dates before 1930 were adjusted to 1930, effectively treating all individuals born before 1930 as having an estimated age of 90 years. Patients were included in the study if they met the inclusion criteria of an estimated age above 64 years and a diagnosis of AD based on the presence of the ICD-10-CM codes G30.1, G30.8, or G30.9. We focused on LOAD because it is a highly polygenic disorder, whereas EOAD is thought to have a distinct etiology and a stronger genetic predisposition [40]. Additionally, only patients with at least one documented comorbidity in addition to AD were included in the analysis. Sex was determined based on the most recent sex assignment recorded in the electronic medical record. UCDDP includes the de-identified records from UCSF, UC Los Angeles (UCLA), UC San Diego (UCSD), UC Davis (UCD), UC Irvine (UCI), and UC Riverside (UCR). As in the UCSF UCHDW dataset, the de-identification process adjusted the estimated age of all individuals 91 years or older to 91 years. The validation cohort was identified using the same criteria from UCDDP, excluding UCSF records, and consisted of UC-Wide clinical data from 2012 to 2024.

Sex as a biological variable

Our study examined both males and females using stratified analyses to identify sex differences in AD subphenotypes. By conducting these analyses, we aimed to clarify potential sex-specific variations in disease characteristics.

METHOD DETAILS

Dimensionality reduction and K-means clustering

To represent de-identified AD patients, one-hot encoding was performed for all diagnoses, excluding those explicitly related to Alzheimer's disease. Patients with no recorded comorbidities other than AD were excluded from the clustering analysis. PCA was used to reduce the dimensionality of the dataset while preserving diagnostic information, with the number of components selected to cover at least 80% of the cumulative variance. K-means clustering was then applied to these PCA components, and the optimal number of clusters was determined based on a combination of silhouette scores and WSS reduction analysis. To visualize the clustering results, patient distributions were projected onto a lower-dimensional space using UMAP.

Comorbidity enrichment analysis for overall trends of each cluster

To examine overall trends within each cluster, enrichment analyses were conducted by comparing the frequency of diagnoses using ICD-10-CM codes between each cluster and all other clusters combined. For each diagnosis, the proportion of patients in the target cluster was compared to the proportion in the other clusters using Fisher's exact test when the sample size was below five or the chi-squared test when the sample size was larger. Statistical significance was determined using a Bonferroni-corrected threshold of p-value < 0.05, and the directionality of associations was determined based on odds ratios. The statistical significance of associations was visualized using volcano plots, with effect sizes and magnitudes, while Manhattan plots (for the overall population) and Miami plots (for sex-stratified populations) illustrated the distribution of p-values across diagnoses. Sex-stratified analyses were performed to assess whether comorbidity patterns differed between male and female patients.

Comorbidity enrichment analysis for cluster- or cluster-sex-specific comorbidities

Cluster-specific comorbidities were identified by visualizing significant comorbidities using UpSet plots [83]. The statistical significance of these comorbidities was displayed in Manhattan plots for the overall population and Miami plots for sex-stratified populations. To further explore the phenotypic characteristics of each cluster, a UMAP visualization was generated using standardized odds ratios of cluster-specific comorbidities. Each dot represents a specific ICD-10-CM code, with its size corresponding to the odds ratio of the significantly associated cluster and its color indicating the respective cluster. The input features for the UMAP projection were derived from a matrix of standardized odds ratios in each cluster, where each row represents an ICD-10-CM code and each column corresponds to a cluster. Sex-stratified analyses were conducted to further investigate differences in comorbidity patterns between males and females.

Sensitivity analysis

To validate the robustness of the clustering and comorbidity enrichment results, the same approach used for the UCSF dataset was applied to the UC-Wide dataset. AD patients were identified using the same inclusion criteria, and dimensionality reduction and K-means clustering were performed as described previously. Due to the de-identification process, all estimated ages above 91 were recorded as 91 years. Cluster-specific comorbidities in the UC-Wide dataset were analyzed using the same statistical methods and visualized with UpSet plots. The relationships between clusters in the UCSF and UC-Wide datasets were examined by comparing cluster-specific comorbidities, and the degree of overlap was assessed using Sankey plots. The coverage rate for UCSF clusters was calculated as the proportion of UCSF cluster-specific comorbidities that were statistically significant in UCSF and remained significant in UC-Wide clusters. The UC-Wide clusters with the highest coverage rates were identified as the best matches for each UCSF cluster. Log-log plots were used to compare the odds ratios of matched clusters between the UCSF and UC-Wide datasets for the overall population and sex-stratified subgroups. Pearson correlation

analysis was performed to assess the linear relationship between the odds ratios, and the correlation coefficient and p-value were obtained to evaluate the statistical significance of the association.

QUANTIFICATION AND STATISTICAL ANALYSIS

All statistical and computational analyses were conducted using Python, except for demographic table generation, which was performed in R. In the demographic table analysis, categorical variables such as sex, race, and death status were compared across clusters using the Chi-squared test, while continuous variables such as age and the number of comorbidities were analyzed using the Kruskal-Wallis rank sum test. When the Kruskal-Wallis test indicated a significant difference among clusters, post-hoc pairwise comparisons were conducted using Dunn's test with Bonferroni correction to identify which specific groups differed significantly.

For comorbidity enrichment analysis, the proportions of patients in each ICD-10-CM code were compared using either Fisher's exact test, if fewer than five patients were present in a given ICD-10-CM code, or the Chi-squared test otherwise. Diagnoses were considered significant if the p-value was below 0.05 after Bonferroni correction, and the directionality of the association was determined using the OR.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT-4o (released on May 13, 2024, by OpenAI) in order to refine language and enhance readability. After using this tool or service, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

1. Jack, C.R., Jr., Bennett, D.A., Blennow, K., Carrillo, M.C., Dunn, B., Haeberlein, S.B., Holtzman, D.M., Jagust, W., Jessen, F., Karlawish, J., et al. (2018). NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. Alzheimers Dement *14*, 535-562. 10.1016/j.jalz.2018.02.018.
2. Uddin, M.S., and Lim, L.W. (2022). Glial cells in Alzheimer's disease: From neuropathological changes to therapeutic implications. Ageing Research Reviews *78*, 101622. https://doi.org/10.1016/j.arr.2022.101622.

3.   Hersi, M., Irvine, B., Gupta, P., Gomes, J., Birkett, N., and Krewski, D. (2017). Risk factors associated with the onset and progression of Alzheimer's disease: A systematic review of the evidence. Neurotoxicology *61*, 143-187. 10.1016/j.neuro.2017.03.006.

4.   Liu, C.C., Liu, C.C., Kanekiyo, T., Xu, H., and Bu, G. (2013). Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. Nat Rev Neurol *9*, 106-118. 10.1038/nrneurol.2012.263.

5.   Miller, Z.A., Ossenkoppele, R., Graff-Radford, N.R., Allen, I.E., Shwe, W., Rosenberg, L., Olguin, D.J., Erkkinen, M.G., Butler, P.M., Spina, S., et al. (2022). Left-handedness, learning disability, autoimmune disease, and seizure history influence age at onset and phenotypical targeting of Alzheimer's disease. medRxiv, 2022.2012.2017.22283307. 10.1101/2022.12.17.22283307.

6.   2024 Alzheimer's disease facts and figures. (2024). Alzheimers Dement *20*, 3708-3821. 10.1002/alz.13809.

7.   Irvine, K., Laws, K.R., Gale, T.M., and Kondel, T.K. (2012). Greater cognitive deterioration in women than men with Alzheimer's disease: a meta analysis. J Clin Exp Neuropsychol *34*, 989-998. 10.1080/13803395.2012.712676.

8.   Ferretti, M.T., Iulita, M.F., Cavedo, E., Chiesa, P.A., Schumacher Dimech, A., Santuccione Chadha, A., Baracchi, F., Girouard, H., Misoch, S., Giacobini, E., et al. (2018). Sex differences in Alzheimer disease — the gateway to precision medicine. Nature Reviews Neurology *14*, 457-469. 10.1038/s41582-018-0032-9.

9.   Aggarwal, N.T., and Mielke, M.M. (2023). Sex Differences in Alzheimer's Disease. Neurol Clin *41*, 343-358. 10.1016/j.ncl.2023.01.001.

10.   Gilsanz, P., Mayeda, E.R., Glymour, M.M., Quesenberry, C.P., Mungas, D.M., DeCarli, C., Dean, A., and Whitmer, R.A. (2017). Female sex, early-onset hypertension, and risk of dementia. Neurology *89*, 1886-1893. doi:10.1212/WNL.0000000000004602.

11.   Rogalski, E.J., Rademaker, A., Harrison, T.M., Helenowski, I., Johnson, N., Bigio, E., Mishra, M., Weintraub, S., and Mesulam, M.M. (2011). ApoE E4 is a susceptibility factor in amnestic but not aphasic dementias. Alzheimer Dis Assoc Disord *25*, 159-163. 10.1097/WAD.0b013e318201f249.

12.   Arnold, M., Nho, K., Kueider-Paisley, A., Massaro, T., Huynh, K., Brauner, B., MahmoudianDehkordi, S., Louie, G., Moseley, M.A., Thompson, J.W., et al. (2020). Sex and APOE ε4 genotype modify the Alzheimer's disease serum metabolome. Nat Commun *11*, 1148. 10.1038/s41467-020-14959-w.

13.   Larson, E.B., Shadlen, M.F., Wang, L., McCormick, W.C., Bowen, J.D., Teri, L., and Kukull, W.A. (2004). Survival after initial diagnosis of Alzheimer disease. Ann Intern Med *140*, 501-509. 10.7326/0003-4819-140-7-200404060-00008.

14.   Xie, J., Brayne, C., and Matthews, F.E. (2008). Survival times in people with dementia: analysis from population based cohort study with 14 year follow-up. BMJ *336*, 258-262. 10.1136/bmj.39433.616678.25.

15.   Santiago, J.A., and Potashkin, J.A. (2023). Biological and Clinical Implications of Sex-Specific Differences in Alzheimer's Disease. Handb Exp Pharmacol *282*, 181-197. 10.1007/164_2023_672.

16.   Geifman, N., Kennedy, R.E., Schneider, L.S., Buchan, I., and Brinton, R.D. (2018). Data-driven identification of endophenotypes of Alzheimer's disease progression: implications for clinical trials and therapeutic interventions. Alzheimer's Research & Therapy *10*, 4. 10.1186/s13195-017-0332-0.

17.   Xu, J., Wang, F., Xu, Z., Adekkanattu, P., Brandt, P., Jiang, G., Kiefer, R.C., Luo, Y., Mao, C., Pacheco, J.A., et al. (2020). Data-driven discovery of probable Alzheimer's disease and related dementia subphenotypes using electronic health records. Learn Health Syst *4*, e10246. 10.1002/lrh2.10246.

18. Tang, A.S., Oskotsky, T., Havaldar, S., Mantyh, W.G., Bicak, M., Solsberg, C.W., Woldemariam, S., Zeng, B., Hu, Z., Oskotsky, B., et al. (2022). Deep phenotyping of Alzheimer's disease leveraging electronic medical records identifies sex-specific clinical associations. Nature Communications *13*, 675. 10.1038/s41467-022-28273-0.

19. Woldemariam, S.R., Tang, A.S., Oskotsky, T.T., Yaffe, K., and Sirota, M. (2023). Similarities and differences in Alzheimer's dementia comorbidities in racialized populations identified from electronic medical records. Communications Medicine *3*, 50. 10.1038/s43856-023-00280-2.

20. Meng, W., Xu, J., Huang, Y., Wang, C., Song, Q., Ma, A., Song, L., Bian, J., Ma, Q., and Yin, R. (2024). Autoencoder to Identify Sex-Specific Sub-phenotypes in Alzheimer's Disease Progression Using Longitudinal Electronic Health Records. medRxiv. 10.1101/2024.07.07.24310055.

21. McAdams-DeMarco, M.A., Bae, S., Chu, N., Gross, A.L., Brown, C.H.t., Oh, E., Rosenberg, P., Neufeld, K.J., Varadhan, R., Albert, M., et al. (2017). Dementia and Alzheimer's Disease among Older Kidney Transplant Recipients. J Am Soc Nephrol *28*, 1575-1583. 10.1681/asn.2016080816.

22. Ruck, J.M., Chu, N.M., Liu, Y., Li, Y., Chen, Y., Mathur, A., Carlson, M.C., Crews, D.C., Chodosh, J., Segev, D.L., and McAdams-DeMarco, M. (2024). Association of Postoperative Delirium With Incident Dementia and Graft Outcomes Among Kidney Transplant Recipients. Transplantation *108*, 530-538. 10.1097/tp.0000000000004779.

23. Evans, D.A., Bennett, D.A., Wilson, R.S., Bienias, J.L., Morris, M.C., Scherr, P.A., Hebert, L.E., Aggarwal, N., Beckett, L.A., Joglekar, R., et al. (2003). Incidence of Alzheimer disease in a biracial urban community: relation to apolipoprotein E allele status. Arch Neurol *60*, 185-189. 10.1001/archneur.60.2.185.

24. Tang, M.X., Stern, Y., Marder, K., Bell, K., Gurland, B., Lantigua, R., Andrews, H., Feng, L., Tycko, B., and Mayeux, R. (1998). The APOE-epsilon4 allele and the risk of Alzheimer disease among African Americans, whites, and Hispanics. Jama *279*, 751-755. 10.1001/jama.279.10.751.

25. Weuve, J., Barnes, L.L., Mendes de Leon, C.F., Rajan, K.B., Beck, T., Aggarwal, N.T., Hebert, L.E., Bennett, D.A., Wilson, R.S., and Evans, D.A. (2018). Cognitive Aging in Black and White Americans: Cognition, Cognitive Decline, and Incidence of Alzheimer Disease Dementia. Epidemiology *29*, 151-159. 10.1097/ede.0000000000000747.

26. Nasr, H., and Scriven, J.M. (2015). Superficial thrombophlebitis (superficial venous thrombosis). Bmj *350*, h2039. 10.1136/bmj.h2039.

27. Décousus, H., Bertoletti, L., and Frappé, P. (2015). Spontaneous acute superficial vein thrombosis of the legs: do we really need to treat? J Thromb Haemost *13 Suppl 1*, S230-237. 10.1111/jth.12925.

28. Di Minno, M.N., Ambrosino, P., Ambrosini, F., Tremoli, E., Di Minno, G., and Dentali, F. (2016). Prevalence of deep vein thrombosis and pulmonary embolism in patients with superficial vein thrombosis: a systematic review and meta-analysis. J Thromb Haemost *14*, 964-972. 10.1111/jth.13279.

29. Simões, A.M.N., Vendramim, P., and Pedreira, M.L.G. (2022). Risk factors for peripheral intravenous catheter-related phlebitis in adult patients. Rev Esc Enferm USP *56*, e20210398. 10.1590/1980-220X-REEUSP-2021-0398en.

30. Undas, A. (2017). Prothrombotic Fibrin Clot Phenotype in Patients with Deep Vein Thrombosis and Pulmonary Embolism: A New Risk Factor for Recurrence. Biomed Res Int *2017*, 8196256. 10.1155/2017/8196256.

31. Czysz A, H.S. (2023). Superficial Thrombophlebitis. StatPearls [Internet].

32. Cortes-Canteli, M., Zamolodchikov, D., Ahn, H.J., Strickland, S., and Norris, E.H. (2012). Fibrinogen and altered hemostasis in Alzheimer's disease. J Alzheimers Dis *32*, 599-608. 10.3233/jad-2012-120820.

33. Cortes-Canteli, M., Paul, J., Norris, E.H., Bronstein, R., Ahn, H.J., Zamolodchikov, D., Bhuvanendran, S., Fenz, K.M., and Strickland, S. (2010). Fibrinogen and beta-amyloid association alters thrombosis and fibrinolysis: a possible contributing factor to Alzheimer's disease. Neuron *66*, 695-709. 10.1016/j.neuron.2010.05.014.

34. Fibrin induces neurotoxic microglia gene programs in neurodegeneration. (2023). Nature Immunology *24*, 1062-1063. 10.1038/s41590-023-01542-w.

35. Merlini, M., Rafalski, V.A., Rios Coronado, P.E., Gill, T.M., Ellisman, M., Muthukumar, G., Subramanian, K.S., Ryu, J.K., Syme, C.A., Davalos, D., et al. (2019). Fibrinogen Induces Microglia-Mediated Spine Elimination and Cognitive Impairment in an Alzheimer's Disease Model. Neuron *101*, 1099-1108.e1096. 10.1016/j.neuron.2019.01.014.

36. Ryu, J.K., Rafalski, V.A., Meyer-Franke, A., Adams, R.A., Poda, S.B., Rios Coronado, P.E., Pedersen, L., Menon, V., Baeten, K.M., Sikorski, S.L., et al. (2018). Fibrin-targeting immunotherapy protects against neuroinflammation and neurodegeneration. Nat Immunol *19*, 1212-1223. 10.1038/s41590-018-0232-x.

37. Bergamaschini, L., Rossi, E., Storini, C., Pizzimenti, S., Distaso, M., Perego, C., De Luigi, A., Vergani, C., and De Simoni, M.G. (2004). Peripheral treatment with enoxaparin, a low molecular weight heparin, reduces plaques and beta-amyloid accumulation in a mouse model of Alzheimer's disease. J Neurosci *24*, 4181-4186. 10.1523/jneurosci.0550-04.2004.

38. Avitan, I., Halperin, Y., Saha, T., Bloch, N., Atrahimovich, D., Polis, B., Samson, A.O., and Braitbard, O. (2021). Towards a Consensus on Alzheimer's Disease Comorbidity? J Clin Med *10*. 10.3390/jcm10194360.

39. Nedelec, T., Couvy-Duchesne, B., Monnet, F., Daly, T., Ansart, M., Gantzer, L., Lekens, B., Epelbaum, S., Dufouil, C., and Durrleman, S. (2022). Identifying health conditions associated with Alzheimer's disease up to 15 years before diagnosis: an agnostic study of French and British health records. The Lancet Digital Health *4*, e169-e178. https://doi.org/10.1016/S2589-7500(21)00275-2.

40. Wingo, T.S., Lah, J.J., Levey, A.I., and Cutler, D.J. (2012). Autosomal recessive causes likely in early-onset Alzheimer disease. Arch Neurol *69*, 59-64. 10.1001/archneurol.2011.221.

41. Sirkis, D.W., Bonham, L.W., Johnson, T.P., La Joie, R., and Yokoyama, J.S. (2022). Dissecting the clinical heterogeneity of early-onset Alzheimer's disease. Mol Psychiatry *27*, 2674-2688. 10.1038/s41380-022-01531-9.

42. Mantyh, W.G., Cochran, J.N., Taylor, J.W., Broce, I.J., Geier, E.G., Bonham, L.W., Anderson, A.G., Sirkis, D.W., Joie, R., Iaccarino, L., et al. (2023). Early-onset Alzheimer's disease explained by polygenic risk of late-onset disease? Alzheimers Dement (Amst) *15*, e12482. 10.1002/dad2.12482.

43. Falgàs, N., Allen, I.E., Spina, S., Grant, H., Piña Escudero, S.D., Merrilees, J., Gearhart, R., Rosen, H.J., Kramer, J.H., Seeley, W.W., et al. (2022). The severity of neuropsychiatric symptoms is higher in early-onset than late-onset Alzheimer's disease. Eur J Neurol *29*, 957-967. 10.1111/ene.15203.

44. Ramey, G.D., Tang, A., Phongpreecha, T., Yang, M.M., Woldemariam, S.R., Oskotsky, T.T., Montine, T.J., Allen, I., Miller, Z.A., Aghaeepour, N., et al. (2025). Exposure to autoimmune disorders is associated with increased Alzheimer's disease risk in a multi-site electronic health record analysis. Cell Rep Med, 101980. 10.1016/j.xcrm.2025.101980.

45. Sundaresan, B., Shirafkan, F., Ripperger, K., and Rattay, K. (2023). The Role of Viral Infections in the Onset of Autoimmune Diseases. Viruses *15*. 10.3390/v15030782.

46. Liao, L. (2015). Evaluation and Management of Neurogenic Bladder: What Is New in China? Int J Mol Sci *16*, 18580-18600. 10.3390/ijms160818580.

47. Gómez-Gómez, M.E., and Zapico, S.C. (2019). Frailty, Cognitive Decline, Neurodegenerative Diseases and Nutrition Interventions. Int J Mol Sci *20*. 10.3390/ijms20112842.

48. Makhnevich, A., Perrin, A., Talukder, D., Liu, Y., Izard, S., Chiuzan, C., D'Angelo, S., Affoo, R., Rogus-Pulia, N., and Sinvani, L. (2024). Thick Liquids and Clinical Outcomes in Hospitalized Patients With Alzheimer Disease and Related Dementias and Dysphagia. JAMA Internal Medicine *184*, 778-785. 10.1001/jamainternmed.2024.0736.

49. Yurtdaş Depboylu, G., Acar Tek, N., Akbulut, G., Günel, Z., and Kamanlı, B. (2023). Functional Constipation in Elderly and Related Determinant Risk Factors: Malnutrition and Dietary Intake. J Am Nutr Assoc *42*, 541-547. 10.1080/27697061.2022.2096150.

50. Seo, D.O., and Holtzman, D.M. (2024). Current understanding of the Alzheimer's disease-associated microbiome and therapeutic strategies. Exp Mol Med *56*, 86-94. 10.1038/s12276-023-01146-2.

51. Kuźniar, J., Kozubek, P., Czaja, M., and Leszek, J. (2024). Correlation between Alzheimer's Disease and Gastrointestinal Tract Disorders. Nutrients *16*. 10.3390/nu16142366.

52. Vogt, N.M., Kerby, R.L., Dill-McFarland, K.A., Harding, S.J., Merluzzi, A.P., Johnson, S.C., Carlsson, C.M., Asthana, S., Zetterberg, H., Blennow, K., et al. (2017). Gut microbiome alterations in Alzheimer's disease. Sci Rep *7*, 13537. 10.1038/s41598-017-13601-y.

53. Cunha, B.A. (2001). Pneumonia in the elderly. Clin Microbiol Infect *7*, 581-588. 10.1046/j.1198-743x.2001.00328.x.

54. Whitson, H.E., Colton, C., El Khoury, J., Gate, D., Goate, A., Heneka, M.T., Kaddurah-Daouk, R., Klein, R.S., Shinohara, M.L., Sisodia, S., et al. (2022). Infection and inflammation: New perspectives on Alzheimer's disease. Brain Behav Immun Health *22*, 100462. 10.1016/j.bbih.2022.100462.

55. Lehrer, S., and Rheinstein, P.H. (2022). Circulating Chlamydia Trachomatis Antigens in Subjects With Alzheimer's Disease. In Vivo *36*, 2650-2653. 10.21873/invivo.12999.

56. Lehrer, S., and Rheinstein, P.H. (2024). RPTOR Is an Alzheimer's Disease Susceptibility Gene Associated with the Risk Factors Body Mass Index and Infectious Encephalitis. J Alzheimers Dis Rep *8*, 715-721. 10.3233/adr-230185.

57. Ukraintseva, S., Duan, M., Simanek, A.M., Holmes, R., Bagley, O., Rajendrakumar, A.L., Yashkin, A.P., Akushevich, I., Tropsha, A., Whitson, H., et al. (2023). Vaccination Against Pneumonia May Provide Genotype-Specific Protection Against Alzheimer's Disease. J Alzheimers Dis *96*, 499-505. 10.3233/jad-230088.

58. Lehrer, S., and Rheinstein, P.H. (2022). Vaccination Reduces Risk of Alzheimer's Disease, Parkinson's Disease and Other Neurodegenerative Disorders. Discov Med *34*, 97-101.

59. Iwai-Saito, K., Sato, K., Fujii, M., and Kondo, K. (2024). Pneumococcal vaccination, but not influenza vaccination, is negatively associated with incident dementia among Japanese older adults: The JAGES 2013–2022 prospective cohort study. Brain, Behavior, and Immunity *120*, 452-463. https://doi.org/10.1016/j.bbi.2024.06.020.

60. Huo, X., and Finkelstein, J. (2024). Pneumococcal Vaccination Lowers the Risk of Alzheimer's Disease: A Study Utilizing Data from the IBM® MarketScan® Database. Stud Health Technol Inform *310*, 961-965. 10.3233/shti231107.

61. Perry, D.C., Brown, J.A., Possin, K.L., Datta, S., Trujillo, A., Radke, A., Karydas, A., Kornak, J., Sias, A.C., Rabinovici, G.D., et al. (2017). Clinicopathological correlations in behavioural variant frontotemporal dementia. Brain *140*, 3329-3345. 10.1093/brain/awx254.

62. Musa, G., Slachevsky, A., Muñoz-Neira, C., Méndez-Orellana, C., Villagra, R., González-Billault, C., Ibáñez, A., Hornberger, M., and Lillo, P. (2020). Alzheimer's

Disease or Behavioral Variant Frontotemporal Dementia? Review of Key Points Toward an Accurate Clinical and Neuropsychological Diagnosis. J Alzheimers Dis *73*, 833-848. 10.3233/jad-190924.

63. Wagner, M., Lorenz, G., Volk, A.E., Brunet, T., Edbauer, D., Berutti, R., Zhao, C., Anderl-Straub, S., Bertram, L., Danek, A., et al. (2021). Clinico-genetic findings in 509 frontotemporal dementia patients. Molecular Psychiatry *26*, 5824-5832. 10.1038/s41380-021-01271-2.

64. Ranasinghe, K.G., Rankin, K.P., Pressman, P.S., Perry, D.C., Lobach, I.V., Seeley, W.W., Coppola, G., Karydas, A.M., Grinberg, L.T., Shany-Ur, T., et al. (2016). Distinct Subtypes of Behavioral Variant Frontotemporal Dementia Based on Patterns of Network Degeneration. JAMA Neurol *73*, 1078-1088. 10.1001/jamaneurol.2016.2016.

65. Wang, Y., Yang, P., Yan, Z., Liu, Z., Ma, Q., Zhang, Z., Wang, Y., and Su, Y. (2021). The Relationship between Erythrocytes and Diabetes Mellitus. J Diabetes Res *2021*, 6656062. 10.1155/2021/6656062.

66. Lowe, G.D. (2001). The relationship between infection, inflammation, and cardiovascular disease: an overview. Ann Periodontol *6*, 1-8. 10.1902/annals.2001.6.1.1.

67. Pankratz, V.S., Roberts, R.O., Mielke, M.M., Knopman, D.S., Jack, C.R., Geda, Y.E., Rocca, W.A., and Petersen, R.C. (2015). Predicting the risk of mild cognitive impairment in the Mayo Clinic Study of Aging. Neurology *84*, 1433-1442. doi:10.1212/WNL.0000000000001437.

68. Gong, J., Harris, K., Peters, S.A.E., and Woodward, M. (2021). Sex differences in the association between major cardiovascular risk factors in midlife and dementia: a cohort study using data from the UK Biobank. BMC Med *19*, 110. 10.1186/s12916-021-01980-z.

69. Chatterjee, S., Peters, S.A.E., Woodward, M., Mejia Arango, S., Batty, G.D., Beckett, N., Beiser, A., Borenstein, A.R., Crane, P.K., Haan, M., et al. (2015). Type 2 Diabetes as a Risk Factor for Dementia in Women Compared With Men: A Pooled Analysis of 2.3 Million People Comprising More Than 100,000 Cases of Dementia. Diabetes Care *39*, 300-307. 10.2337/dc15-1588.

70. Anthopoulos, P.G., Hamodrakas, S.J., and Bagos, P.G. (2010). Apolipoprotein E polymorphisms and type 2 diabetes: a meta-analysis of 30 studies including 5423 cases and 8197 controls. Mol Genet Metab *100*, 283-291. 10.1016/j.ymgme.2010.03.008.

71. Shi, J., Liu, Y., Liu, Y., Li, Y., Qiu, S., Bai, Y., Gu, Y., Luo, J., Cui, H., Li, Y., et al. (2018). Association between ApoE polymorphism and hypertension: A meta-analysis of 28 studies including 5898 cases and 7518 controls. Gene *675*, 197-207. 10.1016/j.gene.2018.06.097.

72. Bennet, A.M., Di Angelantonio, E., Ye, Z., Wensley, F., Dahlin, A., Ahlbom, A., Keavney, B., Collins, R., Wiman, B., de Faire, U., and Danesh, J. (2007). Association of apolipoprotein E genotypes with lipid levels and coronary risk. Jama *298*, 1300-1311. 10.1001/jama.298.11.1300.

73. Riedel, B.C., Thompson, P.M., and Brinton, R.D. (2016). Age, APOE and sex: Triad of risk of Alzheimer's disease. The Journal of Steroid Biochemistry and Molecular Biology *160*, 134-147. https://doi.org/10.1016/j.jsbmb.2016.03.012.

74. Maestre, G., Ottman, R., Stern, Y., Gurland, B., Chun, M., Tang, M.X., Shelanski, M., Tycko, B., and Mayeux, R. (1995). Apolipoprotein E and Alzheimer's disease: ethnic variation in genotypic risks. Ann Neurol *37*, 254-259. 10.1002/ana.410370217.

75. Li, M.L., Song, S.C., Yang, F., Gao, C., Zhou, B., and Wang, Q. (2024). Risk assessment and prevention of urolithiasis in urban areas of Baoding, China. Medicine (Baltimore) *103*, e35880. 10.1097/md.0000000000035880.

76. Stamatelou, K., and Goldfarb, D.S. (2023). Epidemiology of Kidney Stones. Healthcare (Basel) *11*. 10.3390/healthcare11030424.

77. Siener, R., Herwig, H., Rüdy, J., Schaefer, R.M., Lossin, P., and Hesse, A. (2022). Urinary stone composition in Germany: results from 45,783 stone analyses. World J Urol *40*, 1813-1820. 10.1007/s00345-022-04060-w.

78. Miller, A.W., Penniston, K.L., Fitzpatrick, K., Agudelo, J., Tasian, G., and Lange, D. (2022). Mechanisms of the intestinal and urinary microbiome in kidney stone disease. Nat Rev Urol *19*, 695-707. 10.1038/s41585-022-00647-5.

79. Dodiya, H.B., Lutz, H.L., Weigle, I.Q., Patel, P., Michalkiewicz, J., Roman-Santiago, C.J., Zhang, C.M., Liang, Y., Srinath, A., Zhang, X., et al. (2022). Gut microbiota-driven brain Aβ amyloidosis in mice requires microglia. J Exp Med *219*. 10.1084/jem.20200895.

80. Bashir, B., Gulati, M., Vishwas, S., Gupta, G., Dhanasekaran, M., Paudel, K.R., Chellappan, D.K., Anand, K., Negi, P., Singh, P.K., et al. (2025). Bridging gap in the treatment of Alzheimer's disease via postbiotics: Current practices and future prospects. Ageing Res Rev *105*, 102689. 10.1016/j.arr.2025.102689.

81. Jiang, X., Zheng, Y., Sun, H., Dang, Y., Yin, M., Xiao, M., and Wu, T. (2025). Fecal Microbiota Transplantation Improves Cognitive Function of a Mouse Model of Alzheimer's Disease. CNS Neurosci Ther *31*, e70259. 10.1111/cns.70259.

82. Mevcha, A., and Drake, M.J. (2010). Etiology and management of urinary retention in women. Indian J Urol *26*, 230-235. 10.4103/0970-1591.65396.

83. Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R., and Pfister, H. (2014). UpSet: Visualization of Intersecting Sets. IEEE Trans Vis Comput Graph *20*, 1983-1992. 10.1109/tvcg.2014.2346248.

	Cluster 1 N = 328		Cluster 2 N = 538		Cluster 3 N = 815		Cluster 4 N = 1,521		Cluster 5 N = 5,161		Overall N = 8,363		p-val
Sex, *n* (%)													< 0.05
Female	217	(66%)	358	(67%)	514	(63%)	1,083	(71%)	3,143	(61%)	5,315	(64%)	
Male	110	(34%)	180	(33%)	301	(37%)	438	(29%)	2,000	(39%)	3,029	(36%)	
Unknown	<10	(-)	<10	(-)	<10	(-)	<10	(-)	18	(0.3%)	19	(0.2%)	
Race, *n* (%)													< 0.05
White or Caucasian	148	(45%)	248	(46%)	439	(54%)	897	(59%)	3,431	(66%)	5,163	(62%)	
Other/Unknown	45	(14%)	98	(18%)	75	(9.2%)	205	(13%)	886	(17%)	1,309	(16%)	
Asian	108	(33%)	90	(17%)	222	(27%)	109	(7.2%)	347	(6.7%)	876	(10%)	
Black or African American	27	(8.2%)	70	(13%)	65	(8.0%)	180	(12%)	240	(4.7%)	582	(7.0%)	
Native Hawaiian or Other Pacific Islander	<10	(-)	32	(5.9%)	14	(1.7%)	130	(8.5%)	257	(5.0%)	433	(5.2%)	
Death Status, *n* (%)													< 0.05
Alive	218	(66%)	258	(48%)	550	(67%)	912	(60%)	4,342	(84%)	6,280	(75%)	
Deceased	110	(34%)	280	(52%)	265	(33%)	609	(40%)	819	(16%)	2,083	(25%)	
Age (yr), Median (IQR)	89.5	(84.0-91.0)	90.0	(90.0-91.0)	88.0	(82.0-90.0)	91.0	(90.0-91.0)	90.0	(81.0-91.0)	90.0	(84.0-91.0)	< 0.05
Comorbidities, Median (IQR)	313.5	(258.0-379.0)	132.0	(106.0-175.5)	114.0	(80.0-156.0)	47.0	(33.0-65.0)	12.0	(5.0-21.0)	23.0	(9.0-63.0)	< 0.05

Table 1. Patient demographics in UCSF. Medians and interquartile ranges are presented as median (25th percentile-75th percentile). "Comorbidities" refers to "Number of comorbidities," which represents the total count of diagnoses per patient. The category "Unknown" in Sex includes records classified as "Unknown." The category "Other/Unknown" in Race includes records classified as "American Indian or Alaska Native," "Other," "0," "Unknown/Declined," "Unknown," and "Declined." P-values were calculated using Pearson's Chi-squared test for Sex, Race, Death Status, and Location, and Kruskal-Wallis rank sum test for Age and Comorbidities. p-val, p-value.

	Cluster A N = 1,765		Cluster B N = 12,112		Cluster C N = 2,080		Cluster D N = 4,146		Cluster E N = 5,793		Overall N = 25,896		p-val
Sex, n (%)													< 0.05
Female	1,228	(69.6%)	7,414	(61.2%)	1,170	(56.2%)	2,370	(57.2%)	3,854	(66.5%)	16,036	(61.9%)	
Male	537	(30.4%)	4,672	(38.6%)	910	(43.8%)	1,776	(42.8%)	1,938	(33.5%)	9,833	(38.0%)	
Unknown	<10	(-)	26	(0.2%)	<10	(-)	<10	(-)	<10	(-)	27	(0.1%)	
Race, n (%)													< 0.05
White or Caucasian	1,234	(69.9%)	7,828	(64.6%)	1,322	(63.6%)	2,549	(61.5%)	4,085	(70.5%)	17,018	(65.7%)	
Other/Unknown	248	(14.1%)	2,851	(23.5%)	290	(13.9%)	728	(17.6%)	775	(13.4%)	4,892	(18.9%)	
Asian	147	(8.3%)	993	(8.2%)	271	(13.0%)	556	(13.4%)	625	(10.8%)	2,592	(10.0%)	
Black or African American	131	(7.4%)	419	(3.5%)	187	(9.0%)	287	(6.9%)	295	(5.1%)	1,319	(5.1%)	
Native Hawaiian or Other Pacific Islander	<10	(-)	21	(0.2%)	10	(0.5%)	26	(0.6%)	13	(0.2%)	785	(0.3%)	
Death Status, n (%)													< 0.05
Alive	1,221	(69.2%)	6,265	(51.7%)	717	(34.5%)	1,617	(39.0%)	3,367	(58.1%)	13,187	(50.9%)	
Deceased	544	(30.8%)	5,847	(48.3%)	1,363	(65.6%)	2,529	(61.0%)	2,426	(41.9%)	12,709	(49.1%)	
Location, n (%)													< 0.05
Site 1	320	(18.1%)	2019	(16.7%)	315	(15.1%)	590	(14.2%)	1,112	(19.2%)	4,356	(16.8%)	
Site 2	123	(7.0%)	2,629	(21.7%)	185	(8.9%)	1,032	(24.9%)	566	(9.8%)	4,535	(17.5%)	
Site 3	765	(43.3%)	4,211	(34.8%)	1,185	(57.0%)	1,395	(33.6%)	2,845	(49.1%)	10,401	(40.2%)	
Site 4	<10	(-)	85	(0.7%)	<10	(-)	<10	(-)	<10	(-)	92	(0.4%)	
Site 5	557	(31.6%)	3,168	(26.2%)	395	(19.0%)	1,128	(27.2%)	1,264	(21.8%)	6,512	(25.1%)	
Age (yr), Median (IQR)	87.0	(82.0-90.0)	88.0	(81.0-89.0)	90.0	(87.0-90.0)	90.0	(84.0-90.0)	88.0	(82.0-90.0)	89.0	(82.0-90.0)	< 0.05
Comorbidities, Median (IQR)	201.0	(166.0-253.0)	15.0	(7.0-25.0)	153.0	(127.0-185.0)	52.0	(39.0-71.0)	76.0	(57.0-100.0)	40.0	(16.0-88.0)	< 0.05

Table 2. Patient demographics in UC-Wide. Medians and interquartile ranges are presented as median (25th percentile-75th percentile). "Comorbidities" refers to "Number of comorbidities," which represents the total count of diagnoses per patient. The category "Unknown" in Sex includes records classified as "Unknown" and "*Unspecified." The category "Other/Unknown" in Race includes records classified as "American Indian or Alaska Native," "Other Race," and "Unknown." Site1 to Site5 are obfuscated representations of the following institutions: UCLA, UCSD, UCD, University of California, Irvine (UCI), and University of California, Riverside (UCR). P-values were calculated using Pearson's Chi-squared test for Sex, Race, Death Status, and Location, and Kruskal-Wallis rank sum test for Age and Comorbidities. p-val, p-value.

Figure 1. Cohort selection and overview of study design. Alzheimer's disease patients > 64 years with at least one comorbidity were included in this study. (1) K-means clustering was performed following Principal Component Analysis (PCA). (2) Each cluster was characterized based on cluster-specific comorbidities. For validation, (3) the same approach was applied to the UC-Wide cohort, excluding UCSF patients. ICD10-CM, the International Classification of Diseases, 10th Revision, Clinical Modification; PCA, Principal Component Analysis; UC, University of California

Figure 2. UMAP visualizations using PCA components as features. UMAP plots are colored by cluster (**A** and **D**), sex (**B** and **E**), and number of comorbidities (**C** and **F**). Each dot represents a patient. **A–C** correspond to the UCSF cohort, while **D–G** represent the validation cohort in the UC-Wide dataset. UC, University of California

Figure 3. Volcano plot identifies overall trends of each cluster

The volcano plot displays enriched ICD10-CM codes identified using the two-sided Fisher's exact test (for cases < 5) or the Chi-squared test (for cases ≥ 5). Enrichment was determined based on a Bonferroni-corrected p-value < 0.05. Odds ratios were calculated by comparing each specific cluster to all other clusters. **A:** Cluster 1, **B:** Cluster 2, **C:** Cluster 3, **D:** Cluster 4, **E:** Cluster 5. ICD10-CM, the International Classification of Diseases, 10th Revision, Clinical Modification.

Figure 4. UpSet plot identifies significant cluster-specific comorbidities.

UpSet plot illustrating significant cluster-specific comorbidities. Panels represent the overall population (**A:** Positive associations, **B:** Negative associations), females (**C:** Positive associations, **D:** Negative associations), and males (**E:** Positive associations, **F:** Negative associations).

Figure 5. Manhattan plot for cluster-specific significant comorbidities

Manhattan Plot displaying enriched ICD10-CM codes identified using the two-sided Fisher's exact test (for cases < 5) or the Chi-squared test (for cases ≥ 5). Enrichment was determined based on a Bonferroni-corrected p-value < 0.05. **A:** Cluster 1, **B:** Cluster 2, **C:** Cluster 3, **D:** Cluster 4, **E:** Cluster 5. ICD10-CM, the International Classification of Diseases, 10th Revision, Clinical Modification.

Figure 6. UMAP visualizations using odds ratios as features for each cluster. This figure presents a UMAP visualization of ICD-10-CM codes based on their standardized ORs across different clusters. Each dot represents a specific ICD-10-CM code, with the dot size corresponding to the OR of the significantly associated cluster and the color indicating the corresponding cluster. The input features for the UMAP projection were derived from a matrix of standardized ORs of cluster-specific comorbidities. These standardized ORs were calculated by comparing the prevalence of each diagnosis within a specific cluster against all other clusters using Fisher's exact test (for cases < 5) or the Chi-squared test (for cases ≥ 5). A Bonferroni-corrected p-value threshold of 0.05 was applied to identify significantly enriched diagnoses. **A:** Overall population, **B:** Female, **C:** Male. ICD-10-CM, International Classification of Diseases, 10th Revision, Clinical Modification; UMAP, Uniform Manifold Approximation and Projection.

Figure 7. Validation analysis of cluster-specific significant comorbidities.

A Sankey plot illustrating the overlap of significant comorbidities across UCSF and UC-Wide populations within each cluster. The coverage rate represents the proportion of UCSF cluster-specific comorbidities that were statistically significant in UCSF and remained significant in UC-Wide clusters, identifying the most closely matched UC cluster. Number of cluster-specific comorbidities: Cluster 1 (812), Cluster 2 (225), Cluster 3 (97), Cluster 4 (19), Cluster 5 (1,583), Cluster A (1,976), Cluster B (2,731), Cluster C (819), Cluster D (40), Cluster E (19)

B–F Log-log plot showing significant overlapping comorbidities and the concordance of odds ratios for overlapping conditions in the overall population. P-values were calculated from Pearson correlation analysis. UC, University of California.

Figure 8. Miami plot for cluster-specific significant comorbidities. Miami Plot displaying enriched ICD10-CM codes identified using the two-sided Fisher's exact test (for cases < 5) or the Chi-squared test (for cases ≥ 5). Enrichment was determined based on a Bonferroni-corrected p-value < 0.05. ICD10-CM, the International Classification of Diseases, 10th Revision, Clinical Modification. **A:** Cluster 1, **B:** Cluster 2, **C:** Cluster 3, **D:** Cluster 4, **E:** Cluster 5.

Figure 9. Validation analysis of cluster-specific significant comorbidities across sex. Log-log plot illustrating the significant overlapping comorbidities and the concordance of odds ratios for these conditions in the sex-stratified population. The covered rate represents the proportion of UCSF cluster-sex-specific comorbidities that remain significant in both the UCSF and UC-Wide datasets. **A, C, E, G, I**: Cluster 1–Cluster 5 (Female) vs. Matched UC-Wide Cluster (Female), **B, D, F, H, J:** Cluster 1–Cluster 5 (Male) vs. Matched UC-Wide Cluster (Male). P-values were calculated from Pearson correlation analysis. UC, University of California.