Epidemiological insights into paediatric tuberculosis trends in the Western Cape, South Africa

Lauren R. Brown[1], MSc, Mariette Smith[2,3], MSc, Cari van Schalkwyk[1], PhD, Leigh F. Johnson[3], PhD, Vanessa Mudaly[4], MPH, Erika Mohr-Holland[5], MPH, H. Simon Schaaf[6,] MBChB, PhD, James A. Seddon[6,7,] MBBS, PhD, Anneke C. Hesseling[6], MBChB, PhD, James Nuttall[8], MBChB, PhD, Helena Rabie[9], MBChB, PhD, Mary-Ann Davies[3,4], MBChB, PhD, Andrew Boulle[2,3], MBChB, PhD, Karen du Preez[6], MBChB, PhD

Affiliations
[1]South African Centre for Epidemiological Modelling and Analysis, Stellenbosch University, Stellenbosch, South Africa
[2]Provincial Health Data Centre, Western Cape Department of Health, Cape Town, South Africa
[3]Centre for Integrated Data and Epidemiological Research, University of Cape Town, Cape Town, South Africa
[4]Department of Health and Wellness, Provincial Government of the Western Cape, Cape Town, South Africa
[5]City of Cape Town Health Department, Specialised Health, Epidemiology Unit, Cape Town, South Africa
[6]Desmond Tutu TB Centre, Department of Paediatrics and Child Health, Stellenbosch University, Cape Town, South Africa
[7]Department of Infectious Disease, Imperial College London, United Kingdom
[8]Department of Paediatrics and Child Health, University of Cape Town, Cape Town, South Africa
[9]Department of Paediatrics and Child Health, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa

Address correspondence to: Lauren Brown, SACEMA, Stellenbosch University, 19 Jonkershoek Rd, Mostertsdrift, Stellenbosch, 7600 [laurenbrown@sun.ac.za]

Short title: Epidemiological insights into paediatric TB in the Western Cape

Conflict of interest disclosures: The authors have no conflicts of interest relevant to this article to disclose.

Funding/support: This work was supported by the Fogarty International Center of the US National Institutes of Health [K43TW011006 K.d.P] and the Bill & Melinda Gates Foundation [INV063625 L.R.B., C.v.S., and L.F.J.]. These funders had no role in the design and conduct of the study.

Abbreviations: TB = tuberculosis, HIV = Human Immunodeficiency Virus, WC = Western Cape, CAWTB = children and adolescents with TB, PHDC = Provincial Health Data Centre, (E)PTB = (extra)pulmonary TB, (I)LTFU = (initial) loss to follow-up

Article summary
Using comprehensive provincial data, this study captures high TB incidence in young children, geographic variation in diagnosis, care cascade losses, and underreporting in routine surveillance.

What's Known on This subject
TB among children and adolescents remains a major public health concern in high-burden settings like the Western Cape. Improved understanding of local epidemiology and integrated data systems could assist in strengthening surveillance, guiding interventions, and improving outcomes for paediatric TB.

What This Study Adds
Analysis of microbiologically confirmed and clinically diagnosed TB in the Western Cape (2017-2023) revealed high incidence among young children, substantial losses across the care cascade, geographic heterogeneity in diagnosis, and underreporting, highlighting the value of integrated data for improved surveillance.

Contributors statement

Lauren R Brown conceptualized the study, conducted the formal analysis and investigation, designed the methodology, wrote the original draft and edited the final manuscript.

Mariette Smith conceptualized the study, designed the methodology, and extracted the original dataset.

Dr Cari van Schalkwyk and Dr Karen du Preez conceptualized the study, designed the methodology, and reviewed and edited the final manuscript.

Prof Leigh Johnson, Vanessa Mudaly, Erika Mohr-Holland, Prof H. Simon Schaaf, Prof James A. Seddon, Prof Anneke C. Hesseling, Prof James Nuttall, Prof Helena Rabie, Prof Mary-Ann Davies, and Prof Andrew Boulle reviewed and edited the final manuscript.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Abstract

Background and objectives

Paediatric tuberculosis (TB) remains a major public health concern in high-burden settings like the Western Cape (WC), South Africa. We analysed geographic differences in TB burden among children and adolescents, described temporal trends, and quantified gaps in the TB care cascade.

Methods

We analysed TB episodes recorded in the WC Provincial Health Data Centre (PHDC) from 2017-2023, stratified by 5-year age groups, and compared them to adult episodes. We assessed HIV status, drug resistance status, microbiological testing, disease classification, place of diagnosis, and TB treatment outcomes. Reporting gaps were estimated by comparing PHDC-recorded episodes to national notifications. Incidence rates were calculated using mid-year population estimates.

Results

In 2023, TB incidence rates of diagnosis in the WC were 722.4, 189.1, 171.2, and 523.4 per 100,000 population ages 0-4, 5-9, 10-14, and 15-19 years. Children aged 0-4 years accounted for 47.9% of paediatric TB episodes. In the Cape Winelands district in 2023, TB incidence among 0-4-year-olds was double that of adults in the district and 2-4 times higher than 0-4-year-olds in other districts. Among PHDC-recorded episodes, 17.3% were not reported at national level. Treatment success was low, with only 70.3% of diagnosed children and adolescents completing treatment in 2023.

Conclusions

Our findings highlight geographic variation in paediatric TB burden in the WC, emphasizing the need to address local drivers to inform targeted interventions. Gaps in the paediatric TB care cascade remain major concerns. Strengthening integrated data systems beyond TB treatment registers, could improve surveillance, health system planning, and patient outcomes.

Introduction

Childhood tuberculosis (TB) continues to be a global public health priority. In 2023, the

World Health Organization (WHO) estimated that 1.3 million children and young adolescents

(aged 0–14 years) developed TB, accounting for 12% of the global TB burden [1]. Children

under 5 years of age are particularly vulnerable to developing severe forms of TB and

experiencing rapid disease progression compared to adults [2,3]. This risk is exacerbated in

settings like South Africa, where the prevalence of Human Immunodeficiency Virus (HIV)

contributes to the TB burden in children [4]. Diagnosing TB in children remains challenging

due to non-specific clinical symptoms, the difficulty of obtaining respiratory samples, and the

low sensitivity of microbiological tests due to mainly paucibacillary disease, leading to

underdiagnosis and delayed treatment [5]. While South Africa achieved the End TB Strategy's

milestone of reducing total TB incidence by at least 50% by 2025, in 2023 only 64% of the

WHO's estimated TB burden among children and young adolescents under 15 years of age

were notified and treated in the country [1].

A study using routinely collected data from the Electronic TB Register (ETR.Net) found that

children and adolescents with TB (CAWTB) aged 0-19 years old contributed to 22.7% the TB

burden in the Western Cape (WC) province of South Africa in 2011 [6]. However, this study

predated the establishment of the Western Cape Provincial Health Data Centre (PHDC), a

unique data integration system that consolidates clinical information from multiple health

platforms to create a single patient record [7]. The PHDC uses data from the primary health

care information system (PHCIS), patient registration and health management information

system (PREHMIS), and the electronic drug-resistant TB register (EDRWeb) to create TB

episodes [7] and additionally includes evidence relating to TB patient care, including, but not

limited to, TB drug regimens dispensed, laboratory test results, admissions, diagnosis coding

and observational data. The PHDC supports patient care, service delivery, operational

management, and research. Other provinces in South Africa, as well as most low- and

middle-income countries, rely on TB treatment register data that reflects notifications and

treatment initiations rather than actual TB episodes, demonstrating an important reporting gap

[8,9]. Through its episode algorithm, the PHDC enables a more comprehensive analysis of TB

burden than is available in other provinces by providing critical insights into patients'

pathways of care.

Monitoring of the progression from TB diagnosis to treatment initiation, the frequency of

health facility visits, recurrent TB episodes and treatment outcomes by the PHDC provides

information on the TB care cascade [10–12]. The TB care cascade is a framework used to

quantify patient outcomes at various stages of TB care [13], and losses along this care cascade

present a challenge for reducing the TB burden in South Africa [14]. Individuals with TB may

be lost along the care cascade anywhere between accessing healthcare services, receiving test

results, initiating TB treatment, and successfully completing their treatment. Quantifying

these losses can aid in highlighting gaps in health systems, and planning strategies to improve

the quality of patient care.

This study aimed to analyse geographic differences in the burden of TB among children (0-9

years) and adolescents (10-19 years) within the context of the adult epidemic and examine

trends across districts in the WC from 2017 to 2023 using comprehensive data from the

PHDC. Additionally, we sought to quantify gaps in the TB care cascade for these age groups.

Methods

Data sources

We obtained data on the number of TB episodes that were laboratory confirmed or clinically

diagnosed in the WC at district level from the PHDC over a seven-year period (1 January

2017 to 31 December 2023). A summary of differences in the PHDC TB episode algorithm

and notifications reported by other provinces is highlighted in the supplement. Data on TB

episodes were stratified by routine PHDC characteristics which include HIV status, TB drug

resistance status, laboratory testing (whether a diagnostic test was conducted for the episode),

disease classification (pulmonary TB [PTB] with or without extra-pulmonary TB [EPTB] vs

EPTB only), place of diagnosis (at a primary healthcare [PHC] facility or hospital), inclusion

in the TB register and treatment outcomes at the time of data extraction (including successful

treatment, initial loss to follow-up [ILTFU], or treatment loss to follow-up [LTFU], and

death). Deaths occurring in-hospital or those known to the TB programme are recorded, but

there is no routine linkage to vital registration systems and thus some deaths may not be

recorded. Supplementary Table S1 presents definitions of these characteristics. Both episodes

and testing data were extracted in five-year age groups for individuals <20 years old, and for

adults 20 years and over for comparison.

We used mid-year population estimates from the provincial Thembisa model [15] as

denominators to calculate the incidence of TB diagnosis between 2017 and 2023 for each age

group. Thembisa is an integrated demographic and HIV projection model developed for

South Africa, applied at national and provincial levels. To capture incidence of TB diagnosis

at a smaller spatial scale, we used district-level population estimates from the Naomi model

[16], a Bayesian small-area estimation model. Naomi population estimates were available for

June 2017, September 2022, and December 2023 within our seven-year period. To calculate

mid-year population estimates at the district level, we applied district-level proportions from

the Naomi model to Thembisa province-level population estimates. More details can be

found in Supplementary Figure S1.

Epidemiological analysis

We analysed the number of TB episodes at district-, and province-level, focusing on children

(0-4 and 5-9 years) and adolescents (10-14 and 15-19 years), and included adults (≥20 years)

to provide context of the local epidemic. The geographic distribution of TB episodes was

based on the facility with the earliest electronic evidence of TB rather than place of residence.

We further disaggregated TB episodes by characteristics recorded in the PHDC

(supplementary Table S1). We aggregated these characteristics over 2017-2023 in the primary

analysis and provide yearly trends in the supplement. We highlighted spatial differences in

paediatric TB indicators in 2023 to show the most recent data on the epidemic.

We defined incidence as the number of TB episodes recorded in the PHDC divided by

Thembisa mid-year population estimates for 2017-2023 by 5-year age groups. This measure

reflects incidence of diagnosis as opposed to true incidence. Incidence rates are reported per

100,000 population. We analysed gaps in reporting defined as the difference between TB

episodes in the PHDC and treatment initiations captured in the TB treatment register which is

reported to national government (the reporting gap is further described in the supplement).

Ethics statement

Ethics approval was obtained from the Health Research Ethics Committee at Stellenbosch

University (N21/04/041) and the Western Cape Provincial Department of Health and

Wellness (WC_202107_021).

Results

Between 2017 and 2023, a total of 339,573 high-confidence episodes of TB occurred in the WC using the PHDC algorithm. Of these, 40,065 (11.8%) were CAWTB aged 0-14 years. Numbers of TB episodes across age groups and districts over the seven-year period are provided in supplementary Table S2. Incidence declined between the end of 2019 and 2020, coinciding with the COVID-19 pandemic and its associated public health measures [17] (Figure 1). The largest decline was experienced in children aged 0-4 years (by 145.7 per 100,000 population). By the end of 2021, incidence rates across all age groups had returned to expected levels based on pre-COVID-19 trends, followed by an increase in incidence among 0-4 years compared to other age groups in 2022. In 2023, the province-level incidence of diagnosis was estimated at 722.4, 189.1, 171.2, and 523.4 per 100,000 population for individuals aged 0-4, 5-9, 10-14, and 15-19 years, respectively, with the highest incidence estimated in adults (876.8 per 100,000 population). In the Cape Winelands district, the estimated incidence among children aged 0-4 years in 2023 was approximately double that of adults in the district and 2-4 times higher than young children in other districts (Supplementary Table S3).

Figure 1: TB incidence rates in the WC between 2017 and 2023 for different age groups spanning children, adolescents and adults. Time points represent incidence based on cumulative totals at the end of each year. *Incidence refers to the incidence of TB diagnosis as opposed to true TB incidence.

Despite routine HIV testing for CAWTB, a high proportion of TB episodes had unrecorded

HIV status, particularly in those aged 5-9 and 10-14 years (25% and 23.8%, respectively),

over the 7-year period (Table 1). Of the CAWTB who had a known HIV status, 11.1% (5,165

of 46,423) were living with HIV. Microbiological testing for TB was limited among children

aged 0-4 and 5-9 years, with 56.4% and 32.9% of TB episodes, respectively, being diagnosed

in the absence of microbiological testing. District-level comparisons among children under 5

years for 2023 can be found in supplementary Figure S2. Of TB episodes among individuals

aged 0-19 years diagnosed in a hospital, 43% were not recorded in the TB treatment register,

compared to 7.8% of those diagnosed in PHC facilities (supplementary Table S4). Among

CAWTB, 97.3% were treated for drug-susceptible (DS)-TB and 86.3% were classified as

having pulmonary TB, with similar proportions across age groups. TB treatment outcomes

show that ILTFU was 15%, driven largely by the 5-9 and 10-14-year-olds, and treatment

LTFU was 12%. Treatment success was the highest among adolescents (75.2%). Less than

2% of CAWTB are known to have died. The proportions of each characteristic remain

reasonably consistent over time (supplementary Figure S3).

Table 1: Demographic and clinical characteristics of children and adolescents diagnosed with

TB in the Western Cape between 2017 and 2023 (N = 57,614).

	Age 0-4 years	Age 5-9 years	Age 10-14 years	Age 15-19 years	Age 0-19 years
Total episodes*	N = 27,620 (47.9%)	N = 6,427 (11.2%)	N = 6,018 (10.4%)	N = 17,549 (30.5%)	N = 57,614
HIV status					
Negative	21,443 (77.6%)	4,123 (64.2%)	3,927 (65.3%)	11,765 (67%)	41,258 (71.6%)
Positive	1,625 (5.9%)	696 (10.8%)	657 (10.9%)	2,187 (12.5%)	5,165 (9%)
Unrecorded[†]	4,552 (16.5%)	1,608 (25%)	1,434 (23.8%)	3,597 (20.5%)	11,191 (19.4%)
Disease susceptibility type					
DS-TB	27,049 (97.9%)	6,245 (97.2%)	5,836 (97%)	16,920 (96.4%)	56,050 (97.3%)
DR-TB	571 (2.1%)	182 (2.8%)	182 (3%)	629 (3.6%)	1,564 (2.7%)

Microbiological testing for *Mycobacterium tuberculosis*					
Confirmed	5,352 (19.4%)	2,882 (44.8%)	4,438 (73.7%)	14,783 (84.2%)	27,455 (47.7%)
Negative	6,699 (24.3%)	1,431 (22.3%)	943 (15.7%)	1,555 (8.9%)	10,628 (18.4%)
No testing done	15,569 (56.4%)	2,114 (32.9%)	637 (10.6%)	1,211 (6.9%)	19,531 (33.9%)
Disease classification					
EPTB only	1,686 (6.1%)	713 (11.1%)	801 (13.3%)	1,265 (7.2%)	4,465 (7.7%)
PTB[‡]	23,738 (85.9%)	5,187 (80.7%)	4,921 (81.8%)	15,867 (90.4%)	49,713 (86.3%)
Unknown	2,196 (8%)	527 (8.2%)	296 (4.9%)	417 (2.4%)	3,436 (6%)
In TB treatment register					
Yes	22,643 (82%)	4,534 (70.5%)	4,626 (76.9%)	15,851 (90.3%)	47,654 (82.7%)
No	4,977 (18%)	1,893 (29.5%)	1,392 (23.1%)	1,698 (9.7%)	9,960 (17.3%)
Place of diagnosis					
Hospital	9,226 (33.4%)	1,788 (27.8%)	1,511 (25.1%)	3,048 (17.4%)	15,573 (27%)
PHC facility	18,394 (66.6%)	4,639 (72.2%)	4,507 (74.9%)	14,501 (82.6%)	42,041 (73%)
TB treatment outcome					
Successful treatment	19,542 (70.8%)	3,892 (60.6%)	4,048 (67.3%)	13,204 (75.2%)	40,686 (70.6%)
Still on treatment	54 (0.2%)	15 (0.2%)	13 (0.2%)	100 (0.6%)	182 (0.3%)
Died with TB	333 (1.2%)	80 (1.2%)	89 (1.5%)	464 (2.6%)	966 (1.7%)
ILTFU	4,144 (15%)	1,734 (27%)	1,296 (21.5%)	1,461 (8.3%)	8,635 (15%)
Treatment LTFU	3,430 (12.4%)	684 (10.6%)	549 (9.1%)	2,223 (12.7%)	6,886 (12%)
Other[Ω]	117 (0.4%)	22 (0.3%)	23 (0.4%)	97 (0.6%)	259 (0.4%)

DR-TB = drug-resistant TB, DS-TB = drug-susceptible TB, (E)PTB = (extra)pulmonary TB, PHC = primary health care, (I)LTFU = (initial) loss to follow-up

* Supplementary table S5 provides a comparison of these characteristics to adults aged 20 years and over

† More likely to be negative as negative rapid tests for HIV are not always recorded

‡ PTB includes episodes classified as PTB only or PTB and EPTB

Ω Including transferred out of WC (N = 257 for 0-19-year-olds) or unsuccessful treatment (N = 2 for 0-19-year-olds)

There was substantial heterogeneity in paediatric TB indicators across districts in the WC, as shown in Figure 2(A) and 2(B). The proportion of TB episodes in 0–14-year-olds relative to total TB episodes ranged from 9.91% in Cape Town to 20.26% in the Cape Winelands (Figure 2(A)). The ratio of TB episodes in 0–4-year-olds to those in 5–14-year-olds, ranged from 1.73 in Garden Route to 3.57 in the Cape Winelands.

Figure 2: District-level indicators in the WC in 2023. (A) Proportion of TB episodes in 0-14-year-olds relative to total TB episodes, including adults. (B) Ratio of TB episodes in 0–4-year-olds versus 5–14-year-olds. CK = Central Karoo, CPT = City of Cape Town, CW = Cape Winelands, GR = Garden Route, Ov = Overberg, WC = West Coast.

Large gaps exist between diagnosis, treatment initiation, and successful treatment for

CAWTB in 2023 (Figure 3). Of the total episodes among CAWTB, 15.5% did not initiate

treatment after being diagnosed with TB (ILTFU), with the largest loss occurring in the 5–9-

year-olds (26%). Between 26% and 39% of CAWTB did not successfully complete treatment,

attributed to treatment LTFU or death. Gaps in reporting and proportions of individuals LTFU

differed between districts as observed in Figure 4.

Figure 3: Age-stratified TB care cascades for children and adolescents in 2023 in the Western Cape.

The true burden of TB in these age groups has not been adequately estimated in South Africa, represented by the "unknown" first pillar in the cascade. The green bars represent numbers of episodes at each step of the care cascade, including diagnosis, treatment initiation and registration, and successful treatment (including those still on treatment at the time of data extraction). Percentage differences between each pillar are illustrated in grey blocks. Includes children treated for both drug-resistant and drug-susceptible TB.

Between 2017 and 2023, 17.3% of TB episodes among 0-19-year-olds were not included in

the TB treatment register (reporting gap) of which 67.2% (6,698 of 9,960) were diagnosed in

hospital (supplementary Table S4). In 2023, reporting gaps were as high as 18.2% among

CAWTB in Overberg, and below 10% in the Cape Winelands and Central Karoo (Figure 4A).

Combined LTFU (ILTFU and treatment LTFU) among CAWTB was high in all districts, with

West Coast (36%) having the highest proportion and Central Karoo (20.1%) the lowest

(Figure 4B).

Figure 4: District-level losses along the TB care cascade among children and adolescents in 2023. (A) the reporting gap between the PHDC versus the TB treatment register, and (B) combined ILTFU and treatment LTFU.

CK = Central Karoo, CPT = City of Cape Town, CW = Cape Winelands, GR = Garden Route, Ov = Overberg, WC = West Coast.

Discussion

We analysed TB episodes among children and adolescents in the WC between 2017 and

2023. TB incidence was particularly high among children aged 0–4 years, with substantial

variation across districts and age groups. Nearly a third of children and young adolescents

were lost along the TB care cascade, including major reporting gaps between PHDC-recorded

episodes and those reported through the national TB treatment register.

Geographical heterogeneity TB diagnosis was observed across the WC between 2017 and

2023, highlighting the importance of spatial factors when analysing TB burden [18]. In the Cape

Winelands district, the estimated incidence of TB diagnosis among children aged 0-4 years in

2022-2023 was approximately double that of adults in the district and 2-4 times higher than

0-4-year-olds in other districts. This district also had the highest adult TB incidence. One

contributing factor may be variable implementation of national TB screening guidelines for

child contacts under five to assess eligibility for TB preventive treatment (TPT) [19], leading to

inter-district variation in household contact screening. Similar heterogeneity was observed in

the proportion of overall burden among children (Figure 2A). In most districts, this

proportion fell within the expected range of 10% to 20% [20], with the Cape Winelands district

highest at 20.3%, 5.7% higher than the second highest district. This may indicate a localized

epidemic, proactive case detection with a low threshold to start treatment, or overdiagnosis

Children from neighbouring districts may also seek care in Cape Winelands due to its well-

established health services, including a regional specialist TB hospital [21]. Ongoing paediatric

TB research conducted by academic institutions in the Cape Winelands and stronger training

initiatives at healthcare facilities in the district may further contribute to high detection rates

and clinician confidence in diagnosing and initiating TB treatment. The district also had

better cascade outcomes, with lower reporting gaps and loss to follow-up than others,

suggesting strong case management. Further investigation, including patient-level analysis

and clinician engagement, may help explain these geographic differences.

Identifying and quantifying gaps in the TB care cascade for children is critical to improve

child and adolescent TB care. Our findings indicate substantial losses between diagnosis,

treatment initiation, and successful treatment completion, likely driven by multiple factors.

The gap between diagnosis and treatment initiation (ILTFU) is particularly concerning, as

individuals lost at this stage face an elevated risk of mortality [11,12]. Between 2017 and 2023,

17.3% of CAWTB were ILTFU, 67.2% of whom were diagnosed in hospital. Ensuring

completeness of hospital reporting would help address this gap [22]; however, most hospitals

lack TB registers, and diagnosed individuals are referred to PHC facilities for registration and

treatment initiation. Movement between levels of care poses a challenge for continuity of TB

care, especially as children frequently require hospital-based diagnosis due to diagnostic

challenges [5]. This gap is being addressed through the TB treatment action list (TTAL), which

links hospital-diagnosed patients to PHC facilities and prompts recall for treatment.

Interventions such as LINKEDIn [12] have proved successful in reducing ILTFU in two WC

subdistricts and should be considered by TB programs. Overall, 70.6% of CAWTB were

treated successfully, notably lower than success rates in smaller WC studies (78% in 2014 [23];

90% between 2005-2012 [24]) and a national 2016 study (80%) [25]. Treatment LTFU contributed

an additional 12-18%. As no national or sub-national estimates of the true TB burden among

children and adolescents currently exist, we could not quantify the gap between true burden

and diagnosed cases - an important consideration for future research.

In this high TB-burden setting, children and young adolescents under 15 years contributed

11.8% of all TB episodes, consistent with the WHO's estimated 12% for South Africa [1].

Children under 5 years accounted for the majority of TB in children overall (Table 1; Figure

2B). Microbiological confirmation of TB remains challenging in this group due to the

paucibacillary nature of the disease and difficulties in collecting specimens [5]. Over half of

children under 5 diagnosed with TB in our study had no microbiological test for

Mycobacterium tuberculosis. This proportion varied by district (supplement Figure S2) and

may reflect differences in access to diagnostics, staff training, or testing practices, though

specific drivers remain unclear. Further research is needed to understand how resource or

strategy disparities contribute to these differences. The high incidence of TB diagnosis in

young children likely reflects ongoing community transmission, particularly in households

where they may be exposed to infectious adults [26]. This highlights the need for strengthened

TB control measures, including systematic household contact screening in high-risk groups

and strengthened implementation of targeted universal testing for TB (TUTT) and TPT

policies that endorse a test-and-treat approach.

Our data highlight the advantages of an integrated data system, offering a more comprehensive surveillance approach compared to registry-based systems used elsewhere in South Africa and other high TB-burden countries. We observed a reporting gap of 17.3% (9,960 unreported episodes) among CAWTB, emphasizing limitations in relying solely on routine surveillance. Similar underreporting has been found in previous WC studies, ranging from 15% to 38% [8,9,22]. Contributing factors likely include patients being managed exclusively in hospitals, failure to initiate treatment after referral to PHC facilities, and data capture challenges, particularly in smaller clinics without access to TB registers. The PHDC helps address these challenges by integrating data across healthcare levels, improving case ascertainment and patient follow-up. Integrated systems that capture both microbiologically confirmed and clinically diagnosed TB episodes offer substantial advantage for surveillance of paediatric TB. Scaling up similar systems in other settings could enhance routine surveillance and support monitoring and evaluation of child and adolescent TB services.

Our study has several limitations. Population estimates, particularly at the district level, may be unreliable and introduce uncertainty; however, the Thembisa [15] and Naomi [16] model estimates are sufficiently robust to inform government portfolio planning and management. Additionally, we were unable to estimate the incidence of TB diagnosis at the subdistrict level due to the lack of reliable subdistrict population data – an important limitation that should be addressed in future research. Our analysis relied on aggregated data, which allowed for broad observations of key indicators but limited our ability to examine individual-level health-seeking behaviour. Treatment outcome data in the PHDC are based on clinician-recorded entries in PHCIS and PREHMIS, which are vulnerable to documentation errors and subjectivity, especially when follow-up is limited. Incomplete capture of outcomes in TB

hospitals and outpatient settings, where many children are treated, further limits the accuracy

of routine reporting. Additionally, point-of-care HIV tests are often not captured

electronically and, for TB episodes not captured in registers, HIV-negative results may go

unrecorded. The geographic distribution of TB episodes was based on the facility with the

earliest electronic evidence of TB rather than place of residence, which may not fully reflect

where transmission or exposure occurred. Addressing these gaps is critical for improving

paediatric TB surveillance in the Western Cape.

Conclusion

The PHDC provides a unique dataset that comprehensively identifies clinically diagnosed as

well as microbiologically confirmed episodes of TB in children and adolescents in a high TB

burden setting. Our analysis highlights important geographical heterogeneity in paediatric TB

at provincial level, emphasising the need to understand local drivers to inform targeted

interventions. Gaps in the TB care cascade remain a major concern, particularly initial and

treatment loss to follow up as well as the large reporting gap for children diagnosed in-

hospital. Expanding integrated, electronic data systems can greatly strengthen health system

planning and improve patient care.

Data Sharing Statement: Deidentified individual participant data will not be made available.

References

1. World Health Organization. Global tuberculosis report 2024. Geneva, Switzerland: World Health Organization, 2024.
2. Newton SM, Brent AJ, Anderson S, Whittaker E, Kampmann B. Paediatric tuberculosis. Lancet Infect Dis 2008; 8:498–510.
3. Roya-Pabon CL, Perez-Velez CM. Tuberculosis exposure, infection and disease in children: a systematic diagnostic approach. Pneumonia 2016; 8:23.
4. Anyalechi GE, Bain R, Kindra G, et al. Tuberculosis prevalence, incidence and prevention in a South African cohort of children living with HIV. J Trop Pediatr 2022; 68.
5. Gunasekera KS, Vonasek B, Oliwa J, et al. Diagnostic challenges in childhood pulmonary tuberculosis—optimizing the clinical approach. Pathogens 2022; 11:382.
6. Snow K, Hesseling AC, Naidoo P, Graham SM, Denholm J, du Preez K. Tuberculosis in adolescents and young adults: epidemiology and treatment outcomes in the Western Cape. Int J Tuberc Lung Dis 2017; 21:651–657.
7. Boulle A, Heekes A, Tiffin N, et al. Data centre profile: the provincial health data centre of the Western Cape province, South Africa. Int J Popul Data Sci 2019; 4.
8. Preez K Du, Schaaf HS, Dunbar R, et al. Incomplete registration and reporting of culture-confirmed childhood tuberculosis diagnosed in hospital. Public Health Action 2011; 1:19–24.
9. Marais BJ, Hesseling AC, Gie RP, Schaaf HS, Beyers N. The burden of childhood tuberculosis and the accuracy of community-based surveillance data. Int J Tuberc Lung Dis 2006; 10:259–63.
10. Jennings K, Lembani M, Hesseling AC, et al. A decline in tuberculosis diagnosis, treatment initiation and success during the COVID-19 pandemic, using routine health data in Cape Town, South Africa. PLoS One 2024; 19:e0310383.
11. Osman M, Meehan S-A, von Delft A, et al. Early mortality in tuberculosis patients initially lost to follow up following diagnosis in provincial hospitals and primary health care facilities in Western Cape, South Africa. PLoS One 2021; 16:e0252084.
12. Meehan S-A, Hesseling AC, Boulle A, et al. Reducing initial loss to follow-up among people with bacteriologically confirmed tuberculosis: LINKEDin, a quasi-experimental study in South Africa. Open Forum Infect Dis 2024; 11.
13. Subbaraman R, Jhaveri T, Nathavitharana RR. Closing gaps in the tuberculosis care cascade: an action-oriented research agenda. J Clin Tuberc Other Mycobact Dis 2020; 19:100144.
14. Naidoo P, Theron G, Rangaka MX, et al. The South African tuberculosis care cascade: estimated losses and methodological challenges. J Infect Dis 2017; 216:S702–S713.
15. Johnson LF, Meyer-Rath G, Dorrington RE, et al. The effect of HIV programs in South Africa on national HIV incidence trends, 2000–2019. J Acquir Immune Defic Syndr 2022; 90:115–123.
16. Eaton JW, Dwyer☐Lindgren L, Gutreuter S, et al. Naomi: a new modelling tool for estimating HIV epidemic indicators at the district level in sub☐Saharan Africa. J Int AIDS Soc 2021; 24.
17. Dheda K, Perumal T, Moultrie H, et al. The intersecting pandemics of tuberculosis and COVID-19: population-level and patient-level impact, clinical presentation, and corrective interventions. Lancet Respir Med 2022; 10:603–622.
18. Trauer JM, Dodd PJ, Gomes MGM, et al. The importance of heterogeneity to the epidemiology of tuberculosis. Clin Infect Dis 2019; 69:159–166.
19. Department of Health South Africa. Management of tuberculosis in children and adolescents: A clinical guideline for the diagnosis and treatment of drug-susceptible TB in

children and adolescents in South Africa. Available at: https://knowledgehub.health.gov.za. Accessed 4 February 2025.

20. Seddon JA, Jenkins HE, Liu L, et al. Counting children with tuberculosis: why numbers matter. Int J Tuberc Lung Dis 2015; 19:9–16.
21. Moyo S, Verver S, Hawkridge A, et al. Tuberculosis case finding for vaccine trials in young children in high-incidence settings: a randomised trial. Int J Tuberc Lung Dis 2012; 16:185–191.
22. du Preez K, Schaaf HS, Dunbar R, Swartz A, Naidoo P, Hesseling AC. Closing the reporting gap for childhood tuberculosis in South Africa: improving hospital referrals and linkages. Public Health Action 2020; 10:38–46.
23. du Preez K, du Plessis L, O'Connell N, Hesseling AC. Burden, spectrum and outcomes of children with tuberculosis diagnosed at a district-level hospital in South Africa. Int J Tuberc Lung Dis 2018; 22:1037–1043.
24. Osman M, Lee K, Du Preez K, Dunbar R, Hesseling AC, Seddon JA. Excellent treatment outcomes in children treated for tuberculosis under routine operational conditions in Cape Town, South Africa. Clin Infect Dis 2017; 65:1444–1452.
25. Osman M, van Schalkwyk C, Naidoo P, et al. Mortality during tuberculosis treatment in South Africa using an 8-year analysis of the national tuberculosis treatment register. Sci Rep 2021; 11:15894.
26. Seid G, Alemu A, Dagne B, Sinshaw W, Gumi B. Tuberculosis in household contacts of tuberculosis patients in sub-Saharan African countries: a systematic review and meta-analysis. J Clin Tuberc Other Mycobact Dis 2022; 29:100337.

Figure legends

Figure 1: TB incidence rates in the WC between 2017 and 2023 for different age groups spanning children, adolescents and adults. Time points represent incidence based on cumulative totals at the end of each year. *Incidence refers to the incidence of TB diagnosis as opposed to true TB incidence.

Figure 2: District-level indicators in the WC in 2023. (A) Proportion of TB episodes in 0-14-year-olds relative to total TB episodes, including adults. (B) Ratio of TB episodes in 0–4-year-olds versus 5–14-year-olds. CK = Central Karoo, CPT = City of Cape Town, CW = Cape Winelands, GR = Garden Route, Ov = Overberg, WC = West Coast.

Figure 3: Age-stratified TB care cascades for children and adolescents in 2023 in the Western Cape. The true burden of TB in these age groups has not been adequately estimated in South Africa, represented by the "unknown" first pillar in the cascade. The green bars represent numbers of episodes at each step of the care cascade, including diagnosis, treatment initiation and registration, and successful treatment (including those still on treatment at the time of data extraction). Percentage differences between each pillar are illustrated in grey blocks. Includes children treated for both drug-resistant and drug-susceptible TB.

Figure 4: District-level losses along the TB care cascade among children and adolescents in 2023. (A) the reporting gap between the PHDC versus the TB treatment register, and (B) combined ILTFU and treatment LTFU. CK = Central Karoo, CPT = City of Cape Town, CW = Cape Winelands, GR = Garden Route, Ov = Overberg, WC = West Coast.