Hematological Events in Metformin Treated NSCLC Patients Post Chemotherapy: Drug Repurposing Based on Real-World Evidence from Case Controlled Study

Saikat Samadder[1*]

[1*]Corresponding author; Department of Pharmacology, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-Gu, Seoul, Korea.

Email: SaikatSamadder@yuhs.ac; saikat.samadder46@gmail.com

ORCID number: Saikat Samadder (0000-0001-8693-0863)

Author contribution: Saikat Samadder solely contributed all the works related to research and manuscript preparation.

Funding: No grants were obtained for this research.

Aim: In this multi-center case controlled study aimed at metformin repurposing on the basis of survival benefit in non-small cell lung cancer patients compared to non-diabetic patients, based on excessive myelosuppression or myeloupregualtion induced by chemotherapy and adverse events mitigating ability of metformin.

Methods: At the Severance hospitals from 1^{st} Jan 2010 to 1^{st} Jan 2020, (n = 1851) patients were identified from hospital database. Further analysis was performed using propensity score matched to visualize the survival outcome of metformin comparing with non-diabetic control group in non-small cell lung cancer patients. Using propensity score matching of 8 variables, post-chemotherapy days survived by diabetic patients compared with untreated patients shown in this study. Six different chemotherapy induced clinical events related to WBC, platelets, and neutrophils are studied to identify the relative detrimental role of metformin post chemotherapy frequency/cycle stratification.

Results: Total 354/1851 patients identified as diabetic non-small cell lung cancer patients remaining were non-diabetic patients. Post-propensity score matching 354 non-diabetic patients survived 521 days (95% CI 459 - 582), compared to treatment group patients survived for 580 days (95% CI 508 - 651) non-significantly with (*p*-value 0.1). The patients under radiotherapy survived 624 days (95% CI 521 - 726) in 142 control group, compared to that chemoradiation metformin treated patients survived 798 days (95% CI 698 - 967) *p*-value-0.02. Among (n = 136) metformin patients treated with metformin prior to 180 days of chemotherapy survived maximum days post chemotherapy start 2-year hazard ratio 0.42 (95% CI 0.008 – 0.7) *p*-value 0.02 compared to patients received metformin 180 days post chemotherapy. Post propensity score matching and chemotherapy cycle stratification leukocytosis, neutrophilic leukocytosis, and thrombocytopenia events were significantly high (*p*-value < 0.05) causing increased deaths in chemotherapy non-responder compared to responder group. Metformin treatment did not significantly reduce leukopenia, neutropenia, and thrombocytopenia events post chemotherapy compared to control (*p*-value < 0.05).

Conclusion: The overall survival of diabetic patients treated with metformin is not different from untreated patients. Fatalities in treatment and control groups induced by chemotherapy were similar. Prolonged use of metformin reduced hematological event risk in NSCLC patients.

Keywords: First-line chemotherapy, hematological events, non-responders, metformin, drug repurposing, real-world evidence.

INTRODUCTION: Lung Cancer is the second most cause of deaths among all cancers in the global population. As per GLOBOCAN report until 2020 a total of 1.7 million deaths due to lung cancer were recorded each year and 2.2 million newly diagnosed lung cancer were reported [1]. Several medicines were repurposed in past decades to counter this severe disease and to enhance the survival outcome [2, 3]. Anti-inflammatory medicine like celecoxib is presently prescribed for lung cancer patients originally known for the treatment of rheumatoid arthritis and ankylosing spondylitis. Pemetrexed is another best example of drug repurposed for lung cancer [4]. Maintaining the hematological balance within the clinical range was reported to have positive survival effect on cancer patients [5]. At the same time dual mode of actions such as immunosuppression and tumor growth inhibition is necessary to reduce adverse outcome in cancer. Identifying potential immunosuppressive agents similar to pemetrexed and celecoxib, beside immune-activating agents like GM-CSF, G-CSF, are the prime medications used to tackle chemotherapy induced immunopathologies [4, 6]. Events occurring from myelosuppression or hyperactivation may lead to detrimental conditions including death [7-12]. Chemotherapy induced diabetes is a rising concern and one of the longest ongoing debates to find the relationship between diabetes and cancer [12]. Type-II Diabetic patients were reportedly more prone toward various cancers depicting mainly gender, age, and lifestyle as the major confounding factors of disease progression [10].

Metformin, one of the oldest known medicines for diabetes, is known for increasing the risk of myelosuppression and related serious adverse events in long-term users [13]. In several retrospective analysis it was demonstrated that metformin use benefited cancer patient's survival outcome [14, 15]. At the same time metformin was not found to reduce the risk of deaths in chemoradiation patients with advanced lung cancer [16, 17]. Myelosuppression and myeloupregualtion may occur due to ongoing or prolong chemotherapy regimen, or metformin use or due to cancer progression [18, 19, 20]. Therefore, events resulting in deaths of patients post chemotherapy end potentially after 1 to 2 years of standard first-line treatment, monitoring deceased patient's immunopathological events retrospectively would potentially identify the actual cause of deaths in population with lung cancer [20]. Events such as leukopenia, lymphocytopenia, neutropenia, leukocytosis, thrombocytopenia, and thrombocytosis are often caused by cardiac events as well as infections thereby increasing the disease burden [21, 22]. These events are often known as chemotherapy induced adverse events reduces quality of life. Diabetes is induced by cancer progression, chemotherapy, and radiation therapy as well [23].

The main focus of this propensity score matched (PSM) retrospective analysis is to measure and compare the hazard ratio of metformin users and non-users diagnosed with non-small cell lung cancer (NSCLC) for drug repurposing. In this study events and event-to-death occurrence in patients post chemotherapy initiation in metformin patients compared to non-diabetic patients is analyzed. Events such as leukopenia, lymphocytopenia, neutropenia, neutrophilic leukocytosis, thrombocytopenia, and thrombocytosis are compared in overall patients post-chemotherapy initiation to find the risk or benefit from metformin usage. Identifying moderate responders toward chemotherapy and finding the event related deaths is of great importance and are rarely performed for drug repurposing. Here it is further demonstrated using chemotherapy frequency/cycle stratified Kaplan-Meier plot the effect of chemotherapy responding NSCLC patient's overall survival. By calculating the events occurrence rate and mortality rate in control and treatment groups as well in chemotherapy cycle stratified sub-groups, during 1st year and 2nd year post chemotherapy initiation to identify underlying cause of fatalities in diabetic patients. As diabetic patients may endure maximum chemotherapy treatment response compared to non-diabetic cancer patients is presently not well-known necessary for drug repositioning.

MATERIALS AND METHODS

Study design and population: In this present multi-center case controlled study, the real world data (RWD) was obtained from three different branches of Severance hospital consisting of electronic medical record (EMR) of patients receiving treatment for NSCLC. In this database patient information including age, gender, height, weight, body surface area (BSA), chemotherapy, immunotherapy, radiotherapy dose, dates of exposure, various clinical laboratory values obtained from continuous monitoring of hematological factors and patient vitals, various diseases and diagnosis dates, surgery and dates of surgical and radiological procedures are available, various non-chemo drugs, intake dates along with dose are accurately maintained. There is no missing information of patient included in this study.

To evaluate the effect of metformin in diabetic lung cancer patients aged above 19 years of age was included in this study. In hospital database there are 7,532 lung cancer patients identified based on searching this database with international classification of diseases-10 (ICD-10) codes among these patients (n=4,182) died until 30th Jun 2024 received chemotherapy. Patients are excluded based on following basis; missing chemo start date or chemo end date, SCLC and

neuroendocrine lung cancer were excluded, missing WBC counts, neutrophil, RBC, platelet counts, small cell lung cancer, patients diagnosed with neuroendocrine tumor, and patients diagnosed with stage I/II during diagnosis were excluded. Among the deceased patients there are 2688 patients with available WBC counts were selected for analysis. By implementing these exclusion criteria patients are finally included based on chemotherapy treatment provided within 180 days post cancer diagnosis and patients receiving chemotherapy dose above 0.4 mg/m^2 were included. A total of (n=1851) patients were included in baseline, among these patients (n=354) were diagnosed with diabetes pre/post cancer diagnosis. The inclusion and exclusion criteria of patients in this study is shown using flowchart (Figure 1). Clinical baseline patients were selected based on the average count of 10 days pre/post cancer diagnosis allowing to include1851 patients. Post curation of data, the analysis was performed based on 1:5 PSM of 1851 patients including 354 samples in each arms allowing to obtain statistical strength of above 100%. The patients died during 1-year and 2-year of chemotherapy start within 8-days post event occurrence were analyzed post chemotherapy frequency/cycles stratification. This study design allowed to identify the cause of survival benefit or deaths arising from cancer progression, diabetes, chemotherapy, and/or metformin in diabetic and non-diabetic patients. Further stratification of groups as per chemotherapy cycles was applied and similar analytics was performed.

Chemotherapy Dosage: To identify chemotherapy dose, volume of drug injected per cycle was divided from the BSA. The values of each patient's BSA were pre-calculated and stored in database with high compliance for each patient. This allowed to acquire and calculate lung cancer patient total chemotherapy dose in mg/m^2 as total dose for all patients. Total dosage for oral formulation was calculated from the standard dose in mg available as per formulation type, multiplied by daily recommended dose, and duration of the treatment. Therefore, the total chemotherapy dosage was total of both mg and mg/m^2 further pooled for matched patients based on total dosage [25]. The chemotherapeutic dosage was used as a variable in PSM this enabled to match patients with similar efficacy towards chemotherapy. While further stratification of chemotherapy frequency/cycle was performed to identify event occurrence and mortality rates among diabetic and non-diabetic groups.

Propensity Score Matching: To perform PSM using 8 variables were included during the matching processes in RStudio. PSM was matched including gender, age, lung cancer location, stage, total chemo dosage, metformin treated/untreated, surgery, and radiotherapy status. The

lung cancer metformin users ratio 1 were matched with nondiabetic metformin non-users with ratio 5 based on algorithm or values matched available from matched total chemotherapy dose and 6 other variables [24, 25]. Obtaining standardized difference of between -0.1 to 0.1 post-PSM is considered as a sign of balance and was not found (supplementary figure **1-2**). PSM using chemotherapy dosage were rarely performed in previous literatures.

Statistical Analysis: The statistical analysis was based on patients receiving all necessary therapies available for treatment post diagnosis as per patient's requirement and physician's decision. The confounding factors for this analysis are the age of cancer diagnosis, gender, body weight, cancer stage at diagnosis, lung cancer type, surgery, radiotherapy, salvage therapy, and total chemotherapy dosage. Responder and non-responders were analyzed by stratifying the frequency of chemotherapy post-PSM. The overall survival curve of responders and non-responders allowed to understand the impact of metformin post-chemotherapy. Stratification method allowed to obtain efficiency of chemotherapy curve obtained by segregating patients based on chemotherapy frequency/cycles (shown in supplementary figure 5). Observing the change in hematological laboratory values could identify frequency of events and event-to-death within 1-year and 2-year post-chemotherapy start were analyzed in this study, provided significant percentage change from baseline (shown in table **3 & 4)**. Hazard ratio (HR) was calculated by dividing diabetic patients died within 2 years with total number of patients, further dividing the ratio of patients died in Group A (180 pre chemotherapy start) by Group B (180 days pre/post chemotherapy start) and vise-versa for finding HR of Group A Vs Group C (180 days post chemotherapy start). Cox-proportional hazard regression analysis was used to estimate the HR of metformin users compared to non-users with cancer, with 95% confidence interval (CI). Data curation was performed in Microsoft excel, PSM and visualization were done with R (ver. 3.6.1; R Foundation for Statistical Computing, Vienna, Austria). The calculation of 95% confidence interval (CI) and *p*-value was performed online using MedCalc [26].

Event-to-Death Analysis: In the majority of patients event related mortality was observed within 0-7 days after 8 days, the mortality rate reduced. Therefore, the decision to analyze 0-7day mortality rate in patients was adapted. This fact was further confirmed in a separate analysis event to death per day reduced after 8 days (Day 0 to day 7) this confirmed post event 8 days was appropriate for analysis (supplementary figure 3). High mortality rate in the first month might not be specifically due to myelosuppression or other related events

(supplementary figure 4). Standard event occurrence rate was calculated, dividing events in group by total events, event-to-death was calculated, dividing number of fatalities by total number of patients/groups. The normal laboratory range of WBC, platelets, neutrophil cell counts, hematological events leukopenia (> $4x10^3$ cells/ul), lymphocytopenia (< $10x10^3$ cells/ul), neutropenia (> $2x10^3$ cells/ul), neutrophilic leukocytosis (< $7x10^3$ cells/ul), thrombocytopenia (> $150x10^3$ cells/ul), and thrombocytosis (< $400x10^3$ cells/ul) are provided (supplementary table 2).

RESULTS

Baseline Demographics: All deceased chemotherapy treated patients with cancer identified from 1st January 2010 to 1st January 2020 were selected for baseline (n = 1851) patients assessment. Patients survived or died until 30th June 2024 were considered as cut-off date prior to analysis (4.5 years). Among all 1851 were identified for analysis 68.07% male and 31.9% female, mean age at diagnosis was 63 years, mean body weight was 61.6 kg, all 360 (19.4%) patients at stage-III, 1440 (77.5%) were in stage-IV, remaining 51 (2.7%) were found recurred stage at baseline. In 1419/1851 (87.7%) patients were diagnosed with various types of non-small cell lung cancer during the first hospital visit remaining 432 (23.3%) were diagnosed as unspecified. Pulmonary surgical interventions were provided specifically for lung cancer 858 (46.35%) patients, 682 (36.8%) patients received chemo-radiation therapy, 365 patients were found to have undergone salvage therapy. At baseline in 1851 deceased individuals the WBC count was 9.0, RBC count was 4.2, platelet count was 290.7, neutrophil 6.3, albumin was 3.9 during the diagnosis of NSCLC. Post-PSM there was non-significant differences between the variables among groups (refer supplementary table 1). The chemotherapy duration mean days was 45 days less in control group compared to metformin treated patients.

Identifying and providing treatment for immunosuppressed patients prior to initiation of chemotherapy is highly recommended as immunosuppressant or chemotherapy may exacerbate the myelosuppression mainly for metformin patients. It was found that among 545/1851 patients and in matched 708 patients 230 patients received immunosuppressants at some point post diagnosis of cancer. Total 124/230 (53%) were in metformin group and 110/230 (47%) were treated with immunosuppressants in control group. Immunosuppressed 45/230 patients received immunosuppressants before 30 days of first-line chemotherapy that lasted for 6 months, 45 patients possibly experienced myeloupregualtion. Among 49/110 patients in control

arm received immunosuppressants during similar timeline. Post chemotherapy initiation within 1-year 456 patients and 448 patients experienced leukopenia and leukocytosis, respectively. In 411 patients neutropenia, and 523 neutrophilia leukocytosis was distinctly observed. In 437 patients thrombocytopenia and 283 experienced thrombocytosis within 1-year post first-line therapy.

Efficiency of Chemotherapy: Efficiency of chemotherapy was analyzed based on chemotherapy stratified dosage frequency. A small proportion of population died after the treatment start resulting in 380 deaths within 180 days post-chemotherapy initiation and these patients could be regarded as non-responder of chemotherapy. It was important to provide an overall survival curve (supplementary figure 5) to understand the efficiency of chemotherapy, based on stratification of patients as per response, as these patients completed at-least 3 to 6 months of first-line treatment and few presented better survival ability compared to non-responders, irrespective of chemotherapy dosage. Among 708/1851 matched patients responders survived maximum (n =58/708) 1351 days (95% CI 1160 - 1541) compared to non-responders (n = 380/708) 294 days (95% CI 254 - 333), and moderate-responders (n =270/708) survived 741 days (95% CI 658 - 823). PSM ensures partial randomization resulting in non-significant mortality difference between the groups of three sub-population caused by specific events (figure 5).

Benefit from Metformin: Total 354/1851 patients identified as diabetic NSLC patients remaining are non-diabetic patients. Pre-PSM there was significant survival benefit results not shown. Post PSM 354 non-diabetic patients survived 521 days (95% CI 459 - 582), compared to treatment group patients survived for 580 days (95% CI 508 - 651) with non-significant statistical difference *p*-value 0.1. Beyond 2-year survival post-chemo in control group matched patients in 77/354 patients survived mean 1392 days (95% CI 1228 – 1537) and in metformin group 85/354 patients survived 1580 days (95% CI 1418 – 1741) with *p*-value 0.1. Among 174/354 patients mean days survived 670 days (95% CI 569 - 770) received surgery in control group, compared to 176/354 patients in metformin 136 patients survived mean 745 days (95% CI 630 - 859) with non-significant *p*-value 0.1. The patients under radiotherapy survived 624 days (95% CI 521 - 726) post-chemotherapy (n = 142) control group patients, while in (n = 136) metformin treated patients survived 798 days (95% CI 698 - 967) with *p*-value-0.02. Among 136 diabetic patients diagnosed prior to 180 days of chemotherapy survived maximum days post chemotherapy start, 2-year HR 0.42 (95% CI 0.008 – 0.7) with *p*-0.02 compared to

patients receiving chemotherapy and metformin post-chemo. The 2-year mean survival days is 322 in (n=106/142) untreated patients compared to 276 days (n=83/136) in metformin group, the 2-year hazard ratio of metformin Vs control group patient is 0.82 (95% CI 0.02 – 0.24) with p-value 0.01. The non-diabetic chemoradiation recipients survived 692 days (95% CI 576 – 806) post cancer diagnosis compared to diabetic chemoradiation patients survived 926 days (95% CI 768 - 1083) with statistical significance of p-value 0.009.

Myelosuppression: Frequency of leukopenia, neutropenia, and thrombocytopenia was statistically significant in metformin patients (p-value < 0.05) compared to non-diabetic/control group patients post-chemotherapy, fatality was non-significantly high in metformin group. All these three events were chemotherapy induced events (figure 3 & 4**).** The risk of event occurrence rate increased significantly in metformin group for thrombocytopenia events after 1^{st} year of chemotherapy. There is no statistical significance in mortality rate observed between the control and metformin treated myelosuppressed patients within 1^{st} year or 2^{nd} year post chemotherapy start. During 2^{nd} year post-chemotherapy patients in control group experienced leukopenia and thrombocytopenia events rate increased significantly (p-value < 0.05) compared to metformin treated patients. The number of patients experienced events were not significantly different among the groups, the number of patients expired due to events were non-significantly lower in metformin group compared to control group (figure 4). Event-to-death prior to chemotherapy initiation did not occur in either of the groups for all hematological events.

Myeloupregualtion: Total immune-hyperactivation events such as neutrophilic leukocytosis, and thrombocytosis was experienced significantly by metformin group patients (p-value < 0.05) pre/post initiation of chemotherapy. Myeloupregulation was non-significant in both diabetic and non-diabetic patients within 1^{st} year post-initiation of chemotherapy. In the 2^{nd} year thrombocytosis related events occurred significantly in metformin treated patients (p-value < 0.05) compared to control. Leukocytosis and neutrophilic leukocytosis events increased in control group patients (p-value < 0.05) significantly than metformin treated patients. There is no statistical significance in mortality rate caused due to events observed between the control and metformin treated myeloupregulated patients. Although fatalities in metformin group was lower in leukocytosis, and neutrophilic leukocytosis events compared to treatment group except for thrombocytosis event number of fatalities were not different among groups. The number of patients experiencing events, or event-to-death are not statistically significant

among the groups (figure 3 & 4).

Sub-group analysis: In this study 8-day mortality rate in 3 different sub-groups are shown in (**table 5**), sub-groups are divided as per efficiency toward chemotherapy cycles. Event-to-death occurrence rate, caused by leukocytosis, and thrombocytosis were absent in diabetic chemotherapy responder patients. No statistical significance observed in event-to-death between the sub-groups non-responders, moderate-responders, and responders between control Vs metformin groups. Diabetic patients not responding (n = 188) well towards chemotherapy were not at higher risk of death due to events related to myelosuppression and myeloupregulation compared to diabetic responders (n = 32). To confirm this event-to-death in all three chemotherapy groups of control and metformin are added (**table** 6) further divided by the total number of patients, as no statistical significance is observed in mortality rate between the control and treatment groups (figure 5). Total event-to-death in leukocytosis, neutrophilic leukocytosis, and thrombocytosis were significantly high (*p*-value < 0.05) in non-responder compared to responder group (table 6).

DISCUSSION

In a recent randomized controlled trial (RCT), lung cancer patients showed no survival benefit from metformin and chemoradiation compared to patients treated with chemoradiation therapy [16]. In this study there is no overall survival benefit in diabetic NSCLC patients receiving metformin and chemotherapy compared to patients treated with only chemotherapy. This study found diabetic chemoradiation recipients survived significantly longer compared to patients treated with chemotherapy, because most of diabetic patients received radiation therapy at the end stage and here survival was calculated post-chemotherapy start. Kim et al. reported maximum survival benefit in metformin treated squamous cell carcinoma patients, here adenocarcinoma patients are maximum [27]. It is confirmed in this study patients diagnosed with diabetes 6 months in prior to chemotherapy were less prone to fatality receiving chemotherapy, similar retrospective findings previously reported by Lee et al. in Asian diabetic NSCLC patients [28]. Patients responding well toward chemotherapy regimen were less prone to infections and bleeding caused by hematological events. These findings suggest that all responders blood cell sub-population were maintained within moderate range for prolong time post cancer diagnosis. Anti-tumor T-lymphocyte production and activation to eliminate tumor debris post-chemo, from the tumor site and in blood/lymph circulation possibly results in

increased WBC activation [29]. Chemoresistance is one of the main causes of deaths in non-responder NSCLC patients, at the same time results in fluctuations of blood cell counts due to chemotherapy initiation are more likely related to cardiac events or infections [6, 19, 29]. In a prospective study recruiting cervical cancer patients, found leukopenia as one of the leading causes of deaths after grade 2+ leukopenia events [7]. Prognosis of non-responders is one of the major challenges at present, identifying these patients as non-responders in prior to initiation of chemotherapy would benefit the overall survival of these patients. Several studies reported numerous biological markers, and cancer stage to be a good prognostic factor for survival, at present very few studies are available on prognosis of non-responder patients post cancer diagnosis prior to chemotherapy initiation [30]. Early detection of non-responders would allow selection of low dose metronomic chemotherapeutic regimen that would benefit patient's treatment outcome. In this study NSCLC patients lived shorter duration post first-line chemotherapy in non-responders group, and these patients are usually withdrawn from chemotherapeutic regimen due to various adverse events further resulted in cancer progression.

In control group patients, the frequency of event occurrence was significantly low compared to diabetic patients, withdrawal of metformin for short duration might reduce the event occurrence rate, further study is necessary to confirm [27, 31]. Here it is found that a chemotherapy responder or non-responder could be a diabetic or non-diabetic patient. Several hematological events could be distinct in three different stratified chemotherapy frequency/cycle sub-population of NSCLC patients, no significant number of patients were found to be affected severely within stratified groups (table 5). To repurpose a drug the event-to-death caused by chemotherapeutic regimen initiated against cancer should be significantly low in treatment group compared to control, irrespective of chemotherapy response. At present there is no study available to confirm hematological marker assisted drug repurposing, mostly overall survival and disease free survival were major focus of all RCTs (32). Future studies should focus on hematological biomarkers for drug repositioning, mainly WBC counts. Prospectively designed clinical studies recruits non-diabetic cancer patients to study survival benefit from chemotherapeutic regimen concomitantly treated with metformin or other non-chemotherapeutic drugs considered for repositioning against cancer, at the same time effect on hematological markers due to prolong use of metformin in treatment-arm are not monitored (33). This study found that during the 2nd year post chemotherapy the number of events reduced significantly in metformin group for most of the events except thrombocytopenia.

Chemotherapy induced thrombocytosis is a biomarker for bleeding, or severe infections may lead to internal bleeding in patients (34-35). All the event-to-death were chemotherapy induced, and deaths were mainly caused due to infections and bleeding were significantly higher in non-responder group (table 6). Increased WBC and neutrophils counts indicates infection or body's natural attempt to recover chemotherapy induced immunosuppression; neutropenia manifests sepsis, are related to chemotherapy use (36-37).

One of the main limitations of this study is one-to-many (1:M) PSM, this allowed to overcome the limitation of selection bias, and patients were not excluded from analysis post PSM. The types of chemotherapeutic agents are not provided in this study, beside that concomitant use of other drugs, and other comorbidities of patients were not included in the baseline or matched patients. In hospital, patients are generally treated with GM-CSF prior to the start or end of chemotherapy to reduce the risk of decreased WBC counts and related events that was not considered at baseline. In this study patient's economic background, smoking, drinking, and health insurance status are important confounding factors were not taken into consideration. It is shown for the first time that post-PSM, segregating patients on the basis of chemotherapy cycles, it is possible to distinguish the safety of secondary concomitant drug, from adverse events caused by primary therapeutic agents (chemotherapy), as metformin is a long-term prescribed drug. Decades ago, clinicians segregated patients as per chemotherapy response, later was found not feasible for adoption in trials due to lack of randomization (38). Therefore, survival benefit of metformin group compared to control is not shown using Kaplan-Meier plot in responders, moderate responders, and non-responders. Moreover, there is no difference in survival between control and treated patients in these sub-populations.

All together metformin use is safe for diabetic NSCLC patients and the event related to death were not associated with metformin use. During the 1^{st} year post chemotherapy number of events increased but did not contribute to death of diabetic patients. Further study is required to understand the reason behind this trend in metformin treated diabetic patients not found in non-diabetic control patients. In previous studies, mortality was not related to neutropenia or thrombocytopenia in cancer patients treated with metformin [35, 39-40]. In contrast this study found total neutropenic events post chemotherapy were significantly higher than control, but metformin did not contribute to high mortality rate. Total leukopenia, neutropenia, and thrombocytopenia events total occurrence rate were significantly higher post chemotherapy but significantly reduced in the 2^{nd} year, possibly due to prolonged use of metformin possibly

reduced risk of hematological events. This fact could be further confirmed in overall survival benefit of NSCLC patients diagnosed with diabetes (< 180) days prior to chemotherapy initiation. Diabetic patients are often at risk of low platelet count due to high glucose level in blood. In overall diabetic patients metformin could not be considered as a major causative agent of thrombocytopenia or associated deaths in NSCLC patients. Here hematological events associated with chemotherapy or cancer progression resulted in event related deaths of NSCLC patients. Metformin attributed to increased event occurrence rate did not cause increased deaths caused by these events, suggesting metformin use is safe, withdrawal of metformin during event occurrence during cancer treatment regimen may increase therapeutic efficacy. The efficacy of metformin in survival outcome of NSCLC patients is beneficial for patients diagnosed with diabetes and treated prior to chemotherapy initiation. Metformin could be beneficial in other cancer types comorbid with diabetes, significant survival benefit could not be observed in NSCLC patients for metformin to be repositioned against NSCLC.

Acknowledgement

The author would like to thank professor Kyungsoo park (Department of pharmacology, Yonsei University) for providing access to severance hospital database for obtaining patient's electronic health records.

Ethics Statement

I confirm all relevant ethical guidelines have been followed, and any necessary IRB and/or ethics committee approval was obtained. The studies involving human participants were waived by the Institutional review board (IRB) of Yonsei University. Written informed consent form for participation was not required for this study in accordance with the national legislation and the institutional requirements.

Funding

No funding was received for this study.

Conflict of Interest

The author declare that no conflict of interest exists.

Data availability:

Data will be made available upon request, post-approval from Yonsei University IRB.

REFERENCE

1. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021;71(3):209-249. doi:10.3322/caac.21660

2. Dhodapkar M, Zhang AD, Puthumana J, Downing NS, Shah ND, Ross JS. Characteristics of Clinical Studies Used for US Food and Drug Administration Supplemental Indication Approvals of Drugs and Biologics, 2017 to 2019. JAMA Netw Open. 2021 Jun 1;4(6):e2113224. doi: 10.1001/jamanetworkopen.2021.13224.

3. Pushpakom S, Iorio F, Eyers PA, Escott KJ, Hopper S, Wells A, Doig A, Guilliams T, Latimer J, McNamee C, Norris A, Sanseau P, Cavalla D, Pirmohamed M. Drug repurposing: progress, challenges and recommendations. Nat Rev Drug Discov. 2019 Jan;18(1):41-58. doi: 10.1038/nrd.2018.168.

4. Edelman MJ, Wang X, Hodgson L, Cheney RT, Baggstrom MQ, Thomas SP, Gajra A, Bertino E, Reckamp KL, Molina J, Schiller JH, Mitchell-Richards K, Friedman PN, Ritter J, Milne G, Hahn OM, Stinchcombe TE, Vokes EE; Alliance for Clinical Trials in Oncology. Phase III Randomized, Placebo-Controlled, Double-Blind Trial of Celecoxib in Addition to Standard Chemotherapy for Advanced Non-Small-Cell Lung Cancer with Cyclooxygenase-2 Overexpression: CALGB 30801 (Alliance). J Clin Oncol. 2017 Jul 1;35(19):2184-2192. doi: 10.1200/JCO.2016.71.3743.

5. Hao L, Zhang J, Di Y, Tan Z. Prognostic Value of White Blood Cells Detected for the First Time After Adjuvant Chemotherapy in Primary Operable Non-Small Cell Lung Cancer. *Technol Cancer Res Treat.* 2018;17:1533033818802813. doi:10.1177/1533033818802813

6. Skoetz N, Bohlius J, Engert A, Monsef I, Blank O, Vehreschild JJ. Prophylactic antibiotics or G(M)-CSF for the prevention of infections and improvement of survival in cancer patients receiving myelotoxic chemotherapy. *Cochrane Database Syst Rev.* 2015;2015(12):CD007107. Published 2015 Dec 21. doi:10.1002/14651858.CD007107.pub3

7. Bogani G, Sabatucci I, Maltese G, et al. Chemotherapy-related leukopenia as a biomarker predicting survival outcomes in locally advanced cervical cancer. *Eur J Obstet Gynecol Reprod Biol.* 2017;208:41-45. doi:10.1016/j.ejogrb.2016.11.017

8. Huang WM, Cheng HM, Huang CJ, et al. Hemographic indices are associated with mortality in acute heart failure. *Sci Rep.* 2017;7(1):17828. doi:10.1038/s41598-017-17754-8

9. Alenzi EO, Kelley GA. The association of hyperglycemia and diabetes mellitus and the risk of chemotherapy-induced neutropenia among cancer patients: A systematic review with meta-analysis. *J Diabetes Complications.* 2017;31(1):267-272. doi:10.1016/j.jdiacomp.2016.09.006

10. Shahid RK, Ahmed S, Le D, Yadav S. Diabetes and Cancer: Risk, Challenges, Management and Outcomes. *Cancers (Basel).* 2021;13(22):5735. Published 2021 Nov 16. doi:10.3390/cancers13225735

11. Carobbio A, Antonioli E, Guglielmelli P, et al. Leukocytosis and risk stratification assessment in essential thrombocythemia. *J Clin Oncol.* 2008;26(16):2732-2736. doi:10.1200/JCO.2007.15.3569

12. Hillson R. Diabetes and the blood–white cells and platelets. Pract Diabetes. 2015;32(5):159–60.

13. Xiao K, Liu F, Liu J, Xu J, Wu Q, Li X. The effect of metformin on lung cancer risk and survival in patients with type 2 diabetes mellitus: A meta-analysis. *J Clin Pharm Ther.* 2020;45(4):783-792.

doi:10.1111/jcpt.13167

14. Xiao K, Liu F, Liu J, Xu J, Wu Q, Li X. The effect of metformin on lung cancer risk and survival in patients with type 2 diabetes mellitus: A meta-analysis. *J Clin Pharm Ther*. 2020;45(4):783-792. doi:10.1111/jcpt.13167

15. Hung MS, Chuang MC, Chen YC, et al. Metformin Prolongs Survival in Type 2 Diabetes Lung Cancer Patients With EGFR-TKIs. *Integr Cancer Ther*. 2019;18:1534735419869491. doi:10.1177/1534735419869491

16. Tsakiridis T, Pond GR, Wright J, et al. Metformin in Combination with Chemoradiotherapy in Locally Advanced Non-Small Cell Lung Cancer: The OCOG-ALMERA Randomized Clinical Trial. *JAMA Oncol*. 2021;7(9):1333-1341. doi:10.1001/jamaoncol.2021.2328

17. Sayed R, Saad AS, El Wakeel L, Elkholy E, Badary O. Metformin Addition to Chemotherapy in Stage IV Non-Small Cell Lung Cancer: an Open Label Randomized Controlled Study. *Asian Pac J Cancer Prev*. 2015;16(15):6621-6626. doi:10.7314/apjcp.2015.16.15.6621

18. Arrieta O, Varela-Santoyo E, Soto-Perez-de-Celis E, et al. Metformin use and its effect on survival in diabetic patients with advanced non-small cell lung cancer. *BMC Cancer*. 2016;16:633. Published 2016 Aug 12. doi:10.1186/s12885-016-2658-6

19. Arruda-Olson AM, Reeder GS, Bell MR, Weston SA, Roger VL. Neutrophilia predicts death and heart failure after myocardial infarction: a community-based study. *Circ Cardiovasc Qual Outcomes*. 2009;2(6):656-662. doi:10.1161/CIRCOUTCOMES.108.831024

20. Lin JJ, Gallagher EJ, Sigel K, et al. Survival of patients with stage IV lung cancer with diabetes treated with metformin. *Am J Respir Crit Care Med*. 2015;191(4):448-454. doi:10.1164/rccm.201407-1395OC

21. Connolly GC, Khorana AA, Kuderer NM, Culakova E, Francis CW, Lyman GH. Leukocytosis, thrombosis and early mortality in cancer patients initiating chemotherapy. *Thromb Res*. 2010;126(2):113-118. doi:10.1016/j.thromres.2010.05.012

22. Jameus A, Kennedy AE, Thome C. Hematological Changes Following Low Dose Radiation Therapy and Comparison to Current Standard of Care Cancer Treatments. *Dose Response*. 2021;19(4):15593258211056196. Published 2021 Nov 15. doi:10.1177/15593258211056196

23. Sempere-Bigorra M, Julián-Rochina I, Cauli O. Chemotherapy-Induced Neuropathy and Diabetes: A Scoping Review. *Curr Oncol*. 2021;28(4):3124-3138. Published 2021 Aug 19. doi:10.3390/curroncol28040273

24. Shitara K, Ikeda J, Kondo C, et al. Reporting patient characteristics and stratification factors in randomized trials of systemic chemotherapy for advanced gastric cancer. *Gastric Cancer*. 2012;15(2):137-143. doi:10.1007/s10120-011-0083-8

25. Jang HY, Kim I-W and Oh JM (2022) Cysteinyl Leukotriene Receptor Antagonists Associated With a Decreased Incidence of Cancer: A Retrospective Cohort Study. Front. Oncol. 12:858855. doi: 10.3389/fonc.2022.858855

26. MedCalc Software Ltd. Comparison of proportions calculator. Website: https://www.medcalc.org/calc/comparison_of_proportions.php (version 23.2.1) [Accessed Date: 18 Oct, 2024].

27. Kim J, Bae YJ, Lee JW, et al. Metformin use in cancer survivors with diabetes reduces all-cause mortality, based on the Korean National Health Insurance Service between 2002 and 2015. *Medicine (Baltimore)*. 2021;100(11):e25045. doi:10.1097/MD.0000000000025045

28. Lee Y, Joo J, Lee YJ, et al. Randomized phase II study of platinum-based chemotherapy plus controlled diet with or without metformin in patients with advanced non-small cell lung cancer. *Lung Cancer*. 2021;151:8-15. doi:10.1016/j.lungcan.2020.11.011

29. Daniela S. Thommen, T Cell Dysfunction in Cancer. Cancer Cell. 2018 Apr 9;33(4):547-562.

30. Arrieta O, Aviles-Salas A, Orozco-Morales M, et al. Association between CD47 expression, clinical characteristics and prognosis in patients with advanced non-small cell lung cancer. *Cancer Med*. 2020;9(7):2390-2402. doi:10.1002/cam4.2882

31. Kim YJ, Jung SM, Kang J, et al. Risk factors for extended-spectrum beta-lactamase-producing Enterobacteriaceae infection causing septic shock in cancer patients with chemotherapy-induced febrile neutropenia. *Intern Emerg Med*. 2019;14(3):433-440. doi:10.1007/s11739-018-02015-x

32. Nechushtan H, Hamamreh Y, Nidal S, el al. A phase IIb trial assessing the addition of disulfiram to chemotherapy for the treatment of metastatic non-small cell lung cancer. Oncologist. 2015;20(4):366-367. Doi: 10.1634/theoncologist.2014-0424.

33. Yu-OHY, Suissa S. Metformin and Cancer: Solutions to a Real-World Evidence Failure. Diabetes Care. 2023;1;46(5):904-912. Doi: 10.2337/dci22-0047.

34. Gao A, Zhang L, Zhong D. Chemotherapy induced thrombocytopenia: literature review. Discov Oncol. 2023;14(1):10. Doi: 10:1007/s12672-023-00616-3.

35. Kristensen DT, Ovlisen AK, Jakobsen LHK, et al. Metformin Use and risk of myeloproliferative neoplasms: a Danish population-based case-control study. Blood Adv. 2024;8(16):4478-4485. Doi: 10.1182/bloodadvances.20233012266.

36. Yin F, Xi YL, Wang Y, et al. The clinical outcomes and biomarker features of severe sepsis/septic shock with severe neutropenia: a retrospective cohort study. Transl Pediatr. 2021;10(3):464-473. Doi: 10.21037/tp-20-230.

37. Huang X, Nepovimova E, Adam V. et al. Neutrophils in Cancer immunotherapy: friends or foes?. Mol Cancer 23, 107 (2024). Doi: 10.1186/s12943-024-02004-z

38. Gary B Weiss, Harvey Bruce, James A Hoakanson, Comparing Survival of Responders and Nonresponders after treatment: A potential Source of Confusion in Interpreting Cancer Clinical Trials. Controlled Clinical Trials 4:43 (1983)

39. Serageldin MA, Kassem AB, El-kerm Y, Helmy MW, El-Mas MM, El-Bassiouny NA. The effect of Metformin on Chemotherapy-induced Toxicities in Non-diabetic Breast Cancer Patients: A Randomized Controlled Study. Drug Saf. 2023;46(6):587-599. Doi: 10.1007/s40264-023-01305-4

40. Baaten CCFMJ, Moenen FCJI, Henskens YMC, et al. Impaired mitochondrial activity explains platelet dysfunction in thrombocytopenic cancer patients undergoing chemotherapy. Haematologica. 2018;103(9):1557-1567. Doi: 10.3324/haematol.2017.185165.

Figure 1: Study flowchart to identify myelosuppression and myeloupregualtion in Metformin treated/untreated Lung cancer patients. Three sub-groups were stratified based on chemotherapy frequency/cycle post propensity score matching to segregate chemotherapy response.

Variables	Number (%)
Gender	
Male	1260 (68.07)
Female	591 (31.9)
Age	63 (19 - 99)
Weight	61.6 (32 - 110)
Stage	
III	360 (19.4)
IV	1440 (77.5)
recurred	51 (2.7)
Cancer type	
Non-small cell lung cancer	204 (11)
Adenocarcinoma	923 (49.8)
Squamous cell	289 (15.6)
Pancoast	2 (0.1)
Pulmonary blastoma	1 (0.05)
unspecified	432 (23.3)
Therapeutic Intervention	
Surgery	858 (46.35)
Radiotherapy	682 (36.8)
Salvage	365
Blood cell types (mean)	
WBC	9.0
RBC	4.2
Platelet	290.7
Neutrophil	6.3
Albumin	3.9
Diabetic patients	354 (19)
Immunosuppressants	545 (29.4%)

Table 1: Pre-propensity score match baseline characteristics of NSCLC patients (n = 1851).

Metformin Start pre/post Chemo	No. of patients (%)	Mean Days (SD 95% CI)	Hazard ratio
<180 Days pre-chemo (Group A)	23 (8)	1296 (920 95% CI 544 – 1295) *	0.42
180 days pre/post chemo (Group B)	81 (58)	670 (828 95% CI 489 – 850)	2.05
>180 Days post-chemo (Group C)	32 (17)	765 (645 95% CI 535 – 976)	1.67

Table 2: Sub-group survival analysis as per diabetes diagnosis and metformin treatment initiation in NSCLC patients. (*) represents statistical significance (*p*-value < 0.05) compared to Group B & Group C patients.

	Leukopenia	Leukocytosis	Neutropenia	Neutrophilic leukocytosis	Thrombocytopenia	Thrombocytosis
Total events pre/post chemotherapy, events (event %) (n = number of patients)						
Control (n = 354)	2451 (46.25%) (n = 264)	3951 (50.2%) (n = 310)	1704 (46.45%) (n = 244)	5007 (48.2%) (n = 334)	4581 (54.7%)* (n =260)	1596 (47.5%) (n = 202)
Metformin (n = 354)	2848 (53.75%)* (n = 267)	3923 (49.8%) (n = 313)	1964 (53.5%)* (n = 248)	5373 (51.8%)* (n = 337)	3790 (45.3%) (n =270)	1764 (52.5%)* (n = 206)
Total events post chemotherapy initiation, events (event %) (n = number of patients)						
Control (n = 354)	2168 (48.2%) (n =252)	3259 (52.3%)* (n = 282)	1469 (48.2%) (n = 223)	4092 (50.2%) (n = 309)	3358 (47.9%) (n = 245)	1227 (49.7%) (n = 175)
Metformin (n = 354)	2330 (51.8%)* (n = 255)	2972 (47.7%) (n = 287)	1581 (51.8%)* (n = 229)	4064 (49.8%) (n = 320)	3651 (52.1%)* (n = 259)	1239 (50.2%) (n = 174)
Post chemotherapy initiation 1st year, total events (event %) (n = number of patients)						
Control (n = 354)	1456 (49.8%) (n = 228)	1899 (50.8%) (n = 217)	961 (51.3%) (n = 205)	2324 (49.7%) (n = 250)	2029 (48%) (n =213)	761 (47.8%) (n = 143)
Metformin (n = 354)	1468 (50.2%) (n = 228)	1833 (49.2%) (n = 231)	910 (48.7%) (n = 206)	2348 (50.2%) (n = 273)	2196 (52%)* (n = 224)	830 (52.2%) (n = 140)
Post chemotherapy initiation 2nd year, total events (event %) (n = number of patients)						
Control (n = 354)	399 (53.9%)* (n = 67)	508 (56.7%)* (n = 75)	276 (52.5%) (n = 58)	743 (55.7%)* (n = 93)	590 (53%)* (n =72)	195 (45.4%) (n = 41)
Metformin (n = 354)	341 (46.1%) (n = 64)	389 (43.3%) (n =76)	249 (47.4%) (n = 53)	590 (44.3%) (n = 90)	522 (47%) (n = 71)	234 (54.6%)* (n = 47)

Table 3: Hematological events occurrence rate in NSCLC patients. Total events occurred at different timelines post cancer diagnosis, post chemotherapy start, compared between control and treated groups. (*) represents statistical significance (p-value < 0.05).

	Leukopenia	Leukocytosis	Neutropenia	Neutrophilic leukocytosis	Thrombocytopenia	Thrombocytosis
Total event-to-death pre/post chemotherapy, N = number of fatality (Mortality rate %)						
Control (n = 354)	N = 37 (10.4%)	N = 153 (43.2%)	N = 25 (7%)	N = 168 (47.4%)	N = 112 (31.6%)	N = 26 (7.3%)
Metformin (n = 354)	N =29 (8.1%)	N = 143 (40.3%)	N = 21 (5.9%)	N = 173 (48.8%)	N = 119 (33.6%)	N = 23 (6.4%)
Total event-to-death post chemotherapy initiation, n = number of fatality (Mortality rate %)						
Control (n = 354)	N = 37 (10.4%)	N = 153 (43.2%)	N = 25 (7%)	N = 168 (47.4%)	N = 112 (31.6%)	N = 26 (7.3%)
Metformin (n = 354)	N =29 (8.1%)	N = 143 (40.3%)	N = 21 (5.9%)	N = 173 (48.8%)	N = 119 (33.6%)	N = 23 (6.4%)
Post chemotherapy initiation 1st year event-to-death, N = number of fatality (Mortality rate %)						
Control (n = 354)	N =23 (6.4%)	N = 96 (27%)	N = 15 (4.2%)	N = 103 (29%)	N = 65 (18%)	N = 14 (3.9%)
Metformin (n = 354)	N = 18 (5%)	N = 81 (22.8%)	N = 10 (2.8%)	N = 95 (26.8%)	N = 71 (20%)	N = 12 (3.3%)
Post chemotherapy initiation 2nd year, total event-to-death, N = number of fatality (Mortality rate %)						
Control (n = 354)	N = 7 (2%)	N = 32 (9%)	N = 6 (2%)	N = 33 (9.3%)	N = 25 (7%)	N = 6 (2%)
Metformin (n = 354)	N = 4 (1%)	N = 20 (38%)	N = 2 (0.5%)	N = 29 (8.1%)	N = 17 (5%)	N = 6 (2%)

Table 4: 8-day standard mortality rate post event occurrence within 1st year and 2nd year post-chemotherapy initiation in (n = 708) NSCLC diabetic (metformin treated) Vs non-diabetic (untreated/control) patients. No statistical significance observed between the groups.

Events	Non-responder (n = 380)		Moderate-responder (n = 270)		Responder (n = 58)	
(Total patients n = 708)	Metformin (n=188)	Control (n=192)	Metformin (n=134)	Control (n=136)	Metformin (n=32)	Control (n=26)
Leukopenia	14 (7.4%)	20 (10.4%)	5 (3.7%)	9 (6.6%)	3 (9.3%)	1 (3.8%)
Leukocytosis	66 (35.1%)	79 (41.1%)	35 (26.1%)	45 (33%)	0	4 (15.3%)
Neutropenia	7 (3.7%)	13 (6.7%)	3 (2.2%)	7 (5.1%)	2 (6.25%)	1 (3.8%)
Neutrophilic Leukocytosis	78 (41.48%)	82 (42.7%)	45 (33.5%)	51 (37.5%)	1 (3.1%)	3 (11.5%)
Thrombocytopenia	58 (30.8%)	59 (30.7%)	27 (20.1%)	28 (20.5%)	3 (9.3%)	3 (11.5%)
Thrombocytosis	12 (6.3%)	10 (5.2%)	6 (4.4%)	9 (6.6%)	0	1 (3.8%)

Table 5: 8-day standard mortality rate post-chemotherapy in non-responder, moderate-responder and chemotherapy responders within 2 years of treatment initiation. Non-significant statistical difference observed between the three groups.

Events (Total patients n = 708)	Total events Non-responder (n = 380)	Total events Moderate-responder (n = 270)	Total events Responder (n = 58)
Leukopenia	34 (9%)	14 (5%)	4 (7%)
Leukocytosis	145 (38%) *	80 (29%)	4 (7%)
Neutropenia	20 (5%)	10 (4%)	3 (5%)
Neutrophilic Leukocytosis	160 (42%) *	96 (36%)	4 (7%)
Thrombocytopenia	117 (31%) *	55 (20%)	6 (10%)
Thrombocytosis	22 (6%)	15 (6%)	1 (2%)

Table 6: Total event-to-death in non-responder, moderate responder and responder sub-groups. (*) represents statistical significance (*p*-value < 0.05) comparison between non-responder Vs responder.