Who would take part in a pandemic preparedness

cohort study? The role of vaccine-related affective

polarisation: cross-sectional survey

Aziz Mert Ipekci[1,2,3], Eva Maria Hodel[1,3], Maximilian Filsinger[3,4,5], Selina Wegmüller[3,6,7],

Simone Schuller[3,8], Markus Freitag[3,4], Annika Frahsa[1,3], Gilles Wandeler[3,9], Nicola Low[1,3]*

1. Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

2. Graduate School for Health Sciences, University of Bern, Bern, Switzerland

3. Multidisciplinary Center for Infectious Diseases, University of Bern, Bern, Switzerland

4. Institute of Political Science, University of Bern, Bern, Switzerland

5. European School of Political and Social Sciences, Université Catholique de Lille, Lille,

France

6. Department of Clinical Research, CTU Bern, University of Bern, Bern, Switzerland

7. Institute of Philosophy, University of Bern, Bern, Switzerland

8. Department of Clinical Veterinary Medicine, University of Bern, Switzerland.

9. Department of Infectious Diseases, Bern University Hospital, University of Bern, Bern,

Switzerland

* Corresponding author e-mail: nicola.low@unibe.ch (NL)

Abstract

Cohort studies are an important tool for public health research about emerging infectious diseases. Willingness to participate in research is associated with multiple factors. Little is known about the role of people's feelings, especially in the aftermath of the COVID-19 pandemic. Affective polarisation, an affinity for people with similar attitudes to oneself and hostility toward those with opposing views, is a measure of feelings about issues, including COVID-19 vaccination. The aim of this study was to investigate factors associated with willingness to participate in a future household-based cohort study on pandemic preparedness. We did a cross-sectional online survey in the Canton of Bern, Switzerland. We invited a random sample of persons aged 18+ years from 15,000 private households. We asked about willingness to take part and collected data about demographic, social and household characteristics, and affective polarisation related to COVID-19 vaccination. We performed univariable and multivariable analyses, weighted by household size to estimate odds ratios (with 95% confidence intervals, CI). Among responders, 49.8% (95% CI 47.9-51.8, weighted proportion) were willing to take part in a cohort study. In multivariable analysis, higher educational level (adjusted odds ratio 2.48, 95% CI 1.81-3.39) and higher monthly income (1.92, 1.39-2.65) were most strongly associated with higher willingness to participate. Opposition to COVID-19 vaccination was associated with lower willingness to participate (0.53, 0.39-0.72). Affective polarisation modified the relationship between vaccination attitudes and willingness to participate. Compared with non-polarised vaccination supporters, polarised supporters were more willing to participate (marginal adjusted odds ratio 1.51, 1.05-2.16), whereas willingness to participate was lower among both non-polarised (0.53, 0.32-0.86) and affectively polarised (0.26, 0.12-0.56) vaccination opposers. Willingness to participate in a cohort study was moderate. Following the COVID-19 pandemic, addressing affective polarisation, as well as socioeconomic factors, is needed to improve participation in pandemic preparedness research.

Introduction

Cohort studies are an important tool for public health research about emerging infectious diseases and pandemics [1]. At the start of the COVID-19 pandemic in early 2020, policy makers needed to know the proportion of the population who had already been exposed to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) to plan appropriate prevention measures. Researchers in Switzerland were among the first to provide this information, by contacting participants in an existing cohort study on non-communicable diseases and obtaining blood samples for SARS-CoV-2 antibody testing [2]. Seroprevalence was 10%, showing that most people were still susceptible to SARS-CoV-2, guiding the need for ongoing measures to prevent transmission in the population with the minimum of restrictions [2]. In anticipation of the next pandemic, global health experts have proposed that 'pandemic preparedness cohorts' can establish the research infrastructure and then pivot quickly to collect relevant data when a new pandemic pathogen starts to spread [1]. New cohort studies in this field are needed because existing studies, established for other reasons, have not necessarily collected the data needed to improve understanding about behaviours and exposures associated with susceptibility, transmission and immunity of emerging pandemic pathogens.

People's feelings and concerns about public health issues may influence their willingness to participate in research in a new field and should be considered when designing and starting studies [3]. Documented reasons for longstanding declines in participation in population-based research include limited time, declines in altruism and community involvement in general, and mistrust in science [4]. Previous studies have focused on socioeconomic and demographic factors associated with willingness to participate in health research, and on views about issues such as data privacy [4-6]. The novelty and scale of the COVID-19 pandemic may now have affected attitudes towards research because of the unprecedented amounts of new research findings [7], awareness about research [8], and uncertainties about communication, and public trust between scientists, policymakers and the public [9-11].

Political polarisation, according to ideology, partisan affiliation or trust in political institutions,

has characterised responses to many aspects of COVID-19 prevention [12, 13]. The concept

of affective polarisation, defined as an affinity for people with similar attitudes to oneself and

hostility toward those with opposing views, has been used as a quantitative method [14].

Whilst affective polarisation was first defined and studied in the United States of America to

understand [14], the COVID-19 pandemic revealed associations between affective

polarisation and health behaviours such as COVID-19 vaccination uptake, adherence to

pandemic countermeasures or anxiety about COVID-19 [14-18]. Measures of people's

feelings about public health measures such as COVID-19 vaccination, and how they view

others, are more proximate to health-associated behaviours than partisan or ideological

differences. Whether affective polarisation is associated with willingness to take part in

research about pandemics more generally is not well understood. The objectives of this

study were: 1) to investigate sociodemographic factors associated with willingness to

participate in a planned pandemic preparedness cohort study; 2) to explore associations

between COVID-19 vaccination-related affective polarisation and willingness to participate;

and 3) to describe reasons for taking part, or not taking part, in a pandemic preparedness

cohort study.

Materials and methods

We did a cross-sectional survey in the canton of Bern, Switzerland. Bern, population

1,070,898, is the largest of the 26 Swiss cantons geographically, with demographic

characteristics similar to those of Switzerland as a whole. The survey was part of the

preparation for a population-based pandemic preparedness cohort study, BEready ("Bern,

get ready"). The BEready cohort will study whole households [19], including pets [20],

because of the importance of within household transmission of infectious agents [21] and a

One Health approach to understand and prepare for pandemics which may spread between

humans and animals [22]. The Cantonal Ethics Committee in Bern provided an exemption

from ethical approval under the Human Research Act because the survey did not ask for any

personal health-related information (BASEC-Nr: Req-2022-00877). We report this study

following the Strengthening the Reporting of Observational Studies in Epidemiology

guideline for cross-sectional studies (research checklist) (S1 Table).

Study sample

We contacted the Cantonal Administration and Information Office to obtain a random sample

of private household addresses in the canton of Bern. The requested sample of 15,000

households was the same as that of a previous online study about willingness to participate

in personalised health research in Switzerland [5]. We requested a random sample of 3,000

households, each with 1, 2, 3, 4, or 5+ members, oversampling larger households because

their participation is important for a cohort study that aims to investigate household infection

transmission. One person aged 18 or older from each household was randomly chosen as

the primary contact. The selected household members received a letter by surface mail,

which introduced the issue of pandemic preparedness and about potential participation in

research about existing and future infectious diseases. The letter contained a personal

access code and a link to an online questionnaire. On request, they could receive a printed

version of the survey. They were informed that by filling in the questionnaire, they consented

to participate in the study. Invitation letters were sent on 19[th] September 2022, with two

reminders on 14[th] October and 9[th] November 2022. The survey closed on 18[th] December

2022.

Survey development

The survey questions were taken from published national and international questionnaires,

whenever possible, including the Swiss Health Study pilot [23], a nationwide survey about

personalised health research [24], the COVID-19 social mixing survey [25], the International

Social Survey Programme [26] and a study of COVID-19-associated affective polarisation

[15]. The survey was available in German, French, Italian, and English. A convenience

sample of 10 people speaking the languages of the survey tested the questions and we

made changes to improve readability where necessary.

The primary outcome was self-reported willingness to participate in a long-term cohort study

about infectious diseases, measured on a 4-point scale, "yes", "probably yes", "probably no",

or "no". The survey questions covered: demographic characteristics including age and

gender identity, social, and household characteristics, patterns of social mixing and a pre-

defined list of motivations to participate or not participate in cohort studies (S2 Table).

Opinions about COVID-19 vaccination were measured on a scale from zero (complete

rejection) to 10 (complete support). We measured affective polarisation using a feeling

thermometer [27]. First, we asked participants about their feelings towards people who get

vaccinated from –5 ("very cold") to +5 ("very warm"). Then, we asked the same question

about people who do not get vaccinated. The absolute difference between their answers to

the two questions gave the affective polarisation level on a scale from zero to ten [15].

Patient and public involvement

139 We did not involve patients or the public in the development of the questionnaire used in this

140 study. After the survey was completed, we put together a community advisory group of

141 representatives and members of the public, who reviewed the study findings. Their feedback

142 is informing community engagement strategies to optimise enrolment into the BEready

143 cohort study.

Statistical analysis

We used RStudio for all analyses [28]. To account for the oversampling of larger households,

we performed inverse probability weighting (S3 Table) using the survey package (version

4.4-2) [29]. We defined respondents as those who answered at least half of the survey

questions [5] and included respondents who answered the question about the primary

outcome in statistical analyses. We compared responders and non-responders, based on

the two variables available, household size and main language spoken at home. We used

the municipality of residence to map the proportions of responders willing to participate. We

dichotomised three variables (S2 Table, S1 Fig): 1) the primary outcome variable,

willingness to participate (yes/no), combining responses "yes" and "probably yes", and "no"

and "probably no"; 2) opinion about vaccination (opposition, 0-4, support, 6-10). A response

of 5 was considered as neutral and was treated as missing in the main analysis [15];

affective polarisation (not polarised, 0-4, polarised, 5-10). We grouped variables such as

education and income into a smaller number of categories, following conventions used in

previous publications (S2 Table) [25].

We described our study population characteristics using means (and standard deviation, SD)

or median (and interquartile range, IQR) for continuous variables and frequencies and

percentages for categorical variables. We used logistic regression to estimate univariable

odds ratios (OR and 95% confidence intervals, CI) for the association between reported

willingness to participate and exposure variables. We built a multivariable regression model,

based on knowledge available on the topic and findings from the univariable analyses. The

model included age, gender, education, current work situation, nationality, household income,

household size, location, language, opinion about COVID-19 vaccination, and vaccination-

related polarisation. We compared models using likelihood ratio tests and expressed

associations as adjusted odds ratios, aOR, with 95% CI. To explore affective polarisation

related to opinions about COVID-19 vaccination further, we examined the observed

association between respondents' support or opposition to COVID-19 vaccination and

willingness to participate, according to the presence or absence of affective polarisation. We

included an interaction term in the multivariable model and reported marginal odds ratios

[30]. Lastly, we summarised motivations for participation or non-participation in a cohort

study. Participants could choose multiple reasons.

We conducted three sensitivity analyses: 1) we reclassified people with a score of 5 in the

vaccination opinion scale with either supporters of vaccination or as opposing vaccination; 2)

we tested alternative cut-off points for affective polarisation (not polarised, 0-3; polarised, 4

to 6; heavily polarised, 7-10); 3) we varied the definition of a responder by adjusting the

minimum number of answered survey questions required for inclusion (33% and 66%

considered as responders).

Results

Characteristics of the study population

The 15,000 selected households came from 306/335 municipalities in the Canton of Bern. Overall, 3,425/15,000 (22.8%) people from 273 municipalities responded to at least half of the survey and 3,394 (22.6%) responders replied to the question about willingness to participate in a cohort study (S2 Fig). The highest proportion of responders were from 2-person households (29.2%, 877/3,000), with the lowest proportion from households with 5 or more members (11.7%, 351/3,000) (S4 Table). The distribution of respondents, according to the main language spoken in the household, was similar to the numbers invited (S4 Table). Responses were similar among German-speaking (22.8%, 3,074/13,474) and French-speaking households (21.2%, 312/1,473). Among the remaining households, there were 18.2% (2/11) Italian-speaking and 14.6% (6/41) English-speaking respondents. The percentage of missing values was 0.5% to 4.1% for sociodemographic variables. For the questions related to COVID-19 vaccine, 8.5% and 11.8% did not respond. We did not conduct imputation for missing values.

We present results from the weighted analysis; descriptive results in the unweighted analysis were very similar (S5 Table). Table 1 presents the weighted distribution of the participants' characteristics. Amongst responders, there were more women (256,277, 52.3%) than men (229,354, 46.8%), more than a third (182,256, 37.2%) had tertiary education and two-fifths had a household income between 4,500–9,000 Swiss Francs (205,540, 42.0%).

Sociodemographic factors and willingness to participate in a long-term cohort study

Overall, 1,660/3,394 (unweighted proportion 48.9%, weighted proportion 49.8%, 95% CI 47.9–51.8) of responders reported being willing to participate in a long-term study. Willingness to participate was higher near the city of Bern than in rural areas (Table 1, Fig 1). The highest levels of willingness were among individuals with tertiary education (weighted proportion 62.6%) and those earning more than 9,000 Swiss Francs per month (64.9%). The

lowest levels were seen among individuals with less than compulsory education (31.1%) and

those categorised as being opposed to COVID-19 vaccination (32.6%). Similar proportions

of women and men were willing to participate. We asked respondents whether they would be

willing for their children and pets to participate in a long-term study (Table 1). Among

respondents with children, the willingness for their children to participate was lower (40.0%)

than for themselves (49.8%). Among those with pets, willingness to include pets varied by

species, with the highest participation rates for rodents (58.6%), followed by dogs (47.9%),

cats (46.0%), and rabbits (37.9%).

Fig 1. Proportion of respondents willing to participate, by municipality in the canton of

Bern, Switzerland. The map uses: the swissBOUNDARIES3D dataset and the official

directory of towns and cities both published by the Federal Office of Topography swisstopo

(swisstopo.admin.ch); and the lakes and rivers 2025 GEOM TK dataset of Swiss Federal

Statistical Office (bfs.admin.ch) [31, 32].

Table 1. Self-reported willingness to participate in a long-term study about infectious
diseases, weighted denominator.

		Denominator N[1]		Willingness to participate in cohort study, % (95% confidence intervals)
Number of people		489,973	(100%)	49.8 (47.9 - 51.8)
Age group, years	18-29	44,522	(9.1%)	58.5 (52.9 - 64.0)
	30-64	291,949	(59.6%)	52.5 (50.0 - 54.9)
	64+	153,501	(31.3%)	42.2 (38.5 - 46.0)
Gender	Female	256,277	(52.3%)	50.5 (47.8 - 53.2)
	Male	229,354	(46.8%)	49.2 (46.4 - 52.1)
	Other	1,854	(0.4%)	58.7 (27.2 - 85.8)
	No response	2,488	(0.5%)	25.2 (5.1 - 58.4)
Education level	Compulsory education or less	69,135	(14.1%)	31.1 (26.5 - 36.0)
	Upper secondary	225,091	(46.0%)	46.0 (43.1 - 48.9)
	Tertiary	182,256	(37.2%)	62.6 (59.5 - 65.6)
	Other	4,752	(1.0%)	43.9 (23.1 - 66.4)
	No response	8,740	(1.8%)	33.2 (20.3 - 48.2)
Current work situation	Full-time employee	211,250	(43.1%)	54.9 (51.9 - 57.9)
	Part-time employee	97,113	(19.8%)	55.5 (51.5 - 59.5)

	Not employed	151,874	(31.0%)	41.0 (37.3 - 44.7)
	In education	10,546	(2.2%)	60.7 (51.3 - 69.6)
	Other	6,523	(1.3%)	32.1 (18.7 - 48.0)
	No response	12,666	(2.6%)	27.4 (17.3 - 39.3)
Income, Swiss Francs	<4,500	96,560	(19.7%)	43.6 (38.7 - 48.5)
	4,500 – 9,000	205,540	(42.0%)	51.6 (48.5 - 54.6)
	>9,000	105,130	(21.5%)	64.9 (61.3 - 68.5)
	Other	62,679	(12.8%)	32.7 (27.9 - 37.7)
	No response	20,064	(4.1%)	26.9 (19.0 - 35.8)
Household residents, number	1	185,820	(38.0%)	52.6 (48.7 - 56.4)
	2	170,629	(34.8%)	48.2 (45.0 - 51.5)
	3	54,814	(11.2%)	51.1 (47.6 - 54.7)
	4	53,981	(11.0%)	48.5 (45.1 - 51.8)
	5+	24,729	(5.0%)	40.5 (35.7 - 45.3)
Household location[2]	Urban	278,236	(56.9%)	52.3 (49.6 - 55.0)
	Intermediate	118,778	(24.5%)	48.4 (44.6 - 52.3)
	Rural	90,725	(18.5%)	44.3 (40.0 - 48.7)
Nationality	Swiss	415,354	(84.8%)	50.8 (48.6 - 52.9)
	Foreign	69,632	(14.2%)	45.7 (41.1 - 50.3)
	No response	4,987	(1.0%)	28.0 (13.0 - 47.2)
Language spoken at home	German	428,396	(87.4%)	51.1 (48.9 - 53.3)
	French	45,301	(9.2%)	49.6 (43.2 - 55.9)
	Italian	6,604	(1.4%)	49.4 (32.4 - 66.5)
	English	9,671	(2.0%)	54.2 (42.5 - 65.6)
Opinion about vaccination[3]	For vaccination	322,536	(65.8%)	57.7 (55.3 - 60.1)
	Oppose vaccination	83,395	(17.0%)	32.6 (28.4 - 36.9)
	No response	41,576	(8.5%)	33.3 (27.3 - 39.7)
Vaccination related affective polarisation	Polarised	244,371	(49.9%)	56.8 (54.0 - 59.5)
	Not polarised	187,844	(38.3%)	44.6 (41.5 - 47.6)
	No response	57,758	(11.8%)	37.6 (32.3 - 43.1)

Willingness to participate with children			
	Number with children		**Willingness to participate**
On behalf of their children	-	80,605	(100%)	40.0 (37.0 - 42.9)

Willingness to participate with pets			
	Number with pets		**Willingness to participate**
On behalf of their pets	Any pet	134,046	(100%)	46.1 (42.6 - 49.6)
Type of pet[4]	Dogs	43,521	(32.5%)	47.9 (41.8 - 54.1)
	Cats	90,996	(67.9%)	46.0 (41.7 - 50.3)
	Rabbits	7,066	(5.3%)	37.9 (26.9 - 49.9)
	Rodents	6,686	(5.0%)	58.6 (45.2 - 71.2)
	Others	13,575	(10.1%)	47.8 (37.6 - 58.0)

[1] Weighted number of participants
[2] The sum of percentages for the household location variable does not equal 100% due to missing data that were not classified by the Cantonal Administration and Information Office.
[3] The sum of percentages for the opinions on vaccination does not total 100% because

responses marked as "5", representing a neutral position at the midpoint of the scale, were excluded from the analysis.
[4] The sum of the percentages exceeds 100% because some households reported owning multiple types of pets.

After adjustment for all variables included in the model, willingness to participate was more

likely among participants with the highest educational level (aOR 2.48, 95% CI 1.81 - 3.39)

and those with the highest income (aOR 1.92, 1.39 - 2.65); and was less likely among older

participants (aOR 0.99, 0.98 - 0.99 for a one-year increase) and larger households (5 or

more people vs. 1 person, aOR 0.67, 0.44 - 1.01) (Table 2).

Table 2. Factors associated with willingness to participate, univariable and
multivariable logistic regression.

	Univariable model			Multivariable model[1]		
Characteristic	OR[2]	(95% CI)[2]	p-value	aOR[3]	(95% CI)[2]	p-value
Age, per year	0.99	(0.98 - 0.99)	<0.001	0.99	(0.98 - 0.99)	<0.001
Gender			0.346			0.165
Female	1			1		
Male	0.95	(0.81 - 1.11)		0.9	(0.74 - 1.09)	
Other	1.39	(0.39 - 4.93)		3.13	(0.87 - 11.3)	
No response	0.33	(0.09 - 1.24)		0.46	(0.06 - 3.33)	
Education level			<0.001			<0.001
Compulsory education or less	1			1		
Upper secondary education	1.89	(1.47 - 2.43)		1.59	(1.19 - 2.12)	
Tertiary education	3.70	(2.86 - 4.79)		2.48	(1.81 - 3.39)	
Other	1.74	(0.72 - 4.19)		1.57	(0.49 - 5.05)	
No response	1.10	(0.57 - 2.12)		1.61	(0.75 - 3.43)	
Current work situation			<0.001			0.053
Full-time employee	1			1		
Part-time employee	1.03	(0.84 - 1.25)		1.28	(0.98 - 1.67)	
Not employed	0.57	(0.47 - 0.69)		0.93	(0.70 - 1.23)	
In education	1.27	(0.85 - 1.91)		1.91	(1.15 - 3.15)	
Other	0.39	(0.20 - 0.77)		0.85	(0.40 - 0.79)	
No response	0.31	(0.18 - 0.54)		0.92	(0.49 - 1.76)	
Household income, Swiss Francs			<0.001			<0.001
< 4'500	1			1		
4'500 - 9'000	1.39	(1.11 - 1.75)		1.21	(0.93 - 1.59)	
> 9'000	2.40	(1.87 - 3.08)		1.92	(1.39 - 2.65)	

	OR	(95% CI)	p-value	aOR	(95% CI)	p-value
Other	0.62	(0.46 - 0.83)		0.62	(0.44 - 0.87)	
No response	0.47	(0.29 - 0.75)		0.46	(0.26 - 0.83)	
Number of household members			0.002			<0.001
1	1			1		
2	0.84	(0.69 - 1.03)		0.74	(0.59 - 0.95)	
3	0.94	(0.76 - 1.16)		0.65	(0.50 - 0.85)	
4	0.85	(0.69 - 1.04)		0.50	(0.38 - 0.67)	
5≤	0.61	(0.48 - 0.79)		0.67	(0.44 - 1.01)	
House location			0.008			0.601
Urban	1	1		1	1	
Intermediate	0.86	(0.71 - 1.03)		1.03	(0.83 - 1.27)	
Rural	0.73	(0.59 - 0.89)		0.90	(0.71 - 1.14)	
Nationality			0.013			0.791
Swiss	1	1		1	1	
Foreign	0.81	(0.66 - 1.00)		0.90	(0.65 - 1.25)	
No response	0.38	(0.16 - 0.87)		0.84	(0.30 - 2.31)	
Language			0.907			0.729
German	1	1		1	1	
French	0.99	(0.76 - 1.29)		1.11	(0.80 - 1.54)	
Italian	0.99	(0.51 - 1.92)		1.34	(0.70 - 2.53)	
English	1.19	(0.74 - 1.92)		0.90	(0.48 - 1.67)	
Opinion about COVID-19 vaccination			<0.001			<0.001
Support vaccination	1	1		1	1	
Oppose vaccination	0.35	(0.28 - 0.44)		0.53	(0.39 - 0.72)	
No response	0.37	(0.27 - 0.49)		0.34	(0.07 - 1.69)	
Vaccination-related polarisation			<0.001			<0.001
Not polarised	1	1		1	1	
Polarised	1.64	(1.38 - 1.93)		1.51	(1.20 - 1.89)	
No response	0.75	(0.58 - 0.97)		0.88	(0.47 - 1.67)	

[1] Model includes the following variables: Age, gender, education level, current work situation, income, household size, household location, language, nationality, opposition to vaccination, affective polarisation: vaccination opposer (interaction term) and willingness to participate (outcome)

[2] OR, odds ratio, CI, confidence interval

[3] aOR, adjusted odds ratio

COVID-19 vaccination-related factors and willingness to participate

Most respondents expressed opinions in support of COVID-19 vaccination (65.8%) (Table 1, S1 Fig). In both univariable and multivariable analyses, people who had negative opinions about COVID-19 vaccination were less willing than those who expressed support for vaccination to say they would participate in a long-term cohort study (aOR 0.53, 0.39 - 0.72). Amongst all respondents, people with polarised feelings towards others about COVID-19 vaccination were more willing than those defined as non-polarised to participate (aOR 1.51, 1.20 - 1.89) (Table 2). The direction of the association between COVID-19 vaccination opinions and willingness to participate differed between those who showed affective polarisation and those who were non-polarised (Table 3). The increase in willingness to participate among people with polarised opinions was only observed among those who supported COVID-19 vaccination (aOR 1.51, 1.05 - 2.16). People with opinions opposed to COVID-19 vaccination were less willing to participate than non-polarised supporters of vaccination, particularly those who were affectively polarised (aOR 0.26, 0.12 - 0.56). The three sensitivity analyses did not change the results (S6 Table).

Table 3. Effect modification by affective polarisation status of the association between opinion about COVID-19 vaccination and willingness to participate.

Affective polarisation status	Support vaccination		Oppose vaccination	
	N^2 with outcome	aOR[3] (95% CI)	N^2 with outcome	aOR[3] (95% CI)
Not polarised	97,556	1	56,783	0.53 (0.32 - 0.86)
Polarised	216,219	1.51 (1.05 - 2.16)	21,406	0.26 (0.12 - 0.56)

[1] Adjusted for age, education level, current work situation, income, household size, household location, gender, language spoken, foreign status, polarisation status, vaccine support status
[2] Weighted number of participants
[3] Marginal adjusted odds ratio

Reasons for participation or non-participation in research

People who answered "yes", "probably yes", and "probably no" to the willingness to participate question were asked about their reasons to participate in a long-term study

(n=2,776). People who answered "no" were asked for reasons to decline participation

(n=637) (S7 Table). The most common reasons for participation were: "I can contribute to

the health of fellow human beings" (n=1,559, 56%); "I can contribute to better preparation for

the next pandemic" (n=1,134, 41%), followed by answers such as, "I am interested in

research and health" (n=1,031, 37%), "I will get a free health check" (n=713, 26%) and "I will

receive the study results as feedback" (n=682, 25%). The main reasons to decline were: lack

of interest (n=266, 42%); privacy concerns and mistrust (Fig 2, S7 Table).

Fig 2. Reasons for non-participation, amongst 637 respondents who stated that they
would not be willing to take part in a cohort study.

264 Discussion

In this cross-sectional online survey, about half of respondents reported willingness to take

part in a long-term cohort study for pandemic preparedness, with slightly lower proportions

willing for their children or pets to take part. Older adults, individuals with lower levels of

education and income, and those opposing COVID-19 vaccination were less likely to report

that they would participate. Affective polarisation was associated with increased willingness

to participate among those supporting COVID-19 vaccination, whilst people expressing

opposition to vaccination were less willing to participate, particularly those who were

affectively polarised. The main reasons given for participation in research were to help

others and to improve pandemic preparedness, while lack of interest, privacy concerns and

mistrust were the most stated reasons for non-participation.

275 Strengths and limitations

The design of our study has three main strengths. First, the sample was drawn at random

from a population register, with oversampling from larger households, which account for a

smaller proportion of households in the canton of Bern. Second, we asked about affective

polarisation related to feelings about COVID-19 vaccination, in addition to more traditional

sociodemographic factors. Third, we used questions that had been used and/or validated in

other surveys to facilitate comparisons with other studies. There are three main limitations to

the study design. First, the study participants came from a single canton in Switzerland

because this is the setting for the planned BEready cohort study. Whilst this may limit

generalisability of the study findings, the demographic characteristics of the canton of Bern

are similar to those of Switzerland as a whole. Second, for reasons of efficiency, the

questionnaire was primarily designed to be answered online and was available in only four

languages. The survey was, therefore, not accessible to people without ready access to a

computer and the internet, or to those who could not understand one of the survey

languages; only 184 people asked for a paper version of the questionnaire. Our findings may

therefore be biased if factors limiting participation to those with digital access are also

associated with willingness to participate in research. Third, the number of questions in the

survey was restricted since a lengthy questionnaire might have decreased the response rate

[33]. Whilst there is evidence that political ideology and trust in government institutions are

associated with an individual's willingness to participate in scientific research [34, 35], we

were not able to include questions about a broader range of political variables.

Comparison with other studies

Assessments of the level of willingness to participate in public health research need to take

into account the proportion responding to the online survey. When compared with other

online surveys using random sampling approaches, our study's response rate (23%) was

acceptable in comparison with 2.9% for a survey about precision medicine in South Korea

[36], 16% for a pilot study for national cohort study [23] and 34% for a study about

personalised health research in Switzerland [5]. We could not assess the extent of non-

response bias in detail because only two characteristics were available for comparison; the

proportion of responders according to household language, which was similar to that of the

sample as a whole, and household size, with people from larger households less likely to

respond than those from smaller households.

The proportion of survey participants reporting willingness to participate varies between

studies. About 50% of potential participants in our study were willing to participate, which is

comparable to findings from studies on personalised health research in Switzerland (54%)

[5] and the United States (54%) [37], and clinical research in India (42%) [8]. Similarly, a

large European survey involving 32 countries found that 53% of respondents were willing to

provide personal information to a biobank [38]. In Australia, a telephone survey found that

61% of respondents were willing to participate in health research in general, with willingness

increasing when specific information about the study was provided [39]. In Switzerland, 90%

of respondents to an online survey reported willingness to participate in a national health

cohort study [6]. The researchers suggested this high proportion could reflect a strong civic

duty to support public research, particularly when led by the Federal Office of Public Health

[6]. Indeed, altruism was one of the main reasons, stated by 56% of people who said that

they would be willing to take part in a cohort study about pandemic preparedness.

Willingness to take part in future research might also be overestimated, however, given that

they selectively took part in the online survey. Actual participation may also be lower than

intended participation. In studies in high-income countries, including ours, younger people,

those with higher levels of education and/or higher incomes were more likely to say they

would participate. Gender was not strongly associated with willingness to participate; the

small number of respondents reporting a non-binary gender makes interpretation uncertain.

We did not find any other studies about affective polarisation and willingness to participate in

research.

328   Interpretation of the study findings

Our investigation of affective polarisation as a dimension of willingness to participate in

research adds to the knowledge in this field, particularly in the context of designing research

that might investigate preventive interventions, such as vaccination against COVID-19. The

finding that people who support vaccination were more willing to take part in a long-term

cohort study than people who oppose vaccination is in line with general support for research

by "good citizens" [3]. The modifying effect of affective polarisation provides additional

insight. People who supported COVID-19 vaccination were even more motivated to take part

in research if they also held polarised views about other people's opinions. In contrast, both

polarised and non-polarised opponents of vaccination reported lower willingness to

participate than non-polarised supporters of vaccination. Existing research provides

evidence of a link between affective polarisation and reduced vaccination uptake [40, 41],

lower concern about COVID-19, and decreased support for COVID-19 policies [42]. These

findings align with our study if we consider participation in a long-term cohort study as a

health-related behaviour. Additionally, we found that mistrust and concerns about privacy of

health data were among the top reasons for unwillingness to participate. We did not examine

the association between affective polarisation and trust, but other researchers have found

this in relation to vaccination and political opinions [43-45].

Implications for research and public health

The findings from our study have implications for public health research and policy making.

The strong associations between opinions about COVID-19 vaccination, vaccine-related

affective polarisation and willingness to participate in future studies supports a

multidisciplinary approach to research and its communication [9]. The relationship between

affective polarisation and willingness to participate in public health research should also be

extended to different contexts and countries with varying levels of affective polarisation [15]

to examine the generalisability of our findings. Strategies to overcome under-representation

of people who are older, less well educated, have lower incomes and have more negative

opinions about preventive measures in research for pandemic preparedness are also

needed to reduce the potential for participation bias. Community engagement activities could

raise awareness of the importance of research about infectious diseases and pandemic

preparedness and reduce mistrust about sharing health data. Technological advances,

which reduce barriers to participation, such as lack of time, including the use of online

surveys and mailing of self-collected biological samples might also be beneficial. Policy

makers will benefit from future research about potential pandemic threats, despite the

difficulties of achieving and sustaining high levels of participation [1]. Our study examined

willingness to participate in research after the emergency phase of the COVID-19 pandemic,

which might be affected by pandemic fatigue and shifting attitudes to trust. A study in

Switzerland found that affective polarisation in relation to COVID-19 vaccination was less

marked in 2023 than in 2022 [45]. Longer term study of affective polarisation in pandemic

preparedness cohort studies, such as the BEready cohort, should be conducted to

determine whether, in the next pandemic, early-stage solidarity in a crisis increases

participation. Understanding factors affecting willingness to participate could help develop

future strategies to sustain long-term research engagement across diverse populations.

Acknowledgements

373 The authors would like to acknowledge Jean-Benoît Rossel for his contribution to data

374 preparation, Christiane Pelzer for her support in data management, and Martin Samuel

375 Wohlfender for his help with data visualisation.

Author contributions

AMI, EMH, SW, MaxF, MarF, AF, GW, NL conceptualised the questionnaire survey. EMH,

SW administered the project. AMI, EMH, MaxF, NL developed the methodology for analysis.

AMI, EMH, SW curated the data. AMI analysed the data. AMI, EMH, MaxF, NL drafted the

manuscript. AMI, EMH, MaxF, SW, SS, MarF, AF, GW, NL reviewed and edited the

manuscript. NL supervised the study and is guarantor for the manuscript. All authors read

the final manuscript and agreed to submission.

Financial disclosure statement

384 This work was supported by the Multidisciplinary Center for Infectious Diseases, University

385 of Bern (award number MA21 awarded to AF, MarF, NL and the BEready project, no award

386 number, awarded to NL, GW).

Competing interests

AF is a representative for the German Speaking Network of Participatory Health Research.

All other authors declare no competing interests.

References

References

1. Norton A, Sigfrid L, Aderoba A, Nasir N, Bannister PG, Collinson S, et al. Preparing for a

pandemic: highlighting themes for research funding and practice-perspectives from the

Global Research Collaboration for Infectious Disease Preparedness (GloPID-R). BMC

Med. 2020;18(1):273. Epub 20200908. doi: https://doi.org/10.1186/s12916-020-01755-y.

PubMed PMID: 32895055; PubMed Central PMCID: PMCPMC7476792.

2. Stringhini S, Wisniak A, Piumatti G, Azman AS, Lauer SA, Baysson H, et al.

Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-

POP): a population-based study. Lancet. 2020;396(10247):313-9. Epub 20200611. doi:

https://doi.org/10.1016/s0140-6736(20)31304-0. PubMed PMID: 32534626; PubMed

Central PMCID: PMCPMC7289564.

3. Bühler N, Frahsa A, Morand Bourqui R, Von Götz N, Bochud M, Panese F. Between data

providers and concerned citizens: Exploring participation in precision public health in

Switzerland. Public Underst Sci. 2024;33(1):105-20. Epub 20230717. doi:

https://doi.org/10.1177/09636625231183265. PubMed PMID: 37458286; PubMed

Central PMCID: PMCPMC10756011.

4. Galea S, Tracy M. Participation rates in epidemiologic studies. Ann Epidemiol.

2007;17(9):643-53. Epub 20070606. doi:

https://doi.org/10.1016/j.annepidem.2007.03.013. PubMed PMID: 17553702.

5. Brall C, Berlin C, Zwahlen M, Ormond KE, Egger M, Vayena E. Public willingness to

participate in personalized health research and biobanking: A large-scale Swiss survey.

PLoS One. 2021;16(4):e0249141. Epub 20210401. doi:

https://doi.org/10.1371/journal.pone.0249141. PubMed PMID: 33793624; PubMed

Central PMCID: PMCPMC8016315.

6. Bühler N, Frahsa A, Jaramillo NG, Bourqui RM, Nusslé SG, Zuppinger C, et al.

Willingness to participate in a personalized health cohort – insights from the swiss health

study pilot phase. BMC Public Health. 2024;24(1):2140. doi:

https://doi.org/10.1186/s12889-024-19650-z.

7. Ipekci AM, Buitrago-Garcia D, Meili KW, Krauer F, Prajapati N, Thapa S, et al. Outbreaks

of publications about emerging infectious diseases: the case of SARS-CoV-2 and Zika

virus. BMC Medical Research Methodology. 2021;21(1):50. doi:

https://doi.org/10.1186/s12874-021-01244-7.

8. Raj JP, Tomy SK, Jose A, Kashyap A, Kureethara JV, Kallarakal TK. Knowledge and

perceptions about clinical research and its ethical conduct among college students from

non-science background: a representative nation-wide survey from India. BMJ Public

Health. 2024;2(1):e000748. Epub 20240530. doi: https://doi.org/10.1136/bmjph-2023-

000748. PubMed PMID: 40018094; PubMed Central PMCID: PMCPMC11812855.

9. McKee M, Altmann D, Costello A, Friston K, Haque Z, Khunti K, et al. Open science

communication: The first year of the UK's Independent Scientific Advisory Group for

Emergencies. Health Policy. 2022;126(3):234-44. Epub 20220115. doi:

https://doi.org/10.1016/j.healthpol.2022.01.006. PubMed PMID: 35140018; PubMed

Central PMCID: PMCPMC8760632.

10. Cristea F, Weishaar H, Geurts B, Delamou A, Tan MMJ, Legido-Quigley H, et al. A

comparative analysis of experienced uncertainties in relation to risk communication

during COVID19: a four-country study. Globalization and Health. 2022;18(1):66. doi:

https://doi.org/10.1186/s12992-022-00857-x.

11. Heyerdahl LW, Borzykh Y, Lana B, Volkmann A-M, Crusefalk L, Colman E, et al.

Outsciencing the scientists: a cross-sectional mixed-methods investigation of public trust

in scientists in seven European countries. BMJ Public Health. 2023;1(1):e000280. doi:

https://doi.org/10.1136/bmjph-2023-000280.

12. Klösch B, Hadler M, Reiter-Haas M, Lex E. Polarized opinions on Covid-19 and

environmental policy measures. The role of social media use and personal concerns in

German-speaking countries. Innovation: The European Journal of Social Science

Research. 2023:1-24. doi: https://doi.org/10.1080/13511610.2023.2201877.

13. Grimalda G, Murtin F, Pipke D, Putterman L, Sutter M. The politicized pandemic:

Ideological polarization and the behavioral response to COVID-19. European Economic

Review. 2023;156:104472. doi: https://doi.org/10.1016/j.euroecorev.2023.104472.

14. Iyengar S, Lelkes Y, Levendusky M, Malhotra N, Westwood SJ. The Origins and

Consequences of Affective Polarization in the United States. Annual Review of Political

Science. 2019;22(1):129-46. doi: https://doi.org/10.1146/annurev-polisci-051117-073034.

15. Filsinger M, Freitag M. Asymmetric affective polarization regarding COVID-19

vaccination in six European countries. Scientific Reports. 2024;14(1):15919. doi:

https://doi.org/10.1038/s41598-024-66756-w.

16. Gadarian SK, Goodman SW, Pepinsky TB. Partisanship, health behavior, and policy

attitudes in the early stages of the COVID-19 pandemic. Plos one. 2021;16(4):e0249596.

doi: https://doi.org/10.1371/journal.pone.0249596.

17. Schieferdecker D, Joly P, Faas T. Affective Polarization Between Opinion-Based Groups

in a Context of Low Partisan Discord: Measuring Its Prevalence and Consequences.

International Journal of Public Opinion Research. 2024;36(2). doi:

https://doi.org/10.1093/ijpor/edae009.

18. Ipekci AM, Filsinger M, Buitrago-Garcia D, Kobler Betancourt CI, Liwanag HJ, Frahsa A,

et al. Polarization and health-related behaviors and outcomes during the COVID-19

pandemic: a systematic review. SSM - Population Health. 2026;33:101891. doi:

https://doi.org/10.1016/j.ssmph.2025.101891.

19. Verberk JDM, de Hoog MLA, Westerhof I, van Goethem S, Lammens C, Ieven G, et al.

Transmission of SARS-CoV-2 within households: a remote prospective cohort study in

European countries. Eur J Epidemiol. 2022;37(5):549-61. Epub 20220528. doi:

https://doi.org/10.1007/s10654-022-00870-9. PubMed PMID: 35644003; PubMed Central

PMCID: PMCPMC9146817.

20. Alberto-Orlando S, Calderon JL, Leon-Sosa A, Patiño L, Zambrano-Alvarado MN,

Pasquel-Villa LD, et al. SARS-CoV-2 transmission from infected owner to household

dogs and cats is associated with food sharing. Int J Infect Dis. 2022;122:295-9. Epub

20220526. doi: https://doi.org/10.1016/j.ijid.2022.05.049. PubMed PMID: 35643308;

PubMed Central PMCID: PMCPMC9132679.

21. Derqui N, Koycheva A, Zhou J, Pillay TD, Crone MA, Hakki S, et al. Risk factors and

vectors for SARS-CoV-2 household transmission: a prospective, longitudinal cohort study.

The Lancet Microbe. 2023;4(6):e397-e408. doi: https://doi.org/10.1016/s2666-

5247(23)00069-1.

22. Davis MF, Innes GK. The Cat's in the Bag: Despite Limited Cat-to-Cat Severe Acute

Respiratory Syndrome Coronavirus 2 Transmission, One Health Surveillance Efforts Are

Needed. J Infect Dis. 2021;223(8):1309-12. doi: https://doi.org/10.1093/infdis/jiab106.

PubMed PMID: 33605418; PubMed Central PMCID: PMCPMC7928724.

23. Morand Bourqui R, Nusslé SG, von Goetz N, Veys-Takeuchi C, Zuppinger C, Boulez Y,

et al. Towards a Swiss health study with human biomonitoring: Learnings from the pilot

phase about participation and design. PLoS One. 2023;18(7):e0289181. doi:

https://doi.org/10.1371/journal.pone.0289181.

24. Brall C, Berlin C, Zwahlen M, Vayena E, Egger M, Ormond KE. Public preferences

towards data management and governance in Swiss biobanks: results from a nationwide

survey. BMJ Open. 2022;12(8):e060844. Epub 20220826. doi:

https://doi.org/10.1136/bmjopen-2022-060844. PubMed PMID: 36028266; PubMed

Central PMCID: PMCPMC9422864.

25. Reichmuth ML, Heron L, Riou J, Moser A, Hauser A, Low N, et al. Socio-demographic

characteristics associated with COVID-19 vaccination uptake in Switzerland: longitudinal

analysis of the CoMix study. BMC Public Health. 2023;23(1):1523. doi:

https://doi.org/10.1186/s12889-023-16405-0.

26. Issp Research Group. International Social Survey Programme: Social Networks and

Social Resources - ISSP 2017. GESIS Data Archive, Cologne. ZA6980 Data file Version

2.0.0, https://doi.org/10.4232/1.13322; 2019.

27. Druckman JN, Levendusky MS. What Do We Measure When We Measure Affective

Polarization? Public Opinion Quarterly. 2019;83(1):114-22. doi:

https://doi.org/10.1093/poq/nfz003.

28. RStudio: Integrated Development Environment for R Version 2024.4.2.764. 2024.

29. Lumley T. Survey: analysis of complex survey samples. R package version 4.2. 2023 doi:

https://doi.org/10.18637/jss.v009.i08.

30. Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect

modification and interaction. International journal of epidemiology. 2012;41(2):514-20.

doi: https://doi.org/10.1093/ije/dyr218.

31. Federal Office of Topography. swissBOUNDARIES3D. Federal Office of Topography

swisstopo; 2025 https://www.swisstopo.admin.ch/en/landscape-model-

swissboundaries3d.

32. Federal Statistical Office. GEOM TK (Geometrische Grundlagen für die thematische

Kartografie) 2025. Federal Statistical Office (FSO); 2025

https://www.bfs.admin.ch/bfs/de/home/statistiken/regionalstatistik/kartengrundlagen/basis

geometrien.assetdetail.33807959.html.

33. Rolstad S, Adler J, Rydén A. Response Burden and Questionnaire Length: Is Shorter

Better? A Review and Meta-analysis. Value in Health. 2011;14(8):1101-8. doi:

https://doi.org/10.1016/j.jval.2011.06.003.

34. Gabel M, Gooblar J, Roe CM, Selsor NJ, Morris JC. Political Ideology, Confidence in

Science, and Participation in Alzheimer Disease Research Studies. Alzheimer Dis Assoc

Disord. 2018;32(3):179-84. doi: https://doi.org/10.1097/wad.0000000000000244.

PubMed PMID: 29351092; PubMed Central PMCID: PMCPMC6051930.

35. Onyeaka H, Weber DB, Chido-Amajuoyi O, Muoghalu C, Amonoo HL. The influence of

political ideology on clinical trial knowledge, invitation, and participation among adults in

the United States. Clin Trials. 2023;20(6):708-13. Epub 20230622. doi:

https://doi.org/10.1177/17407745231178790. PubMed PMID: 37345562; PubMed

Central PMCID: PMCPMC10741248.

36. Kim H, Kim HR, Kim S, Kim E, Kim SY, Park H-Y. Public attitudes toward precision

medicine: a nationwide survey on developing a national cohort program for citizen

participation in the republic of korea. Frontiers in Genetics. 2020;11:283. doi:

https://doi.org/10.3389/fgene.2020.00283.

37. Kaufman DJ, Baker R, Milner LC, Devaney S, Hudson KL. A Survey of U.S Adults'

Opinions about Conduct of a Nationwide Precision Medicine Initiative® Cohort Study of

Genes and Environment. PLOS ONE. 2016;11(8):e0160461. doi:

https://doi.org/10.1371/journal.pone.0160461.

38. Gaskell G, Gottweis H, Starkbaum J, Gerber MM, Broerse J, Gottweis U, et al. Publics

and biobanks: Pan-European diversity and the challenge of responsible innovation. Eur J

Hum Genet. 2013;21(1):14-20. Epub 20120606. doi:

https://doi.org/10.1038/ejhg.2012.104. PubMed PMID: 22669414; PubMed Central

PMCID: PMCPMC3522201.

39. Glass DC, Kelsall HL, Slegers C, Forbes AB, Loff B, Zion D, et al. A telephone survey of

factors affecting willingness to participate in health research surveys. BMC Public Health.

2015;15(1):1017. doi: https://doi.org/10.1186/s12889-015-2350-9.

40. Hierro LÁ, Patiño D, Atienza P, Garzón AJ, Cantarero D. The effect of altruism on

COVID-19 vaccination rates. Health Economics Review. 2023;13(1):2. doi:

https://doi.org/10.1186/s13561-022-00415-6.

41. Dolman AJ, Fraser T, Panagopoulos C, Aldrich DP, Kim D. Opposing views: associations

of political polarization, political party affiliation, and social trust with COVID-19

vaccination intent and receipt. Journal of Public Health. 2023;45(1):36-9. doi:

https://doi.org/10.1093/pubmed/fdab401.

42. Druckman JN, Klar S, Krupnikov Y, Levendusky M, Ryan JB. Affective polarization, local

contexts and public opinion in America. Nat Hum Behav. 2021;5(1):28-38. Epub

20201123. doi: https://doi.org/10.1038/s41562-020-01012-5. PubMed PMID: 33230283.

43. Torcal M, Thomson ZA. Social trust and affective polarization in Spain (2014–19).

Electoral Studies. 2023;81:102582. doi: https://doi.org/10.1016/j.electstud.2023.102582.

44. Rekker R, Harteveld E. Understanding factual belief polarization: the role of trust, political

sophistication, and affective polarization. Acta Politica. 2022:1-28. doi:

https://doi.org/10.1057/s41269-022-00265-4.

45. Filsinger M, Freitag M. Divided by vaccination? Evaluating the intergroup conflict

between pro-and anti-vaccination groups in the post-pandemic era. Humanities and

Social Sciences Communications. 2025;12(1):1-12. doi: https://doi.org/10.1057/s41599-

024-04016-y.

Supporting information

S1 Fig. Distribution of raw data from variables which were dichotomised

S2 Fig. Study participants flow chart

S1 Table. STROBE checklist for cross-sectional studies

S2 Table. Descriptive table of variables included in the analysis

S3 Table. Weighting levels and represented households

S4 Table. Responders of the survey by household size and language

S5 Table. Willingness to participate in a long-term cohort study, unweighted denominators
and proportions

S6 Table. Sensitivity analysis

S7 Table. Reasons to participate or decline participation in a long-term study, unweighted

Willingness to participate, proportion

1.00
0.75
0.50
0.25
0.00

✕ Bern city

No household selected

Canton of Bern in Switzerland

Grouped reasons for non–participation in the study

	Number (%)
Not interested	283 (44.4%)
Privacy concerns	271 (42.5%)
Mistrust	252 (39.6%)
No time / Inconvenient	222 (34.9%)
Other reasons / No response	144 (22.6%)
No personal benefit	108 (17%)