Modelling a potential zoonotic spillover event of H5N1 influenza

Philip Cherian[1], Gautam I. Menon[1,2,3*]

1 Department of Physics, Ashoka University, Sonepat, Haryana, India
2 Department of Biology, Trivedi School of Biological Sciences, Ashoka University, Sonepat, Haryana, India
3 Center for Climate Change and Sustainability, Ashoka University, Sonepat, Haryana, India

[*] gautam.menon@ashoka.edu.in (GIM)

Abstract

Background   Highly Pathogenic Avian Influenza (HPAI) is a prominent candidate for a future human pandemic arising from a zoonotic spillover event. Its best-known strain is H5N1, with South- or South-East Asia a likely location for an initial outbreak. Such an outbreak would be initiated through a primary event of bird-to-human infection, followed by sustained human-to-human transmission. Early interventions would require the extraction, integration and interpretation of epidemiological information from the limited and noisy case data available at outbreak onset.

Methods   We studied the implications of a potential zoonotic spillover of H5N1 influenza into humans. Our simulations used *BharatSim*, an agent-based model framework designed primarily for the population of India, but which can be tuned easily for others. We considered a synthetic population representing farm-workers (primary contacts) in a farm with infected birds. These primary contacts transfer infections to secondary (household) contacts, from where the infection spreads further. We simulate outbreak scenarios in such a setting, accounting for the network structure of human contacts and the stochasticity of the infection process. We further simulated multiple interventions, including bird-culling, quarantines, and vaccinations.

Findings   We show how limited, noisy data for primary and secondary infections can be used to estimate epidemiological transmission parameters, such as the basic reproductive ratio R_0, in realistic settings. We describe the impact of early interventions (bird-culling, quarantines, and vaccination), taken together or separately, in slowing or terminating the outbreak.

Interpretation   An individual-based model allows for the most granular description of the bird-human spillover and subsequent human-to-human transmission for the case of H5N1. Such models can be contextualised to individual communities across varied geographies, given representative contact networks. We show how such models allow for the systematic real-time exploration of policy measures that could constrain disease-spread, as well as guide a better understanding of disease epidemiology.

Research in context

Evidence before this study   To identify similar modelling studies, we searched PubMed using the search terms "("HPAI" OR "H5N1" OR "avian influenza") AND ("zoonosis" OR "zoonotic") AND ("modeling" OR "modelling")". Only a few mathematical modelling studies assessed the risk of human infection from novel avian influenzas. None of these studies examined potential sustained human-to-human transmission. We identified studies that simulated pathogen transmission within realistic poultry production and distribution networks, but they did not use these simulations to estimate disease parameters. The available evidence suggesting the potential of zoonotic spillover of avian influenza supports the inclusion of such parameters into transmission models.

Added value of this study   South- and South-East Asia have in recent years recorded a number of significant outbreaks of avian influenza in birds, accompanied by spillover events into non-poultry species. Within the past three months, India has recorded 34 epicentres of infection across 8 states, raising concerns about the possibility of a pandemic driven by an adaptation of the virus to infectious spread between humans. Our agent-based simulation framework can be used to tune policy measures for a potential outbreak of H5N1 influenza, as well as to calibrate both epidemiological parameters and the impact of interventions.

Implications of all the available evidence   Apart from providing realistic models for the consecutive events of a bird-human spillover infection followed by human-human transmission for the case of H5N1 influenza, our models can describe the impact of a variety of policy measures aimed at stemming a potential HPAI-initiated pandemic in humans. These include bird culling, quarantines and vaccination drives. We show that that timing of such measures is crucial if they are at all to be effective: culling of infected birds and quarantining of primary contacts should happen well before the force of infection peaks.

1   Introduction

A likely scenario for a future Disease X [1, 2] associates it with a viral infection in humans, initiated through a zoonotic spillover event. Such an infection would transmit efficiently via a respiratory route. A strong candidate for such a disease is highly pathogenic avian influenza (HPAI), commonly called "bird-flu" [3–5].

The most prominent example of HPAI is H5N1 influenza, a disease very largely confined to birds, but with a demonstrated ability to infect both terrestrial and marine mammals [4, 6–8]. There have, so far, only been isolated reports of bird-to-human transmission, either directly or via household pets. There are no confirmed reports of human-to-human transmission [9]. However, in the few known cases of human infection, case fatality rates of more than 30% have been suggested [10, 11], pointing to the importance of planning for a future outbreak.

The initial stages of any outbreak of a novel disease are inevitably marked by uncertainty as to the epidemiological parameters that characterise its spread [12]. The epidemiology here must account for two bottlenecks [13]. The first is the ability of the virus to cause infections in human hosts, while the second describes the barriers to subsequent human-to-human transmission. Once the initial threshold of viral adaptability is breached, the potential to cause a pandemic depends on the efficiency of human-to-human transmission.

The usefulness of models is that they allow for the reconstruction of potential disease trajectories from which, by comparison, such epidemiological parameters and their consequences can be inferred. A variety of disease models can be used to describe an outbreak. They typically differ in the scale of their description. The model may be defined at the level of groups or categories of individuals (susceptible or infected, as in conventional compartmental models) or at the level of single individuals, where individual-level variation and networks of potential contacts leading to infection must be accounted for. A number of models have been used to describe avian influenza and its transmission both between birds and into mammals [14–26].

Here, since the early stages of an outbreak are marked by a small number of infections, each of them occurring through a stochastic infection event, any useful analysis must account for multiple alternative disease trajectories describing transmission between individuals, a task for which agent-based simulations are best suited. In this paper, we simulate potential scenarios of an H5N1 outbreak in humans. Our simulations use BharatSim, an agent-based simulation framework developed for India [27]. Agent-based models are the most granular of models since they are defined at the level of single individuals and the networks of their interactions. While BharatSim's capabilities extend to simulations of up to 50 million individual agents, a much smaller population of 10,000 agents is sufficient for us to be able to address the early stages of an outbreak.

BharatSim uses a synthetic population of considerable complexity, including social and economic information obtained from combining a variety of surveys and census information using refined AI/ML techniques [27]. Much of this complexity is irrelevant to the questions we ask here, so we choose simpler representations, arguing that these should generalize across multiple LMIC geographies as well. Our simulations work with networks of homes, workplaces (including schools) and the interactions between primary, secondary and tertiary contacts across them.

The ability to describe a range of different disease trajectories while also potentially accommodating individual-level heterogeneity is a feature of the simulation methodology described in this paper. We use results from these simulations to extract information about parameters governing epidemiological transmission, such as the basic reproductive ratio and the secondary attack rate, across a range of scenarios. Our simulation methods allow us to explore targeted interventions and synergies between them, for example isolating and quarantining populations vulnerable to infection [28].

2 Methods

Our simulations of a potential H5N1 epidemic originating in a spillover to humans use an agent-based approach for disease-spread in the human population. This disease is seeded through infected birds within a single location (a farm).

2.1 Model

A schematic of our model, implemented within BharatSim, is shown in Fig. 1(A). This describes a farm with infected birds and a set of primary contacts. These primary contacts have secondary contacts at their homes.

The individuals in our model are part of larger networks. They move between different locations during a single day, reconfiguring their contact networks. BharatSim assumes that any individual above the age of 18 works at a defined 'workplace', while the remaining are children who go to schools. Agents spend 12 hours at their homes, interacting with their household contacts. The remaining 12 hours are spent at their workplaces or schools, where they interact with a different set of contacts. One specific

workplace is designated as a poultry farm, from where the infection is initiated, with individuals working in this farm being primary contacts.

Fig 1. (A) Schematic illustrating primary and secondary contacts of infected birds associated with a single poultry farm. (B) A representative time-series of daily incidence in the case of a spillover event for both primary (crosses) and secondary cases (circles) at the onset of the outbreak. Also shown is the number of infected birds in the farm, directly proportional to the force of infection.

2.2   Disease progression

The infection of primary contacts, humans in direct contact with infected birds, is described through a force of infection. This is depicted in Fig 1B (the smooth curve). This force of infection in the farm is taken to be proportional to the fraction of infected birds in that location. We choose parameters of transmission among birds such that the force of infection arising from the birds peaks at approximately 12 days from the start of our simulations. The first primary case is detected close to the peak of the force of infection, about 10 days after the initiation of the infection in this run. In practice, information would be available, at least in principle, about the fraction of birds which die (containing information about to the force of infection and how it changes in time, the curve) and the time-line of cases in humans (circles and crosses).

Fig. 2 shows the disease progression in our model for H5N1. Disease states are described using the notation of conventional compartmental models. Infection among birds is assumed to follow a well-mixed continuous SIR model, given their high density. Susceptible birds (S_B) can become infected (I_B) before dying from the disease and transitioning to the D_B compartment.

For humans, even though they are described as individual agents, we use the *notation* of an SEIR model to classify their disease states. At any given time, each individual can be in one of four possible states: (S)usceptible, (E)xposed, (I)nfectious, and (R)emoved. The disease states are shown in the Fig. 2. Disease is assumed to be symptomatic. We assume that recovered individuals are immune to further infection. Our model can be generalised to include the possibility of asymptomatic transmission.

Human agents transition stochastically between disease compartments after a probabilistically drawn duration (or "sojourn time"). BharatSim allows modellers the flexibility of choosing these sojourn times from different probability distributions. We draw these times from lognormal distributions, whose properties are specified further in Section 2.3.

2.3 Parameters

Our model community consists of 10,000 individuals, constructed to simulate a catchment area for the given poultry farm. (Since our interest in the trajectory of initial infections, ranging between 0 and 100 typically, this choice is sufficient.) Agents employed by this farm are primary contacts. The remaining network of individuals represents (i) their household members (in addition to other, secondary, contacts), and (ii) work- or school-contacts of their household members (tertiary contacts). Our parameter choices are detailed in Table 1.

Par.	Description	Range of values	Comment
β_{BH}	Bird-human transmission probability	$(1, 5, 10) \times 10^{-6}$	Correspond to $\sim 20, \sim 70$, and $\sim 90\%$ probabilities of at least one spillover event.
R_0^B	R_0 for bird SIR model	2.3	Corresponds to a peak at ~ 15 days after infection initialised.
N_{pop}	Human population size	10,000	Primary contacts: 162, secondary contacts: 805
τ_E	Latent period for human infection	lognormal(3,1)	Drawn from mean of 3 days and standard deviation of 1.
τ_I	Infectious period for human infection	lognormal(7,1)	Drawn from mean of 7 days and standard deviation of 1.
β_{HH}	Human-human transmission parameter	$(0.05, 0.5)$	Correspond to $R_0 \in (0, 3)$.
CD	Culling day	$(10, 20, 600)$	The FOI due to birds is set to zero post $t =$ CD.
QT	Quarantine threshold	2, 10	Threshold number of infectious agents for lockdown of primary contact households.
DVR	Daily vaccination rate	0%, 1%, 5% of N_{pop}	Number of vaccines available per day.
VE	Vaccine efficacy	0%, 50%, 100%	Reduces relative risk of infection by VE.

Table 1. Parameter choices for our model. Our simulations are run using multiple parameter sets. Those used in this paper correspond to the parameter choices above.

We begin our simulations with zero infected individuals in the human population, but with a small number of infected birds in the poultry farm. Susceptible birds get infected at some rate β_{BB}, and eventually die. At every time-step, individual agents who work in the farm are assumed to come in contact with these infected birds. The fraction of infected birds thus contributes to the force of infection within the farm location.

For human-to-human transmission to occur at all, a pre-requisite is an initial spillover event. To account for this, we weight the infected fraction of birds by a "relative risk of spillover", β_{BH}, which we vary in the range $\beta_{BH} \in (10^{-6}, 10^{-5})$. This corresponds to between 20% and 90% of the simulation runs leading to at least one spillover event. Once such an event occurs, the disease is assumed to spread amongst the human population through direct contact. At any given time, each location within our simulation (home, work, school) is considered to be a fully-connected network of individuals. However, the number of individuals an agent comes in contact with changes with time as they move between these locations.

Susceptible individuals in each location experience a force of infection given by

$$\mathcal{F}(\ell) \equiv \text{Force of infection at location } \ell = \beta_{HH} \times \left(\frac{I_\ell}{N_\ell} \right), \qquad (1)$$

where β_{HH} is the human-human transmission parameter, N_ℓ is the total number of individuals in that location, and I_ℓ is the number of infected individuals in that location. In an interval of time Δt, "Susceptible" individuals transition to the "Exposed" compartment with probability $\mathcal{F}\Delta t$.

For β_{HH}, we choose values in a range that corresponds to a basic reproductive ratio $R_0 \in (0,3)$. (This must be back-calculated given the trajectory of infections for a given β_{HH}, as we detail below.) The distributions of sojourn times in each compartment (shown in Fig 2) are uncertain at the outbreak onset. We draw them from lognormal distributions, choosing parameter values that should be largely representative [12]. In itself, this choice of distribution provides a considerable improvement on conventional compartmental models, where the sojourn times in each compartment are constrained to be exponentially distributed. These parameter choices will have to be refined as more information about the outbreak becomes available. Our simulation framework can easily account for updated parameter values.

Fig 2. Description of the disease model. The birds are assumed to follow a well-mixed compartmental variant of the SIR model where they are infected and eventually die. The total number of birds contributes to the force of infection at the farm. Individual agents can be in any one of the four states (S)usceptible, (E)xposed, (I)nfected, or (R)emoved. An agent working at the farm may be infected by these birds, after which they transition to the "Exposed" state, and eventually become infectious and can infect their secondary (household) contacts who can, in turn, infect other contacts in the population.

2.4 Interventions

We consider three specific interventions and study their impacts on the disease trajectory.

Culling At a specific day, all the birds in the farm are culled. Beyond this time, no further infections can arise from bird-human interactions. Any new cases after this date occur solely due to human-human interactions. This is modelled in our simulations by setting the force of infection arising from birds to zero, as shown in panels (C) and (E) of Fig 3.

Vaccination We consider a vaccination drive that begins one week after the detection of the first case in the population. Vaccines are restricted to primary and secondary contacts, and have the effect of reducing the agents' susceptibility to the disease. Given that little is known about how such vaccines might function, we choose to assume that vaccination does not alter the infectivity of an individual who contracts the disease, vaccinated or not.

We consider a daily vaccination rate (DVR) of 1% of the population (which translates to 100 vaccines available for use every day). We choose vaccine efficacies of 50% and 100%, which corresponds to the reduction in susceptibility of a vaccinated agent. (A vaccine efficacy of 100% means that an individual is completely immune to the disease.) The effect of the vaccine is immediate.

We have also performed simulation runs for a DVR of 5%, and with a vaccine drive that begins two weeks after the first detected cases. These results are shown in 6.

Quarantine When the number of cases reaches a threshold value (which we choose as either 2 or 10 individuals in our simulations, to explore a range of possible scenarios), all households of primary contacts are quarantined, such that no individual is allowed to leave their household to their workplaces or schools.

2.5 Computing R_0 and the secondary attack rate

From our simulations, we compute two specific quantities: the basic reproductive ratio R_0 and the secondary (or household) attack rate (SAR).

2.5.1 Reproductive ratio R_0

The basic reproductive ratio is defined as the average number of people infected by a single infectious person in the background of a population of completely susceptible people. Its value, in an averaged sense, is dependent on the mean number of contacts per unit time, the probability that an infection will result from an encounter with an infected contact over the period of the interaction, and the time over which an individual remains infected. Stochasticity enters through the network structure of interactions and the non-deterministic nature of the transmission of infection.

Our agent-based simulations take both of these into account when determining R_0. The advantage of agent-based models is that these different sources of stochasticity can be tuned independently of each other, in contrast to purely statistical models.

To compute R_0 from our simulations, we record all chains of infection. Whenever an individual is infected in a specific location, an agent is drawn at random from the potential "infectors" at that location. For a given value of β_{HH}, we run our simulations for a period of 21 days from the date of spillover. This period is chosen so that a sufficient number of infectious individuals recover, while the number of susceptibles in the accessible population can be assumed to be more or less constant. At the end of this

period, we compute the average number of people infected by a person who has recovered from the disease.

2.5.2 Secondary or household attack risk (SAR)

The secondary attack risk measures the difference between community transmission of illness versus transmission of illness in a household and is a useful indicator of infectivity at outbreak onset. It is calculated as the proportion of infected household or family contacts [29]:

$$\text{SAR} = \frac{\text{Number of infected secondary cases in primary-case households}}{\text{Total number in those households} - \text{Number of primary cases}}. \tag{2}$$

Consider an outbreak in which 5 "primary" cases are detected. After one incubation period, 10 new cases develop in the households of these primary contacts. If the combined total number of individuals in these households was 30, then the secondary attack rate is calculated as

$$\text{SAR} = \frac{10}{30 - 5} = 40\%. \tag{3}$$

As with our computation of R_0, we restrict our computation of the secondary attack risk to cases within the first 21 days of the first recorded case. In this time, it is unlikely that *all* households containing primary contacts have an infection. As a result, the secondary attack risk uses only those households in which one primary case has been detected. Additionally, in our computation of the SAR, we do not differentiate between infections in primary contacts that arise due to interaction with the FOI, and those that are due to interactions with other infected primary contacts.

3 Results

We begin our simulation with no infected humans and a small fraction of infected birds. The infection spreads rapidly amongst the birds and potentially infecting the farm workers (the primary contacts). In this section we study the effect of different bird-human interaction probabilities, culling days, and quarantine strategies on the disease trajectory.

3.1 Distribution of primary cases

In Fig 3A, we show the force of infection arising from the spread of disease in the bird population. In the absence of culling the force of infection follows the dashed line. The solid line shows the force of infection when all birds are culled on three specific days following the onset of the infection.

In Fig 3B we show the distribution of primary cases for three values of the human-bird infectivity. For the earliest culling dates, the distribution of primary cases decreases monotonically. The most likely outcome is that there are no primary cases.

For later culling dates (and shown in the later panels) the distribution of primary cases peaks at 1 or 2 individuals, but also has a long tail, where up to 7 or 8 may be infected. As β_{BH} increases, the peak of this distribution shifts outward, while its spread becomes larger.

3.2 Reproductive ratio and secondary attack rate

In an agent-based simulation, the basic reproductive ratio is not simply specified in terms of the model parameters, because it depends on the assumed network structure.

Fig 3. The distribution of primary cases. Each row represents a different culling date, as shown in the left panels (A), (C), and (E). The solid curve represents the number of infected birds in the farm, which abruptly goes to zero in the case of culling. The dashed grey curve represents the number of infected birds in the absence of culling. The right panels (B), (D), and (F), show the probability distributions of the number of primary cases for these culling dates, with the different colours representing different values of β_{BH}, the bird-human interaction probability.

To account for this, we compute R_0 directly from the distribution of secondary infections. Figs 4 and 5 show R_0 for different values of the human-human transmission coefficient (β_{HH}), where the error-bars arise from the inherent stochasticity of the transmission process: these graphs are obtained by averaging over multiple stochastic runs. In principle, one might have to average over different network structures as well,

but here we assume that a well-constructed synthetic population should naturally improve on the conventional approximation of a well-mixed population generally used in traditional compartmental models.

In Figs. 4B and 5B, we show the secondary attack rate as a function of β_{HH}, and show how the SAR may be used to infer R_0. Given an estimated value of the SAR, we can compute the corresponding value of β_{HH} and hence the basic reproductive ratio R_0 by simply reading off the curve. Both quantities are monotonic in the human-human transmission parameter β_{HH}. While the structure of these curves depends in detail on the assumptions we make, such as for the parameters of the lognormal distribution assumed for sojourn times in disease states, further refinements are possible as more clinical information is available to be assimilated during the early course of the outbreak.

In Appendix S2 we show the epidemic curves with a variety of interventions for five different values of β_{HH} that correspond to an $R_0 > 1$. In addition to the no-intervention scenario, we also show the epidemic curves for a daily vaccination rate of 1% and a vaccine efficacy of 100%, and quarantining scenarios where the threshold is 10 and 2 individuals respectively.

Fig 4. Calibrating the human-human transmission rate. (A) The mean reproductive ratio as a function of the human-human transmission β_{HH}. We see that in our model, $\beta_{HH} \approx 0.25$ represents the threshold $R_0 = 1$, beyond which the epidemic takes off among humans in the case of no interventions. The introduction of vaccinations causes the critical threshold of $R_0 = 1$ to shift rightwards towards higher values of β_{HH}. (B) The secondary attack risk (SAR) among secondary contacts, evaluated as described in Equation (2). We see that the introduction of vaccinations does not alter the secondary attack risk. The error bars in both plots represent the variation (one standard deviation) across 500 different stochastic runs.

.

3.3 Conclusions

The results of Fig. 3 illustrates both the impact of increasing the probability of a spillover event as well as the effect of culling on the distribution of primary human cases. We find that culling has a large effect on this distribution, provided it occurs sufficiently early, before the peak in the force of infection arising from the infected birds. In this case, the most likely scenario is that the infection will not spill over, irrespective of bird-human spillover β_{BH}. For later culling dates, the most likely scenario involves one

Fig 5. Effects of quarantining strategies. (A) The mean reproductive ratio as a function of the human-human transmission β_{HH} in the case of quarantining homes of farmers when a threshold of infections is reached. As in Fig. 4, $\beta_{HH} \approx 0.25$ represents the threshold $R_0 = 1$ in the case of no interventions. However, if farmers and their households are quarantined when either 2 or 10 infected cases are detected, this can lead to a reduction in the number of cases within primary and secondary contacts. (B) The secondary attack rate (SAR) among household contacts of farmers. The quarantining of farmer households greatly reduces the SAR, although it does not necessarily reduce the spread of infection in the population unless it is done early. As can be seen, using a threshold number of cases of 2, for example, leads to an epidemic with an R_0 consistently below 1, meaning the epidemic never takes off. The error bars in both plots represent variation across 500 different stochastic runs.

.

or more primary cases, a number that depends strongly on β_{BH}. This argues for early culling as an important strategy to curb further spillover events.

In Figs. 4 and 5, we show how the secondary attack risk can be used to infer the basic reproductive ratio, a central quantity in determining the pandemic potential of our outbreak. The reproductive ratio rises smoothly with β_{HH}. In Fig. 4, we show the results with a vaccination drive. Introducing vaccinations causes the critical threshold of $R_0 = 1$ to shift towards higher β_{HH}, by an amount determined by the vaccine efficacy. We further find that the secondary attack risk is left relatively unchanged by the vaccination drive. This is due in part to our assumption that vaccination does not reduce the infectivity of an individual, only their relative risk of infection.

In Fig. 5, we show a similar graph but for quarantining primary and secondary contacts at different threshold values. As described above, when the number of active cases in the population exceeds a threshold of either 2 or 10 cases, the households of all primary contacts are locked down, and these agents are not allowed to travel to their workplace or school locations.

We find that a threshold of 10 infected individuals produces both an R_0 and an SAR that is comparable to the case of no interventions. We attribute this to the fact that the time taken for 10 active cases to be present in the population is much larger than the 21 day interval used to compute the SAR, which is typically a quantity calculated early in

an outbreak. In the case of a threshold of 2 infected individuals, we see a larger SAR. This is because households of primary contacts are quarantined early on, and our models do not account for any reduction in spread within households. Consequently, infected primary contacts have a much higher probability of transmitting the disease to the secondary contacts in the households as they spend more time within the house.

Nevertheless, the reproductive ratio remains consistently below $R_0 = 1$, meaning that the epidemic does not take off, since the disease is not allowed to enter the tertiary contact network due to the quarantine. This is verified in Appendix S3, where we compute the tertiary attack rate explicitly.

Thus, we find that while quarantining households of primary contacts has by far the largest effect in reducing the spread of the disease, this must be done very soon after the first case is detected, failing which the disease will enter the tertiary contact network and a more aggressive lockdown strategy would be required. The error bars in both plots represent variation across 500 different stochastic runs.

We have also conducted runs combining these different interventions, as shown in Appendix S3, to show how the combined effects of vaccinations and quarantining primary and secondary contacts can affect these epidemiological parameters.

4   Discussion

In this paper we showed how computational models can describe the sequential stages of a zoonotic spillover. We showed that a recently developed, ultra-large-scale agent-based simulation model devised for an LMIC context, *BharatSim*, could be used to study the impact of various policy interventions in the case of an epidemic of a spillover event of HPAI. For concreteness we assumed an initial spillover event of H5N1 from birds to humans, followed by sustained human-to-human transmission. Our model describes the two-step nature of outbreak initiation, showing how crucial epidemiological parameters governing transmission can be calibrated given data for the distribution of the number of primary and secondary cases at early times.

It is in the very early stages of an outbreak that control measures make the most difference. Once community transmission takes over, cruder public-health measures such as lockdowns, compulsory masking, and large-scale vaccination drives are the only options left. At this stage, the large number of cases ensures that stochastic effects should play a smaller role and conventional compartmental models should provide appropriate guidance. However, projecting the trajectory of a disease from initial fragmentary information available at outbreak onset is a far harder task. It in this setting that the methods described here are expected to be most useful [28].

Conventional models that calculate the reproductive ratio from case data attempt to reproduce the observed trajectory of infections using statistical assumptions [30, 31]. This assumption collapses many independent sources of stochasticity, including network effects and the random nature of the transmission of infection. An agent-based model allows one to separate these different contributions. Because many parallel scenarios can be explored in real-time, and the models can be adapted to the measures that are actually implemented, such methods provide powerful input to policy [32].

We considered the initial stages of an outbreak, reasoning that confining ourselves to simulating 10,000 individuals overall should be sufficient. The world's fastest growing poultry market are South- and South-East Asia [33, 34]. One feature of poultry farms in these regions is a coexistence of small family-holdings with large-scale factory-farmed poultry. Since large factory farms are extensively monitored, the more interesting situation for us is an intermediate one. We ignore stochastic effects in the bird population in this limit.

The range of estimated epidemiological disease parameters in humans is very broad,

given the sparsity of data. We used the recent meta analysis of Ref. [12] to fix values for the incubation and infectious periods. Changes in these within the estimated ranges will not affect our qualitative results.

Our simulations account for different interventions. These include (i) the culling of all birds in the farm, (ii) the quarantining of primary and secondary contacts once a threshold of cases is crossed, and (iii) a vaccination drive where primary and secondary contacts are targeted. We find that culling birds is effective, provided no primary infection has occurred. The earlier birds are culled, the larger the probability that a spillover can be prevented. Even if an infection of a primary case occurs, onward infections are limited if cases are isolated and their household contacts quarantined. However, if tertiary contacts are infected, establishing control becomes impossible unless far more stringent measures are applied, including a total lockdown.

Our methods allow us to explore a range of possibly applicable policy measures. Understanding the scope of possible interventions in advance of a potential outbreak strengthens the ability of a public health infrastructure to respond. To the best of our knowledge, no other work exists which has attempted to capture the consequences of such a two-step spillover event occurring in a realistic population, especially in the LMIC context.

Our simulations can be run in real time, responding to initial reports of cases. They can be tuned further as more information is collected. Further generalizations include the incorporation of asymptomatic infections and of delays in reporting, the possibility of a bird to intermediate mammal to human chain of spillovers, the inclusion of serological data in real time, and a more refined description of the consequences of vaccination drives, including model descriptions of vaccine efficacy. These are the subject of ongoing work.

5   Contributors

Both authors did the literature search, design and implementation of the study, and wrote the original draft. Both authors had final responsibility for the decision to submit for publication.

6   Acknowledgements

The authors are grateful for ongoing support from the Mphasis F1 Foundation. BharatSim development was supported by the Bill and Melinda Gates Foundation, Grant No: R/BMG/PHY/GMN/20, as well as by the Mphasis F1 Foundation. GIM thanks Sindura Ganapati for ongoing discussions. The authors acknowledge the use of computational facilities provided by the Centre for Bioinformatics and Computational Biology, as well as the One-Health initiative of the Center for Climate Change and Sustainability at Ashoka University. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Simpson S, Kaufmann MC, Glozman V, Chakrabarti A. Disease X: accelerating the development of medical countermeasures for the next pandemic. The Lancet Infectious Diseases. 2020;20(5):e108–e115. doi:10.1016/S1473-3099(20)30123-7.

2. Sikkema RS, Koopmans MPG. Preparing for Emerging Zoonotic Viruses. Encyclopedia of Virology. 2021; p. 256–266. doi:10.1016/B978-0-12-814515-9.00150-8.

3. Al-Tawfiq JA, Tirupathi R, Temsah MH. Feathered fears: Could avian H5N1 influenza be the next pandemic threat of disease X? New Microbes and New Infections. 2024;59:101416. doi:10.1016/j.nmni.2024.101416.

4. Mostafa A, Naguib MM, Nogales A, Barre RS, Stewart JP, García-Sastre A, et al. Avian influenza A (H5N1) virus in dairy cattle: origin, evolution, and cross-species transmission. mBio. 2024;15(12):e02542–24. doi:10.1128/mbio.02542-24.

5. Peiris JSM, de Jong MD, Guan Y. Avian Influenza Virus (H5N1): a Threat to Human Health. Clinical Microbiology Reviews. 2007;20(2):243–267. doi:10.1128/CMR.00037-06.

6. Pardo-Roa C, Nelson MI, Ariyama N, Aguayo C, Almonacid LI, Gonzalez-Reiche AS, et al. Cross-species and mammal-to-mammal transmission of clade 2.3.4.4b highly pathogenic avian influenza A/H5N1 with PB2 adaptations. Nature Communications. 2025;16(1):2232. doi:10.1038/s41467-025-57338-z.

7. Campbell AJ, Brizuela K, Lakdawala SS. mGem: Transmission and exposure risks of dairy cow H5N1 influenza virus. mBio. 2025;16(3):e02944–24. doi:10.1128/mbio.02944-24.

8. Wang X, Yu H, Ma Y, Zhang P, Wang X, Liang J, et al. The novel H10N3 avian influenza virus acquired airborne transmission among chickens: an increasing threat to public health. mBio. 2024;16(2):e02363–24. doi:10.1128/mbio.02363-24.

9. Fok B. H5N1 influenza viral lineages beginning to evade human immunological defenses; 2025. Available from: https://www.the-microbiologist.com/news/h5n1-influenza-viral-lineages-beginning-to-evade-human-immunological-defenses/5472.article.

10. Li FCK, Choi BCK, Sly T, Pak AWP. Finding the real case-fatality rate of H5N1 avian influenza. Journal of Epidemiology & Community Health. 2008;62(6):555–559. doi:10.1136/jech.2007.064030.

11. Lai S, Qin Y, Cowling BJ, Ren X, Wardrop NA, Gilbert M, et al. Global epidemiology of avian influenza A H5N1 virus infection in humans, 1997-2015: a systematic review of individual case data. The Lancet Infectious Diseases. 2016;16(7):e108–e118. doi:10.1016/S1473-3099(16)00153-5.

12. Ward J, Lambert JW, Russell TW, Azam JM, Kucharski AJ, Funk S, et al.. Estimates of epidemiological parameters for H5N1 influenza in humans: a rapid review; 2024. Available from: https://www.medrxiv.org/content/10.1101/2024.12.11.24318702v2.

13. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic spillover. Nature Reviews Microbiology. 2017;15(8):502–510. doi:10.1038/nrmicro.2017.45.

14. Beaunée G, Deslandes F, Vergu E. Inferring ASF transmission in domestic pigs and wild boars using a paired model iterative approach. Epidemics. 2023;42:100665. doi:10.1016/j.epidem.2023.100665.

15. Cecilia H, Métras R, Fall AG, Lo MM, Lancelot R, Ezanno P. It's risky to wander in September: Modelling the epidemic potential of Rift Valley fever in a Sahelian setting. Epidemics. 2020;33:100409. doi:10.1016/j.epidem.2020.100409.

16. Cecilia H, Arnoux S, Picault S, Dicko A, Seck MT, Sall B, et al. Dispersal in heterogeneous environments drives population dynamics and control of tsetse flies. Proceedings of the Royal Society B: Biological Sciences. 2021;288(1944):20202810. doi:10.1098/rspb.2020.2810.

17. Cecilia H, Vriens R, Wichgers Schreur PJ, De Wit MM, Métras R, Ezanno P, et al. Heterogeneity of Rift Valley fever virus transmission potential across livestock hosts, quantified through a model-based analysis of host viral load and vector infection. PLOS Computational Biology. 2022;18(7):e1010314. doi:10.1371/journal.pcbi.1010314.

18. Cristancho-Fajardo L, Vergu E, Beaunée G, Arnoux S, Ezanno P. Learning and strategic imitation in modelling farmers' dynamic decisions on bovine viral diarrhoea vaccination. Veterinary Research. 2022;53(1):102. doi:10.1186/s13567-022-01112-2.

19. Ezanno P, Arnoux S, Joly A, Vermesse R. Rewiring cattle trade movements helps to control bovine paratuberculosis at a regional scale. Preventive Veterinary Medicine. 2022;198:105529. doi:10.1016/j.prevetmed.2021.105529.

20. Guyver-Fletcher G, Gorsich E, Jewell CP, Tildesley MJ. Controlling Foot-and-Mouth Disease: Vaccination is More Important than Movement Bans. A Simulation Study in the Republic of Turkey; 2024. Available from: https://www.ssrn.com/abstract=4886121.

21. Hayes B, Vergne T, Rose N, Mortasivu C, Andraud M. A multi-host mechanistic model of African swine fever emergence and control in Romania; 2023. Available from: https://zenodo.org/doi/10.5281/zenodo.10174922.

22. Hobbelen PHF, Elbers ARW, Werkman M, Koch G, Velkers FC, Stegeman A, et al. Estimating the introduction time of highly pathogenic avian influenza into poultry flocks. Scientific Reports. 2020;10(1):12388. doi:10.1038/s41598-020-68623-w.

23. Johnson RM, Stopard IJ, Byrne HM, Armstrong PM, Brackney DE, Lambert B. Investigating the dose-dependency of the midgut escape barrier using a mechanistic model of within-mosquito dengue virus population dynamics. PLOS Pathogens. 2024;20(4):e1011975. doi:10.1371/journal.ppat.1011975.

24. Ssematimba A, Malladi S, Hagenaars TJ, Bonney PJ, Weaver JT, Patyk KA, et al. Estimating within-flock transmission rate parameter for H5N2 highly pathogenic avian influenza virus in Minnesota turkey flocks during the 2015 epizootic. Epidemiology and Infection. 2019;147:e179. doi:10.1017/S0950268819000633.

25. Tiensin T, Nielen M, Vernooij H, Songserm T, Kalpravidh W, Chotiprasatintara S, et al. Transmission of the Highly Pathogenic Avian Influenza Virus H5N1 within Flocks during the 2004 Epidemic in Thailand. The Journal of Infectious Diseases. 2007;196(11):1679–1684. doi:10.1086/522007.

26. Vergne T, Gubbins S, Guinat C, Bauzile B, Delpont M, Chakraborty D, et al. Inferring within-flock transmission dynamics of highly pathogenic avian influenza H5N8 virus in France, 2020. Transboundary and Emerging Diseases. 2021;68(6):3151–3155. doi:10.1111/tbed.14202.

27. Cherian P, Kshirsagar J, Neekhra B, Deshkar G, Hayatnagarkar H, Kapoor K, et al. BharatSim: An agent-based modelling framework for India. PLOS Computational Biology. 2024;20(12):e1012682. doi:10.1371/journal.pcbi.1012682.

28. Chowell G, Sattenspiel L, Bansal S, Viboud C. Mathematical models to characterize early epidemic growth: A Review. Physics of life reviews. 2016;18:66–97. doi:10.1016/j.plrev.2016.07.005.

29. Centers for Disease Control and Prevention. Principles of Epidemiology, Lesson 3: Measures of Risk; 2023. Available from: https://archive.cdc.gov/www_cdc_gov/csels/dsepd/ss1978/lesson3/section2.html.

30. Breban R, Vardavas R, Blower S. Theory versus Data: How to Calculate R0? PLoS ONE. 2007;2(3):e282. doi:10.1371/journal.pone.0000282.

31. Pellis L, Ball F, Trapman P. Reproduction numbers for epidemic models with households and other social structures. I. Definition and calculation of R0. Mathematical Biosciences. 2012;235(1):85–97. doi:10.1016/j.mbs.2011.10.009.

32. Nespeca V, Comes T, Brazier F. A Methodology to Develop Agent-Based Models for Policy Support Via Qualitative Inquiry. Journal of Artificial Societies and Social Simulation. 2023;26(1):10.

33. Global Ag Media. South, Southeast Asia to play pivotal role in global poultry market growth; 2024. Available from: https://www.thepoultrysite.com/news/2024/04/south-southeast-asia-to-play-pivotal-role-in-global-poultry-market-growth.

34. Mulder ND, Pan C, Beisly C, McCracken C, Yanaguizawa W, Gidley-Baird A, et al.. Poultry Quarterly; 2025. Available from: https://www.rabobank.com/knowledge/q011332991-global-poultry-quarterly-q2-2025.

Supporting information

Appendix S1 Calibrating the bird-human interaction parameter. We show how we calibrate the bird-human interaction parameter β_{BH} using the spillover probability.

Appendix S2 Epidemic curves with different interventions. We compare the different epidemic curves for different values of β_{BH}, β_{HH}, and different intervention strategies: no interventions, a vaccine drive with a daily vaccination rate of 1%, and quarantining all primary and secondary contacts when the number of infected agents in the population is greater than 10 or 2 respectively.

Appendix S3 Synergies of different intervention strategies. We compare the epidemiological parameters R_0, SAR, and TAR for different combinations of interventions.