Using step selection functions to analyse human mobility using telemetry data in infectious disease epidemiology: a case study of leptospirosis

Pablo Ruiz Cuenca1 , Fabio Neves Souza2,3 , Roberta Coutinho do Nascimento3 , Ariane Goncalves da Silva2 , Max T. Eyre4 , Juliet O. Santana2,3 , Daiana Santos de Oliveira2,3 , Emile Victoria Ribeiro de Souza2 , Fabiana Almerinda G. Palma2 , Diogo César de Carvalho Santiago2 , Priscyla dos Santos Ribeiro5,6 , Priscilla Elizabeth Ferreira dos Santos2, Hussein Khalil7, Jonathan M. Read1 , Cleber Cremonese2 , Federico Costa*2,3,8 , and Emanuele Giorgi*1

1 Centre for Health Informatics, Computing, and Statistics (CHICAS), Lancaster Medical School, Lancaster University, United Kingdom

2 Institute of Collective Health, Federal University of Bahia, Salvador, Bahia, Brazil

3 Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Ministério da Saúde, Salvador, Bahia,Brazil

4 Environmental Health Group, London School of Hygiene and Tropical Medicine, London,United Kingdom

5 Federal University of Bahia, National Institute of Science and Technology in Interdisciplinary and Transdiciplinary Studies in Ecology and Evolution, Salvador, Brazil

6 Amsterdam University Medical Centre, Leptospirosis Reference Center, Medical Microbiology and Infection Control, Amsterdam , Netherlands

7 Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sciences (SLU), Sweden

8 Department of Epidemiology of Microbial Diseases, School of Public Health, Yale University, New Haven, CT, United States

* These authors contributed equally to this work.

Abstract

Human movement plays a critical role in the transmission of infectious diseases, especially those with environmental drivers like leptospirosis—a zoonotic bacterial infection linked to mud and water contact. Using GPS loggers, we collected detailed telemetry data to understand how fine-scale movements can be analysed in the context of an infectious disease. We recruited individuals living in urban slums in Salvador, Brazil to analyse how they interact with environmental risk factors such as domestic rubbish piles, open sewers, and a local stream. We aimed to identify differences in movement patterns inside the study areas by gender, age, and leptospirosis serological status. Step-selection functions, a spatio-temporal model used in animal movement ecology, estimated selection coefficients to represent the likelihood of movement toward specific environmental factors. With 128 participants wearing GPS devices for 24 to 48 hours, recording locations every 35 seconds during active daytime hours, we segmented movements into morning, midday, afternoon, and evening. Our results suggested women moved closer to the central stream and farther from open sewers compared to men, while serologically positive individuals avoided open sewers. This study introduces a novel method for analysing human telemetry data in infectious disease research.

Keywords: Leptospirosis, human movement, GPS, urban health, infectious diseases, zoonosis

Introduction

GPS loggers are a growing tool for capturing both human and animal movements.[1,2] These small devices can be worn by individuals and record locations at regular preset time intervals. Compared to other methods of collecting human movements, such as cell tower traffic or Google Location History which are suited for analysing large-scale mobility,[3,4] these devices can capture very fine-scale movements. These data are

crucial in quantifying exposure within complex environments, where terrain can change
rapidly. Furthermore, movements recorded by GPS loggers can be assigned to specific
individuals. This allows linkage between individual socio-demographic factors and the
data collected, especially convenient when performing epidemiological analyses. Other
methods for measuring human mobility are inherently anonymous and do not allow this
connection to be made. An important challenge when using GPS loggers is that they
rely on individual compliance for carrying the device at all times, an issue which is
overcome by the other methods mentioned above.

The analysis of human telemetry data is an emerging field of research in epidemiology.
Whilst previous methods have advanced this area of research, improvements could be
made. For example, the methods used by Owers et al.[1] to assess the relationship
between urban slum residents' movements and the risk of leptospirosis infection were
able to analyse differences between genders, but did not consider other important
socio-demographic factors. In another study, Fornace et al.[2] used GPS loggers to
assess human exposure to mosquito vectors of *Plasmodium knowlesi* malaria and
environmental factors associated with this. Various individual-level factors were
included in the analyses performed in this paper, questioning how these could affect
participants' movements. However, by not including comparisons of possible choices an
individual could have made, this study could not determine how the environment may
have influenced movement.

Leptospirosis is a zoonotic bacterial infectious disease with strong environmental
drivers. It has been estimated to cause over 1 million yearly human cases worldwide,
leading to 58 900 deaths.[5] Rats are the main reservoir of the disease, shedding bacteria
in their urine.[6] Human infection is associated with exposure to contaminated waters and
soils.[6–8]Evidence shows that in urban slum settings, men have a higher infection risk
than women.[9] This has been attributed to differences in behaviours, especially in how
individuals move through their communities, rather than a biological differences. Indeed,
there is evidence that men tend to visit much larger areas during their daily journeys
than women.[1]

Exactly where people are most exposed to high leptospirosis contamination, and

therefore where infection is most likely to occur, has not been investigated. Previous

studies have focused on the assessment of the peri-domiciliary environment and its

associations to infection risk.[7–9] However, these analyses assume people are mostly

exposed to infection risk in this area and ignore the exposure that individuals may incur

when they move further away from their households. Furthermore, people's movement

patterns may differ depending on individual socio-demographic factors which could in

turn affect their risk of exposure. If individuals traverse highly contaminated areas where

the risk of exposure is heightened, it can lead to an increased risk of infection. This is

particularly important in environmentally heterogeneous areas, such as urban slums,

where the landscape can change drastically in small spaces. Technological advances

now allow us to record and analyse fine-scale movements to understand how these

may affect infection risk.

In this paper, we developed a modelling framework to understand how telemetry data

can be used to identify and quantify determinants of human movements, adapting

methods from animal movement ecology. We present a novel method for analysing

telemetry data to estimate environmental selection as individuals move through their

urban communities. This method is applied in a low-income urban setting in Salvador,

Brazil, and is used to examine how individuals interact with various key points in their

surrounding environment . Furthermore, we analyse if there are any differences in

movements inside the study areas between genders, ages and leptospirosis serological

status . This method of analysis overcomes limitations from other studies by, firstly,

specifically modelling choice of movement in relation to environmental factors and,

secondly, incorporating multiple socio-demographic factors which allows regression

relationships to be jointly adjusted for these.

Methods

Study areas

This study was nested in a prospective cohort study taking place across Salvador,

Brazil.[10] Salvador is the third largest city in Brazil, located in the north-eastern region of

the country and has a tropical climate. The study areas are considered urban slums

(locally called `favelas`). They were selected for a number of reasons: firstly, they all

have similar demographic and socio-economic factors within their populations;

secondly, they all have a stream running through the centre of the community, which is

considered contaminated; and thirdly, there is a high burden of leptospirosis in these

populations.

All four study areas are small, with an approximate size of 0.03 km2. They are located

across the outskirts of Salvador (Figure 1). The communities have very heterogeneous

environments, with rapid changes in both land cover and slope. Buildings in these

communities have been built with limited or no urban planning. They can be of varying

quality, ranging from gated areas with multiple dwellings protected from rain and

flooding to single brick buildings with informal entryways.

[**Figure 1** - Map showing location of each study area in Salvador. Each area includes

symbology for stream (blue line), open sewer points (purple diamond), and domestic

rubbish piles (orange triangle).]

Individual characteristics

The eligibility criteria for inclusion in the study was: individuals who (1) had been living

at one of the study areas for at least 6 months, (2) slept there at least 3 nights a week,

(3) were at least 18 years old and (4) gave written consent.[10] Participants were asked to

answer a baseline survey which collected their demographic, social and economic

characteristics, including age and gender. A blood sample was taken from each

participant to determine serological evidence for *Leptospira* infection using the

microscopic agglutination test (MAT), the standard test used for leptospirosis

diagnosis.[6] In this analysis, a MAT showing antibodies with a titre >1:50 against any

Leptospira serovar was considered a positive result. Further details about the laboratory

work carried out are available in Supplementary Material I. The location of their

household was recorded and georeferenced by the research team.

Participants who were already enrolled in the cohort study were recruited to take part in

the movement analysis study. At the time of recruitment, we found no published

scientific studies detailing how to perform sample size calculations for research using

GPS data in humans. Therefore, we opted to use convenience sampling instead. A

target of 30 people per study area, balanced by gender and blind to their serological

status, was chosen for this study.

GPS Data

Individuals who consented to take part in this study were asked to wear GPS loggers for

continuous periods of up to 48 hours, which could be repeated. The GPS loggers used

were i-got U GT-600, set to record their location every 35 seconds. We used the

manufacturer's software to programme the devices. Data were collected between March
and November 2022.

Once the GPS telemetry data was collected, participants' recorded locations were
cleaned so as to retain only relocations within the study area boundaries that were
recorded between 5 am and 9 pm. This period generally corresponds to an individual's
active hours. Interactions with environmental factors outside of the study area
boundaries could not be considered in the analysis because high-resolution
environmental data outside of the study areas was not available. Individuals with less
than 50 relocations within the study area were excluded from the analysis to ensure
good model convergence. Details of these excluded individuals can be found in
Supplementary Material I.

Environmental Data

This analysis focused on three environmental factors: community stream, open sewers
and domestic rubbish piles. The latter factor represented areas where rats were more
likely to be found, whilst the other factors represented risks of having close contact with
Leptospira contaminated muds or waters. The location of these different points of
interest in the study area were mapped by trained research teams.

These environmental factors were included in analyses in two ways: using distance
rasters and buffer rasters. A 1 meter resolution raster was created for each
environmental factor by calculating the nearest distance for each pixel to the reference
points. The buffer rasters, one for each factor, were created using a 20 meter buffer
around each reference point. The size of this buffer was decided after visiting the study
areas and represented an area within which it could be considered a strong interaction
with the point of interest. All pixels within this buffer were assigned a value of 1, whilst
those outside were given a value of 0. Buffers were used to understand the effect of the
immediate vicinity of each reference point on movement behaviours. Buffer rasters were
also created for each individual's household location, with a 10 meter buffer around

each location.This represented space within and immediately outside each house. This
buffer size accounted for the size of dwellings in these study areas.

Movement Analysis

The analysis was performed in two phases (Figure 2). Firstly, each individual's data was
analysed alongside the environmental factors. This phase created a set of
parameters—called selection coefficients—for each individual. These selection
coefficients were specific to each of the environmental factors. In the second phase, the
selection coefficient for a particular environmental factor was analysed across the study
population. This phase incorporated the individual characteristics for each participant:
gender, age and *Leptospira* serological status. These phases are detailed below. All
analyses were carried out in R, version 4.2.1,[11] using tools from tidyverse.[12] Specific
movement analyses were carried out using package amt.[13]

[**Figure 2** - Schematic diagram showing what data sources are used in which model,

and how models are linked with each other. The blue sections represent phase one, the

individual-level models, whilst the orange section represent phase two, the population-

level model.]

Phase 1: Individual-level model

Drawing from the current methodological developments in animal movement ecology,

we used step-selection functions to characterise individuals' movement behaviours in

relation to the environmental factors described above. Step-selection functions are a

type of movement analysis method that fall under the Resource Selection umbrella.

They can also be classified as spatio-temporal point process models.[14] In these models,

an individual's location at time point i (μ_i) is conditioned on the previous location it was

in (μ_{i-1}), the selection coefficients of the environment (β) and the available space the

individual could have travelled to (θ).

$$[\mu_i|\mu_{i-1}, \beta, \theta] \equiv \frac{g(x(\mu_i), \beta) f(\mu_i|\mu_{i-1}, \Delta_i, \theta)}{\int g(x(\mu), \beta) f(\mu|\mu_{i-1}, \Delta_i, \theta) \, d\mu}$$

Step-selection functions have two important components: the availability function ($f(\dots)$)

and the selection function ($g(\dots)$). The availability function defines the available space

that an individual could move inside of within a set of space and time constraints. The

selection function specifies how the individual responds to the environmental factors

that are close to them when choosing their path, creating a set of selection coefficients

for each factor—or resource–included in the model. These selection coefficients are

specific to a given individual. This latter component is the focus of our analysis, whilst

the former availability function was pre-defined using the empirical data.

The availability function was fitted separately to each recorded location. The step
lengths and turning angles between consecutive steps were used to parametrise
movement characteristics for an individual (Figure 3A). Using these characteristics, a
group of available steps (Figure 3B, grey dots) was created for each used step (Figure
3B, black dots). These represented locations that were consistent with human
movements that an individual could have travelled to but chose not to. A total of 100
available steps were created for each used step (Figure 3B).

[**Figure 3** - Descriptive diagram of step-selection functions. A: step lengths (sl) and
turning angles (θ) are used to characterise an individual's movements. B: these
parameters are used to create a set of available steps (grey dots) for every used step
(black dots).]

Each individual's telemetry data was analysed by time periods within daytime active
hours. These were periods of 4 hours, representing morning (05:00 – 09:00), midday
(09:00 – 13:00), afternoon (13:00 – 17:00) and evening (17:00 – 21:00) activities.
Movements across the whole daytime period were also analysed (05:00 – 21:00). This
analysis was performed to examine the effects of circular journeys, when people travel
to and back from a same place using a very similar route. By looking at specific time
periods, we hoped to capture one-way journeys. As with the full day analysis, any

individuals with less than 50 relocations within the period of analysis were removed from
the models.

A conditional logistic regression was used to estimate the selection coefficients for each
of the environmental variables for a given individual. A separate model was used for
each time period.

$$logit(p) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \beta_4 x_4 + \beta_5 x_5 \beta_6 x_6 + \beta_7 x_7 + \beta_8 x_8 + \alpha_{stratum_j}$$

The model estimated the odds of a step being used compared to it being available and
unused (p), with a logit transformation ($logit(p)$). The first three variables included in
the model ($x_1 - x_3$) represented the different environmental factors (central stream,
open sewers and domestic rubbish piles) and their corresponding selection coefficients
($\beta_1 - \beta_3$). Distance rasters and buffer rasters were included in separate models. The
household buffer rasters were included in the next variable (x_4). The following three
variables ($x_5 - x_7$) represent the movement characteristics of the individual: the step
length (sl), the natural logarithm of the step lengths ($log(sl)$) and the cosine of the
turning angle ($cos(\theta)$). These are the same movement characteristics used to create the
set of available steps. The final variable included (x_8) was the hour within which each
step was recorded. The model was stratified by each used step ($\alpha_{stratum_i}$), where j
represents each used step and its associated available steps. This model estimates a
selection coefficient for each of the environmental factors of interest, conditioned on all
other environmental factors, the individual's household location, the individual's
movement characteristics and the hour of the day. These selection coefficients can be
interpreted as the likelihood of moving into a specific environmental condition whilst
keeping other environmental factors, movement characteristics and hour of the day
constant. For distance rasters, the selection coefficient represents odds of moving
further away from the reference point. For buffer rasters, it represents the odds of
moving inside of the 20 meter buffer of each reference point.

272 Phase 2: Population level model

To assess movement differences between individual characteristics, a population-level
linear regression model was used. Separate models were created for each of the three

environmental factors, using their corresponding selection coefficients as the outcome,

and for each time period (whole daytime period, morning, midday, afternoon and

evening). We used two main group of models: (1) those assessing differences between

genders and ages, which were conditioned on both of these variables and the study

area; (2) those assessing differences between *Leptospira* antibody statuses, which

were conditioned on gender, age and study area. The shared equation for each of the

models is defined as follows:

$$\hat{\beta}_k = \gamma_0 + \gamma_1 x_1 + \gamma_2 x_2 + \gamma_3 x_3 + \gamma_4 x_4 + Z_k$$

In these models, the outcome was the estimated selection coefficient ($\hat{\beta}$) for each

environmental factor (k). The first two variables, x_1 and x_2, represented gender (taking

values 0 for male and 1 for female), and age, used as a continuous variable. The third

variable, x_3, represented *Leptospira* antibody status, as a binary variable taking values

1 for a positive test and 0 otherwise. As mentioned previously, a positive result was

defined as a positive MAT result for any *Leptospira* serovar. The final variable in the

model, x_4, represented the study area, included to adjust for any unmeasured

differences between study areas. The error term, Z_k, captured the residuals from the

model, which also accounted for any variation between individuals which was not

measured as well as the sampling error inherent to the estimates of the selection

coefficients. To account for variation in the standard errors of the selection coefficients,

the variance of Z_k was defined as w_k/τ^2, where w_k is the estimated variance of $\hat{\beta}_k$

which was used to account for the heterogeneity in the estimate of $\hat{\beta}_k$.

Results

Descriptive statistics

There were a total of 130 individuals who consented to take part in this movement

study. Of these, 2 individuals were removed from further analysis due to not having

sufficient re-locations within the study area boundaries during the 5 am to 9 pm period.

They were both male, older than 50 and tested negative for leptospirosis serology. The

remaining 128 individuals represented 11.7% of the sample population from the parent

study (n = 1086). Of the participants in the movement study, 59 (46.0%) were female

and their ages ranged from 18 to 83, with a median age of 38 and mean age of 39.5 (sd

= 15.5). There were 13 individuals (10.2%) who tested positive for *Leptospira*

antibodies. Although these proportions were very similar to those present in the larger

sample population from the parent study, the individuals in the movement analysis

skewed female and older (Table 1).

Table 1 – Summary table comparing parent study participants and movement study participants

	Parent study participants		Movement study participants	
	n = 1086	*% / mean*	*n = 128*	*% / mean*
Study area				
1 – Nova Sussuarana	297	27.3%	32	25.0%
2 – Arenoso	246	22.7%	28	21.9%
3 – Jardim Santo Inacio	278	25.6%	35	27.3%
4 – Calabetao	265	24.4%	33	25.7%
Gender				
Female	454	41.8%	59	46.1%
Male	632	58.2%	69	53.9%
Age (mean +/- SD)	32.2	+/- 19.7	39.4	+/- 15.4
***Leptospira* antibody status**				
Positive	94	8.7%	13	10.2%
Negative	992	91.3%	115	89.8%

The majority of individuals spent most of their recorded time during their active daytime
hours within their study area boundaries. The percentage of recorded time spent within
the study area boundaries ranged from 4% to 100%. The mean percentage was 80%,
with a median of 91% and a standard deviation of 25%. Females spent less time within
the boundaries than men (females: mean = 76%, sd = 28%; males: mean = 83%, sd =
22%). Individuals who had antibodies against *Leptospira* spent the same time within the
study area boundaries as individuals with no antibodies (positive: mean = 83%, sd =
26%; negative: mean = 80%, sd = 25%).

The maximum values for the different environmental distance rasters varied across the
four study areas. The maximum distance to open sewers was lowest in study area 3
and highest in study area 2 (1: 199 m; 2: 235m; 3: 80 m; 4: 208 m). Similarly, the
maximum distance to domestic rubbish piles was lowest in study area 3 and highest in
study area 2 (1: 214 m; 2: 363 m; 3: 153 m; 4: 247 m). These differences are attributed
to the number of open sewer points and domestic rubbish piles within each study area.
The maximum distance to the central stream was highest in study area 1 and lowest in
study area 3 (1: 217 m; 2: 209 m; 3: 94 m; 4: 172 m).

Similarly, the proportion of tracked time spent within the various environmental buffers
varied across characteristics. These can be found in table 2. There were significant
differences between characteristics and within characteristics, represented by the high
standard deviations. More detailed descriptive statistics are available in Supplementary
Material I.

Table 2: Proportion of tracked time (full day period, 9 am to 5 pm) spent within each buffer. Mean (Standard Deviation)

	River buffer	Open sewer buffer	Domestic rubbish buffer

Total		0.54 (0.33)	0.33 (0.31)	0.07 (0.15)
Area	1 - Nova Sussuarana	0.54 (0.29)	0.43 (0.29)	0.15 (0.18)
	2 - Arenoso	0.65 (0.25)	0.28 (0.29)	0.03 (0.13)
	3 - Jardim Santo Inacio	0.36 (0.35)	0.41 (0.31)	0.08 (0.18)
	4 - Calabetao	0.67 (0.31)	0.21 (0.28)	0.01 (0.02)
Gender	Female	0.56 (0.32)	0.37 (0.32)	0.07 (0.15)
	Male	0.53 (0.34)	0.30 (0.29)	0.06 (0.16)
Leptospirosis serological status	Negative	0.55 (0.33)	0.35 (0.31)	0.07 (0.16)
	Positive	0.48 (0.29)	0.25 (0.30)	0.02 (0.05)

Movement analysis

338 The results from the movement analysis are presented in the odds scale. A positive
339 value represents higher odds of moving towards an increasing value for each raster. As
340 described previously, for distance rasters this is interpreted as moving further away from
341 the point of reference (Table 3), whilst for buffer rasters this is interpreted as moving
342 into the 20 meter buffer area for each point of reference (Table 4).

Table 3 - Estimated differences (γ) in selection coefficients (ß) for each environmental factor using distance-based rasters. Values >1 represent increasing distance from points of reference

	Community Stream		Open Sewers		Domestic Rubbish Piles	
	Estimate	*95% CI*	*Estimate*	*95% CI*	*Estimate*	*95% CI*
Gender *						
Male	(Ref)	-	(Ref)	-	(Ref)	-
Female	0.98	0.97, 0.99	1.04	1.02, 1.06	0.99	0.98, 1.01
Age **						
Per year increase	1.00	1.00, 1.00	1.00	1.00, 1.00	1.00	1.00, 1.00
Leptospira serological status ‡						

Negative	(Ref)	-	(Ref)	-	(Ref)	-
Positive	0.99	0.96, 1.01	1.03	1.00, 1.07	1.00	0.98, 1.02

* Adjusted for age and study area

** Adjusted for gender and study area. Values represent increases by one year of age

‡ Adjusted for gender, age and study area

We found no differences in how individuals moved in regards to the distance to the
central stream by age (OR: 1.00; 95% CI: 1.00, 1.00; p = 0.697) or *Leptospira* antibody
status (OR: 0.99; 95% CI: 0.96, 1.01; p = 0.273). Similarly, movements relative to the 20
meter buffer for the central stream were the same across ages (OR: 1.00; 95% CI: 1.00,
1.01; p = 0.280) and across *Leptospira* serological status (OR: 0.89; 95% CI: 0.67, 1.19;
p = 0.433). There was evidence that women moved closer to the stream than men, even
after accounting for the effects of age, study area and the location of their households
(OR: 0.98; 95% CI: 0.97, 0.99; p = 0.003). This effect was more pronounced in the
analysis of the 20 meter buffered area (OR: 1.22, 95% CI: 1.02, 1.46; p = 0.026).

Table 4 - Estimated differences (γ) in selection coefficients (β) for each environmental factor using 20 meter buffers around each point of reference. Values >1 represent movement within the buffer zone for each point of reference.

	Community Stream		Open Sewers		Domestic Rubbish Piles	
	Estimate	*95% CI*	*Estimate*	*95% CI*	*Estimate*	*95% CI*
Gender *						
Male	(Ref)	-	(Ref)	-	(Ref)	-
Female	1.22	1.02, 1.46	0.95	0.80, 1.14	0.92	0.66, 1.27
Age **						
Per year increase	1.00	1.00, 1.00	0.99	0.98, 1.00	1.00	0.99, 1.01
Leptospira serological status ‡						
Negative	(Ref)	-	(Ref)	-	(Ref)	-
Positive	0.89	0.67, 1.19	0.64	0.47, 0.87	0.85	0.48, 1.49

* Adjusted for age and study area

** Adjusted for gender and study area. Values represent increases by one year of age

‡ Adjusted for gender, age and study area

As with the above, there was no evidence of different movement behaviours relative to

distance to open sewers by age (OR 1.00; 95% CI: 1.00, 1.00; p = 0.572) or *Leptospira*

antibody status (OR: 1.03; 95% CI: 1.00, 1.07; p = 0.054). Women were found to move

further away from open sewers compared to men (OR 1.04; 95% CI: 1.02, 1.06; p <

0.001). When analysing movements relative to the 20 meter buffer around open sewers,

we found no evidence of differences between genders (OR: 0.95; 0.80, 1.14; p = 0.580).

We found evidence of a small tendency to move outside of the 20 meter buffer around

open sewers as people aged, although the effect could be considered negligible (OR:

0.99; 95% CI: 0.98, 1.00; p = 0.003). We also found evidence of a strong inclination for

people with *Leptospira* antibodies to move outside of the buffers around open sewers,

compared to people with no antibodies (OR: 0.64; 95% CI: 0.47, 0.87; p = 0.005).

Our analysis showed no evidence of different movement behaviours relative to the

distance to rubbish piles across genders (OR: 0.99; 95% CI: 0.98, 1.01; p = 0.280),

ages (OR: 1.00; 95% CI: 1.00, 1.00; p = 0.466) or *Leptospira* antibody statuses (OR:

1.00; 95% CI: 0.98, 1.02; p = 0.760). We also found no evidence when analysing

movements relative to the 20 meter buffer around rubbish piles across genders (OR:

0.92; 95% CI: 0.66, 1.27; p = 0.600), ages (OR: 1.00; 95% CI: 0.99, 1.01; p = 0.989) or

Leptospira antibody statuses (OR: 0.80; 95% CI: 0.44, 1.49; p = 0.482).

It is important to highlight that the effect sizes of the selection coefficients for the

distance based rasters (Table 3) are very small and could be considered negligible. This

may be linked to the spatial scale used, as these values represent increases of 1 meter.

A coarser scale may have produced larger effect sizes that may have been easier to

conceptualise. However, given the focus on fine-scale movement, we decided to keep

this spatial scale for the analysis.

Analysis by time periods

Movements were subdivided into four time periods: morning (5 am -- 9 am), midday (9

am -- 1 pm), afternoon (1 pm -- 5 pm) and evening (5 pm -- 9 pm). The demographic

characteristics of all individuals removed from analyses for having less than 50

relocations within a specific time period can be found in Supplementary Material I. The

interactions with the environmental factors were similar to those reported for whole day
activities, although there were some key differences (Figure 4).

We found no differences in movements relative to the central stream as people aged or
between *Leptospira* antibody status across the four periods. Women still moved closer
to the central stream than men across all periods. We also saw that women had a
higher tendency to move within the 20 meter buffer for the stream compared to men
across all periods.

Movement in relation to distance to open sewer points and their respective 20 meter
buffers showed no difference across all four periods. The strength of the selection effect
seen in serological positive individuals for moving outside of the 20 meter buffer varied,
with stronger effects seen in the morning and evening periods.

Domestic rubbish piles did not appear to have an effect on movement differences

between ages or *Leptospira* antibody status across all periods. We found women

moved outside of the 20 meter buffer zone more than men during the morning period

only. Otherwise, no notable differences were seen.

[**Figure 4** - Graph showing results of final analyses: A) results for distance based
rasters, values above 1 interpreted as increasing distance to points of reference; B)
results for 20 meter buffer based rasters, values above 1 show movement within buffer
zones. Each horizontal band represents a specific time period (right hand side y-axis
label): all day (5 am – 9 pm, Tables 2 and 3), morning (5 am – 9 am), midday (9 am – 1
pm), afternoon (1pm – 5 pm) and evening (5 pm – 9 pm). All data points include their
corresponding 95% confidence intervals, some of which are too narrow to show up
clearly.]

Discussion

Our study aimed to apply a novel methodology to the area of human mobility analysis in
infectious disease epidemiology, focusing on leptospirosis in four urban slums in
Salvador, Brazil. We assessed movements in relation to central streams, open sewer
points, and domestic rubbish piles and observed changes throughout the day using
step-selection functions. These are a modelling approach which we have taken and
adapted from animal movement ecology. Our findings showed that step selection
functions could be an effective method to identify movement behaviours. To understand
how the results could be described in the context of infectious disease epidemiology, we
have explained our interpretation of the findings, including strengths and limitations.
However, it is important to highlight that, given this is a novel methodology, the
evidence we present is not conclusive and further research is required.

The results suggested no movement differences between *Leptospira* antibody statuses
or ages concerning the distances to stream, open sewer points, or domestic rubbish
piles. Our findings consistently showed that women tended to move closer to the central
stream and farther from open sewer points than men, adjusted for age and study area.
We also found that women had a tendency to move within the 20 meter buffer of the

central streams compared to men, and that seropositive individuals were more inclined

to move outside of the buffer zone for open sewers compared to seronegative

individuals. Movement patterns did not vary significantly throughout the day. Previous

research indicates that men in similar communities perceive themselves as less

vulnerable to leptospirosis compared to women.[15] Additionally, a knowledge, attitudes

and practices analysis showed that men have lower scores for both knowledge and

attitudes towards leptospirosis and its associated risks.[16] Our findings align with these

studies, suggesting that women may avoid open sewers due to perceived risks, while

men may not share these perceptions. Social areas, which may have gender

differences, also contribute to different movement behaviours. One might conclude that

the stream is used for gendered chores such as washing clothes. However, this is not

the case in our study areas. Following discussion with residents, we know that they

perceive the stream as highly contaminated and avoid using its waters for cleaning or

other household chores.

Our results contrast those reported by Owers et al.[1], who found no differences in space

use between genders after using GPS loggers to analyse individuals' movements. This

discrepancy could be explained by the differences in length of time being analysed.

Owers et al. were only able to analyse data collected over 24 hour periods, whereas our

analysis was longer and included data collected over periods of up to 48 hours, which

could be repeated. The contrasting results could also be attributed to the different

populations studied. Although overall these populations resided in very similar

communities in Salvador, they could have different characteristics that affect movement

behaviours.

Our findings regarding the interactions with rubbish piles may be explained by various

reasons. There is evidence that proximity to rubbish piles does not drive *Leptospira*

seropositivity in similar areas to those used in our analysis.[15] Whilst this proximity does

increase rat sightings, this reduced effect on infection risk could lead individuals to

disregard the locations of rubbish piles when choosing their travel paths. Another

possible explanation is that there may be an unmeasured environmental variable that is

interacting with the distance to rubbish piles, which needs further investigation. For

example, violence could be interacting with where rubbish accumulates. We discuss
violence further in a paragraph below.

The evidence showing *Leptospira* positive individuals avoiding open sewers was
surprising. Although we were expecting to see an effect in the opposite direction,
showing individuals with *Leptospira* antibodies interacting closely with open sewers,
there are a few possible explanations for our findings. If individuals with antibodies are
also actively infected, they could be symptomatic and therefore alter their behaviour to
avoid high risk areas. Alternatively, individuals with antibodies could be more aware of
risks due to previous infections and display more protective behaviours than people who
have not had any previous infections.

During informal conversations with community residents, it became clear that violence
plays a key role in individual's decisions on where they go. Violence in these
communities is perceived as hyper-local, restricted to one corner or small square within
the communities. It is unclear what drives this perception, but nevertheless it is an
important factor that could be accounted for. Further research is required to develop
methods that can capture these perceptions in spatial formats that could be
incorporated into similar movement studies. Age did not affect movement choices,
suggesting consistent perceptions of environmental risks or stable use of urban spaces
across ages.

We expected different movement patterns at various times of day, anticipating circular
journeys (an individual going somewhere and back again on the same route). However,
our results showed consistent movement patterns, possibly due to the analysis period's
length or other unmeasured factors modulating movements. Our results could also be
indicative of evidence that strictly circular journeys through these communities, where
individuals are travelling through the exact same path for both journeys, are not
common, and that movement interactions with urban surroundings do not vary
throughout the day.

To our knowledge, this is the first study that uses step-selection functions to model
movement behaviours in the context of human infectious disease epidemiology. This
method has provided quantitative evidence that there may be differences in how men

and women move through their communities, strengthening the argument that the

variation in leptospirosis exposure and infection risk between genders is due to

behavioural differences rather than physiological differences. Additionally, we show that

individuals consider environmental features differently when moving through their

communities. Highlighting the effects of these variables on movement would not have

been possible with the approaches previously used to model human movement. Our

approach provides a better understanding of how individuals relate to their surrounding

urban environment and how they interact with features that could increase risk of

leptospirosis.

Several important limitations must be highlighted. This study involves a relatively small

sample of a larger population, slightly skewed towards older women compared to the

parent study. There are few individuals testing positive for *Leptospira* antibodies. As a

result, the findings are biased towards the more represented individuals, limiting their

generalisability. Additionally, all participants are from specific areas in Salvador, which

may further limit the generalisability to similar contexts. Further research is needed to

develop appropriate study designs using these methods, including how many

individuals should be recruited. The small number of *Leptospira*-positive individuals also

makes the estimation for the effect of this characteristic more difficult. We would also

like to restate that a positive antibody response to any *Leptospira* serovar does not

indicate active infection. A positive result merely indicates that the individual has been

infected at some stage, either symptomatically or asymptomatically, and has produced

an immune response. Information on the timing of the infection could instead be a

variable showing a stronger association with movement. Another important limitation is

that we did not collect data on behaviours. If risky or protective behaviours, such as the

use of closed footware, had been available at the appropriate temporal resolution (e.g.

hourly intervals), these could have been included in the step selection functions and

could have shown significant associations. Although these are important limitations

which require cautious interpretation of results, they do not detract from the value of

exploring this novel methodology in this context. This methodology also provides a

crucial starting point for exploring how movement characteristics can differ between

individuals in these environments.

Step-selection functions also have limitations that must be considered. While these

methods can model the choice of moving in a specific direction, they do not account for

the initial distance from the individual. For instance, an individual moving towards the

central stream from far away will have a high selection coefficient for this environmental

factor, which does not indicate their starting distance. This is important because

environmental risk factors cease to provide risk beyond a certain distance. This

limitation was overcome by using buffer zones around specific points of interest, but it is

crucial to highlight to correctly interpret all results. Similarly, step-selection functions do

not quantify how long an individual spent within this high-risk distance. Additionally,

these models have some underlying assumptions that may be violated in this study.

Step-selection functions assume each step is independent, conditioned on the previous

step. This can be violated by circular journeys. Although we attempted to account for

these by analysing specific periods of the day, a higher temporal resolution of analysis

may be needed if circular journeys are still present within each period. Another

assumption is that movement is smooth through the environment. In urban

environments this may not hold true, as street layouts may force sharp corners in

movements. The effect of violating this assumption is not immediately clear and

requires further methodological research to understand its significance. Finally, we

assumed that by including movement characteristics (step lengths and turning angles)

into our models, we were accounting for goal-oriented behaviour. These assumptions

need to be considered in future studies that attempt to use step-selection functions to

analyse human mobility.

Despite these limitations, this study has several valuable strengths. By including steps

an individual could have taken but did not (i.e., available steps, grey dots in Figure 3B),

the models allow us to estimate choice. Additionally, the models use each individual's

movement characteristics to create these available steps, resulting in a realistic

representation of movement behaviours. This creates more realistic estimates of

environmental interactions than those created using existing methods.

Another significant strength is the specificity of the individual-level and population-level

models. First, the population-level linear regression models allow multiple individual

characteristics to be included, producing results that can be adjusted as needed.

Although not considered in this study, these models also provide flexibility in the type of

variable interactions that can be specified, allowing for non-linear effects if necessary.

Second, the individual-level conditional logistic regression models are conditioned for all

included variables. This enables the estimation of the selection coefficients for each

environmental factor after adjusting for potential confounders. This is particularly useful

in our case, as open sewer points are often close to the central stream in all study areas

(Figure 1).

Overall, we believe this method is a useful tool in analysing human mobility in the

context of infectious disease epidemiology. This modelling approach could also be used

in other areas of research which analyse human movements and choice relating to

surrounding environmental features, such as urban planning. A major benefit of step

selection functions is the use of rasters, which provide flexibility when investigating

environmental features. Creative uses of rasters could provide interesting questions and

results. Although the focus of these models is looking at choice in space, the methods

could also be adapted to analyse choice in time (e.g. are there temporal variables that

affect when a rat enters a household).

To conclude, we provide a worked example of how to use step-selection functions to

analyse human movements in the field of infectious disease epidemiology. This

highlights the usefulness of adapting methods from other fields to answer questions that

would otherwise be difficult to answer with the existing methodology. By doing so, we

develop a better understanding of environmental interactions and how to leverage the

large data sets provided by GPS loggers. Although our focus was leptospirosis, these

methods can be adapted to model the exposure to any disease where movement and

the environment play an important role.

Ethics

Ethical approval for this study was obtained from the ethics committee at the Collective Health Institute, Federal University of Bahia (CEP/ISC/UFBA) under number CAEE 32361820.7.0000.5030, and the national research ethics committee (CONEP) linked to the Brazilian Ministry of Health under approval number 4.235.251. All participants involved in the study provided written informed consent before data collection.

Authors' contributions

P.R.C., F.N.S., F.C., and E. G.: conceptualization, formal analysis, investigation, methodology, data curation, software, visualization, writing—original draft, writing—review and editing; J.R., C.C. , F.C. and E.G.: Supervision; F.C.: funding acquisition; M.T.E., J. R. and C.C.: formal analysis, writing—original draft, writing—review and editing; J.O.S, D.S.O., R.C.N., A.G.S., E.V.R.S., F.A.G, D.C.C.S and P.S.R: investigation, field team, laboratory, data curation, writing—review and editing.

Funding

This work was supported by the Wellcome Trust, NIHR/Wellcome Global health Partnership (218987/Z/19/Z) to F.C.; F.N.S. received a research scholarship from the Brazilian National Research Council (CNPq:150142/2024-2). PRC is in receipt of a studentship from the Medical Research Council, United Kingdom. ME was supported through a Reckitt Global Hygiene Institute fellowship. JMR acknowledges support from National Institute of Allergy and Infectious Diseases (Grant number 1R01AI160780-01).

Acknowledgments

We would like to thank all residents from the study areas, without whom this work would not have been able to be completed. The authors would also like to specially thank the GIC (Grupo Impulsor Comunitario) for their support and warmth.

References

1. Owers, K. A. *et al.* Fine-scale GPS tracking to quantify human movement patterns and exposure to leptospires in the urban slum environment. *PLoS Negl. Trop. Dis.* **12**, e0006752 (2018).

2. Fornace, K. M. *et al.* Local human movement patterns and land use impact exposure to zoonotic malaria in Malaysian Borneo. *eLife* **8**, e47602 (2019).

3. Kraemer, M. U. G. *et al.* Mapping global variation in human mobility. *Nat. Hum. Behav.* **4**, 800–810 (2020).

4. Moncayo-Unda, M. G., Van Droogenbroeck, M., Saadi, I. & Cools, M. A longitudinal analysis of the COVID-19 effects on the variability in human activity spaces in Quito, Ecuador. *J. Transp. Geogr.* **113**, 103735 (2023).

5. Costa, F. *et al.* Global Morbidity and Mortality of Leptospirosis: A Systematic Review. *PLoS Negl. Trop. Dis.* **9**, e0003898 (2015).

6. Adler, B. & de la Peña Moctezuma, A. *Leptospira* and leptospirosis. *Vet. Microbiol.* **140**, 287–296 (2010).

7. Johnson, M. A. S. *et al.* Environmental Exposure and Leptospirosis, Peru - Volume 10, Number 6—June 2004 - Emerging Infectious Diseases journal - CDC. doi:10.3201/eid1006.030660.

8. Reis, R. B. *et al.* Impact of Environment and Social Gradient on Leptospira Infection in Urban Slums. *PLoS Negl. Trop. Dis.* **2**, e228 (2008).

9. Hagan, J. E. *et al.* Spatiotemporal Determinants of Urban Leptospirosis Transmission: Four-Year Prospective Cohort Study of Slum Residents in Brazil. *PLoS Negl. Trop. Dis.* **10**, e0004275 (2016).

10. Cremonese, C. *et al.* Simplified sewerage to prevent urban leptospirosis

transmission: a cluster non-randomised controlled trial protocol in disadvantaged

urban communities of Salvador, Brazil. *BMJ Open* **13**, e065009 (2023).

11. R Core Team. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing (2024).

12. Wickham, H. & RStudio. tidyverse: Easily Install and Load the 'Tidyverse'. (2023).

13. Signer, J. *et al.* amt: Animal Movement Tools. (2024).

14. Fieberg, J., Signer, J., Smith, B. & Avgar, T. A 'How to' guide for interpreting

parameters in habitat-selection analyses. *J. Anim. Ecol.* **90**, 1027–1043 (2021).

15. Khalil, H. *et al.* Poverty, sanitation, and Leptospira transmission pathways in

residents from four Brazilian slums. *PLoS Negl. Trop. Dis.* **15**, e0009256 (2021).

16. Palma, F. A. G. *et al.* Why is leptospirosis hard to avoid for the impoverished?

Deconstructing leptospirosis transmission risk and the drivers of knowledge,

attitudes, and practices in a disadvantaged community in Salvador, Brazil. *PLOS*

Glob. Public Health **2**, e0000408 (2022).

Supplementary Material I

Descriptive Statistics

Telemetry data

The mean number of hours of telemetry data provided by an individual was 13.3 hours, with a
standard deviation of 13.5 hours. The mean number of locations recorded by the GPS loggers
was 2767 points (SD = 1947.2). There were no differences in the number of hours or number of
locations recorded by gender, age or leptospirosis antibody status. There were notable
differences in the number of hours recorded and the number of locations by study area. Study
area 1 (NVS) had the lowest number of hours recorded (mean = 5.6 hours, SD = 5.6), whilst all
other areas had similar hours recorded (area 2: mean = 15.0, SD = 11.4; area 3: mean = 10.9, SD
= 14.0; area 4; mean = 20.7, SD = 15.3). The mean number of locations recorded were all similar
across all study areas (area 1: mean = 2048, SD = 1206; area 2: mean = 2831, SD = 1302; area 3:
mean = 2992, SD = 2737; area 4: mean = 3107, SD = 1761).

 Figure 1 – ...eriods (x

659   axis), separated into each of the four study areas. Overlapping areas represent multiple days.

660   Vertical bars represent 5 am (left hand bar) and 9 pm (right hand bar), the period of analysis.

Excluded individuals

664   Table 1 – Demographic details of excluded individuals due to having less than 50 relocations

ID (anonymised)	Relocation below 50	Period	Gender	Age group	Leptospirosis serological status
60	TRUE	05-21	Male	50-54	Neg
91	TRUE	05-21	Male	>55	Neg
15	TRUE	05-09	Female	45-49	Neg
60	TRUE	05-09	Male	50-54	Neg
81	TRUE	05-09	Female	50-54	Neg
91	TRUE	05-09	Male	>55	Neg
108	TRUE	05-09	Female	50-54	Neg
109	TRUE	05-09	Male	20-24	Neg
128	TRUE	05-09	Male	25-29	Neg
129	TRUE	05-09	Male	40-44	Pos
15	TRUE	09-13	Female	45-49	Neg
24	TRUE	09-13	Female	50-54	Neg
60	TRUE	09-13	Male	50-54	Neg
70	TRUE	09-13	Male	35-39	Neg
71	TRUE	09-13	Female	35-39	Neg
76	TRUE	09-13	Male	35-39	Neg
91	TRUE	09-13	Male	>55	Neg

108	TRUE	09-13	Female	50-54	Neg
109	TRUE	09-13	Male	20-24	Neg
128	TRUE	09-13	Male	25-29	Neg
129	TRUE	09-13	Male	40-44	Pos
24	TRUE	13-17	Female	50-54	Neg
60	TRUE	13-17	Male	50-54	Neg
71	TRUE	13-17	Female	35-39	Neg
76	TRUE	13-17	Male	35-39	Neg
91	TRUE	13-17	Male	>55	Neg
5	TRUE	17-21	Male	35-39	Neg
7	TRUE	17-21	Male	50-54	Neg
18	TRUE	17-21	Male	30-34	Neg
22	TRUE	17-21	Male	45-49	Neg
24	TRUE	17-21	Female	50-54	Neg
27	TRUE	17-21	Male	30-34	Neg
30	TRUE	17-21	Female	>55	Neg
60	TRUE	17-21	Male	50-54	Neg
71	TRUE	17-21	Female	35-39	Neg
91	TRUE	17-21	Male	>55	Neg
114	TRUE	17-21	Female	40-44	Neg

Serological data

670 Serologically positive individuals were equally distributed across ages and genders, although the

671 oldest male included in the analysis was also serologically positive (Figure 2).

[Figure 2 – Distribution of Leptospirosis antibody status (serological status) by gender and age.]

There were also no significant skews in the household characteristics relative to the

environmental factors being analyzed (Figure 3).

[FIGURE 3 – Distribution of nearest distance to each of the environmental factors being

analysed (central stream, open sewer points and domestic rubbish piles) by serological status and

study area. NA represents rest of households in study area that did not take part in movement

analysis.]

Laboratory work

All samples were tested using the MAT test, the reference test for serological diagnosis of

leptospirosis, as designated by the WHO. The diagnostic panel used included the following

serovars:

- *L. kirschneri* serovar Cynopteri strain 3522C

- *L. kirschneri* serovar Grippothyphosa strain Duyster

- *L. interrogans* serovar Canicola strain H. Ultrech

- *L. interrogans* serovar Autumnlais strain Akiyami A

- *L. borgspetersenii* serovar Ballum strain MUS 127

- *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (locally isolated in 1996)

- *L. interrogans* serovar Copenhageni strain Fiocruz LV3954