Using step selection functions to analyse human mobility using telemetry data in infectious disease epidemiology: a case study of leptospirosis

Pablo Ruiz Cuenca1 , Fabio Neves Souza2,3 , Roberta Coutinho do Nascimento3 , Ariane Goncalves da Silva2 , Max T. Eyre4 , Juliet O. Santana2,3 , Daiana Santos de Oliveira2,3 , Emile Victoria Ribeiro de Souza2 , Fabiana Almerinda G. Palma2 , Diogo César de Carvalho Santiago2 , Priscyla dos Santos Ribeiro5,6 , Jonathan M. Read1 , Cleber Cremonese2 , Federico Costa*2,3,7 , and Emanuele Giorgi*1

1 Centre for Health Informatics, Computing, and Statistics (CHICAS), Lancaster Medical School, Lancaster University, United Kingdom

2 Institute of Collective Health, Federal University of Bahia, Salvador, Bahia, Brazil

3 Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Ministério da Saúde, Salvador, Bahia,Brazil

4 Environmental Health Group, London School of Hygiene and Tropical Medicine, London,United Kingdom

5 Federal University of Bahia, National Institute of Science and Technology in Interdisciplinary and Transdiciplinary Studies in Ecology and Evolution, Salvador, Brazil

6 Amsterdam University Medical Centre, Leptospirosis Reference Center, Medical Microbiology and Infection Control, Amsterdam , Netherlands

7 Department of Epidemiology of Microbial Diseases, School of Public Health, Yale University, New Haven, CT, United States

* These authors contributed equally to this work.

Abstract

Human movement plays a critical role in the transmission of infectious diseases, especially those with environmental drivers like leptospirosis—a zoonotic bacterial infection linked to mud and water contact. Using GPS loggers, we collected detailed telemetry data to understand how fine-scale movements can be analysed in the context of an infectious disease. We recruited individuals living in urban slums in Salvador, Brazil to analyse how they interact with environmental risk factors such as domestic rubbish piles, open sewers, and a local stream. We aimed to identify differences in movement patterns inside the study areas by gender, age, and leptospirosis serological status. Step-selection functions, a spatio-temporal model used in animal movement ecology, estimated selection coefficients to represent the likelihood of movement toward specific environmental factors. With 128 participants wearing GPS devices for 24 to 48 hours, recording locations every 35 seconds during active daytime hours, we segmented movements into morning, midday, afternoon, and evening. Our results suggested women moved closer to the central stream and farther from open sewers compared to men, while serologically positive individuals avoided open sewers. This study introduces a novel method for analysing human telemetry data in infectious disease research.

Keywords: Leptospirosis, human movement, GPS, urban health, infectious diseases, zoonosis

Introduction

GPS loggers are a growing tool for capturing both human and animal movements.[1,2] These small devices can be worn by individuals and record locations at regular preset time intervals. Compared to other methods of collecting human movements, such as cell tower traffic or Google Location History which are suited for analysing large-scale mobility,[3,4] these devices can capture very fine-scale movements. These data are crucial in quantifying exposure within complex environments, where terrain can change rapidly. Furthermore, movements recorded by GPS loggers can be assigned to specific individuals. This allows linkage between individual socio-demographic factors and the data collected, especially convenient when performing

epidemiological analyses. Other methods for measuring human mobility are inherently

anonymous and do not allow this connection to be made. An important challenge when using

GPS loggers is that they rely on individual compliance for carrying the device at all times, an

issue which is overcome by the other methods mentioned above.

The analysis of human telemetry data is an emerging field of research in epidemiology. Whilst

previous methods have advanced this area of research, improvements could be made. For

example, the methods used by Owers et al.[1] to assess the relationship between urban slum

residents' movements and the risk of leptospirosis infection were able to analyse differences

between genders, but did not consider other important socio-demographic factors. In another

study, Fornace et al.[2] used GPS loggers to assess human exposure to mosquito vectors of

Plasmodium knowlesi malaria and environmental factors associated with this. Various

individual-level factors were included in the analyses performed in this paper, questioning how

these could affect participants' movements. However, by not including comparisons of possible

choices an individual could have made, this study could not determine how the environment may

have influenced movement.

Leptospirosis is a zoonotic bacterial infectious disease with strong environmental drivers. It has

been estimated to cause over 1 million yearly human cases worldwide, leading to 58 900 deaths.[5]

Rats are the main reservoir of the disease, shedding bacteria in their urine.[6] Human infection is

associated with exposure to contaminated waters and soils.[6–8]Evidence shows that in urban slum

settings, men have a higher infection risk than women.[9] This has been attributed to differences in

behaviours, especially in how individuals move through their communities, rather than a

biological differences. Indeed, there is evidence that men tend to visit much larger areas during

their daily journeys than women.[1]

Exactly where people are most exposed to high leptospirosis contamination, and therefore where

infection is most likely to occur, has not been investigated. Previous studies have focused on the

assessment of the peri-domiciliary environment and its associations to infection risk.[7–9]

However, these analyses assume people are mostly exposed to infection risk in this area and

ignore the exposure that individuals may incur when they move further away from their

households. Furthermore, people's movement patterns may differ depending on individual socio-

demographic factors which could in turn affect their risk of exposure. If individuals traverse

highly contaminated areas where the risk of exposure is heightened, it can lead to an increased
risk of infection. This is particularly important in environmentally heterogeneous areas, such as
urban slums, where the landscape can change drastically in small spaces. Technological
advances now allow us to record and analyse fine-scale movements to understand how these may
affect infection risk.

In this paper, we developed a modelling framework to understand how telemetry data can be
used to identify and quantify determinants of human movements, adapting methods from animal
movement ecology. We present a novel method for analysing telemetry data to estimate
environmental selection as individuals move through their urban communities. This method is
applied in a low-income urban setting in Salvador, Brazil, and is used to examine how
individuals interact with various key points in their surrounding environment . Furthermore, we
analyse if there are any differences in movements inside the study areas between genders, ages
and leptospirosis serological status . This method of analysis overcomes limitations from other
studies by, firstly, specifically modelling choice of movement in relation to environmental
factors and, secondly, incorporating multiple socio-demographic factors which allows regression
relationships to be jointly adjusted for these.

Methods

Study areas

This study was nested in a prospective cohort study taking place across Salvador, Brazil.[10]
Salvador is the third largest city in Brazil, located in the north-eastern region of the country and
has a tropical climate. The study areas are considered urban slums (locally called `favelas`).
They were selected for a number of reasons: firstly, they all have similar demographic and socio-
economic factors within their populations; secondly, they all have a stream running through the
centre of the community, which is considered contaminated; and thirdly, there is a high burden
of leptospirosis in these populations.

All four study areas are small, with an approximate size of 0.03 km2. They are located across the
outskirts of Salvador (Figure 1). The communities have very heterogeneous environments, with

rapid changes in both land cover and slope. Buildings in these communities have been built with

limited or no urban planning. They can be of varying quality, ranging from gated areas with

multiple dwellings protected from rain and flooding to single brick buildings with informal

entryways.

[**Figure 1** - Map showing location of each study area in Salvador. Each area includes symbology

for stream (blue line), open sewer points (purple diamond), and domestic rubbish piles (orange

triangle).]

Individual characteristics

The eligibility criteria for inclusion in the study was: individuals who (1) had been living at one

of the study areas for at least 6 months, (2) slept there at least 3 nights a week, (3) were at least

18 years old and (4) gave written consent.[10] Participants were asked to answer a baseline survey

which collected their demographic, social and economic characteristics, including age and

gender. A blood sample was taken from each participant to determine serological evidence for

Leptospira infection using the microscopic agglutination test (MAT), the standard test used for
leptospirosis diagnosis.[6] In this analysis, a MAT showing antibodies with a titre >1:50 against
any *Leptospira* serovar was considered a positive result. Further details about the laboratory
work carried out are available in Supplementary Material I. The location of their household was
recorded and georeferenced by the research team.

Participants who were already enrolled in the cohort study were recruited to take part in the
movement analysis study. At the time of recruitment, we found no published scientific studies
detailing how to perform sample size calculations for research using GPS data in humans.
Therefore, we opted to use convenience sampling instead. A target of 30 people per study area,
balanced by gender and blind to their serological status, was chosen for this study.

GPS Data

Individuals who consented to take part in this study were asked to wear GPS loggers for
continuous periods of up to 48 hours, which could be repeated. The GPS loggers used were i-got
U GT-600, set to record their location every 35 seconds. We used the manufacturer's software to
programme the devices. Data were collected between March and November 2022.

Once the GPS telemetry data was collected, participants' recorded locations were cleaned so as
to retain only relocations within the study area boundaries that were recorded between 5 am and
9 pm. This period generally corresponds to an individual's active hours. Interactions with
environmental factors outside of the study area boundaries could not be considered in the
analysis because high-resolution environmental data outside of the study areas was not available.
Individuals with less than 50 relocations within the study area were excluded from the analysis to
ensure good model convergence. Details of these excluded individuals can be found in
Supplementary Material I.

Environmental Data

This analysis focused on three environmental factors: community stream, open sewers and
domestic rubbish piles. The latter factor represented areas where rats were more likely to be

found, whilst the other factors represented risks of having close contact with *Leptospira*

contaminated muds or waters. The location of these different points of interest in the study area

were mapped by trained research teams.

These environmental factors were included in analyses in two ways: using distance rasters and

buffer rasters. A 1 meter resolution raster was created for each environmental factor by

calculating the nearest distance for each pixel to the reference points. The buffer rasters, one for

each factor, were created using a 20 meter buffer around each reference point. The size of this

buffer was decided after visiting the study areas and represented an area within which it could be

considered a strong interaction with the point of interest. All pixels within this buffer were

assigned a value of 1, whilst those outside were given a value of 0. Buffers were used to

understand the effect of the immediate vicinity of each reference point on movement behaviours.

Buffer rasters were also created for each individual's household location, with a 10 meter buffer

around each location.This represented space within and immediately outside each house. This

buffer size accounted for the size of dwellings in these study areas.

Movement Analysis

The analysis was performed in two phases (Figure 2). Firstly, each individual's data was

analysed alongside the environmental factors. This phase created a set of parameters—called

selection coefficients—for each individual. These selection coefficients were specific to each of

the environmental factors. In the second phase, the selection coefficient for a particular

environmental factor was analysed across the study population. This phase incorporated the

individual characteristics for each participant: gender, age and *Leptospira* serological status.

These phases are detailed below. All analyses were carried out in R, version 4.2.1,[11] using tools

from tidyverse.[12] Specific movement analyses were carried out using package amt.[13]

[**Figure 2** - Schematic diagram showing what data sources are used in which model, and how

models are linked with each other. The blue sections represent phase one, the individual-level

models, whilst the orange section represent phase two, the population-level model.]

Phase 1: Individual-level model

Drawing from the current methodological developments in animal movement ecology, we used

step-selection functions to characterise individuals' movement behaviours in relation to the

environmental factors described above. Step-selection functions are a type of movement analysis

method that fall under the Resource Selection umbrella. They can also be classified as spatio-

temporal point process models.[14] In these models, an individual's location at time point i (μ_i) is

conditioned on the previous location it was in (μ_{i-1}), the selection coefficients of the

environment (β) and the available space the individual could have travelled to (θ).

$$[\mu_i|\mu_{i-1}, \beta, \theta] \equiv \frac{g(x(\mu_i), \beta)f(\mu_i|\mu_{i-1}, \Delta_i, \theta)}{\int g(x(\mu), \beta)f(\mu|\mu_{i-1}, \Delta_i, \theta)\, d\mu}$$

Step-selection functions have two important components: the availability function ($f(...)$) and the selection function ($g(...)$). The availability function defines the available space that an individual could move inside of within a set of space and time constraints. The selection function specifies how the individual responds to the environmental factors that are close to them when choosing their path, creating a set of selection coefficients for each factor—or resource–included in the model. These selection coefficients are specific to a given individual. This latter component is the focus of our analysis, whilst the former availability function was pre-defined using the empirical data.

The availability function was fitted separately to each recorded location. The step lengths and turning angles between consecutive steps were used to parametrise movement characteristics for an individual (Figure 3A). Using these characteristics, a group of available steps (Figure 3B, grey dots) was created for each used step (Figure 3B, black dots). These represented locations that were consistent with human movements that an individual could have travelled to but chose not to. A total of 100 available steps were created for each used step (Figure 3B).

[**Figure 3** - Descriptive diagram of step-selection functions. A: step lengths (sl) and turning angles (θ) are used to characterise an individual's movements. B: these parameters are used to create a set of available steps (grey dots) for every used step (black dots).]

Each individual's telemetry data was analysed by time periods within daytime active hours.
These were periods of 4 hours, representing morning (05:00 – 09:00), midday (09:00 – 13:00),
afternoon (13:00 – 17:00) and evening (17:00 – 21:00) activities. Movements across the whole
daytime period were also analysed (05:00 – 21:00). This analysis was performed to examine the
effects of circular journeys, when people travel to and back from a same place using a very
similar route. By looking at specific time periods, we hoped to capture one-way journeys. As
with the full day analysis, any individuals with less than 50 relocations within the period of
analysis were removed from the models.

A conditional logistic regression was used to estimate the selection coefficients for each of the
environmental variables for a given individual. A separate model was used for each time period.

$$logit(p) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \beta_4 x_4 + \beta_5 x_5 \beta_6 x_6 + \beta_7 x_7 + \beta_8 x_8 + \alpha_{stratum_j}$$

The model estimated the odds of a step being used compared to it being available and unused
(p), with a logit transformation $(logit(p))$. The first three variables included in the model
$(x_1 - x_3)$ represented the different environmental factors (central stream, open sewers and
domestic rubbish piles) and their corresponding selection coefficients $(\beta_1 - \beta_3)$. Distance rasters
and buffer rasters were included in separate models. The household buffer rasters were included
in the next variable (x_4). The following three variables $(x_5 - x_7)$ represent the movement
characteristics of the individual: the step length (sl), the natural logarithm of the step lengths
$(log(sl))$ and the cosine of the turning angle $(cos(\theta))$. These are the same movement
characteristics used to create the set of available steps. The final variable included (x_8) was the
hour within which each step was recorded. The model was stratified by each used step
$(\alpha_{stratum_i})$, where j represents each used step and its associated available steps. This model
estimates a selection coefficient for each of the environmental factors of interest, conditioned on
all other environmental factors, the individual's household location, the individual's movement
characteristics and the hour of the day. These selection coefficients can be interpreted as the
likelihood of moving into a specific environmental condition whilst keeping other environmental
factors, movement characteristics and hour of the day constant. For distance rasters, the selection

243   coefficient represents odds of moving further away from the reference point. For buffer rasters, it
244   represents the odds of moving inside of the 20 meter buffer of each reference point.

Phase 2: Population level model

To assess movement differences between individual characteristics, a population-level linear
regression model was used. Separate models were created for each of the three environmental
factors, using their corresponding selection coefficients as the outcome, and for each time period
(whole daytime period, morning, midday, afternoon and evening). We used two main group of
models: (1) those assessing differences between genders and ages, which were conditioned on
both of these variables and the study area; (2) those assessing differences between *Leptospira*
antibody statuses, which were conditioned on gender, age and study area. The shared equation
for each of the models is defined as follows:

$$\hat{\beta}_k = \gamma_0 + \gamma_1 x_1 + \gamma_2 x_2 + \gamma_3 x_3 + \gamma_4 x_4 + Z_k$$

In these models, the outcome was the estimated selection coefficient ($\hat{\beta}$) for each environmental
factor (k). The first two variables, x_1 and x_2, represented gender (taking values 0 for male and 1
for female), and age, used as a continuous variable. The third variable, x_3, represented
Leptospira antibody status, as a binary variable taking values 1 for a positive test and 0
otherwise. As mentioned previously, a positive result was defined as a positive MAT result for
any *Leptospira* serovar. The final variable in the model, x_4, represented the study area, included
to adjust for any unmeasured differences between study areas. The error term, Z_k, captured the
residuals from the model, which also accounted for any variation between individuals which was
not measured as well as the sampling error inherent to the estimates of the selection coefficients.
To account for variation in the standard errors of the selection coefficients, the variance of Z_k
was defined as w_k/τ^2, where w_k is the estimated variance of $\hat{\beta}_k$ which was used to account for
the heterogeneity in the estimate of $\hat{\beta}_k$.

269 Results

270 Descriptive statistics

There were a total of 130 individuals who consented to take part in this movement study. Of
these, 2 individuals were removed from further analysis due to not having sufficient re-locations
within the study area boundaries during the 5 am to 9 pm period. They were both male, older
than 50 and tested negative for leptospirosis serology. The remaining 128 individuals represented
11.7% of the sample population from the parent study (n = 1086). Of the participants in the
movement study, 59 (46.0%) were female and their ages ranged from 18 to 83, with a median
age of 38 and mean age of 39.5 (sd = 15.5). There were 13 individuals (10.2%) who tested
positive for *Leptospira* antibodies. Although these proportions were very similar to those present
in the larger sample population from the parent study, the individuals in the movement analysis
skewed female and older (Table 1).

Table 1 – Summary table comparing parent study participants and movement study participants

	Parent study participants		Movement study participants	
	n = 1086	*% / mean*	*n = 128*	*% / mean*
Study area				
1 – Nova Sussuarana	297	27.3%	32	25.0%
2 – Arenoso	246	22.7%	28	21.9%
3 – Jardim Santo Inacio	278	25.6%	35	27.3%
4 – Calabetao	265	24.4%	33	25.7%
Gender				
Female	454	41.8%	59	46.1%
Male	632	58.2%	69	53.9%
Age (mean +/- SD)	32.2	+/- 19.7	39.4	+/- 15.4
Leptospira **antibody status**				
Positive	94	8.7%	13	10.2%

Negative	992	91.3%	115	89.8%

The majority of individuals spent most of their recorded time during their active daytime hours within their study area boundaries. The percentage of recorded time spent within the study area boundaries ranged from 4% to 100%. The mean percentage was 80%, with a median of 91% and a standard deviation of 25%. Females spent less time within the boundaries than men (females: mean = 76%, sd = 28%; males: mean = 83%, sd = 22%). Individuals who had antibodies against *Leptospira* spent the same time within the study area boundaries as individuals with no antibodies (positive: mean = 83%, sd = 26%; negative: mean = 80%, sd = 25%).

The maximum values for the different environmental distance rasters varied across the four study areas. The maximum distance to open sewers was lowest in study area 3 and highest in study area 2 (1: 199 m; 2: 235m; 3: 80 m; 4: 208 m). Similarly, the maximum distance to domestic rubbish piles was lowest in study area 3 and highest in study area 2 (1: 214 m; 2: 363 m; 3: 153 m; 4: 247 m). These differences are attributed to the number of open sewer points and domestic rubbish piles within each study area. The maximum distance to the central stream was highest in study area 1 and lowest in study area 3 (1: 217 m; 2: 209 m; 3: 94 m; 4: 172 m).

Similarly, the proportion of tracked time spent within the various environmental buffers varied across characteristics. These can be found in table 2. There were significant differences between characteristics and within characteristics, represented by the high standard deviations. More detailed descriptive statistics are available in Supplementary Material I.

Table 2: Proportion of tracked time (full day period, 9 am to 5 pm) spent within each buffer. Mean (Standard Deviation)

	River buffer	Open sewer buffer	Domestic rubbish buffer
Total	0.54 (0.33)	0.33 (0.31)	0.07 (0.15)
Area 1 - Nova Sussuarana	0.54 (0.29)	0.43 (0.29)	0.15 (0.18)

	2 - Arenoso	0.65 (0.25)	0.28 (0.29)	0.03 (0.13)
	3 - Jardim Santo Inacio	0.36 (0.35)	0.41 (0.31)	0.08 (0.18)
	4 - Calabetao	0.67 (0.31)	0.21 (0.28)	0.01 (0.02)
Gender	Female	0.56 (0.32)	0.37 (0.32)	0.07 (0.15)
	Male	0.53 (0.34)	0.30 (0.29)	0.06 (0.16)
Leptospirosis serological status	Negative	0.55 (0.33)	0.35 (0.31)	0.07 (0.16)
	Positive	0.48 (0.29)	0.25 (0.30)	0.02 (0.05)

Movement analysis

The results from the movement analysis are presented in the odds scale. A positive value represents higher odds of moving towards an increasing value for each raster. As described previously, for distance rasters this is interpreted as moving further away from the point of reference (Table 3), whilst for buffer rasters this is interpreted as moving into the 20 meter buffer area for each point of reference (Table 4).

Table 3 - Estimated differences (γ) in selection coefficients (β) for each environmental factor using distance-based rasters. Values >1 represent increasing distance from points of reference

	Community Stream		Open Sewers		Domestic Rubbish Piles	
	Estimate	*95% CI*	*Estimate*	*95% CI*	*Estimate*	*95% CI*
Gender *						
Male	(Ref)	-	(Ref)	-	(Ref)	-
Female	0.98	0.97, 0.99	1.04	1.02, 1.06	0.99	0.98, 1.01
Age **						
Per year increase	1.00	1.00, 1.00	1.00	1.00, 1.00	1.00	1.00, 1.00
Leptospira serological status ‡						
Negative	(Ref)	-	(Ref)	-	(Ref)	-
Positive	0.99	0.96, 1.01	1.03	1.00, 1.07	1.00	0.98, 1.02

* Adjusted for age and study area

** Adjusted for gender and study area. Values represent increases by one year of age

 ‡ Adjusted for gender, age and study area

We found no differences in how individuals moved in regards to the distance to the central

stream by age (OR: 1.00; 95% CI: 1.00, 1.00; p = 0.697) or *Leptospira* antibody status (OR:

0.99; 95% CI: 0.96, 1.01; p = 0.273). Similarly, movements relative to the 20 meter buffer for

the central stream were the same across ages (OR: 1.00; 95% CI: 1.00, 1.01; p = 0.280) and

across *Leptospira* serological status (OR: 0.89; 95% CI: 0.67, 1.19; p = 0.433). There was

evidence that women moved closer to the stream than men, even after accounting for the effects

of age, study area and the location of their households (OR: 0.98; 95% CI: 0.97, 0.99; p = 0.003).

This effect was more pronounced in the analysis of the 20 meter buffered area (OR: 1.22, 95%

CI: 1.02, 1.46; p = 0.026).

Table 4 - Estimated differences (γ) in selection coefficients (β) for each environmental factor using 20 meter buffers around each point of reference. Values >1 represent movement within the buffer zone for each point of reference.

	Community Stream		Open Sewers		Domestic Rubbish Piles	
	Estimate	*95% CI*	*Estimate*	*95% CI*	*Estimate*	*95% CI*
Gender *						
Male	(Ref)	-	(Ref)	-	(Ref)	-
Female	1.22	1.02, 1.46	0.95	0.80, 1.14	0.92	0.66, 1.27
Age **						
Per year increase	1.00	1.00, 1.00	0.99	0.98, 1.00	1.00	0.99, 1.01
Leptospira serological status ‡						
Negative	(Ref)	-	(Ref)	-	(Ref)	-
Positive	0.89	0.67, 1.19	0.64	0.47, 0.87	0.85	0.48, 1.49

* Adjusted for age and study area

** Adjusted for gender and study area. Values represent increases by one year of age

‡ Adjusted for gender, age and study area

As with the above, there was no evidence of different movement behaviours relative to distance

to open sewers by age (OR 1.00; 95% CI: 1.00, 1.00; p = 0.572) or *Leptospira* antibody status

(OR: 1.03; 95% CI: 1.00, 1.07; p = 0.054). Women were found to move further away from open

sewers compared to men (OR 1.04; 95% CI: 1.02, 1.06; p < 0.001). When analysing movements

relative to the 20 meter buffer around open sewers, we found no evidence of differences between

genders (OR: 0.95; 0.80, 1.14; p = 0.580). We found evidence of a small tendency to move

outside of the 20 meter buffer around open sewers as people aged, although the effect could be

considered negligible (OR: 0.99; 95% CI: 0.98, 1.00; p = 0.003). We also found evidence of a

strong inclination for people with *Leptospira* antibodies to move outside of the buffers around

open sewers, compared to people with no antibodies (OR: 0.64; 95% CI: 0.47, 0.87; p = 0.005).

Our analysis showed no evidence of different movement behaviours relative to the distance to

rubbish piles across genders (OR: 0.99; 95% CI: 0.98, 1.01; p = 0.280), ages (OR: 1.00; 95% CI:

1.00, 1.00; p = 0.466) or *Leptospira* antibody statuses (OR: 1.00; 95% CI: 0.98, 1.02; p = 0.760).

We also found no evidence when analysing movements relative to the 20 meter buffer around

rubbish piles across genders (OR: 0.92; 95% CI: 0.66, 1.27; p = 0.600), ages (OR: 1.00; 95% CI:

0.99, 1.01; p = 0.989) or *Leptospira* antibody statuses (OR: 0.80; 95% CI: 0.44, 1.49; p = 0.482).

It is important to highlight that the effect sizes of the selection coefficients for the distance based

rasters (Table 3) are very small and could be considered negligible. This may be linked to the

spatial scale used, as these values represent increases of 1 meter. A coarser scale may have

produced larger effect sizes that may have been easier to conceptualise. However, given the

focus on fine-scale movement, we decided to keep this spatial scale for the analysis.

Analysis by time periods

Movements were subdivided into four time periods: morning (5 am -- 9 am), midday (9 am -- 1

pm), afternoon (1 pm -- 5 pm) and evening (5 pm -- 9 pm). The demographic characteristics of

all individuals removed from analyses for having less than 50 relocations within a specific time

period can be found in Supplementary Material I. The interactions with the environmental

factors were similar to those reported for whole day activities, although there were some key

differences (Figure 4).

We found no differences in movements relative to the central stream as people aged or between

Leptospira antibody status across the four periods. Women still moved closer to the central

stream than men across all periods. We also saw that women had a higher tendency to move

within the 20 meter buffer for the stream compared to men across all periods.

Movement in relation to distance to open sewer points and their respective 20 meter buffers

showed no difference across all four periods. The strength of the selection effect seen in

serological positive individuals for moving outside of the 20 meter buffer varied, with stronger

effects seen in the morning and evening periods.

Domestic rubbish piles did not appear to have an effect on movement differences between ages

or *Leptospira* antibody status across all periods. We found women moved outside of the 20 meter

buffer zone more than men during the morning period only. Otherwise, no notable differences

were seen.

[**Figure 4** - Graph showing results of final analyses: A) results for distance based rasters, values

above 1 interpreted as increasing distance to points of reference; B) results for 20 meter buffer

based rasters, values above 1 show movement within buffer zones. Each horizontal band

represents a specific time period (right hand side y-axis label): all day (5 am – 9 pm, Tables 2

and 3), morning (5 am – 9 am), midday (9 am – 1 pm), afternoon (1pm – 5 pm) and evening (5

pm – 9 pm). All data points include their corresponding 95% confidence intervals, some of

which are too narrow to show up clearly.]

Discussion

Our study aimed to apply a novel methodology to the area of human mobility analysis in

infectious disease epidemiology, focusing on leptospirosis in four urban slums in Salvador,

Brazil. We assessed movements in relation to central streams, open sewer points, and domestic

rubbish piles and observed changes throughout the day using step-selection functions. These are

a modelling approach which we have taken and adapted from animal movement ecology. Our

findings showed that step selection functions could be an effective method to identify movement

behaviours. To understand how the results could be described in the context of infectious disease

epidemiology, we have explained our interpretation of the findings, including strengths and

limitations. However, it is important to highlight that, given this is a novel methodology, the

evidence we present is not conclusive and further research is required.

The results suggested no movement differences between *Leptospira* antibody statuses or ages

concerning the distances to stream, open sewer points, or domestic rubbish piles. Our findings

consistently showed that women tended to move closer to the central stream and farther from

open sewer points than men, adjusted for age and study area. We also found that women had a

tendency to move within the 20 meter buffer of the central streams compared to men, and that

seropositive individuals were more inclined to move outside of the buffer zone for open sewers

compared to seronegative individuals. Movement patterns did not vary significantly throughout

the day. Previous research indicates that men in similar communities perceive themselves as less

vulnerable to leptospirosis compared to women.[15] Additionally, a knowledge, attitudes and

practices analysis showed that men have lower scores for both knowledge and attitudes towards

leptospirosis and its associated risks.[16] Our findings align with these studies, suggesting that

women may avoid open sewers due to perceived risks, while men may not share these

perceptions. Social areas, which may have gender differences, also contribute to different

movement behaviours. One might conclude that the stream is used for gendered chores such as

washing clothes. However, this is not the case in our study areas. Following discussion with

residents, we know that they perceive the stream as highly contaminated and avoid using its

waters for cleaning or other household chores.

Our results contrast those reported by Owers et al.[1], who found no differences in space use

between genders after using GPS loggers to analyse individuals' movements. This discrepancy

could be explained by the differences in length of time being analysed. Owers et al. were only

able to analyse data collected over 24 hour periods, whereas our analysis was longer and

included data collected over periods of up to 48 hours, which could be repeated. The contrasting

results could also be attributed to the different populations studied. Although overall these

populations resided in very similar communities in Salvador, they could have different

characteristics that affect movement behaviours.

Our findings regarding the interactions with rubbish piles may be explained by various reasons.

There is evidence that proximity to rubbish piles does not drive *Leptospira* seropositivity in

similar areas to those used in our analysis.[15] Whilst this proximity does increase rat sightings,

this reduced effect on infection risk could lead individuals to disregard the locations of rubbish

piles when choosing their travel paths. Another possible explanation is that there may be an

unmeasured environmental variable that is interacting with the distance to rubbish piles, which

needs further investigation. For example, violence could be interacting with where rubbish

accumulates. We discuss violence further in a paragraph below.

The evidence showing *Leptospira* positive individuals avoiding open sewers was surprising.

Although we were expecting to see an effect in the opposite direction, showing individuals with

Leptospira antibodies interacting closely with open sewers, there are a few possible explanations

for our findings. If individuals with antibodies are also actively infected, they could be

symptomatic and therefore alter their behaviour to avoid high risk areas. Alternatively,

individuals with antibodies could be more aware of risks due to previous infections and display

more protective behaviours than people who have not had any previous infections.

During informal conversations with community residents, it became clear that violence plays a

key role in individual's decisions on where they go. Violence in these communities is perceived

as hyper-local, restricted to one corner or small square within the communities. It is unclear what

drives this perception, but nevertheless it is an important factor that could be accounted for.

Further research is required to develop methods that can capture these perceptions in spatial

formats that could be incorporated into similar movement studies. Age did not affect movement

choices, suggesting consistent perceptions of environmental risks or stable use of urban spaces

across ages.

We expected different movement patterns at various times of day, anticipating circular journeys

(an individual going somewhere and back again on the same route). However, our results showed

consistent movement patterns, possibly due to the analysis period's length or other unmeasured

factors modulating movements. Our results could also be indicative of evidence that strictly

circular journeys through these communities, where individuals are travelling through the exact

same path for both journeys, are not common, and that movement interactions with urban

surroundings do not vary throughout the day.

To our knowledge, this is the first study that uses step-selection functions to model movement

behaviours in the context of human infectious disease epidemiology. This method has provided

quantitative evidence that there may be differences in how men and women move through their

communities, strengthening the argument that the variation in leptospirosis exposure and

infection risk between genders is due to behavioural differences rather than physiological

differences. Additionally, we show that individuals consider environmental features differently

when moving through their communities. Highlighting the effects of these variables on

movement would not have been possible with the approaches previously used to model human

movement. Our approach provides a better understanding of how individuals relate to their
surrounding urban environment and how they interact with features that could increase risk of
leptospirosis.

Several important limitations must be highlighted. This study involves a relatively small sample
of a larger population, slightly skewed towards older women compared to the parent study. There
are few individuals testing positive for *Leptospira* antibodies. As a result, the findings are biased
towards the more represented individuals, limiting their generalisability. Additionally, all
participants are from specific areas in Salvador, which may further limit the generalisability to
similar contexts. Further research is needed to develop appropriate study designs using these
methods, including how many individuals should be recruited. The small number of *Leptospira*-
positive individuals also makes the estimation for the effect of this characteristic more difficult.
We would also like to restate that a positive antibody response to any *Leptospira* serovar does
not indicate active infection. A positive result merely indicates that the individual has been
infected at some stage, either symptomatically or asymptomatically, and has produced an
immune response. Information on the timing of the infection could instead be a variable showing
a stronger association with movement. Another important limitation is that we did not collect
data on behaviours. If risky or protective behaviours, such as the use of closed footware, had
been available at the appropriate temporal resolution (e.g. hourly intervals), these could have
been included in the step selection functions and could have shown significant associations.
Although these are important limitations which require cautious interpretation of results, they do
not detract from the value of exploring this novel methodology in this context. This methodology
also provides a crucial starting point for exploring how movement characteristics can differ
between individuals in these environments.

Step-selection functions also have limitations that must be considered. While these methods can
model the choice of moving in a specific direction, they do not account for the initial distance
from the individual. For instance, an individual moving towards the central stream from far away
will have a high selection coefficient for this environmental factor, which does not indicate their
starting distance. This is important because environmental risk factors cease to provide risk
beyond a certain distance. This limitation was overcome by using buffer zones around specific
points of interest, but it is crucial to highlight to correctly interpret all results. Similarly, step-

selection functions do not quantify how long an individual spent within this high-risk distance.

Additionally, these models have some underlying assumptions that may be violated in this study.

Step-selection functions assume each step is independent, conditioned on the previous step. This

can be violated by circular journeys. Although we attempted to account for these by analysing

specific periods of the day, a higher temporal resolution of analysis may be needed if circular

journeys are still present within each period. Another assumption is that movement is smooth

through the environment. In urban environments this may not hold true, as street layouts may

force sharp corners in movements. The effect of violating this assumption is not immediately

clear and requires further methodological research to understand its significance. Finally, we

assumed that by including movement characteristics (step lengths and turning angles) into our

models, we were accounting for goal-oriented behaviour. These assumptions need to be

considered in future studies that attempt to use step-selection functions to analyse human

mobility.

Despite these limitations, this study has several valuable strengths. By including steps an

individual could have taken but did not (i.e., available steps, grey dots in Figure 3B), the models

allow us to estimate choice. Additionally, the models use each individual's movement

characteristics to create these available steps, resulting in a realistic representation of movement

behaviours. This creates more realistic estimates of environmental interactions than those created

using existing methods.

Another significant strength is the specificity of the individual-level and population-level

models. First, the population-level linear regression models allow multiple individual

characteristics to be included, producing results that can be adjusted as needed. Although not

considered in this study, these models also provide flexibility in the type of variable interactions

that can be specified, allowing for non-linear effects if necessary. Second, the individual-level

conditional logistic regression models are conditioned for all included variables. This enables the

estimation of the selection coefficients for each environmental factor after adjusting for potential

confounders. This is particularly useful in our case, as open sewer points are often close to the

central stream in all study areas (Figure 1).

Overall, we believe this method is a useful tool in analysing human mobility in the context of

infectious disease epidemiology. This modelling approach could also be used in other areas of

research which analyse human movements and choice relating to surrounding environmental
features, such as urban planning. A major benefit of step selection functions is the use of rasters,
which provide flexibility when investigating environmental features. Creative uses of rasters
could provide interesting questions and results. Although the focus of these models is looking at
choice in space, the methods could also be adapted to analyse choice in time (e.g. are there
temporal variables that affect when a rat enters a household).

To conclude, we provide a worked example of how to use step-selection functions to analyse
human movements in the field of infectious disease epidemiology. This highlights the usefulness
of adapting methods from other fields to answer questions that would otherwise be difficult to
answer with the existing methodology. By doing so, we develop a better understanding of
environmental interactions and how to leverage the large data sets provided by GPS loggers.
Although our focus was leptospirosis, these methods can be adapted to model the exposure to
any disease where movement and the environment play an important role.

Ethics

524 Ethical approval for this study was obtained from the ethics committee at the Collective Health
525 Institute, Federal University of Bahia (CEP/ISC/UFBA) under number CAEE
526 32361820.7.0000.5030, and the national research ethics committee (CONEP) linked to the
527 Brazilian Ministry of Health under approval number 4.235.251. All participants involved in the
528 study provided written informed consent before data collection.

Authors' contributions

P.R.C., F.N.S., F.C., and E. G.: conceptualization, formal analysis, investigation, methodology,
data curation, software, visualization, writing—original draft, writing—review and editing; J.R.,
C.C. , F.C. and E.G.: Supervision; F.C.: funding acquisition; M.T.E., J. R. and C.C.: formal
analysis, writing—original draft, writing—review and editing; J.O.S, D.S.O., R.C.N., A.G.S.,
E.V.R.S., F.A.G, D.C.C.S and P.S.R: investigation, field team, laboratory, data curation,
writing—review and editing.

Funding

This work was supported by the Wellcome Trust, NIHR/Wellcome Global health Partnership (218987/Z/19/Z) to F.C.; F.N.S. received a research scholarship from the Brazilian National Research Council (CNPq:150142/2024-2). PRC is in receipt of a studentship from the Medical Research Council, United Kingdom. ME was supported through a Reckitt Global Hygiene Institute fellowship. JMR acknowledges support from National Institute of Allergy and Infectious Diseases (Grant number 1R01AI160780-01).

Acknowledgments

We would like to thank all residents from the study areas, without whom this work would not have been able to be completed. The authors would also like to specially thank the GIC (Grupo Impulsor Comunitario) for their support and warmth.

References

1. Owers, K. A. *et al.* Fine-scale GPS tracking to quantify human movement patterns and exposure to leptospires in the urban slum environment. *PLoS Negl. Trop. Dis.* **12**, e0006752 (2018).

2. Fornace, K. M. *et al.* Local human movement patterns and land use impact exposure to zoonotic malaria in Malaysian Borneo. *eLife* **8**, e47602 (2019).

3. Kraemer, M. U. G. *et al.* Mapping global variation in human mobility. *Nat. Hum. Behav.* **4**, 800–810 (2020).

4. Moncayo-Unda, M. G., Van Droogenbroeck, M., Saadi, I. & Cools, M. A longitudinal analysis of the COVID-19 effects on the variability in human activity spaces in Quito, Ecuador. *J. Transp. Geogr.* **113**, 103735 (2023).

559 5. Costa, F. *et al.* Global Morbidity and Mortality of Leptospirosis: A Systematic Review.

560   *PLoS Negl. Trop. Dis.* **9**, e0003898 (2015).

561 6. Adler, B. & de la Peña Moctezuma, A. *Leptospira* and leptospirosis. *Vet. Microbiol.* **140**,

562   287–296 (2010).

563 7. Johnson, M. A. S. *et al.* Environmental Exposure and Leptospirosis, Peru - Volume 10,

564   Number 6—June 2004 - Emerging Infectious Diseases journal - CDC.

565   doi:10.3201/eid1006.030660.

566 8. Reis, R. B. *et al.* Impact of Environment and Social Gradient on Leptospira Infection in

567   Urban Slums. *PLoS Negl. Trop. Dis.* **2**, e228 (2008).

568 9. Hagan, J. E. *et al.* Spatiotemporal Determinants of Urban Leptospirosis Transmission: Four-

569   Year Prospective Cohort Study of Slum Residents in Brazil. *PLoS Negl. Trop. Dis.* **10**,

570   e0004275 (2016).

571 10. Cremonese, C. *et al.* Simplified sewerage to prevent urban leptospirosis transmission: a

572   cluster non-randomised controlled trial protocol in disadvantaged urban communities of

573   Salvador, Brazil. *BMJ Open* **13**, e065009 (2023).

574 11. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

575   Statistical Computing (2024).

576 12. Wickham, H. & RStudio. tidyverse: Easily Install and Load the 'Tidyverse'. (2023).

577 13. Signer, J. *et al.* amt: Animal Movement Tools. (2024).

578 14. Fieberg, J., Signer, J., Smith, B. & Avgar, T. A 'How to' guide for interpreting parameters in

579   habitat-selection analyses. *J. Anim. Ecol.* **90**, 1027–1043 (2021).

580 15. Khalil, H. *et al.* Poverty, sanitation, and Leptospira transmission pathways in residents from

581   four Brazilian slums. *PLoS Negl. Trop. Dis.* **15**, e0009256 (2021).

582   16. Palma, F. A. G. *et al.* Why is leptospirosis hard to avoid for the impoverished?

583   Deconstructing leptospirosis transmission risk and the drivers of knowledge, attitudes, and

584   practices in a disadvantaged community in Salvador, Brazil. *PLOS Glob. Public Health* **2**,

585   e0000408 (2022).

Supplementary Material I

Descriptive Statistics

Telemetry data

The mean number of hours of telemetry data provided by an individual was 13.3 hours, with a standard deviation of 13.5 hours. The mean number of locations recorded by the GPS loggers was 2767 points (SD = 1947.2). There were no differences in the number of hours or number of locations recorded by gender, age or leptospirosis antibody status. There were notable differences in the number of hours recorded and the number of locations by study area. Study area 1 (NVS) had the lowest number of hours recorded (mean = 5.6 hours, SD = 5.6), whilst all other areas had similar hours recorded (area 2: mean = 15.0, SD = 11.4; area 3: mean = 10.9, SD = 14.0; area 4; mean = 20.7, SD = 15.3). The mean number of locations recorded were all similar across all study areas (area 1: mean = 2048, SD = 1206; area 2: mean = 2831, SD = 1302; area 3: mean = 2992, SD = 2737; area 4: mean = 3107, SD = 1761).

604　Figure 1 –　　　　　　　　　　　　　　　　　　　　　　　　　　　eriods (x

605 axis), separated into each of the four study areas. Overlapping areas represent multiple days.

606 Vertical bars represent 5 am (left hand bar) and 9 pm (right hand bar), the period of analysis.

Excluded individuals

610 Table 1 – Demographic details of excluded individuals due to having less than 50 relocations

ID (anonymised)	Relocation below 50	Period	Gender	Age group	Leptospirosis serological status
60	TRUE	05-21	Male	50-54	Neg
91	TRUE	05-21	Male	>55	Neg
15	TRUE	05-09	Female	45-49	Neg
60	TRUE	05-09	Male	50-54	Neg
81	TRUE	05-09	Female	50-54	Neg
91	TRUE	05-09	Male	>55	Neg
108	TRUE	05-09	Female	50-54	Neg
109	TRUE	05-09	Male	20-24	Neg
128	TRUE	05-09	Male	25-29	Neg
129	TRUE	05-09	Male	40-44	Pos
15	TRUE	09-13	Female	45-49	Neg
24	TRUE	09-13	Female	50-54	Neg
60	TRUE	09-13	Male	50-54	Neg
70	TRUE	09-13	Male	35-39	Neg
71	TRUE	09-13	Female	35-39	Neg
76	TRUE	09-13	Male	35-39	Neg
91	TRUE	09-13	Male	>55	Neg

108	TRUE	09-13	Female	50-54	Neg
109	TRUE	09-13	Male	20-24	Neg
128	TRUE	09-13	Male	25-29	Neg
129	TRUE	09-13	Male	40-44	Pos
24	TRUE	13-17	Female	50-54	Neg
60	TRUE	13-17	Male	50-54	Neg
71	TRUE	13-17	Female	35-39	Neg
76	TRUE	13-17	Male	35-39	Neg
91	TRUE	13-17	Male	>55	Neg
5	TRUE	17-21	Male	35-39	Neg
7	TRUE	17-21	Male	50-54	Neg
18	TRUE	17-21	Male	30-34	Neg
22	TRUE	17-21	Male	45-49	Neg
24	TRUE	17-21	Female	50-54	Neg
27	TRUE	17-21	Male	30-34	Neg
30	TRUE	17-21	Female	>55	Neg
60	TRUE	17-21	Male	50-54	Neg
71	TRUE	17-21	Female	35-39	Neg
91	TRUE	17-21	Male	>55	Neg
114	TRUE	17-21	Female	40-44	Neg

Serological data

616 Serologically positive individuals were equally distributed across ages and genders, although the
617 oldest male included in the analysis was also serologically positive (Figure 2).
618
619

645 [Figure 2 – Distribution of Leptospirosis antibody status (serological status) by gender and age.]
646

There were also no significant skews in the household characteristics relative to the

environmental factors being analyzed (Figure 3).

[FIGURE 3 – Distribution of nearest distance to each of the environmental factors being

analysed (central stream, open sewer points and domestic rubbish piles) by serological status and

study area. NA represents rest of households in study area that did not take part in movement

analysis.]

Laboratory work

All samples were tested using the MAT test, the reference test for serological diagnosis of

leptospirosis, as designated by the WHO. The diagnostic panel used included the following

serovars:

• *L. kirschneri* serovar Cynopteri strain 3522C

• *L. kirschneri* serovar Grippothyphosa strain Duyster

• *L. interrogans* serovar Canicola strain H. Ultrech

• *L. interrogans* serovar Autumnlais strain Akiyami A

• *L. borgspetersenii* serovar Ballum strain MUS 127

• *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (locally isolated in 1996)

• *L. interrogans* serovar Copenhageni strain Fiocruz LV3954