**Title:** Forecastability of infectious disease time series: are some seasons and pathogens intrinsically more difficult to forecast?

**Authors:** Lauren A. White[1] (ORCiD: 0000-0003-3852-5374), Tomás M. León[1] (ORCiD: 0000-0002-7633-677X)

[1]California Department of Public Health, Center for Infectious Diseases, Division of Communicable Disease Control

**Abstract:**
For infectious disease forecasting challenges, individual model performance typically varies across space and time. This phenomenon raises the question: are there properties of the target time series that contribute to a particular season, location, or disease being more difficult to forecast? Here we characterize a time series' future predictability using a forecastability metric that calculates the spectral density of the time series. Forecastability of syndromic influenza hospital admissions for the state of California varied widely across seasons and was positively correlated with peak burden. Next, using archived U.S. state and national forecasts targeting laboratory-confirmed COVID-19 and influenza hospital admissions, we investigated the relationship between forecastability and: (i) population size of the forecasting target, and (ii) forecast performance as measured by mean absolute error, weighted interval score (WIS), and scaled relative WIS. Forecastability increased with increasing population size of the forecasting target, and forecasting performance generally improved with higher forecastability when controlling for population size across scales. These preliminary results support the idea that some targets and respiratory virus seasons may be inherently more difficult to forecast and could help explain inter-seasonal variation in model performance.

**Keywords:** forecastability, Shannon entropy, spectral density, infectious disease forecasting, COVID-19, influenza

**Introduction**

There have been numerous collaborative forecasting efforts across multiple years and diseases including influenza, COVID-19, dengue, and Ebola (1–4). These forecasting challenges or "hubs" provide unified submission targets for modeling teams, evaluate the performance of individual model predictions, and synthesize predictions from multiple submitting teams into a unified, ensemble model. Historically, individual models fail to outperform such ensembles consistently through time (2,5). Forecast performance also tends to decline during periods of rapid change, and certain forecasting targets have been more difficult to forecast than others, e.g., COVID-19 cases compared to COVID-19 deaths (5,6). However, it remains unclear which constituent factors of a model contribute to higher performance, nor is it uncommon for individual model performance to vary within and across seasons and locations (5). Here we ask the inverse question about the underlying forecasting target itself: is there something intrinsic about the qualities of infectious disease time series that make them more or less challenging to forecast?

Other disciplines have used various measures of entropy to characterize the complexity of time series and their propensity for future predictability (7–10). "Forecastability" describes the future predictability—or the inherent uncertainty or surprise—in the signal of a stationary time series. One proposed metric for measuring forecastability calculates the Shannon entropy of the spectral density of the time series (11). This metric is invariant to shifting or scaling, with values near 0% corresponding to white noise and values near 100% corresponding to a single sinusoidal curve (Figure 1). For example, S&P 500 returns which have minimal autocorrelation may have values near 1%, while mean temperature time series with significant autocorrelation lags expected at six and twelve months might have a forecastability closer to 35% (11). This metric does not interact with forecast specifications like model type, forecast horizon, or loss function—depending only on the target time series itself (11). Although there are potentially several ways to define forecastability, we will hereafter use the term "forecastability" to refer to the definition outlined in Goerg (2013). While the 2020 West Nile Virus Forecast Challenge assessed uncertainty in the ensemble model forecast of binned case counts using Shannon entropy (12), we believe this is the first application of a forecastability metric to continuous infectious disease time series.

In the infectious disease forecasting literature, quantile forecasts are commonly evaluated with scoring metrics like mean absolute error (MAE) and weighted interval score (WIS). MAE describes the absolute difference between the forecasted value and the observed value, while WIS also accounts for the dispersion of the forecast's prediction interval. Both of these metrics scale with the absolute size of their target (13). This naturally gives more weight to locations with a higher disease burden. In contrast, applying a log-transformation prior to evaluation means that forecasts are scored based on relative error instead of absolute error, giving equal weight to location targets with smaller population sizes (14). Model performance is commonly evaluated against a baseline model which propagates incidence forward based on values from the prior time point with uncertainty based on changes in incidence observed in prior time intervals (2).

**Figure 1.** Correspondence between example time series (first column), their autocorrelation values (second column) and spectral density in the frequency domain (third column). Time series increase in their "forecastability" from top to bottom, ranging from purely random (first row) to perfectly sinusoidal (fourth row). The second row presents influenza admissions per 100K for the state of Maryland and the third row corresponds to influenza admissions per 100K for the entire United States. The x-axis of the signal column corresponds to days since mandatory reporting for influenza hospital admissions was required beginning February 2, 2021. For consistency, all sample time series are differenced once prior to calculating the forecastability score.

Other studies have found that forecasting performance improves for larger jurisdictions –e.g., states vs. counties (6) and larger vs. smaller counties (15). We further hypothesized that: (1) as population size increases, forecastability of the time series should also increase; and (2) as forecastability increases, the WIS of the natural log transformed forecast should decrease (i.e., forecast performance should improve with higher forecastability, when controlling for population size). We explored the relationship between population size of the representative time series and forecastability: (1) at a county and state-level for California for syndromic influenza hospital admissions data from the California Department of Healthcare Access and Information (HCAI); (2) at a U.S. state and national level for laboratory-confirmed

hospital admissions data for COVID-19 and influenza derived from Health and Human Services (HHS) Patient Impact and Hospital Capacity Data System/National Hospital Safety Network (NHSN) (16). With the latter, we investigated the relationship between forecastability and forecast performance at the state and national scales using archived forecasts from the COVID-19 Forecast Hub (https://covid19forecasthub.org/) and FluSight Forecast challenge (https://github.com/cdcepi/FluSight-forecast-hub)—two collaborative forecasting initiatives that have targeted laboratory-confirmed hospital admissions for COVID-19 and influenza respectively (5,17).

## Methods

### *Data*

California county and state level syndromic influenza hospitalization data were derived from HCAI for the 2000-2022 respiratory virus seasons. State and national, laboratory-confirmed, COVID-19 and influenza hospitalization admission data were obtained from HHS/NHSN via Delphi COVIDcast for the 2022-2024 respiratory virus seasons (16,18). Seasons were classified by the standard influenza season definition of MMWR week 40 to week 39—specifically, October 2, 2022- September 30, 2023 for the 2022-2023 season and October 1, 2023- April 30, 2024 for the 2023-2024 season, where voluntary reporting continued after April 30, 2024. To assess sensitivity to the cut off points of the season definition, an alternative season definition from July 1, 2022-July 30, 2023 & July 1, 2023- April 30, 2024 was also explored.

State and national population size estimates were taken from the 2021 US Census Bureau estimates (19). California county population estimates were taken from 2020 California Department of Finance estimates (20).

To identify the predominant subtype for historical California influenza, the most frequently sequenced subtype was taken from historic Respiratory Laboratory Network (RLN) and clinical sentinel laboratory surveillance data (21).

Historical forecasts for state and national COVID-19 and influenza targets were obtained from the COVID-19 Forecast and the FluSight Forecast challenge, respectively, for the 2022-2023 and 2023-2024 respiratory virus seasons (5,17). Specifically, we focused on a comparison of baseline and ensemble models, as well a few select models that were submitted consistently to both forecasting hubs across both seasons (e.g., CMU-TimeSeries, PSI-DICE, SGroup-RandomForest, UMass-trends_ensemble). Ensemble models from the modeling hubs combine all available models for a given target date and location and usually provide the most robust performance (2,5). In contrast, baseline models offer a reference point for model performance by propagating historical values and uncertainty forward (2).

### *Forecastability*

The forecastability of a stationary timeseries $x_t$ is more formally defined as:

$\Omega(y_t) = 1 - \frac{-\int_{-\pi}^{\pi} f_x(\lambda) \log f_x(\lambda) d\lambda}{\log(2\pi)} \epsilon [0,1]$ where $f_x(\lambda)$ is the normalized spectral density of $x_t$ or $\int_{-\pi}^{\pi} f_x(\lambda) d\lambda = 1$ (11). Time series were assessed for stationarity with both the Kwiatkowski-Phillips-Schmidt-Shin (KPSS) and the Augmented Dickey-Fuller (ADF) tests at an alpha = 0.01. For comparability

across geographies and seasons, time series were differenced once before calculating the forecastability score. The metric of forecastability was calculated using the Omega function of the foreCA package (11).

### Forecast evaluation

Forecasts and their corresponding truth data were first transformed via the natural logarithm to control for the differing magnitude in burden across target locations (14). Point forecast error was assessed with mean absolute error (MAE), which takes the absolute difference between a model's median point predictions $\hat{y}_{i=1:N}$ and the set of observed outcomes $y_{i=1:N}$ for a given model, location, and season such that $MAE = \frac{1}{N}\sum_{i=1}^{N}|y_i - \hat{y}_i|$

Probabilistic forecast accuracy was assessed with the weighted interval score (WIS), which is a metric that compiles the performance of a quantile or interval-based forecasts over a range of prediction intervals (13). The single, interval score for a forecast $F$ with outcome $y$ for prediction interval $(1 - \alpha) \times 100\%$ is given by:

$$IS_\alpha(F, y) = (u - l) + \frac{2}{\alpha}(l - y) \times \mathbf{1}(y < l) + \frac{2}{\alpha}(y - u) \times \mathbf{1}(y > u)$$

Where $\mathbf{1}$ is an indicator function, $u$ is the upper prediction limit and $l$ is the lower prediction limit. The score resolves into three components: $u - l$ represents a penalty incurred for width or dispersion of the prediction interval, $\frac{2}{\alpha}(l - y) \times \mathbf{1}(y < l)$ conveys a penalty for underprediction, and $\frac{2}{\alpha}(y - u) \times \mathbf{1}(y > u)$ conveys a penalty for overprediction.

The WIS then sums intervals scores prediction intervals at different levels $(1 - \alpha_1) < (1 - \alpha_2) < \cdots < (1 - \alpha_K)$ such that:

$$WIS_{\alpha_{\{0:K\}}}(F, y) = \frac{1}{K + \frac{1}{2}}\left(w_0|y - m| + \sum_{k=1}^{K}\{w_k IS_{\alpha_k}(F, y)\}\right)$$

Where $K$ is the total number of quantiles included in the WIS, $m$ is the predictive median forecast value, and $w_k$ is a non-negative, unnormalized weight, often set to: $w_k = \frac{\alpha_k}{2}$ and $w_0 = \frac{1}{2}$ (13).

In this case, both MAE and WIS values approximate relative error since these were computed on the natural logarithm of the forecasts and their respective targets (14).

Lastly, models were scored using relative skill, $\theta_i$, which computes model rankings by taking the geometric mean of all possible pairwise comparisons across all models, as given by:

$$\theta_i = \left(\sum_{m=1}^{M} \theta_{i,m}\right)^{1/M}$$

where $\theta_{i,m}$ represents the mean score ratio of model $i$ and model $m$ out of all possible models $M$. Because this metric is based on pairwise comparisons, it only produces a score when models contribute to the same target for the same date(s) and can therefore help to control for forecast missingness (22). Here we present scaled relative WIS results which normalizes model performance relative to a baseline model; relative WIS values below one indicate that the given model performs better than the baseline model, whereas value greater than one suggest a model performs worse than the baseline model on average.

MAE, mean WIS, and scaled relative risk were calculated for each location, season, and model combination across all four week horizons using the scoringutils package in R (22). All analyses were completed in R (23).

**Results**

When looking at California county and state-level syndromic influenza hospitalizations, for which more seasons of data were available, some respiratory virus seasons were more "forecastable" than others, with the greatest differences occurring for locations with larger population sizes (Fig 2A). For example, forecastability ranged from 19.5% to 41.6% for the state of California for the 2021-2022 and 2017-2018 seasons respectively. Across seasons from 2000-2022, there was a positive relationship between forecastability and both cumulative burden and peak burden of each respective season with an adjusted R-squared of 0.56 and 0.81 respectively (Fig 2B, Supplemental Fig 1). The season with highest peak weakly admissions, cumulative burden, and forecastability was an H3N2 season (Fig 2B, Supplemental Fig 1). Two other H3N2 majority seasons (i.e., 2003-2004, 2005-2006) were outliers with higher-than-expected forecastability given their peak burden. The two seasons bridging the 2009 swine influenza pandemic (2008-2010) were outliers with lower-than-expected forecastability given their burden.

For U.S. state and national laboratory confirmed admissions, forecastability of the time series increased with target population size for both COVID-19 and influenza across U.S. states and nationally (Fig 3A). This pattern was generally of the same order of magnitude for both COVID-19 and influenza, and similar across both the 2022-2023 & 2023-2024 seasons (Fig 3A). As seen in prior studies (6,15), forecast performance generally improved for more populous targets, i.e., for natural log transformed forecasts, MAE and WIS on the log scale decreased with increasing population size for the ensembles (Fig 3B-D) and most individual models (Supplemental Fig 2). Except for the 2022-2023 influenza season, MAE and WIS for the ensemble on the log scale also decreased with higher forecastability, i.e., forecast performance improved with higher forecastability (Fig 3B &C, Table 1). For example, for COVID-19 in 2022-2023 season, forecastability values ranged from 10% (Idaho) to 30% (United States) with corresponding decreases in relative error of MAE from 25% to 21% and relative error of WIS from 26% to 14% for the ensemble. In contrast, baseline model performance showed no such relationship with forecastability, with non-significant or positive slope estimates (Fig 3B & C, Table 1).

Across both COVID-19 and influenza ensembles, rates of change of MAE per unit forecastability ranged from -0.0046 to -0.0086, i.e., every one percent unit increase in forecastability yielded a ~0.46-0.86% decrease in MAE relative error (Table 1). Rates of change of WIS per unit forecastability ranged from -0.0016 to -0.0062 for a corresponding decrease in WIS relative error of ~0.13-0.62% per unit forecastability (Table 1). Relationships between the ensembles' scaled relative skill and forecastability were significant for both seasons and both diseases (Fig 3C, Table 1) with increases in relative skill ranging from -0.0048 to -0.011 per unit of forecastability. Overall, these findings were robust to alternative definitions of seasonal timing (Supplemental Figure 3, Supplemental Table 1).

For lab-confirmed COVID-19 and influenza hospital admissions, the metric of forecastability was very susceptible to data reporting frequency (Fig 4). When compared to a time series of rolling weekly admissions with a daily reporting frequency, time series down sampled to weekly reporting frequency had a decreased forecastability score. While this observed effect increased with population size of the measured time series, population size explained less of the variance in forecastability for the weekly vs. the daily sampled timeseries (Fig 4).

**Figure 2.** Examples of variation in forecastability by season for syndromic HCAI influenza hospitalizations. **(A)** The relationship between forecastability and population size for the state of California and constituent counties by respiratory virus season, derived from HCAI data from 2017-2022 (2020-2021 excluded because of the COVID-19 pandemic) and fitted with a generalized additive model (GAM) for visualization; **(B)** forecastability vs. the natural log of peak influenza weekly admissions for the state of California derived from syndromic HCAI data from 2001-2022 and fitted with a linear regression ($\beta$= 7.49, p < 0.001). The year labels in the plot correspond to the first year of the MMWR season, e.g., 2001 corresponds to the 2001-2002 respiratory virus season.

**Figure 3.** The relationship between population size, forecastability, and forecast performance across two respiratory virus seasons for laboratory-confirmed (HHS/NHSN) COVID-19 and influenza admissions at the U.S. state and national scales. **(A)** Forecastability (Ω) vs. the natural logarithm of population size. Forecast performance for the baseline and ensemble models as measured by: **(B)** MAE and **(C)** mean WIS and **(D)** scaled relative skill for the ensemble model.

**Figure 4.** Comparison of forecastability scores vs. natural logarithm of population size for daily vs. weekly reporting (columns) for COVID-19 vs. influenza laboratory-confirmed hospital admissions (rows) from state and national HHS/NHSN data for the 2022-2023 and 2023-2024 seasons.

**Table 1.** Slope estimates from linear model fits for forecastability (Ω) vs. forecast performance for the ensemble and baseline models targeting laboratory-confirmed (HHS/NHSN) COVID-19 and influenza admissions at the U.S. state and national scales as shown in Figure 2. Values are shown with three significant figures. Bolded rows are those statistically significant at a p = 0.05 threshold.

| Metric | Disease | Model | Season | Estimate (beta) | Standard error | Statistic | p-value |
|--------|---------|-------|--------|-----------------|----------------|-----------|---------|
| MAE | COVID-19 | Baseline | **2022-2023** | **-0.0101** | **0.00237** | **-4.25** | **9.46e-5** |
| | | | **2023-2024** | **-0.00589** | **0.00257** | **-2.29** | **0.0264** |
| | | Ensemble | **2022-2023** | **-0.00857** | **0.00207** | **-4.13** | **1.38e-4** |
| | | | **2023-2024** | **-0.00464** | **0.00216** | **-2.15** | **0.0367** |
| | Influenza | Baseline | 2022-2023 | -0.00428 | 0.00389 | -1.1 | 0.276 |
| | | | **2023-2024** | **-0.00491** | **0.00205** | **-2.4** | **0.0202** |
| | | Ensemble | 2022-2023 | -0.00636 | 0.00326 | -1.95 | 0.0566 |
| | | | **2023-2024** | **-0.00691** | **0.00173** | **-3.99** | **2.17e-4** |
| WIS | | Baseline | 2022-2023 | -5.81E-04 | 0.00222 | -0.261 | 0.795 |
| | | | **2023-2024** | **-0.00369** | **0.00183** | **-2.01** | **0.0496** |

| | | | | | | |
|---|---|---|---|---|---|---|
| | COVID-19 | Ensemble | **2022-2023** | **-0.00627** | **0.00125** | **-5.01** | **7.25e-6** |
| | | | **2023-2024** | **-0.00419** | **0.00132** | **-3.18** | **0.00253** |
| | Influenza | Baseline | **2022-2023** | **0.00522** | **0.00247** | **2.11** | **0.04** |
| | | | 2023-2024 | -1.66E-04 | 0.00137 | -0.121 | 0.904 |
| | | Ensemble | 2022-2023 | -0.00167 | 0.0021 | -0.793 | 0.432 |
| | | | **2023-2024** | **-0.00418** | **0.00113** | **-3.7** | **5.38e-4** |
| Scaled relative skill | COVID-19 | Ensemble | **2022-2023** | **-0.0115** | **0.00232** | **-4.97** | **8.11e-6** |
| | | | **2023-2024** | **-0.00487** | **0.00221** | **-2.2** | **0.0325** |
| | Influenza | | **2022-2023** | **-0.0101** | **0.0013** | **-7.75** | **4.1e-10** |
| | | | **2023-2024** | **-0.0104** | **0.00131** | **-7.95** | **1.99e-10** |

**Discussion**

Using a simple signal processing metric to characterize the predictability of infectious disease time series, we showed that there are differences in the measured forecastability of time series across diseases (i.e., COVID-19 vs. influenza, Fig 3A) and across respiratory virus seasons for a given disease (Fig 2). Moreover, these observed differences in measured forecastability correlated with resulting forecast performance. Our results echoed prior work that found a relationship between population size of the forecast target and forecast performance (Fig 3B & C, Supplementary Fig 2) (6,15), but our work further suggests that forecastability increases with population size (Fig 2A, Fig 3A), and forecast performance improves for time series with higher forecastability scores when controlling for population size (Fig 3D). Notably, scaled relative skill of the ensemble against the baseline model improved with increasing forecastability of the target time series across all four combinations of season and disease (Fig 3D), and these gains came from improvements in ensemble performance rather than changes in the baseline model performance, which remained relatively invariant to the time series' forecastability (Fig 3C, Table 1).

Interestingly, there appears to be a strong relationship between peak seasonal burden and forecastability (Fig 2B), which may point to seasons with more singular, sharper peaks having higher forecastability per this particular metric. This finding contrasts with the general pattern that forecasts struggle most during periods of rapid change, such as during the growth phase or at the epidemic peak (5,6). Although anecdotal, patterns in forecastability across seasons matches what might be expected based on epidemiological understanding: forecastability was lower than expected for disrupted influenza seasons (e.g., during the 2009 swine influenza pandemic and in the wake of the COVID-19 pandemic) (Fig 2B). This observed variability in forecastability across seasons could help explain why there has been such variability in both model rankings across years and the ability (or inability) of models able to outperform the baseline model for any given season (5).

The patterns of forecastability observed here raise a couple of questions about the practical implementation of forecasting challenges for infectious diseases. The first is that forecastability is strongly affected by data frequency (e.g., daily vs. weekly data), with gains for higher resolution temporal data increasing with population size (Fig 4). This may matter practically since there are often changes in infectious disease data continuity through time in terms of targets, reporting frequency, and

reporting definitions (e.g., influenza shifting from influenza like illness (ILI) to confirmed hospitalizations during the COVID-19 pandemic, changes in NHSN reporting requirements from daily to weekly in November of 2024). It also highlights that forecasting a completely new signal (e.g., FluSight 2022-2023) remains challenging regardless of the features of the time series.

The second notable pattern is that the there is a consistent plateau of forecastability at about 10% for smaller geographies, regardless of the disease, and that certain seasons were more or less predictable according to the forecastability metric (Figure 2A ,3A). Since the baseline model showed no significant relationship with forecastability (Fig 3C,D), this raises the question of whether forecasting challenges should also be: (1) controlling for some version of signal predictability based on geographic scale or (2) controlling for season predictability post-hoc when scoring forecasts. The latter might matter more for comparing model performance across seasons than within a season. Finally, although there has been increased interest in providing local level forecasts in the wake of the COVID-19 pandemic (15), this also raises questions about infectious disease surveillance for smaller geographies and the inherent feasibility of providing real time infectious disease modeling intelligence at these scales.

Forecastability is an appealing metric because it simplifies a complex time series into a single value; however, the interpretation of this dimension reduction remains challenging. Since forecastability is measured in the frequency domain, it is sometimes difficult to understand which qualities of a time series directly translate to a more forecastable signal. Another limitation of this analysis is that we measured the forecastability of an entire season, while forecast metrics presented here are mean scores of numerous individual forecasts through time—each occurring at time points with imperfect/incomplete information for the season up to that given date. This analysis also does not address potential underlying differences in clinical reporting patterns or biases across geographies and for different diseases. For example, providers might be less likely to test for and diagnose influenza during off season months, or with increased usage of multiplex testing, influenza detections may increase incidentally as a result.

There are numerous other ways to measure or categorize the properties of time series and future work could explore these metrics in an infectious disease context (9,10). Other methods that consider more local behavior of the time series at the time of the forecast (e.g., shapelet analysis, method of analogues) might also address this limitation of a global time series metric being applied to a summation of local forecast scores (24,25). For example, although COVID-19 and influenza had roughly similar relationship and scale between population size and forecastability (Fig 3A), COVID-19 had noticeably lower relative WIS across both seasons compared to influenza (Fig 3D)—suggesting that this metric is not capturing all features relevant to forecasting performance. We hope this preliminary analysis sparks further interest into delineating the qualities of infectious disease time series themselves that may help further explain forecasting performance across diseases and through time.

**Ethics**

The California Health and Human Services Agency Committee for the Protection of Human Subjects (CPHS) has determined that this research (project number 2024-210) is classified as exempt under the federal Common Rule. This decision is issued under the California Health and Human Services Agency's Federal wide Assurance #00000681 with the Office of Human Research Protections (OHRP).

**Data availability**

Code and underlying data to reproduce the analyses are available at:
https://github.com/cdphmodeling/forecastability.

Laboratory-confirmed hospital admissions data are publicly available data sets through HHS/NHSN for state: https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/g62h-syeh/about_data and facility-level time series: https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/anag-cw7u/about_data

Syndromic influenza hospitalization data derived from the California Department of Healthcare Access and Information (HCAI) are available upon a data request: https://hcai.ca.gov/data/request-data/

**Authors contributions**

L.W.: conceptualization, formal analysis, methodology, validation, visualization, writing—original draft, writing—review and editing; T.L.: conceptualization, data curation, methodology, supervision, writing—review and editing.

**Conflict of interest declaration**

The authors declare no competing interests.

**Funding**

This work was supported by the California Department of Public Health. The findings and conclusions in this article are those of the author(s) and do not necessarily represent the views or opinions of the California Department of Public Health or the California Health and Human Services Agency.

This study used the California Patient Discharge Dataset. The interpretation and reporting of these data are the sole responsibility of the authors. The authors acknowledge the California Department of Healthcare Access and Information for compilation of these data.

This work was funded by Centers for Disease Control and Prevention, Epidemiology and Laboratory Capacity for Infectious Diseases, Cooperative Agreement Number 6 NU50CK000539.

**Acknowledgements**

The authors thank members of the CDPH Modeling and Advanced Analytics team including Héctor Sánchez Castellanos, Mugdha Thakur, Natalie Linton, Phoebe Lu, and Brent Siegel for conversations and insights that improved these analyses.

**References**

1. Biggerstaff M, Alper D, Dredze M, Fox S, Fung ICH, Hickmann KS, et al. Results from the centers for disease control and prevention's predict the 2013-2014 Influenza Season Challenge. BMC Infect Dis. 2016 Jul 22;16:357.

2. Cramer EY, Ray EL, Lopez VK, Bracher J, Brennen A, Castro Rivadeneira AJ, et al. Evaluation of individual and ensemble probabilistic forecasts of COVID-19 mortality in the United States. Proc Natl Acad Sci U S A. 2022 Apr 12;119(15):e2113561119.

3.  Johansson MA, Apfeldorf KM, Dobson S, Devita J, Buczak AL, Baugher B, et al. An open challenge to advance probabilistic forecasting for dengue epidemics. Proc Natl Acad Sci U S A. 2019 Nov 26;116(48):24268–74.

4.  Viboud C, Sun K, Gaffey R, Ajelli M, Fumanelli L, Merler S, et al. The RAPIDD ebola forecasting challenge: Synthesis and lessons learnt. Epidemics. 2018 Mar;22:13–21.

5.  Mathis SM, Webber AE, León TM, Murray EL, Sun M, White LA, et al. Evaluation of FluSight influenza forecasting in the 2021-22 and 2022-23 seasons with a new target laboratory-confirmed influenza hospitalizations [Internet]. medRxiv; 2023 [cited 2024 Jan 30]. p. 2023.12.08.23299726. Available from: https://www.medrxiv.org/content/10.1101/2023.12.08.23299726v1

6.  Lopez VK, Cramer EY, Pagano R, Drake JM, O'Dea EB, Adee M, et al. Challenges of COVID-19 Case Forecasting in the US, 2020–2021. PLOS Comput Biol. 2024 May 6;20(5):e1011200.

7.  Ponce-Flores M, Frausto-Solís J, Santamaría-Bonfil G, Pérez-Ortega J, González-Barbosa JJ. Time Series Complexities and Their Relationship to Forecasting Performance. Entropy. 2020 Jan;22(1):89.

8.  Garland J, James R, Bradley E. Model-free quantification of time-series predictability.

9.  Kang Y, Hyndman RJ, Smith-Miles K. Visualising Forecasting Algorithm Performance using Time Series Instance Spaces. 2016;

10. Papacharalampous G, Tyralis H, Pechlivanidis IG, Grimaldi S, Volpi E. Massive feature extraction for explaining and foretelling hydroclimatic time series forecastability at the global scale. Geosci Front. 2022 May 1;13(3):101349.

11. Goerg GM. Forecastable Component Analysis. In: Proceedings of the 30 th International Conference on Machine Learning. Atlanta, Georgia; 2013.

12. Holcomb KM, Mathis S, Staples JE, Fischer M, Barker CM, Beard CB, et al. Evaluation of an open forecasting challenge to assess skill of West Nile virus neuroinvasive disease prediction. Parasit Vectors. 2023 Jan 12;16(1):11.

13. Bracher J, Ray EL, Gneiting T, Reich NG. Evaluating epidemic forecasts in an interval format. Pitzer VE, editor. PLOS Comput Biol. 2021 Feb 12;17(2):e1008618.

14. Bosse NI, Abbott S, Cori A, van Leeuwen E, Bracher J, Funk S. Scoring epidemiological forecasts on transformed scales. PLOS Comput Biol. 2023 Aug 29;19(8):e1011393.

15. White LA, McCorvie R, Crow D, Jain S, León TM. Assessing the accuracy of California county level COVID-19 hospitalization forecasts to inform public policy decision making. BMC Public Health. 2023 Apr 28;23(1):782.

16. U.S. Department of Health & Human Services. COVID-19 Reported Patient Impact and Hospital Capacity by State Timeseries (RAW) [Internet]. 2024. Available from: https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/g62h-syeh/

17. Cramer EY, Huang Y, Wang Y, Ray EL, Cornell M. The United States COVID-19 Forecast Hub dataset. Sci Data. 2022;9(1):462.

18. Reinhart A, Brooks L, Jahja M, Rumack A, Tang J, Agrawal S, et al. An open repository of real-time COVID-19 indicators. Proc Natl Acad Sci. 2021 Dec 21;118(51):e2111452118.

19. U.S. Census Bureau. Population Estimate, July 1, 2021 [Internet]. [cited 2024 Dec 9]. Available from: https://api.census.gov/data/2021/pep/population/variables.html

20. State of California Department of Finance. California 2020 Population by Age [Internet]. [cited 2024 Dec 9]. Available from: https://dof.ca.gov/forecasting/demographics/

21. California Department of Public Health (CDPH) Influenza Surveillance Program. Historic Influenza and Other Respiratory Disease Surveillance Reports [Internet]. [cited 2025 Feb 7]. Available from: https://www.cdph.ca.gov/Programs/CID/DCDC/pages/immunization/flu-reports.aspx

22. Bosse NI, Gruson H, Cori A, Leeuwen E van, Funk S, Abbott S. Evaluating Forecasts with scoringutils in R. arXiv [Internet]. 2022; Available from: https://arxiv.org/abs/2205.07090

23. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2021. Available from: https://www.R-project.org/

24. Srivastava A. DTW+S: Shape-based Comparison of Time-series with Ordered Local Trend [Internet]. arXiv; 2023 [cited 2024 Nov 1]. Available from: http://arxiv.org/abs/2309.03579

25. Viboud C, Boëlle PY, Carrat F, Valleron AJ, Flahault A. Prediction of the Spread of Influenza Epidemics by the Method of Analogues. Am J Epidemiol. 2003 Nov 15;158(10):996–1006.