Title: Forecastability of infectious disease time series: are some seasons and pathogens

intrinsically more difficult to forecast?

Short title: Forecastability of infectious disease time series

Authors: Lauren A. White[1] (ORCiD: 0000-0003-3852-5374), Tomás M. León[1] (ORCiD: 0000-

0002-7633-677X)

[1]California Department of Public Health, Center for Infectious Diseases, Division of

Communicable Disease Control

Abstract:

For infectious disease forecasting challenges, individual model performance typically varies

across space and time. This phenomenon raises the question: are there properties of the target

time series that contribute to a particular season, location, or disease being more difficult to

forecast? Here we characterize a time series' future predictability using a forecastability metric

that calculates the spectral density of the time series. Forecastability of syndromic influenza

hospital admissions for the state of California varied widely across seasons and was positively

correlated with peak burden. Next, using archived U.S. state and national forecasts targeting

laboratory-confirmed COVID-19 and influenza hospital admissions, we investigated the

relationship between forecastability and: (i) population size of the forecasting target, and (ii)

forecast performance as measured by mean absolute error, weighted interval score (WIS), and

scaled relative WIS. Forecastability increased with increasing population size of the forecasting

target, and forecasting performance generally improved with higher forecastability when

controlling for population size across scales. These preliminary results support the idea that some

targets and respiratory virus seasons may be inherently more difficult to forecast and could help

explain inter-seasonal variation in model performance.

Keywords: forecastability; predictability; Shannon entropy; spectral density; infectious disease

modeling; forecasting; COVID-19; influenza

Author summary: Could intrinsic properties of an epidemiological time series help explain why

a particular season, location, or disease is more difficult to predict in the future? To answer this

question, this analysis uses a measure of a time series' future predictability called

"forecastability," which describes the inherent uncertainty or surprise in the signal. Influenza and

COVID-19 hospital admissions had higher forecastability scores for locations with larger

population sizes, possibly due to larger counts leading to smoother time series. At the same time,

forecasting performance generally improved for time series with higher forecastability scores

when controlling for population size, suggesting that this metric is helpful for understanding ease

of forecasting. These preliminary results support the idea that some epidemiological targets and

respiratory virus seasons may be inherently more difficult to forecast and could help explain why

forecasting model performance changes across different respiratory virus seasons.

Introduction

There have been numerous collaborative forecasting efforts across multiple years and diseases including influenza, COVID-19, dengue, and Ebola (1–4). These forecasting challenges or "hubs" provide unified submission targets for modeling teams, evaluate the performance of individual model predictions, and synthesize predictions from multiple submitting teams into a unified, ensemble model. Historically, individual models fail to outperform such ensembles consistently through time (2,5). Forecast performance also tends to decline during periods of rapid change, and certain forecasting targets have been more difficult to forecast than others, e.g., COVID-19 cases compared to COVID-19 deaths (5,6). However, it remains unclear which constituent factors of a model contribute to higher performance, nor is it uncommon for individual model performance to vary within and across seasons and locations (5). Here we ask the inverse question about the underlying forecasting target itself: is there something intrinsic about the qualities of infectious disease time series that make them more or less challenging to forecast?

In the infectious disease forecasting literature, quantile forecasts are commonly evaluated with scoring metrics like mean absolute error (MAE) and weighted interval score (WIS). MAE describes the absolute difference between the forecasted value and the observed value, while WIS also accounts for the dispersion of the forecast's prediction interval. Both of these metrics scale with the absolute size of their target (7). This naturally gives more weight to locations with a higher disease burden. In contrast, applying a log-transformation prior to evaluation means that forecasts are scored based on relative error instead of absolute error, giving equal weight to location targets with smaller population sizes (8). Model performance is commonly evaluated

against a baseline model which propagates incidence forward based on values from the prior

time point with uncertainty based on changes in incidence observed in prior time intervals (2).

"Forecastability" describes the future predictability—or the inherent uncertainty or surprise—in

the signal of a stationary time series. Other disciplines have used various measures of entropy to

characterize the complexity of time series and their propensity for future predictability (9–12).

Scarpino and Petri proposed permutation entropy as a way of measuring the predictability

limitations of infectious disease time series (13). Predictability decreased with increasing time

series length, illustrating that more time series history did not necessarily correlate with

increased predictability according to that specific metric. This study also found that different

diseases had unique slopes of predictability vs. time series length, suggesting that permutation

entropy was responding to specific signatures in each disease time series. Mills et al. 2025

calculated permutation entropy of a rolling window of COVID-19 weekly incident cases and

compared with time-varying forecast performance to understand how predictability affects

forecast value for decision making (14). In the context of decision making, the ensemble model

generally had greater relative performance especially at longer forecast horizons, even during

periods of lower predictability.

Another proposed metric for measuring forecastability calculates the Shannon entropy of the

spectral density of the time series (15). In the realm of forecasting evaluation, the 2020 West

Nile Virus Forecast Challenge used Shannon entropy to assess uncertainty in the ensemble

model forecast of binned case counts (16). This metric is invariant to shifting or scaling, with

values near 0% corresponding to white noise and values near 100% corresponding to a single

sinusoidal curve (Figure 1). For example, S&P 500 returns which have minimal autocorrelation

may have values near 1%, while mean temperature time series with significant autocorrelation

lags expected at six and twelve months might have a forecastability closer to 35% (15). This

metric does not interact with forecast specifications like model type, forecast horizon, or loss

function—depending only on the target time series itself (15). Although there are potentially

several ways to define forecastability, we will hereafter use the term "forecastability" to refer to

the definition outlined in Goerg (2013).

Figure 1. Correspondence between example time series (first column), their autocorrelation

values (second column) and spectral density in the frequency domain (third column). The red

text in the third column corresponds to a forecastability score of the original signal. Time series

increase in their "forecastability" from top to bottom, ranging from purely random (first row) to

perfectly sinusoidal (fourth row). The second row presents weekly influenza admissions per

100K for the state of Maryland and the third row corresponds to weekly influenza admissions per

100K for the entire United States. The x-axis of the signal column represents days since

mandatory reporting for influenza hospital admissions was required beginning February 2, 2021.

The autocorrelation and spectral densities were calculated for the entire time series displayed in

the first column, and for ease of visual display, autocorrelation lags at intervals of 14 days were

plotted with a maximum lag of 365 days displayed. For consistency, all sample time series are

differenced once prior to calculating the forecastability score.

Other studies have found that forecasting performance improves for larger jurisdictions –e.g.,

states vs. counties (6) and larger vs. smaller counties (17). We further hypothesized that: (1) as

population size increases, forecastability of the time series should also increase; and (2) as

forecastability increases, the WIS of the natural log transformed forecast should decrease (i.e.,

forecast performance should improve with higher forecastability, when controlling for

population size). We explored the relationship between population size of the representative time

series and forecastability: (1) at a county and state-level for California for syndromic influenza

hospital admissions data from the California Department of Healthcare Access and Information

(HCAI); (2) at a U.S. state and national level for laboratory-confirmed hospital admissions data

for COVID-19 and influenza derived from Health and Human Services (HHS) Patient Impact

and Hospital Capacity Data System/National Hospital Safety Network (NHSN) (18). With the

latter, we investigated the relationship between forecastability and forecast performance at the

state and national scales using archived forecasts from the COVID-19 Forecast Hub

(https://covid19forecasthub.org/) and FluSight Forecast challenge

(https://github.com/cdcepi/FluSight-forecast-hub)—two collaborative forecasting initiatives that

have targeted laboratory-confirmed hospital admissions for COVID-19 and influenza

respectively (5,19). We believe this is the first application linking this forecastability metric to

infectious disease forecast performance across diseases and seasons.

Methods

Data

California county and state level syndromic influenza hospitalization data were derived from

HCAI for the 2000-2022 respiratory virus seasons. State and national, laboratory-confirmed,

COVID-19 and influenza hospitalization admission data were obtained from HHS/NHSN via

Delphi COVIDcast for the 2022-2024 respiratory virus seasons (18,20). Seasons were classified

by the standard influenza season definition of MMWR week 40 to week 39—specifically,

October 2, 2022- September 30, 2023 for the 2022-2023 season and October 1, 2023- April 30,

2024 for the 2023-2024 season, where only voluntary reporting continued after April 30, 2024.

To assess sensitivity of results to the season timing, an alternative season definition from July 1,

2022-July 30, 2023 & July 1, 2023- April 30, 2024 was also explored.

State and national population size estimates were taken from the 2021 US Census Bureau

estimates (21). California county population estimates were taken from 2020 California

Department of Finance estimates (22).

To identify the predominant influenza subtype for historical California seasons, the most

frequently sequenced subtype was taken from historic Respiratory Laboratory Network (RLN)

and clinical sentinel laboratory surveillance data (23).

Historical forecasts for state and national COVID-19 and influenza targets were obtained from

the COVID-19 Forecast and the FluSight Forecast challenge, respectively, for the 2022-2023 and

2023-2024 respiratory virus seasons (5,19). Specifically, we focused on a comparison of baseline

and ensemble models, as well a few select models that were submitted consistently to both

forecasting hubs across both seasons (e.g., CMU-TimeSeries, PSI-DICE, SGroup-

RandomForest, UMass-trends_ensemble). Ensemble models from the modeling hubs combine all

available models for a given target date and location and usually provide the most robust

performance (2,5). In contrast, baseline models offer a reference point for model performance by

propagating historical values and uncertainty forward (2).

Forecastability

Time series can be analyzed in the frequency domain through spectral analysis to reveal the

constituent frequencies of a signal. This is reflected by the relationship between the

autocovariance function and the spectral density, which are reciprocal Fourier pairs of one

another. The autocovariance function of a stationary timeseries x_t is given by:

155
$$\gamma_x(h) = \mathbb{E}[(x_t - \mu_x)(x_{t-k} - \mu_x)]$$

156 Where $\mathbb{E}$ is the expectation operator, h is the lag value that takes on any integer value, and $\mu_x =$

157 $\mathbb{E}[x_t]$ or the mean of x_t. The spectral density for the same process, x_t, for a given frequency, ω,

158 is achieved by taking the Fourier transform of the autocovariance γ_x:

159
$$f_x(\omega) = \sum_{h=-\infty}^{h=\infty} \gamma_x(h)e^{-2\pi i \omega h}$$

160 Where $i = \sqrt{-1}$.

161 Shannon entropy measures the inherent uncertainty of a probability distribution for a discrete

162 random variable X for set χ:

165
$$H(X) = -\sum_{x \in \chi} p(x) \log(p(x))$$

163 where $p(x)$ is the probability of state x being observed in set χ. For a continuous random

164 variable, Shannon entropy is approximated by differential entropy:

166
$$H(X) = -\int_{\mathbb{X}} f(x) \log f(x)\, dx$$

167 for a continuous random variable X with probability density function $f(x)$ and support $\mathbb{X} \in \mathbb{R}$.

168 Combining the concepts of spectral density and differential entropy, the forecastability of a

169 stationary timeseries x_t is more formally defined as:

170
$$\Omega(x_t) = 1 - \frac{-\int_{-\pi}^{\pi} f_x(\omega) \log f_x(\omega)\, d\lambda}{\log(2\pi)} \in [0,1]$$

where $f_x(\omega)$ is the normalized spectral density of x_t or $\int_{-\pi}^{\pi} f_x(\omega)d\omega = 1$ (15). Here $\log(2\pi)$

represents the expected differential entropy value of a white noise signal, and so $\Omega(x_t)$ would

equal zero if and only if x_t corresponded to true white noise.

Time series were assessed for stationarity with both the Kwiatkowski-Phillips-Schmidt-Shin

(KPSS) and the Augmented Dickey-Fuller (ADF) tests at an alpha = 0.01. For comparability

across geographies and seasons, time series were differenced once before calculating the

forecastability score. The metric of forecastability was calculated using the Omega function of

the foreCA package (15).

Forecast evaluation

Forecasts and their corresponding truth data were first transformed via the natural logarithm to

control for the differing magnitude in burden across target locations (8). Following Bosse et al.

(2023) and general convention, we added small positive quantity (a =1) to the truth and

predictions data prior to scoring, to account for potential values of zero while maintaining a

monotonic transformation (8). Point forecast error was assessed with mean absolute error

(MAE), which takes the absolute difference between a model's median point predictions $\hat{y}_{i=1:N}$

and the set of observed outcomes $y_{i=1:N}$ for a given model, location, and season such

that $MAE = \frac{1}{N}\sum_{i=1}^{N}|y_i - \hat{y}_i|$

Probabilistic forecast accuracy was assessed with the weighted interval score (WIS), which is a

metric that compiles the performance of a quantile or interval-based forecasts over a range of

prediction intervals (7). The single, interval score for a forecast F with outcome y for prediction

interval $(1 - \alpha) \times 100\%$ is given by:

$$IS_\alpha(F, y) = (u - l) + \frac{2}{\alpha}(l - y) \times \mathbb{1}(y < l) + \frac{2}{\alpha}(y - u) \times \mathbb{1}(y > u)$$

Where $\mathbb{1}$ is the indicator function, u is the upper prediction limit and l is the lower prediction

limit. The score resolves into three components: $u - l$ represents a penalty incurred for width or

dispersion of the prediction interval, $\frac{2}{\alpha}(l - y) \times \mathbb{1}(y < l)$ conveys a penalty for underprediction,

and $\frac{2}{\alpha}(y - u) \times \mathbb{1}(y > u)$ conveys a penalty for overprediction.

The WIS then sums intervals scores prediction intervals at different levels $(1 - \alpha_1) < (1 - $

$\alpha_2) < \cdots < (1 - \alpha_K)$ such that:

199
$$WIS_{\alpha_{\{0:K\}}}(F, y) = \frac{1}{K + \frac{1}{2}} \left(w_0 |y - m| + \sum_{k=1}^{K} \{w_k IS_{\alpha_k}(F, y)\} \right)$$

Where K is the total number of quantiles included in the WIS, m is the predictive median

forecast value, and w_k is a non-negative, unnormalized weight, often set to: $w_k = \frac{\alpha_k}{2}$ and $w_0 = \frac{1}{2}$

(7).

In this case, both MAE and WIS values approximate relative error since these were computed on

the natural logarithm of the forecasts and their respective targets (8).

Lastly, models were scored using relative skill, θ_i, which computes model rankings by taking the

geometric mean of all possible pairwise comparisons across all models, as given by:

207
$$\theta_i = \left(\sum_{m=1}^{M} \theta_{i,m} \right)^{1/M}$$

where $\theta_{i,m}$ represents the mean score ratio of model i and model m out of all possible models M.

Because this metric is based on pairwise comparisons, it only produces a score when models

contribute to the same target for the same date(s) and can therefore help to control for forecast

missingness (24). Here we present scaled relative WIS results which normalizes model

performance relative to a baseline model; relative WIS values below one indicate that the given

model performs better than the baseline model, whereas value greater than one suggest a model

performs worse than the baseline model on average. MAE, mean WIS, and scaled relative risk

were calculated for each location, season, and model combination across all four week horizons

using the scoringutils package in R (24). All analyses were completed in R (25).

Ethics Statement

The California Health and Human Services Agency Committee for the Protection of Human

Subjects (CPHS) has determined that this research (project number 2024-210) is classified as

exempt under the federal Common Rule. This decision is issued under the California Health and

Human Services Agency's Federal wide Assurance #00000681 with the Office of Human

Research Protections (OHRP).

Results

When looking at California county and state-level syndromic influenza hospitalizations, for

which more seasons of data were available, some respiratory virus seasons were more

"forecastable" than others, with the greatest differences occurring for locations with larger

population sizes (Fig 2A). For example, forecastability ranged from 19.5% to 41.6% for the state

of California for the 2021-2022 and 2017-2018 seasons respectively. Across seasons from 2000-

2022, there was a positive relationship between forecastability and both cumulative burden and

peak burden of each respective season with an adjusted R-squared of 0.56 and 0.81 respectively

(Fig 2B, Supplemental Fig 1). The season with highest peak weakly admissions, cumulative

burden, and forecastability was an H3N2 season (Fig 2B, Supplemental Fig 1). Two other H3N2

majority seasons (i.e., 2003-2004, 2005-2006) were outliers with higher-than-expected

forecastability given their peak burden. The two seasons bridging the 2009 swine influenza

pandemic (2008-2010) were outliers with lower-than-expected forecastability given their burden.

For U.S. state and national laboratory confirmed admissions, forecastability of the time series

increased with target population size for both COVID-19 and influenza across U.S. states and

nationally (Fig 3A). This pattern was generally of the same order of magnitude for both COVID-

19 and influenza, and similar across both the 2022-2023 & 2023-2024 seasons (Fig 3A). As seen

in prior studies (6,17), forecast performance generally improved for more populous targets, i.e.,

for natural log transformed forecasts, MAE and WIS on the log scale decreased with increasing

population size for the ensembles (Fig 3B-D) and most individual models (Supplemental Fig 2).

Except for the 2022-2023 influenza season, MAE and WIS for the ensemble on the log scale also

decreased with higher forecastability, i.e., forecast performance improved with higher

forecastability (Fig 3B &C, Table 1). For example, for COVID-19 in 2022-2023 season,

forecastability values ranged from 10% (Idaho) to 30% (United States) with corresponding

decreases in relative error of MAE from 25% to 21% and relative error of WIS from 26% to 14%

for the ensemble. In contrast, baseline model performance showed no such relationship with

forecastability, with non-significant or positive slope estimates (Fig 3B & C, Table 1).

Across both COVID-19 and influenza ensembles, rates of change of MAE per unit

forecastability ranged from -0.0046 to -0.0086, i.e., every one percent unit increase in

forecastability yielded a ~0.46-0.86% decrease in MAE relative error (Table 1). Rates of change

of WIS per unit forecastability ranged from -0.0016 to -0.0062 for a corresponding decrease in

WIS relative error of ~0.13-0.62% per unit forecastability (Table 1). Relationships between the

ensembles' scaled relative skill and forecastability were significant for both seasons and both

diseases (Fig 3C, Table 1) with increases in relative skill ranging from -0.0048 to -0.011 per unit

of forecastability. Overall, these findings were robust to alternative definitions of seasonal timing

(Supplemental Figure 3, Supplemental Table 1).

For lab-confirmed COVID-19 and influenza hospital admissions, the metric of forecastability

was very susceptible to data reporting frequency (Fig 4) and the amount of data smoothing (Fig

262 5). When compared to a time series of rolling weekly admissions with a daily reporting

263 frequency, time series down sampled to weekly reporting frequency had a decreased

264 forecastability score. While this observed effect increased with population size of the measured

265 time series, population size explained less of the variance in forecastability for the weekly vs. the

266 daily sampled time series (Fig 4). The relationship between forecastability and the amount of

267 data smoothing exhibited asymptotic dynamics—beginning with lower forecastability scores for

268 daily reporting that improved to a plateau as the length of the rolling sum window increased (Fig

269 5).

Figure 2. Examples of variation in forecastability by season for syndromic HCAI influenza

hospitalizations. **(A)** The relationship between forecastability and population size for the state of

California and constituent counties by respiratory virus season, derived from HCAI data from

2017-2022 (2020-2021 excluded because of the COVID-19 pandemic) and fitted with a

generalized additive model (GAM) for visualization; **(B)** forecastability vs. the natural log of

peak influenza weekly admissions for the state of California derived from syndromic HCAI data

from 2001-2022 and fitted with a linear regression (β= 7.49, p < 0.001). The year labels in the

 plot correspond to the first year of the MMWR season, e.g., 2001 corresponds to the 2001-2002

 respiratory virus season.

Figure 3. The relationship between population size, forecastability, and forecast performance

across two respiratory virus seasons for laboratory-confirmed (HHS/NHSN) COVID-19 and

influenza admissions at the U.S. state and national scales. **(A)** Forecastability (Ω) vs. the natural

logarithm of population size. Forecast performance for the baseline and ensemble models as

measured by: **(B)** MAE and **(C)** mean WIS and **(D)** scaled relative skill for the ensemble model.

Figure 4. Comparison of forecastability scores vs. natural logarithm of population size for daily

vs. weekly reporting (columns) for COVID-19 vs. influenza laboratory-confirmed hospital

admissions (rows) from state and national HHS/NHSN data for the 2022-2023 and 2023-2024

seasons.

Figure 5. Exploration of the relationship between the amount of signal smoothing and

forecastability score. (A-C) Time series of incident influenza hospital admissions with daily

reporting vs. 28-day or 56-day rolling sums for the state of Maryland. The x-axis of the signal

column represents days since mandatory reporting for influenza hospital admissions was

required beginning February 2, 2021. (D) Forecastability scores calculated on incident influenza

hospitalization time series aggregated by the days of aggregation designated in the x-axis. The

trend line is a Generalized Additive Model (GAM) fit on forecastability as a function of the

rolling sum window (days) with a cubic spline basis function.

303   **Table 1.** Slope estimates from linear model fits for forecastability (Ω) vs. forecast performance

304   for the ensemble and baseline models targeting laboratory-confirmed (HHS/NHSN) COVID-19

305   and influenza admissions at the U.S. state and national scales as shown in Figure 2. Values are

306   shown with three significant figures. Bolded rows are those statistically significant at a p = 0.05

307   threshold.

Metric	Disease	Model	Season	Estimate (beta)	Standard error	Statistic	p-value
MAE	COVID-19	Baseline	**2022-2023**	**-0.0101**	**0.00237**	**-4.25**	**9.46e-5**
			2023-2024	**-0.00589**	**0.00257**	**-2.29**	**0.0264**
		Ensemble	**2022-2023**	**-0.00857**	**0.00207**	**-4.13**	**1.38e-4**
			2023-2024	**-0.00464**	**0.00216**	**-2.15**	**0.0367**
	Influenza	Baseline	2022-2023	-0.00428	0.00389	-1.1	0.276
			2023-2024	**-0.00491**	**0.00205**	**-2.4**	**0.0202**
		Ensemble	2022-2023	-0.00636	0.00326	-1.95	0.0566
			2023-2024	**-0.00691**	**0.00173**	**-3.99**	**2.17e-4**
WIS	COVID-19	Baseline	2022-2023	-5.81E-04	0.00222	-0.261	0.795
			2023-2024	**-0.00369**	**0.00183**	**-2.01**	**0.0496**
		Ensemble	**2022-2023**	**-0.00627**	**0.00125**	**-5.01**	**7.25e-6**
			2023-2024	**-0.00419**	**0.00132**	**-3.18**	**0.00253**
	Influenza	Baseline	**2022-2023**	**0.00522**	**0.00247**	**2.11**	**0.04**
			2023-2024	-1.66E-04	0.00137	-0.121	0.904
			2022-2023	-0.00167	0.0021	-0.793	0.432

		Ensemble	**2023-2024**	**-0.00418**	**0.00113**	**-3.7**	**5.38e-4**
Scaled relative skill	COVID-19	Ensemble	**2022-2023**	**-0.0115**	**0.00232**	**-4.97**	**8.11e-6**
			2023-2024	**-0.00487**	**0.00221**	**-2.2**	**0.0325**
	Influenza		**2022-2023**	**-0.0101**	**0.0013**	**-7.75**	**4.1e-10**
			2023-2024	**-0.0104**	**0.00131**	**-7.95**	**1.99e-10**

Discussion

Using a simple signal processing metric to characterize the predictability of infectious disease time series, we showed that there are differences in the measured forecastability of time series across diseases (i.e., COVID-19 vs. influenza, Fig 3A) and across respiratory virus seasons for a given disease (Fig 2). Moreover, these observed differences in measured forecastability correlated with resulting forecast performance. Our results echoed prior work that found a relationship between population size of the forecast target and forecast performance (Fig 3B & C, Supplementary Fig 2) (6,17), but our work further suggests that forecastability increases with population size (Fig 2A, Fig 3A), and forecast performance improves for time series with higher forecastability scores when controlling for population size (Fig 3D). Notably, scaled relative skill of the ensemble against the baseline model improved with increasing forecastability of the target time series across all four combinations of season and disease (Fig 3D), and these gains came from improvements in ensemble performance rather than changes in the baseline model performance, which remained relatively invariant to the time series' forecastability (Fig 3C, Table 1). This observed relationship between forecastability and population size reflects general

intuition, i.e., that larger populations should have signals closer to the mean field, with

corresponding weaker high frequency components, resulting in a higher forecastability score.

Interestingly, there appears to be a strong relationship between peak seasonal burden and

forecastability (Fig 2B), which may point to seasons with more singular, sharper peaks having

higher forecastability per this particular metric. However, this finding contrasts with the general

pattern that forecasts struggle most during periods of rapid change, such as during the growth

phase or at the epidemic peak (5,6). Although anecdotal, patterns in forecastability across

seasons matches what might be expected based on epidemiological understanding: forecastability

was lower than expected for disrupted influenza seasons (e.g., during the 2009 swine influenza

pandemic and in the wake of the COVID-19 pandemic) (Fig 2B). This also follows intuition, as

one would expect that periods of lower overall burden would exhibit more epidemic

stochasticity, while peak periods would better approximate mean field behavior. This observed

variability in forecastability across seasons could help explain why there has been such

variability in both model rankings across years and the ability (or inability) of models able to

outperform the baseline model for any given season (5).

The patterns of forecastability observed here raise a couple of questions about the practical

implementation of forecasting challenges for infectious diseases. The first is that forecastability

is strongly affected by data frequency (e.g. daily vs. weekly) and amount of data smoothing (e.g.,

daily vs. weekly rolling sum). Although the increase in forecastability observed with population

size is less evident with lower frequency data (Fig 4) and forecastability has a non-linear

relationship with the amount of data smoothing (Fig 5), we were not able to tie these observation

directly to forecast performance. Daily hospital admissions data have been publicly available at

certain times for COVID-19 and influenza (and were briefly a formal target for COVID-19), but

we do not know which models were training on daily vs. weekly data. This may matter

practically since there are often changes in infectious disease data continuity through time in

terms of targets, reporting frequency, and reporting definitions (e.g., influenza shifting from

influenza like illness (ILI) to confirmed hospitalizations during the COVID-19 pandemic,

changes in NHSN reporting requirements from daily to weekly in November of 2024). It also

highlights that forecasting a completely new signal (e.g., FluSight 2022-2023) remains

challenging regardless of the features of the time series.

The second notable pattern is that there is a consistent plateau of forecastability at about 10% for

smaller geographies, regardless of the disease, and that certain seasons were more or less

predictable according to the forecastability metric (Figure 2A, 3A). Since the baseline model

showed no significant relationship with forecastability (Fig 3C, D), this raises the question of

whether forecasting challenges should also be: (1) controlling for some version of signal

predictability based on geographic scale or (2) controlling for season predictability post-hoc

when scoring forecasts. The latter might matter more for comparing model performance across

seasons than within a season. Finally, although there has been increased interest in providing

local level forecasts in the wake of the COVID-19 pandemic (17), this also raises questions about

infectious disease surveillance for smaller geographies and the inherent feasibility of providing

real time infectious disease modeling intelligence at these scales.

Forecastability is an appealing metric because it simplifies a complex time series into a single

value; however, the interpretation of this dimension reduction remains challenging. Since

forecastability is measured in the frequency domain, it is sometimes difficult to understand

which qualities of a time series directly translate to a more forecastable signal. Another

limitation of this analysis is that we measured the forecastability of an entire season, while

forecast metrics presented here represent mean scores of numerous individual forecasts through

time—each occurring at time points with imperfect/incomplete information for the season up to

that given date. This metric only reflects the underlying infectious disease time series itself,

which ignores the other potential covariates that could form the complete information set of a

forecast (e.g., the value of knowing hospitalizations when forecasting deaths) (14). This analysis

also does not address potential underlying differences in clinical reporting patterns or biases

across geographies and for different diseases. For example, providers might be less likely to test

for and diagnose influenza during off season months, or with increased usage of multiplex

testing, influenza detections may increase incidentally as a result.

There are numerous other ways to measure or categorize the properties of time series and future

work could explore these metrics in an infectious disease context (11,12). Other methods that

consider more local behavior of the time series at the time of the forecast (e.g., shapelet analysis,

method of analogues) might also address this limitation of a global time series metric being

applied to a summation of local forecast scores (26,27). For example, although COVID-19 and

influenza had roughly similar relationship and scale between population size and forecastability

(Fig 3A), COVID-19 had noticeably lower relative WIS across both seasons compared to

influenza (Fig 3D)—suggesting that this metric is not capturing all features relevant to

forecasting performance. This work could contribute to improved selection of baseline models

for scoring forecasts (e.g., (28)), better cross-seasonal comparisons of forecasting hub results,

and an increased understanding of how forecasts contribute to decision making at different

thresholds of uncertainty and predictability (14). We hope this analysis sparks further research

into delineating the qualities of infectious disease time series themselves—or the underlying

"information set" of forecasts—that may help further explain forecasting performance across

seasons and diseases.

Data availability

Code and underlying data to reproduce the analyses are available at:

https://github.com/cdphmodeling/forecastability.

Laboratory-confirmed hospital admissions data are publicly available data sets through

HHS/NHSN for state: https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-

Hospital-Capa/g62h-syeh/about_data and facility-level time series:

https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/anag-

cw7u/about_data

Syndromic influenza hospitalization data derived from the California Department of Healthcare

Access and Information (HCAI) contain individual level patient data and so are only available

upon a data request: https://hcai.ca.gov/data/request-data/

Authors contributions

406 L.W.: conceptualization, formal analysis, methodology, validation, visualization, writing—

407 original draft, writing—review and editing; T.L.: conceptualization, data curation, methodology,

408 supervision, writing—review and editing.

Conflict of interest declaration

410 The authors declare no competing interests.

Funding

This work was supported by the California Department of Public Health. The findings and conclusions in this article are those of the author(s) and do not necessarily represent the views or opinions of the California Department of Public Health or the California Health and Human Services Agency.

This study used the California Patient Discharge Dataset. The interpretation and reporting of these data are the sole responsibility of the authors. The authors acknowledge the California Department of Healthcare Access and Information for compilation of these data.

This work was funded by Centers for Disease Control and Prevention, Epidemiology and Laboratory Capacity for Infectious Diseases, Cooperative Agreement Number 6 NU50CK000539.

Acknowledgements

The authors thank members of the CDPH Modeling and Advanced Analytics team including Héctor Sánchez Castellanos, Mugdha Thakur, Natalie Linton, Phoebe Lu, and Brent Siegel for conversations and insights that improved these analyses.

References

1. Biggerstaff M, Alper D, Dredze M, Fox S, Fung ICH, Hickmann KS, et al. Results from the Centers for Disease Control and Prevention's Predict the 2013-2014 Influenza Season Challenge. BMC Infect Dis. 2016 Jul 22;16:357.

2. Cramer EY, Ray EL, Lopez VK, Bracher J, Brennen A, Castro Rivadeneira AJ, et al. Evaluation of individual and ensemble probabilistic forecasts of COVID-19 mortality in the United States. Proc Natl Acad Sci U S A. 2022 Apr 12;119(15):e2113561119.

3. Johansson MA, Apfeldorf KM, Dobson S, Devita J, Buczak AL, Baugher B, et al. An open challenge to advance probabilistic forecasting for dengue epidemics. Proc Natl Acad Sci U S A. 2019 Nov 26;116(48):24268–74.

4. Viboud C, Sun K, Gaffey R, Ajelli M, Fumanelli L, Merler S, et al. The RAPIDD ebola forecasting challenge: Synthesis and lessons learnt. Epidemics. 2018 Mar;22:13–21.

5. Mathis SM, Webber AE, León TM, Murray EL, Sun M, White LA, et al. Evaluation of FluSight influenza forecasting in the 2021-22 and 2022-23 seasons with a new target laboratory-confirmed influenza hospitalizations [Internet]. medRxiv; 2023 [cited 2024 Jan 30]. p. 2023.12.08.23299726. Available from: https://www.medrxiv.org/content/10.1101/2023.12.08.23299726v1

6. Lopez VK, Cramer EY, Pagano R, Drake JM, O'Dea EB, Adee M, et al. Challenges of COVID-19 case forecasting in the US, 2020–2021. PLOS Comput Biol. 2024 May 6;20(5):e1011200.

7. Bracher J, Ray EL, Gneiting T, Reich NG. Evaluating epidemic forecasts in an interval

format. Pitzer VE, editor. PLOS Comput Biol. 2021 Feb 12;17(2):e1008618.

8. Bosse NI, Abbott S, Cori A, van Leeuwen E, Bracher J, Funk S. Scoring epidemiological

forecasts on transformed scales. PLOS Comput Biol. 2023 Aug 29;19(8):e1011393.

9. Ponce-Flores M, Frausto-Solís J, Santamaría-Bonfil G, Pérez-Ortega J, González-Barbosa JJ.

Time Series Complexities and Their Relationship to Forecasting Performance. Entropy. 2020

Jan;22(1):89.

10. Garland J, James R, Bradley E. Model-free quantification of time-series predictability.

11. Kang Y, Hyndman RJ, Smith-Miles K. Visualising Forecasting Algorithm Performance

using Time Series Instance Spaces. 2016;

12. Papacharalampous G, Tyralis H, Pechlivanidis IG, Grimaldi S, Volpi E. Massive feature

extraction for explaining and foretelling hydroclimatic time series forecastability at the

global scale. Geosci Front. 2022 May 1;13(3):101349.

13. Scarpino SV, Petri G. On the predictability of infectious disease outbreaks. Nat Commun.

2019 Feb 22;10(1):898.

14. Mills C, Irons NJ, Tsui JLH, Sparrow S, Carvalho LM, Kucharski AJ, et al. From metric to

action: An evaluation framework to translate infectious disease forecasts into policy

decisions [Internet]. medRxiv; 2025 [cited 2025 Aug 7]. p. 2025.07.20.25331802. Available

from: https://www.medrxiv.org/content/10.1101/2025.07.20.25331802v1

15. Goerg GM. Forecastable Component Analysis. In: Proceedings of the 30 th International

Conference on Machine Learning. Atlanta, Georgia; 2013.

16. Holcomb KM, Mathis S, Staples JE, Fischer M, Barker CM, Beard CB, et al. Evaluation of

an open forecasting challenge to assess skill of West Nile virus neuroinvasive disease

prediction. Parasit Vectors. 2023 Jan 12;16(1):11.

17. White LA, McCorvie R, Crow D, Jain S, León TM. Assessing the accuracy of California

county level COVID-19 hospitalization forecasts to inform public policy decision making.

BMC Public Health. 2023 Apr 28;23(1):782.

18. U.S. Department of Health & Human Services. COVID-19 Reported Patient Impact and

Hospital Capacity by State Timeseries (RAW) [Internet]. 2024. Available from:

https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-

Capa/g62h-syeh/

19. Cramer EY, Huang Y, Wang Y, Ray EL, Cornell M. The United States COVID-19 Forecast

Hub dataset. Sci Data. 2022;9(1):462.

20. Reinhart A, Brooks L, Jahja M, Rumack A, Tang J, Agrawal S, et al. An open repository of

real-time COVID-19 indicators. Proc Natl Acad Sci. 2021 Dec 21;118(51):e2111452118.

21. U.S. Census Bureau. Population Estimate, July 1, 2021 [Internet]. [cited 2024 Dec 9].

Available from: https://api.census.gov/data/2021/pep/population/variables.html

22. State of California Department of Finance. California 2020 Population by Age [Internet].

[cited 2024 Dec 9]. Available from: https://dof.ca.gov/forecasting/demographics/

23. California Department of Public Health (CDPH) Influenza Surveillance Program. Historic

Influenza and Other Respiratory Disease Surveillance Reports [Internet]. [cited 2025 Feb 7].

Available from: https://www.cdph.ca.gov/Programs/CID/DCDC/pages/immunization/flu-

reports.aspx

24. Bosse NI, Gruson H, Cori A, Leeuwen E van, Funk S, Abbott S. Evaluating Forecasts with

scoringutils in R. arXiv [Internet]. 2022; Available from: https://arxiv.org/abs/2205.07090

25. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna,

Austria: R Foundation for Statistical Computing; 2021. Available from: https://www.R-

project.org/

26. Srivastava A. DTW+S: Shape-based Comparison of Time-series with Ordered Local Trend

[Internet]. arXiv; 2023 [cited 2024 Nov 1]. Available from: http://arxiv.org/abs/2309.03579

27. Viboud C, Boëlle PY, Carrat F, Valleron AJ, Flahault A. Prediction of the Spread of

Influenza Epidemics by the Method of Analogues. Am J Epidemiol. 2003 Nov

15;158(10):996–1006.

28. Stapper M, Funk S. Mind the Baseline: The Hidden Impact of Reference Model Selection on

Forecast Assessment [Internet]. 2025 [cited 2025 Aug 26]. Available from:

http://medrxiv.org/lookup/doi/10.1101/2025.08.01.25332807