Retrospective impact evaluation of indoor residual spraying in Côte d'Ivoire using

routine malaria surveillance data: the role of contemporaneous covariates and data

quality

Christian Selinger (1, 2, *), Kouakou David Appeti (3), Sumaiyya Thawer (1, 2), Georgina

Angoa (5), Serge A Aïmain (3), Serge B Assi (3, 4), Emilie Pothin (1, 2), Emily Hilton (6)

Antoine M Tanoh (3)

1 Swiss Tropical and Public Health Institute, Allschwil, Switzerland

2 University of Basel, Basel, Switzerland

3 Programme National de Lutte Contre le Paludisme, Abidjan, Côte d'Ivoire

4 Institut Pierre Richet (IPR), Institut National de la Santé Publique (INSP), Bouaké, Côte

d'Ivoire

5 Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire

6 PMI VectorLink Project, PATH, Seattle, Washington, USA

*corresponding author: christian.selinger@swisstph.ch

Objectives: Indoor residual spraying (IRS) was introduced in Côte d'Ivoire in 2020. We seek

to identify potential factors that could compromise the quality of routine malaria incidence

and determine how impact quantification of IRS on malaria incidence depends on data

quality by emphasizing robust counterfactuals with contemporaneous covariates.

Methods: Using mixed-effect models and correlation analysis we compare data from two

sources (routine surveillance and re-digitized registries). To quantify decrease in incidence

post intervention, we apply Bayesian structural time series analysis with posterior feature

selection to determine impact confounders.

Results: The presence of IRS, test positivity and reporting rates are significant factors

impacting data quality. For high-quality data, we estimate a 12% resp. 19% decrease in

incidence rate one year after the intervention, while for low-quality data estimates are 30%

resp. 42%. Important covariates for counterfactual modeling are control district incidence,

reporting and test positivity rates.

Conclusions: Improvements in data quality by scrutinizing test positivity rates and health

facility reporting can help avoid overly optimistic impact estimations for IRS. Incidence control

districts remain inevitable in order to factor out impact confounders.

Introduction

After two decades of steadily declining malaria mortality from 2000 to 2019, global progress in the fight against malaria has recently stalled and, in some regions, reversed (1). In Côte d'Ivoire, Malaria remains a leading cause of morbidity and mortality, accounting for around 33% of outpatient visits at health facilities (2). Vector control interventions including indoor residual spraying (IRS) and insecticide-treated nets (ITNs) remain important cornerstones of malaria prevention (3). Côte d'Ivoire included IRS in its National Malaria Strategic Plans 2016–2020 and 2021–2025 in order to reduce malaria burden in high transmission districts (2). IRS was introduced in two districts in 2020 and carried out annually until 2022. After 2022, IRS was stopped due to loss of funding through the US President's Malaria Initiative (PMI), despite the evidence of its impact on clinically relevant outcomes in 'real-world' settings (4). Compared to randomized control trials (RCTs), real-world evidence from routine surveillance systems is less costly to produce, specific to the local context and of practical use to decision-makers. On the other hand, such analyses do not typically use any control groups or counterfactuals which is crucial to all retrospective impact evaluations (5–7). Counterfactuals should represent the temporal evolution of the outcome measure in the hypothetical absence of the intervention. Clinical impact of vector control can be estimated using confirmed malaria cases. Every month, such data are reported to a centralized database based upon facility-level paper registries using the District Health Information Software (DHIS2) system. The National Malaria Control Program (NMCP) of Côte d'Ivoire has implemented the DHIS2 database as early as 2015 with only recently improved consistency and validity (8). In a project aiming at assessing the impact of IRS, PMI in collaboration with the NMCP re-digitized key malaria indicators from health facility consultation registers during the years 2019-2022 of four districts: Nassian and Sakassou with IRS campaigns conducted in 2020&2021, and Dabakala and Beoumi as control districts. Based on this data (henceforth referred to as PMI data), a comprehensive impact evaluation estimated a significant reduction in clinical incidence following the IRS campaigns (9).

Complementary to the impact quantification undertaken in (9), our analysis has two main objectives. First, we seek to identify potential factors contributing to the differences between data reported into DHIS2 and the data re-digitized by PMI, assuming that DHIS2 data are generally of lower quality and compromised at the level of data entry and compilation. Second, we utilize a model-based approach to interrogate how differences in data quality alter intervention impact quantification. We emphasize building counterfactuals incorporating a large number of contemporaneous covariates relying on Bayesian structural time series analysis. This technique has been suggested as a viable alternative to interrupted time series analysis for health impact assessment (10,11) with clinical, routine and remote-sensed data

(12–14) and probabilistically combines covariates such as rainfall, vegetation cover,

temperature, but also routine health surveillance quality and most importantly reported

malaria incidence from control districts to factor out possible confounders of IRS impact.

The results of this analysis should help promote critical use of routine malaria surveillance

data for decision-making. Intensified data validation and audits combined with analytics that

allow identifying potential causal confounders should further add to the existing body of

evidence on how IRS affects malaria incidence in the 'real' world. In light of increasingly

limited resources for malaria surveillance, treatment and prevention, the importance of high

quality and appropriately contextualized data for decision making to National Malaria Control

Programs cannot be overestimated.

Methods

Data sources and quality assessment

We extracted and post-processed routine surveillance data for malaria from the DHIS2
database in collaboration with the NMCP of Côte d'Ivoire. In the country, the reporting follows
the hierarchy of referral in the sanitary pyramid, i.e. health facilities in the periphery with
recently added community health workers (CHW) and local general hospitals (Hôpital
Général, HG) refer to regional hospitals (Centre Hospitalier Régional, CHR) who refer to
university hospitals (Centre Hospitalier Universitaire, CHU). The nested spatial organization
consists of 113 health districts within 26 health regions that report to the Ministry of Public
Hygiene and Universal Health Coverage at the national level. We considered clinical malaria
cases of all ages reported at the level of health districts. These monthly data were entered
into DHIS2 from paper registries in health facilities, HG, and CHR by local health workers
and validated by the NMCP. As for CHW and CHU, inconsistency in reporting across districts
and time, prohibited further consideration for this analysis. Since presumed cases (i.e.
without confirmed tests) have been reported into the DHIS2 database only since 2021, we
considered the difference between suspected and tested cases as presumed cases
whenever the latter were not directly available in the database.

In preparation of the impact evaluation for IRS in the districts Nassian and Sakassou with
control districts Dabakala and Beoumi, PMI decided to collect data directly from health facility
registries in the four study districts due to insufficient data quality (15). Patient data were
abstracted from consultation registries ("registres de consultations curatives") for the period
September 2018 to April 2022, we refer to details of inclusion criteria to recently published
study results (9).

In order to control for factors unrelated to the intervention, we also included environmental,
entomological and surveillance covariates into the causal impact analysis (see
Supplementary Table 1). Remote-sensed data (rainfall, average daily temperature, leaf
cover, average daily humidity) was extracted from the ERA5 database (16), aggregated
monthly at the health-district level. Entomological covariates (monthly indoor and outdoor
biting rates and monthly entomological inoculation rates) were extracted from the PMI
VectorLink annual entomological reports for the years 2019-2021 (17–19) for the four
districts of interest. To take into account changes in the health surveillance we also derived
from the DHIS2 database the following district-level monthly quantities: health facility

reporting rate (short: hf_reporting), monthly testing rate (short: testing), test positivity rate

(short: tpr_total), the number of active health facilities (short: N_hf_active) and coverage of

intermittent preventative therapy for pregnant women for the first trimester (short:

ipt1_coverage). Although the latter quantity concerns only a small proportion of the

population at risk, we consider it as a general proxy for access to health care. We normalized

and scaled all covariates prior to further analysis.

Incidence rate calculations with adjustments

To adjust incidence (to both DHIS2 and PMI crude data), we used the following

epidemiological indicators. First, we extracted monthly confirmed malaria cases reported and

validated at the level of health districts in the DHIS2 system or by PMI respectively. Second,

we used presumed cases (short: pr) from the DHIS2 database which are defined as cases

with malaria-like symptoms without having been tested. Third, we calculated the test

positivity rate (short: tpr) as the ratio of confirmed over tested cases. Here, we assumed that

cases were tested with either blood spot or rapid diagnostic test (RDT). For the reporting

rate, we considered for each health district the ratio of months within a year, where health

facilities reported into the DHIS2 system. Finally, the health care seeking rate (short: s) was

extracted from the 2016 Malaria Indicator Survey (20) and the 2021 Demographic and Health

Survey (DHS) report (21) estimates of health care seeking for children with fever made at the

regional level. Sakassou, Beoumi and Dabakala are located in the Center-North region and

Nassian in the North East region. We interpolated these values linearly between the years.

We then calculated adjustments to incidence rates for both data sources DHIS2 and PMI

using the same quantities as detailed in Supplementary Table 2. District-level population

data were extracted from the 2021 census data with 3% annual growth rate extrapolation.

We calculated incidence rate as the ratio of adjusted incidence over district population.

Multiple imputation of missing data

In the period of interest, we have detected 6.7% of missing data among all contemporaneous

covariates across the four health districts (see Supplementary Figure 1). We used the R

package Amelia (22) to perform multiple imputations of missing data for a subset of

contemporaneous variables (biting_rate_hcl_outdoor, biting_rate_hcl_indoor, EIR_ibpn), with

health district as cross-sectional variable and the month during the year as temporal variable

to account for seasonality. Based on the rule-of-thumb we performed 7 imputations for 7% of

missing data. We averaged imputed data points and used them to replace missing data in

the original data set. In addition, we also imputed two incidence points for the Beoumi district,

which had suspicious outliers.

Mixed effect models to analyze difference between data sources

In order to identify associations between difference in incidence rates recorded in the DHIS2
and PMI data sets, we performed a mixed-effect linear model analysis (23) on the difference
of log-transformed incidence data with a random intercept for the health district. We used a
backward stepwise model selection approach (24) to eliminate sequentially factors from two
models (see Supplementary Tables 3&4): *model1* had only health system covariates (IRS
intervention, hf_reporting_rate, N_hf_active, tpr_total, ipt1_coverage), *model2* had in addition
also environmental variables (rainfall, temperature, leaf, humidity, biting rates, EIR) without
and with one month lag obtained from cross-correlation analyses. For the resulting
parsimonious model, we determined the coefficient estimators and two-sided p-values.

Correlation analysis between incidence rates and contemporaneous variables

173 We performed a Pearson correlation analysis between monthly incidence rates and
174 contemporaneous variables for lags of up to one month in the past. This analysis was
175 performed across health districts and per incidence data source.

Bayesian structural time series analysis for impact quantification

Bayesian structural times series analysis (25,26) is an extension of classical time series
analysis techniques such as auto-regression moving average models. The method integrates
elements of state-space models (i.e. observation and hidden state parameters are estimated
concurrently) and Bayesian feature selection (e.g. through spike-and-slab priors) and
provides forecasting of time series with high-dimensional time-dependent covariates. We
utilize this method to construct counterfactuals for intervention impact quantification. For
monthly outcome measures (malaria incidence rates or EIR) we implement local linear and
seasonal effects, and use contemporaneous regressors for lagged time series that are
independent from the incidence rates (e.g. rainfall, temperature). In order to train the pre-
intervention model we use either the incidence data in the IRS districts (control: "none"),
incidence data in both IRS and non-IRS districts (control: "incidence), or all
contemporaneous covariates including incidence from all districts (control: "all covariates)
but excluding entomological covariates, since those are supposed to be statistically
dependent on incidence. Similarly, for the impact analysis with EIR, we use EIR data in IRS
districts only (control: "none"), EIR from both IRS and non-IRS districts (control: "EIR"), or all
contemporaneous covariates except incidence covariates, since those are supposed to be
statistically dependent on EIR.

From the Bayesian feature selection for models with contemporaneous covariates, we
extract marginalized posterior distribution of regression parameters to quantify how often a
regressor has been included in the model, but also its magnitude and whether 95% of the
probability mass was either positive or negative ("consistent regressors"). Causal impact is

determined by the difference of predicted incidence in the absence of the intervention (i.e.

from the counterfactual) and the actually recorded data in presence of the intervention,

averaged over one-year after the intervention deployment. We consider causality as

established if in less than 5% of 10000 simulated counterfactuals we observe incidence

inferior to the observed incidence with IRS.

Results

Factors influencing the difference in incidence between DHIS2 and PMI data

After adjusting crude incidence of malaria cases from DHIS2 and PMI data sources (Table 2, adjustment 2), we observed a range between 15 and 100 monthly incidence points per 1000 population between September 2018 and April 2022 with peaks occurring generally between July and October (Figure 1). Two consecutive months for the control district of Sakassou had zero incidence recorded in the DHIS2 data. We considered these as missing data and imputed them using multiple imputation algorithms (see Methods). PMI incidence rates were generally higher than those based on DHIS2 data, and Sakassou followed more consistently the rainfall seasonality trends than Nassian. The impact analysis focused on incidence data until August 2021, date of the start of the second deployment of IRS. Analyzing the difference between incidence rates from the two data sources including model selection for a comprehensive set of contemporaneous and categorical variables (see Supplementary Table 2&3) suggested that there were several statistically significant factors explaining the difference. We identified the presence of the first IRS intervention, increased IPT1 coverage and test positivity rates to be statistically associated with a decrease in difference ($P<0.05$) Table 1.

Correlations between incidence and contemporaneous variables

For the correlation analysis between monthly incidence rates and contemporaneous variables, we considered both data from the same month and with a one-month lag across all four districts before and after the intervention (Figure 2). We noticed that entomological variables were highly correlated amongst each other (e.g. 0.57 and 0.53 Pearson correlation coefficient between EIR and indoor or outdoor biting rates). Biting rates also positively correlated with malaria incidence for the PMI data (0.53 and 0.57 correlation coefficient). Strikingly, none of the entomological variables had significant correlations with malaria incidence data from DHIS2. On the other hand, DHIS2 incidence data significantly correlated with rainfall and humidity, albeit with smaller coefficients than PMI data. A similar pattern was observed for correlations between incidence and health surveillance yield and quality covariates (number of health facilities active, health facility reporting rates) which had significant correlations with PMI incidence data (0.27, and -0.32 resp.), whereas these same covariates did not correlate with DHIS2 incidence data. Only test positivity rates positively correlated with incidence data from both sources.

Quantification of causal impact of IRS towards malaria incidence

Based on our findings from the correlation analysis, we included all except the entomological
covariates into our Bayesian structural time series analysis (see Methods) as these were
causally dependent on the intervention. We trained a model on incidence data prior to the
intervention including contemporaneous variables and utilized the resulting model estimates
to predict a counterfactual for the Nassian and Sakassou district during the intervention
period. Finally, to measure impact, we compared the counterfactual incidence with actually
recorded incidence data. Using the incidence in the IRS districts alone during the model
training (Figure 3, control=none), we obtained an estimated average decrease in incidence
during the first year post intervention of 30% (95%CI: 11-43%) for Nassian and 36% (95%CI:
20-48%) for Sakassou using DHIS2 data. Incidence decrease according to PMI data was not
significant. Including incidence data from the control districts (Dabakala for Nassian and
Beoumi for Sakassou) to the pre-intervention model training resulted in significant impact of
12% (95%CI: 3-19%) resp. 19% (95%CI: 10-27%) for PMI data and 30% (95%CI: 19-38%)
resp. 43% (95%CI: 34-50%) for DHIS2 data. Considering all covariates besides those related
to entomology yielded an estimated impact for PMI data of 9% (not significant) for Nassian
and 23% (95%CI: 12-31%) for Sakassou. No significant impact for DHIS2 data was
determined in that case. The large error estimates from the credence intervals suggest a
high variability in posterior variable selection.

Contemporaneous covariates influencing the counterfactual

Utilizing the Bayesian structural time series model, we obtained regression coefficients for
each of the contemporaneous covariates. Posterior distributions from the Bayesian updating
stem from spike and slab priors (see Methods), such that not all coefficients were necessarily
selected in each model run. We focused on the marginalized regression coefficients that
consistently either had negative or positive values for 95% of the model runs in order to
quantify covariates influencing the counterfactual, and consequently the evaluated impact.
E.g. a consistently positive regression coefficients would mean increased incidence for the
counterfactual for an increase in the covariates, such that the overall intervention impact
would be have been overestimated without the covariate, and vice versa for negative
regression coefficients.

Considering empirical distributions of posterior regression coefficients (Figure 4), we
observed that incidence from the control district had positive estimates across datasets and
health districts with high probability of inclusion for the model selection process. Health
facility reporting rates were consistently associated with negative coefficients for PMI data,
suggesting that for periods with increased health facility reporting during the intervention, one
would have estimated a decrease in the counterfactual and consequently a decreased
impact of IRS. For DHIS2 on the other hand, test positivity rate, number of health facilities

active and rainfall had positive coefficients, such that an increase in these factors might be

the main driver of the overestimated impact of IRS (Figure 3). We stress that the posterior

distribution of covariate regression coefficients used for the impact quantification are high-

dimensional and not independent from each other such that the underlying causality might

rather be a combination of many factors than just those mentioned above.

Quantification of causal impact of IRS towards entomological inoculation rates

Utilizing the same approach as for the causal impact analysis towards incidence, we also

investigated causal impact towards EIR. This time we excluded malaria incidence as

covariates, as these were not independent from EIR rates. We observed a strong and

statistically significant causal impact of IRS on the decrease of EIR at the order of 70%

averaged during one year post intervention (Figure 5). Among consistent regressors, only

EIR values from control districts were chosen with high inclusion probability (Figure not

shown).

Discussion

With improved quality and availability of routine surveillance data for malaria, retrospective
impact analysis becomes an important tool to bridge the gap from efficacy in clinical trials to
effectiveness at operationally relevant scales such as health districts (15). For NMCPs,
retrospective impact quantification is highly relevant as it provides context-specific
information to evaluate program efficiency and to plan future investments. In the present
study, we interrogate clinical incidence data extracted for the first time in its full breadth from
the DHIS2 routine surveillance system in Côte d'Ivoire (8). Established since 2015, this
system has since seen major improvements (e.g. including new indicators, capturing
community health workers and private sector) but is still underused to provide analytics for
decision support. Côte d'Ivoire has rolled out a pilot of indoor residual spraying in 2020 and
2021 in two health districts thanks to support from PMI. Concurrently with this intervention,
PMI has also collaborated with the NMCP to re-digitize health facility data from paper
registries to quantify the impact of IRS. Our study specifically focuses on quantifying how
data quality impedes impact quantification and how a Bayesian framework could inform
causality between covariates and impact indicators. We summarize and contextualize our
four main findings.

First, correlation analysis with remote-sensed temporal covariates such as rainfall or biting
rates revealed that DHIS2 data is not following expected patterns of seasonality whereas
PMI data does. Likewise, we observed highly significant correlation of PMI incidence with
indicators of surveillance quality such as reporting rates and number of active health
facilities, whereas DHIS2 does not. We conclude that DHIS2 data very likely does not
faithfully represent incidence dynamics at monthly scales and that coarser temporal
aggregation would be better suited for analysis with such data. IRS deployment and test
positivity rates are the factors most strongly associated with decreasing gaps between PMI
and DHIS2 incidence data across several statistical models. We suggest that viability and
sensitivity of tests and their reporting should be further improved in line with observations
from key-informant interviews (8).

Second, although using a completely different method, our results of 12% resp. 19%
incidence decrease one year post intervention for PMI data are in good agreement with a
recently published analysis (9), which reports 16% resp. 15.8% reduction in Nassian resp.
Sakassou. Using DHIS2 data, our statistical models estimated much higher impact of 30%

reps. 42%, which is close to 44% and 47% DHIS2-based estimates for IRS in Ghana and in

Uganda (27,28). This further highlights the robustness of our results and the importance of

the data source.

Third, following the Bayesian model averaging approach, test positivity rates, incidence from

control districts and reporting rates of health facilities were identified as consistent

regressors. Increased test positivity and incidence in control districts during the intervention

period would increase counterfactual incidence and thus not taking these covariates into

account would overestimate the impact of IRS. Likewise, increased facility reporting rates

would decrease counterfactual incidence and not considering these would lead to

underestimated impact.

Fourth, a separate analysis focused on EIR data, which to our knowledge has only been

used to choose insecticides (29) but not for IRS impact quantification before. Our result of a

strong causal impact of IRS of 70% decrease corroborates estimations of more than 80%

vector mortality on walls published in end-of-spray reports (30,31). Transmission model

studies (32) suggest that a 90% decrease in EIR in high endemic areas such as Nassian and

Sakassou would yield a 38% decrease in clinical incidence under active case detection.

Although causality between EIR and incidence remains elusive, our results indicate that

entomological surveillance data could be used as surrogates for intervention impact.

This study has several limitations. The biggest hurdle to generalize our findings is data

quality. We have extracted routine surveillance data from the DHIS2 database with the help

of the monitoring and evaluation division at the NMCP. Despite upfront database validation

work by the NMCP and data post-processing, our correlation analysis showed that temporal

inconsistencies in the malaria incidence data remain and hamper fine-tuned time series

analysis. The improvement in the PMI database concerned mainly electronic data entry, and

our comparative analysis clearly points towards improved data quality, e.g. in terms of

expected patterns of seasonality.

Although our Bayesian analysis considered many contemporaneous covariates, the

underlying causality graph remains elementary. We focused on the relationship between

each covariate and the incidence outcome and did not model the putative causal

relationships between covariates.

Incidence time series from the control district were consistently identified as key to achieve

high accuracy for the counterfactual model training in the intervention district. This finding

suggests that incidence measurements from control districts are vital for counterfactual

model parameter learning as they encode unobserved variables and processes. Relying

solely on surrogate data that is independent from the intervention (e.g. rainfall, vegetation

index) but does not contain incidence measurements would be insufficient to construct

counterfactuals that de-convolute causal impact on malaria transmission.

From the study limitations arise also several opportunities to improve the operability of

routine data for both evaluation and strategic planning. Routine correlation analysis with

remote-sensed data and outlier analysis could help single out spurious signals and guide the

deployment of data quality improvement measures akin to those described in PMI's re-

digitization protocol. Overestimating the impact of particular interventions comes to additional

expenditures, which could be invested in strengthening the routine surveillance system at the

level of data entry. Climate-related changes in patterns of vector dynamics and clinical

malaria incidence call for systematic use of remote-sensed data and summary statistics of

extreme events as contemporaneous variables to more faithfully estimate intervention

impact. The main open question that arises from our study is the problem of causality within

the complex dynamic process of disease transmission of malaria. Ideally, a causal diagram

would single out known mechanism, such as the impact of rainfall on the infectious vector

population, which should be causally related to disease incidence in the following month.

Learning causal diagrams from high-quality data and using more complex observational

models (e.g. over-dispersion and zero-inflation) could further improve the Bayesian

inference.

In conclusion, we highlight that investments to improve data quality of routine surveillance by

focusing on indicators such as test positivity and health facility reporting are key to avoiding

overly optimistic estimations of the impact of IRS on clinical incidence in areas with high

malaria prevalence such as Sakassou and Nassian. The careful choice of control districts for

such analyses remains inevitable in order to factor out potential intervention confounders.

Acknowledgements

We acknowledge the Global Fund, especially Paul Hacopian and Maria Walusimbi, for

initializing the retrospective analysis based on DHIS2 malaria surveillance data. We thank

Sarah Burnett from PMI for sharing previously published data produced through the

collaboration between PATH and the National Malaria Control Program of Côte d'Ivoire.

This work has been funded through the service contract N°UCP-FM/2022/347 from the

Ministère de la Santé, de l'Hygiène Publique et de la Couverture Maladie Universelle de Côte

d'Ivoire.

List of abbreviations

PMI President's Malaria Initiative

NMCP National Malaria Control Program

DHIS2 District health information system

DHS Demographic and Health Survey

ITN Insecticide-treated nets

IRS Indoor residual spraying

RDT Rapid diagnostic test

EIR Entomological inoculation rate

RCT randomized control trial

IPT1 Intermittent preventive therapy, first dose

Figures and Tables

Figure 1: Adjusted monthly incidence rates of confirmed malaria cases in the IRS-treated

districts of Nassian and Sakasso (solid lines) and their respective control district (dashed

lines). The color code refers to the data source (DHIS2 in orange, PMI in black).

| coefficient | model1 | | model2 | 420 |
	Estimate	p-value (t-test)	estimate	p-value (t-test)
(Intercept)	-0.33003	< 2e-16	-0.34043	< 2e-16
IRSTRUE	**-0.3556**	**8.62E-07**	**-0.29326**	**9.32E-05**
tpr_total	**-0.07673**	**0.00384**	**-0.09364**	**0.000661**
hf_reporting_rate	n.s.	n.s.	**0.063**	**0.032704**
ipt1_coverage	n.s.	n.s.	**-0.06111**	**0.047551**
EIR_ibpn	NA	NA	0.04535	0.087861

421 Table 1: Results table of best log-linear mixed-effect model of incidence difference between

422 DHIS2 and PMI data identified after model selection. *model1* considered only health system

423 related covariates, whereas *model2* also included entomological and environmental

424 covariates.

Figure 2: Correlation between contemporaneous variables (and their one-month lag) and incidence rates. The color gradient represents positive (red) and negative (blue) correlations, crossed-out fields are correlations that do not meet the significance criteria (Pearson p-value<0.05).

Figure 3: Causal impact analysis for the 2020 IRS campaign in Nassian and Sakassou. The points show the mean of the posterior distribution, the error bars represent the 95% credence interval of the posterior of the average change in incidence during 1 year post intervention. Colors distinguish data sources, shapes refer to whether only the incidence rate time series itself has been used (none), or incidence of the respective control districts (incidence) or in addition other contemporaneous variables (all_covariates). For the latter, we excluded entomological covariates (EIR_ibnp, biting_rate_hcl_outdoor, biting_rate_hcl_indoor), as these were not independent of the intervention and thus not suitable as controls.

Figure 4: Posterior distribution of regression coefficients (with possible one-month lags) from the Bayesian causal impact analysis that have 95% of the mass either above or below zero.

Positive coefficients indicate an increase in counterfactual incidence for a unit increase of the

regressor, thus an overestimation of intervention impact if the regressor were not taken into

account. The numbers in the graph are inclusion probabilities in the model over 10000 runs

following the spike and slab prior.

Figure 5: Causal impact analysis for the 2020 IRS campaign in Nassian and Sakassou. The

points show the mean of the posterior distribution, the error bars represent the 95% credence

interval of the posterior of the average change in entomological inoculation rate during 1 year

post intervention. The control level "all covariates" in the plot refers to using all covariates in

the intervention districts Nassian and Sakassou (except malaria incidence, see methods), the

level "EIR" refers to using EIR values from control districts Beoumi and Dabakala in addition

to EIR values in the intervention districts. The level "none" refers to using EIR data only from

the intervention districts Nassian and Sakassou.

References

1.	Global Malaria Program WHO Geneva. World Malaria Report 2023. World Health Organization;

2.	Programme National de Lutte contre le Paludisme - PNLP. Plan Stratégique National de Plaidoyer en Matière de la Lutte contre le Paludisme en Côte d'Ivoire 2018-2023. Johns Hopkins University; 2019.

3.	Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature. 2015 Oct 1;526(7572):207–11.

4.	Pryce J, Medley N, Choi L. Indoor residual spraying for preventing malaria in communities using insecticide-treated nets - Pryce, J - 2022 | Cochrane Library. [cited 2024 Sep 9]; Available from: https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012688.pub3/full

5.	Hilton ER, Rabeherisoa S, Ramandimbiarijaona H, Rajaratnam J, Belemvire A, Kapesa L, et al. Using routine health data to evaluate the impact of indoor residual spraying on malaria transmission in Madagascar. BMJ Glob Health. 2023 Jul;8(7):e010818.

6.	Chanda E, Coleman M, Kleinschmidt I, Hemingway J, Hamainza B, Masaninga F, et al. Impact assessment of malaria vector control using routine surveillance data in Zambia: implications for monitoring and evaluation. Malaria Journal. 2012 Dec 29;11(1):437.

7.	Epstein A, Namuganga JF, Nabende I, Kamya EV, Kamya MR, Dorsey G, et al. Mapping malaria incidence using routine health facility surveillance data in Uganda. BMJ Global Health. 2023 May 1;8(5):e011137.

8.	Selinger C, Angoa G. Analyse rétrospective 2012-2022 des activités de lutte contre le paludisme en Côte d'Ivoire. unpublished report; 2023.

9.	Hilton ER, Gning-Cisse N, Assi A, Eyakou M, Koffi J, Gnakou B, et al. Reduction of malaria case incidence following the introduction of clothianidin-based indoor residual spraying in previously unsprayed districts: an observational analysis using health facility register data from Côte d'Ivoire, 2018–2022. BMJ Glob Health. 2024 Mar;9(3):e013324.

10.	Bernal JL, Cummins S, Gasparrini A. Interrupted time series regression for the evaluation of public health interventions: a tutorial. Int J Epidemiol. 2017 Feb;46(1):348–55.

11.	Turner SL, Forbes AB, Karahalios A, Taljaard M, McKenzie JE. Evaluation of statistical methods used in the analysis of interrupted time series studies: a simulation study. BMC Medical Research Methodology. 2021 Aug 28;21(1):181.

12.	Gianacas C, Liu B, Kirk M, Tanna GLD, Belcher J, Blogg S, et al. Bayesian structural time series, an alternative to interrupted time series in the right circumstances. Journal of Clinical Epidemiology. 2023 Nov 1;163:102–10.

13.	Liu J, Spakowicz DJ, Ash GI, Hoyd R, Ahluwalia R, Zhang A, et al. Bayesian structural time series for biomedical sensor data: A flexible modeling framework for

evaluating interventions. PLoS Computational Biology [Internet]. 2021 Aug [cited 2024 Sep
9];17(8). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8412351/

14.   Midelet A, Bailly S, Borel JC, Hy RL, Schaeffer MC, Baillieul S, et al. Bayesian
Structural Time Series With Synthetic Controls for Evaluating the Impact of Mask Changes in
Residual Apnea-Hypopnea Index Telemonitoring Data. IEEE Journal of Biomedical and
Health Informatics. 2022 Oct;26(10):5213–22.

15.   Burnett SM, Davis KM, Assefa G, Gogue C, Hinneh LD, Littrell M, et al. Process and
Methodological Considerations for Observational Analyses of Vector Control Interventions in
Sub-Saharan Africa Using Routine Malaria Data. 2025 Jan 7 [cited 2025 Mar 4]; Available
from: https://www.ajtmh.org/view/journals/tpmd/112/1_Suppl/article-p17.xml

16.   Hersbach H, Bell B, Berrisford P, Hirahara S, Horányi A, Muñoz-Sabater J, et al. The
ERA5 global reanalysis. Quart J Royal Meteoro Soc. 2020 Jul;146(730):1999–2049.

17.   The PMI VectorLink Project. The PMI VectorLink Côte d'Ivoire 2019 Annual
Entomological Report, January 2019–December 2019. 2020.

18.   The PMI VectorLink Project. The PMI VectorLink Côte d'Ivoire 2020 Annual
Entomological Report, January 2020–December 2020. Rockville, MD. The PMI VectorLink
Project, Abt Associates.; 2021.

19.   The PMI VectorLink Project. The PMI VectorLink Côte d'Ivoire 2021 Annual
Entomological Report, January 2021–December 2021. Rockville, MD. The PMI VectorLink
Project, Abt Associates.; 2022.

20.   Institut National de la Statistique - INS, Ministère du Plan et du Développement -
MPD, Programme National de Lutte contre le Paludisme - PNLP, Ministère de la Santé et de
l'Hygiène Publique - MSHP, ICF. République de Côte d'Ivoire Enquête de Prevalence
Parasitaire du Paludisme et de l'Anémie (EPPA-CI) 2016 [Internet]. Abidjan, Côte d'Ivoire:
INS/MPD/PNLP/MSHP and ICF; 2016. Available from:
http://dhsprogram.com/pubs/pdf/FR330/FR330.pdf

21.   Institut National de la Statistique, ICF. Côte d'Ivoire Enquête Démographique et de
Santé 2021 Rapport final [Internet]. Rockville, Maryland, USA et la Côte d'Ivoire: INS et ICF;
2023. Available from: https://www.dhsprogram.com/pubs/pdf/FR385/FR385.pdf

22.   Honaker J, King G, Blackwell M. Amelia II: A Program for Missing Data. J Stat Soft.
2011 Dec 12;45(7):1–47.

23.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. J Stat Soft. 2015 Oct 7;67(1):1–48.

24.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear
Mixed Effects Models. J Stat Soft. 2017 Dec 6;82(13):1–26.

25.   Scott SL, Varian HR. 4. Bayesian Variable Selection for Nowcasting Economic Time
Series [Internet]. Goldfarb A, Greenstein SM, Tucker CE, editors. Economic Analysis of the
Digital Economy. Chicago: University of Chicago Press; 2015. p. 119–36. Available from:
https://doi.org/10.7208/9780226206981-007

26.   Brodersen KH, Gallusser F, Koehler J, Remy N, Scott SL. INFERRING CAUSAL
IMPACT USING BAYESIAN STRUCTURAL TIME-SERIES MODELS. The Annals of Applied
Statistics. 2015;9(1):247–74.

27.   Gogue C, Wagman J, Tynuv K, Saibu A, Yihdego Y, Malm K, et al. An observational
analysis of the impact of indoor residual spraying in Northern, Upper East, and Upper West
Regions of Ghana: 2014 through 2017. Malaria Journal. 2020 Jul 11;19(1):242.

28.   Ronald M, Humphrey W, Adoke Y, Jean-Pierre VG. Impact of population based
indoor residual spraying in combination with mass drug administration on malaria incidence
and test positivity in a high transmission setting in north eastern Uganda. Malaria Journal.
2023 Dec 13;22(1):378.

29.   Kouassi BL, Edi C, Ouattara AF, Ekra AK, Bellai LG, Gouaméné J, et al.
Entomological monitoring data driving decision-making for appropriate and sustainable
malaria vector control in Côte d'Ivoire. Malar J. 2023 Jan 12;22(1):14.

30.   The PMI VectorLink Project. The PMI VectorLink Côte d'Ivoire 2020 End-of-Spray
Report: August 10–September 12, 2020. Rockville, MD. The PMI VectorLink Project, Abt
Associates Inc.;

31.   The PMI VectorLink Project. The PMI VectorLink Côte d'Ivoire 2021 End-of-Spray
Report August 2–September 4, 2021. Rockville, MD. The PMI VectorLink Project, Abt
Associates.;

32.   Cameron E, Battle KE, Bhatt S, Weiss DJ, Bisanzio D, Mappin B, et al. Defining the
relationship between infection prevalence and clinical incidence of Plasmodium falciparum
malaria. Nat Commun. 2015 Sep 8;6(1):8170.

560
561

 Supplementary Material

 Supplementary Figure 1: Missing data of contemporaneous covariates in the four districts of

 the study.

Variable name	Description	Source	monthly	yearly
Rainfall	Total precipitation in m	ERA5	X	
Temperature	2m temperature in degree Celsius	ERA5	X	
Leaf	Leaf area index, low vegetation	ERA5	X	
Humidity	2m dewpoint temperature in degree Celsius	ERA5	X	
biting_rate_hcl_indoor	Indoor biting rate of An. gambiae and funestus combined	PMI-Vectorlink	X	
biting_rate_hcl_outdoor	Outdoor biting rate of An. gambiae and funestus combined	PMI-Vectorlink	X	
EIR_bpn	Infectious bites per night per person	PMI-Vectorlink	X	
N_hf_active	Number of health facilities reporting cases during a month per 1000 population	DHIS2	X	

hf_reporting_rate	Number of consecutive months reporting cases per months from first report until end of year	DHIS2		X
Testing	Number of tested (RDT or blood spot) per suspected cases	DHIS2	X	
tpr_total	Number of positive tested per total number of tested (RDT or blood spot)	DHIS2	X	
ipt1_coverage	Number of IPT doses administered during first trimester ANC visit per total number of first trimester ANC visits	DHIS2		X

Supplementary Table 1: Description of covariates used in the causal impact analysis by source and periodicity.

Adjustment	Description	Details	Formula	Data sources
0	crude incidence	uncomplicated & complicated cases	N0	DHIS2, PMI
1	presumed (**pr**) & test postivity rate (**tpr**)	Add cases that have not been confirmed (i.e. presumed) by assuming that their chance to test positive is as **tpr**	N1=N0+pr*tpr	DHIS2
2	reporting rate (**rr**)	Take into account the annual reporting rate	N2=N1/rr	DHIS2
3	health care seeking (**s**)	Add to incidence those cases with odds of having not seeked care (1-s)/s	N3=N2+N2*(1-s)/s	DHS 2021

Supplementary Table 2: Adjustment of incidence rate for presumed (i.e. untested) cases, reporting rate and health care seeking applied to both DHIS2 and PMI crude incidence.

regression coefficient	eliminated at step…	p-value (F test)
IRS:hf_reporting_rate	1	0.969484
IRS:N_hf_active	2	0.940614
N_hf_active	3	0.847019
IRS:tpr_total	4	0.758068
IRS:ipt1_coverage	5	0.616031
IRS:testing	6	0.599217
testing	7	0.360217
hf_reporting_rate	8	0.056271
ipt1_coverage	9	0.180847
IRS	0	8.62E-07
tpr_total	0	0.003838

577 Supplementary Table 3: Model selection of log-linear mixed-effect models of the difference
578 between DHIS2 and PMI incidence using backward step-wise reduction on *model1* (health
579 system covariates only)

regression coefficient	eliminated at step…	p-value (F test)
IRS:N_hf_active	1	0.931301
IRS:rainfall	2	0.92298
IRS:biting_rate_hcl_indoor	3	0.884611
Rainfall	4	0.84202
IRS:EIR_ibpn	5	0.80147
IRS:biting_rate_hcl_outdoor	6	0.822069
N_hf_active	7	0.802543
IRS:ipt1_coverage	8	0.642298
IRS:hf_reporting_rate	9	0.632453
IRS:tpr_total	10	6.88E-01
IRS:testing	11	0.66586
Testing	12	0.632313
biting_rate_hcl_indoor	13	0.542917
biting_rate_hcl_outdoor	14	0.521993
IRS:humidity	15	0.404982
IRS:leaf	16	0.648124
Leaf	17	0.501763

Humidity	18	0.32239
IRS:temperature	19	0.211798
Temperature	20	0.392839
IRS	0	9.32E-05
EIR_ibpn	0	0.087861
tpr_total	0	0.000661
hf_reporting_rate	0	0.032704
ipt1_coverage	0	0.047551

581   Supplementary Table 4: Model selection of log-linear mixed-effect models of the difference

582   between DHIS2 and PMI incidence using backward step-wise reduction *model2* (health

583   system covariates as well as environmental variables)