The age and sex dynamics of heterosexual HIV transmission in Zambia: an HPTN 071 (PopART) phylogenetic and modelling study

M. Hall*[1,2,3], W. Probert*[3], L Abeler-Dörner[2,3], C. Wymant[2,3], F. Di Lauro[2,3], X. Xi[4], R. Sauter[3], T. Golubchik[3,5], D. Bonsall[2,3], M. Pickles[6,7], A. Cori[6], J. Bwalya[8], S. Floyd[9], N. Mandla[10], K. Shanaube[8], B. Yang[10], P. Bock[10], D. Donnell[11], M.K. Grabowski[12], D. Pillay[13], O. Ratmann[4], S. Fidler[14], H. Ayles[1,8], R. Hayes[9], C. Fraser[2,3], the PANGEA-HIV consortium and the HPTN 071 (PopART) study team

[1] Department of Clinical Research, London School of Hygiene and Tropical Medicine, London, UK

[2] Pandemic Sciences Institute, University of Oxford, Oxford, UK

[3] Big Data Institute, University of Oxford, Oxford, UK

[4] Department of Mathematics, Imperial College, London, UK

[5] Sydney Infectious Diseases Institute, School of Medical Sciences, Faculty of Medicine and Health, University of Sydney, Camperdown, NSW, Australia

[6] MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College, London, UK

[7] HIV Prevention Trials Network Modelling Centre, Imperial College, London, UK

[8] Zambart, Lusaka, Zambia

[9] Department of Infectious Disease Epidemiology and International Health, London School of Hygiene and Tropical Medicine, London, UK

[10] Desmond Tutu TB Centre, Stellenbosch University, Cape Town, South Africa

[11] Fred Hutchinson Cancer Research Center, Seattle, US

[12] Bloomberg School of Public Health, Johns Hopkins University, Baltimore, US

[13] Division of Infection and Immunity, University College London, London, United Kingdom

[14] Department of Infectious Disease, Faculty of Medicine, Imperial College, London, UK

* These authors contributed equally

ABSTRACT

While much progress has been made in reducing the incidence of HIV-1 infection in

sub-Saharan Africa in recent years, bringing the epidemic to an end will require

identification of the demographic groups that continue to contribute to transmission.

Pathogen phylogenetics and individual-based mathematical models (IBMs) of

transmission are approaches that enable researchers to explore such questions.

Here, we used both methods to characterise the ages and sexes of the individuals

involved in heterosexual transmission in the context of the HPTN 071 (PopART) trial

in Zambia.

The two methods produced largely concordant results, strengthening confidence in

both. A principal finding was that when the age gap in transmission (the difference of

ages between the two individuals) was stratified by recipient age, the largest

differences were for the youngest female recipients and the smallest for the

youngest males. For women under 21 this stood at a male 9.87 years older (95% CI:

8.02 - 11.59) in the phylogenetics, compared to 6.93 (95% HDI: 6.56 - 7.32) in the

IBM. As the age of female recipients increased, this gap decreased towards parity.

Conversely, the under-21 male recipients saw the smallest gaps with the female

older by 0.14 years (95% CI: -2.95 - 3.23) in the phylogenetics and 1.38 years (95%

HDI: 0.98 - 1.68) in the IBM. As the age of male recipients decreased, this gap

steadily increased. The consequence of this pattern is that transmission to new age

cohorts first entering into sexual activity is driven predominantly by male-to-female

transmission.

We also showed that targeting interventions at younger adults captures most of the

benefit of population-wide approaches. We used the IBM to simulate the PopART

universal testing and treatment intervention into the future, showing that effective

treatment of under-35-year-olds would account for 94.3% (95% HDI: 65.8% -

126.6%) of the reduction in incidence by 2039 that would be achieved by treating the

entire population, while effective treatment of under-35 men accounts for 60% (95%

HDI: 23.2% - 92.1%).

Finally, we simulated a one-year cessation of ART treatment for the whole

population, which resulted in an immediate increase in both incidence and the

average age at transmission of both sources and recipients. The magnitude of this

was 4.6 years (95% HDI: 2.17 - 6.24) for female recipients, 5.3 (95% HDI: 2.74 -

7.09) for male recipients, 5.24 (95% HDI: 2.78 - 6.97) for female sources, and 6.04

(95% HDI: 2.92 - 8.09) for male sources. These changes would be slow to reverse

even after ART was restored.

These findings indicate that substantial reductions in HIV incidence can be achieved

through intensified testing and treatment of individuals aged under 35, and in

particular young men, a group that drives the infection of younger women and for

whom engagement with care remains disproportionately low.

INTRODUCTION

The widespread rollout of HIV testing, antiretroviral treatment (ART), and voluntary

medical male circumcision (VMMC) in combination prevention programmes has

resulted in considerable declines in both AIDS deaths and new HIV infections (1).

ART is highly effective in preventing HIV transmission as well as clinical illness (2),

and recent dolutegravir-containing regimens have reset the clock with regards to

antiretroviral resistance, whose prevalence had recently been increasing (3).

However, questions remain concerning in which demographic groups transmissions

are still occurring and how prevention interventions can best reach the groups that

contribute most to new infections; and also whether, in the light of rapid population

growth in sub-Saharan Africa (4), these groups will grow as proportions of the

population.

Heterogeneities in the epidemic are well known (5–8), including variations in HIV

acquisition risk by age and sex. Age and sex dynamics of heterosexual sexual

relationships in SSA, regardless of HIV status, have also been explored (9–11),

usually finding the male partner to be on average several (five or less) years older

than the female. Less is understood about which age and sex groups are most likely

to transmit HIV, and the age gap between sources and recipients of transmission.

This is partially because it is harder to identify sources of transmission than it is to

identify recipients. Sources can be identified using traditional epidemiological survey

methods, including recruitment of household pairs into a study (12), characterisation

of the sexual contacts of recipients by contact tracing (13) or interviewing those

recipients (5,14). These can be intensive endeavours, suffering from both recall (15)

and social desirability (16) bias.

Identifying heterogeneities in HIV transmission by age and sex may be critical in

informing the next steps of HIV prevention. For example, testing programmes

typically yield fewer new diagnoses for men than for women, with consequences for

the number of men linked to care. However, programmes aiming to end new

heterosexual infections cannot ignore or de-emphasise the role of men in

transmission, and must concern themselves with the male population. The risk of

being a source of HIV infection will not be evenly distributed amongst males. If

subpopulations disproportionately driving new infections can be identified, then the

focus and impact of interventions could be enhanced by tailoring services to meet

those men's needs.

Survey-based research into heterogeneities in age and sex in HIV transmission in

SSA has produced varied and sometimes contradictory results. A consistent

observation, in many studies conducted during the last two decades, has been that a

large age gap with a partner is associated with increasing HIV prevalence or

acquisition risk in (usually young) women (9,17). However, some recent studies did

not observe a simple increase in risk with increasing age of the male partner

(5,7,14,18–20), suggesting that a new epidemic environment with widespread ART

use has changed transmission dynamics (7). In this environment, partnerships most

likely to result in HIV infection would be with someone who is old enough to have

acquired HIV, but has yet to be diagnosed and started on ART. The effect would be

to reduce the average age of sources of transmission, and in turn reduce the gap

between source and recipient ages. The most recent studies, however, did find the

original association (21,22). Studies of this sort in men are much less common, but

those that have been conducted show a similar pattern; older studies have the risk of

HIV acquisition increasing for men with older or same-age partners when compared

to those with younger partners (9), but with this phenomenon disappearing or

becoming less straightforward in more recent work (14,23). Regardless of recipient

sex, characterisation of the demographics of sources in this study design is generally

rudimentary, with even their ages often simply derived from recipient interviews.

Individual-based mathematical models (IBMs) (24–26) and pathogen phylogenetics

(27,28) are alternative methods to surveys for describing characteristics of sources

of transmission. IBMs model entire transmission networks amongst simulated

individuals with the characteristics of interest. In pathogen genomics, recent

methodological developments have shown that "deep sequence" analysis of within-

host viral genetic diversity can identify pairs of individuals whose HIV infections likely

represent direct transmission, as well as the direction of that transmission (29–31).

For example the method *phyloscanner (27)*, validated using data on known sexual

partnerships (32–34), has been used to investigate transmission patterns through

the identification of transmission pairs.

A previous IBM study (35) predicted that increasing ART rollout would, due to

reduced force of infection, have an "ageing" effect on the epidemic, with the average

age of both sources and recipients of transmission expected to increase over time.

Amongst phylogenetic studies, de Oliveira et al (6) suggested a cycle of transmission

in South Africa in which women under 25 are infected by markedly older men, while

female-to-male transmission occurred in older individuals of similar ages.

Significantly, they did not have access to a method to reconstruct the direction of

transmission using individual-level data, and instead assumed that it was determined

by the prevalence gradient in the relevant communities (36). A large age gap (a

mean of 9.6 years) was also observed in heterosexual transmission in Switzerland

(37). More recently Magosi et al. (8) used more sophisticated methods to infer

direction of transmission (38) and did not find a similar asymmetry in age differences

in Botswana. Instead, they found the age gap in transmission pairs to be small, albeit

with large uncertainty. Monod et al (39), working with data from the Rakai

Community Cohort Study in Uganda, identified a significant upward shift in the age of

both partners in male-to-female transmissions between 2003 and 2018, leading to a

declining proportion of new infections occurring amongst women aged 15-24 and an

increasing one amongst those aged 25-34.

The early months of 2025 saw a sudden suspension of the President's Emergency

Plan for AIDS Relief (PEPFAR). The programme was responsible for 49% of

international spending, and 34% of all spending, on HIV/AIDS in SSA during 2024

(40). While funding has not entirely ended, as of the time of writing its future still

remains highly uncertain (https://www.nytimes.com/2025/07/23/health/pepfar-

shutdown.html). Any suspension, even if temporary, of such a crucial component of

existing HIV care in SSA would likely have a profound impact on health outcomes in

the region. ART prevents both progression to AIDS and onward transmission of the

virus. As a result, a cessation or pause in viral suppression for the millions of HIV-

positive individuals who have relied on treatment provided under PEPFAR would

result in not only a significant increase in the number of individuals progressing to

AIDS, but also in incidence of new HIV infections. Modelling studies have predicted

that:

● In Zimbabwe persistent cuts would lead to a 78% increase in HIV incidence in
Zimbabwe by 2030 (41),

● In seven sub-Saharan African countries, a 90-day funding freeze would result
in between 35,000 and 103,000 new infections, depending on modelling
scenario (42)

● In low and middle income countries globally, cuts in aid spending coupled with
the withdrawal of PEPFAR would cause between 4,43 and 10.75 million
deaths between 2025 and 2026 (43).

There is an urgent need to understand the epidemiology of HIV in SSA as we move

into a new era of reduced international funding of HIV programmes (44).

Here, we provide a comprehensive characterisation of the age- and sex-specific

transmission dynamics during the large HPTN 071 (PopART) trial in Zambia using

two independent methods: an IBM and pathogen genomics. We use these to

investigate the potential impact of future interventions focussed on suppressing

transmissions from inferred source populations. We conclude that there is a

substantial opportunity to achieve substantial reductions in HIV incidence by

improving testing, treatment, and ART adherence of men aged 25-34. We also

investigate the effect of an immediate pause of one year of ART provision on these

dynamics, finding that it would cause a sudden and dramatic increase in the age of

infection for both sexes.

RESULTS

Data

HPTN 071 (PopART) was a cluster-randomised trial of a combination HIV prevention package involving universal testing and treatment in 21 communities, comprising one million individuals, in Zambia and South Africa. It was conducted between 2014 and 2018 (45,46). The intervention involved annual household visits from community health workers offering HIV counselling and rapid testing, followed by linkage to care for HIV-positive clients. The trial had three arms, with arms A and B testing variations of the package and arm C serving as a control arm; for full details see the trial paper (46). To estimate intervention impact, HIV incidence was measured in a cohort of around 2,000 randomly selected individuals in each community, called the Population Cohort (PC). The IBM 'PopART-IBM' was developed as part of the trial to simulate the epidemic and interventions to help understand trial findings and extrapolate them to different settings and timescales (47,48). The IBM was calibrated to various sources of age- and sex-stratified data from the trial. In simulating the whole epidemic, the IBM generates a large number of transmission pairs on a timescale spanning the entire HIV pandemic; for comparability with the phylogenetically identified pairs (described next), from these we selected for analysis a subset which would be eligible to be sampled as part of the phylogenetic study. Across the 1000 replicates of the simulation (see Methods) there were a mean of 213,542 transmissions with recipients in this set, with a 95% highest density interval (HDI) of (191,652 - 231,239).

An ancillary study to the main trial—the HPTN 071-02 (PopART) Phylogenetics Study—was conducted as part of PopART in the Zambian PopART communities only. The study used phylogenetics to identify and characterise sources of HIV transmission in these communities, thus informing modelling of the impact of different intervention scenarios. For this purpose, viral genomic deep-sequencing was performed on samples collected over the course of the trial from both HIV-positive individuals in the PC and HIV-positive individuals attending health-care facilities in the study communities who were not on ART at the time of attendance.

While enrollment in the former was capped at 45 years of age, there was no upper

age limit for inclusion in the latter. Analysing the resulting sequence data with

phyloscanner, we identified 355 opposite-sex transmission pairs with high

confidence in the direction of transmission.

The transmission pairs identified phylogenetically and those simulated from the IBM

give us two independent means of characterising transmission during the trial.

Two independent methods agree on sex and age differences in transmission

Male-to-female transmissions made up 56% (198/355) of the phylogenetically

identified pairs and 57.9% of the IBM pairs (95% HDI: 56.3% – 59.7%). This is in

contrast to males making up, at the start of 2017 in the IBM, 41.6% (95% HDI: 39.9%

- 43.3%) of infected individuals and 44% (95% HDI: 42.2% - 45.8%) of virally

unsuppressed individuals. To estimate the age at transmission of both partners in

each phylogenetic pair, we used an estimate for the time of infection of the recipient

derived from the HIV-phyloTSI method (49), together with the dates of birth of both

partners. The age, sex, and other characteristics of the phylogenetic pairs are

summarised in table 1. Figure 1 shows the distribution of source and recipient ages

at the time of transmission, by sex, as estimated by both the IBM and the

phylogenetic data. Table 2 shows the mean ages, by sex and direction of

transmission in the pairs identified using both methods. Note that the difference in

methodologies dictates that the phylogenetics estimate is given as a frequentist

mean and confidence interval (CI), while the IBM is a posterior mean of means

(PMM), together with the 95% HDI for the mean. The uncertainty intervals are not

directly comparable. A posterior predictive check of the phylogenetic means against

the posterior indicates evidence against the phylogenetics sample coming from the

IBM posterior for the female-to-male age gap (see the last column of table 2).

Estimates of the distribution of transmission age gaps (source minus recipient) are

shown in figure 2 and summarised in table 2. For phylogenetics there was weak

evidence of a difference in the gap separating the partners' ages between male-to-

female and female-to-male pairs (*t*-test *p*=0.058). In the IBM, the gap was 2.44 years

(PMM; 95% HDI: 2.11 - 2.83). A more detailed illustration of the transmission

dynamics can be seen as a who-aquires-infection-from-whom (WAIFW) matrix in

figure 3.

The age gap increases with recipient age in female-to-male pairs and decreases in

male-to-female pairs

The IBM provides a full transmission tree from which the complete dynamics of

transmission can be explored; as inclusion in the dataset is automatic, biased

sampling is not an issue. With the phylogenetic results providing independent

support for its accuracy, we proceeded to explore its outputs in more detail. Figure 4

shows alluvial plots for the transmission history of individuals who became infected

with HIV in the year 2016 in the best-fitting IBM simulation. Orange and green flows

show viral flow to younger and older groups respectively. For female-to-male

transmissions we see that these flows are almost equal, whereas male-to-female

transmission are overwhelmingly orange, i.e. to younger ages. Thus this shows that

there is a clear shift in age for male-to-female transmissions which is absent for

female-to-male transmissions, acting to "rejuvenate" the epidemic.

As this finding seemed contrary to the fairly consistent central estimates of the age

gap described above, we stratified the gap by recipient age (figure 5a,b). This

showed distinctly different patterns for the two directions. In female-to-male

transmission pairs the average age gap for the youngest recipients actually had the

female partner older, but this rapidly reverses as recipient age increases, with the

gap decreasing in a roughly linear fashion. For male-to-female pairs there is also a

decline in the gap with increasing recipient age, but the decline levels out in a

fashion appearing to be non-linear. These trends are similar in the phylogenetics

data (figure 5c,d), although the magnitude of the age gap for the youngest female

recipients is larger, with the mean gap for recipient women aged under 21 years

being 9.87 years (95% CI: 8.02 - 11.59), compared to a PMM of 6.93 years (95%

HDI: 6.56 - 7.32) in the IBM. The equivalent numbers for young men are a female

partner 0.14 years (95% CI: -2.95 - 3.23) older from the phylogenetics and 1.38

years (95% HDI: 0.98-1.68) from the IBM.

Transmissions from an undiagnosed source involve younger individuals

Using the IBM, we stratified the age and sex variables according to the diagnostic

status of the source. A posterior mean of 73.8% (95% HDI: 71%-76.3%) had an

undiagnosed source. This was 70.7% (95% HDI: 67.2%-73.6%) of female sources

and 76% (95% HDI: 73%-78.8%) of males. Figure 6 shows variation in this

proportion by age, with the largest proportions occurring in the youngest age groups

for both sexes.

We also examined age and sex dynamics by diagnostic status (table 3). Both

individuals involved in these transmissions were on average older when the source

had received an HIV diagnosis, regardless of sex. Male sources, female sources,

and male recipients were all a mean of around three years older when the source

was diagnosed. Female recipients, however, showed a smaller increase in age, with

the difference in ages between diagnosed and undiagnosed being 1.45 years (PMM;

95% HDI: 1.06-1.94). This combination meant that there was no significant

difference in age gaps by source diagnostic status for female-to-male couples, with

the PMM difference being 0.044 years and the 95% HDI overlapping zero (-0.12 -

0.2) but one existing for male-to-female pairs (PMM 1.09, 95% HDI 0.9 - 1.27).

Age patterns remain stable over time despite steep increases in ART coverage

We next investigated whether a shift in age patterns in the IBM occurred as a result

of increasing use of ART (7). Figure 7 shows posterior trends in source age, age

gap, and proportion of individuals on ART over time for the period 2005–2020. For

female-to-male transmissions, a small decrease was seen in both source ages and

the age gap. The PMM female source age was 29 (95% HDI: 28.1 - 30) years in

2005 and 27.4 (95% HDI: 26.1 - 28.7) in 2020, while the age gap had the male 3.07

years older (PMM; 95% HDI: 2.68 - 3.46) in 2005 and 2.63 (95% HDI: 2.23 - 3.06) in

2020. In male-to-female transmissions, source ages declined from 33.5 years (PMM;

95% HDI: 32.5 - 34.3) in 2005 to 32.5 (PMM; 95% HDI: 31.5 - 33.4) in 2012, but then

levelled off, with the mean being 32.5 (95% HDI: 31.3 - 33.6) in 2020. Age gaps were

stable between 2005 and 2012, going from 5.26 years (PMM; 95% HDI: 4.93 - 5.59)

to 5.21 (PMM; 95% HDI: 4.85 - 5.53), but then rose, reaching 5.48 (PMM; 95% HDI:

5.06 - 5.91) by 2020.

Rollout of the PopART intervention to individuals aged 35 or under is projected to

reduce incidence by 67% by 2040

We used the IBM to quantify the effect of targeting future interventions at different

demographic groups. We started from the model state of the entire population of the

control communities (those which were not subject to any part of the trial intervention

and received standard of care) at the end of 2019 (47). We considered alternative

hypothetical scenarios where the PopART intervention was used as a proxy to

deliver real-world intensive linkage to care programmes to specific demographic

subgroups. We focus a) on younger age groups, since our data shows that they are

more likely to be unsuppressed than older age groups living with HIV and b) on

younger men, who were shown to drive the rejuvenation of the epidemic (figure 4).

We modelled the effect of four different scenarios, starting in the year 2020:

1. Applying the PopART intervention to the entire population
2. Applying the PopART intervention to only the population aged between 18 and 34, with standard of care for other age groups
3. Applying the PopART intervention to only the male population aged between 18 and 34 , with standard of care for other male age groups and females
4. Standard of care

Figure 8 shows the effects of these four scenarios on HIV incidence, by two-year

period from 2018–2019 (prior to the introduction of the intervention). The posterior

mean incidence in 2018–2019 was 0.8 cases per 100 person-years at risk (95% HDI:

0.592 – 1.01). By 2038–2039, overall incidence was projected to decline to 0.43

(95% HDI: 0.2 – 0.693) even in the absence of any PopART intervention. The

PopART intervention, when provided to the entire population, further reduces this to

0.12 (95% HDI: 0.001 – 0.25). However, the effect of deploying it only to individuals

aged 18-34 has an only slightly smaller effect, with a posterior mean incidence of

0.14 (95% HDI: 0.024 – 0.27), while providing it only to men aged 18-34 gave an

incidence of 0.24 (95% HDI: 0.067 – 0.42). By the end of 2039, the former

intervention achieved 94.3% (95% HDI: 65.8% - 126.6%) of the reduction between

the "standard of care" scenario and full deployment (the stochastic nature of the IBM

meaning that the 18-34 intervention could outperform the full intervention in some

cases), and the latter 60.0% (95% HDI: 23.2% - 92.1%). The 5.78% (95% HDI:

4.7%-7.1%) of the population that remained virally unsuppressed in 2018-19 would

be reduced to 0.38% (95% HDI: 0.072% - 0.76%) by 2038-39 under the full

intervention, to 0.62% (95% HDI: 0.023% - 1.04%) when deployed only to those

aged 18-34 and and to 1.17% (95% HDI: 0.45% - 1.85%) when deployed only to

males 18-34.

Modelling a one-year gap in ART provision

Finally, again starting with the model state in the control communities, we modelled a

one-year complete pause in ART provision for the year 2025. Following this, re-

engagement with care took place following a subsequent positive test; no

deployment of additional resources or targeted interventions to re-integrate

individuals with care was modelled. We examined the effect of this cessation in ART

on age and sex dynamics, which can be seen in figure 9. The pause is responsible

for a substantial increase in incidence (fig 9a), with a posterior mean of 2067 (95%

HDI: 817 - 6708) new cases in 2024 increasing to 6086 (95% HDI: 3463 - 10172) in

2025. Regarding age and sex dynamics (figure 9b), the immediate effect was an

increase in the ages of all those involved in HIV transmission. The change in the

ages of all individuals involved in transmissions from 2024 to 2025 was 4.6 years

(PMM; 95% HDI: 2.17 - 6.24) for female recipients, 5.3 (PMM; 95% HDI: 2.74 - 7.09)

for male recipients, 5.24 (PMM; 95% HDI: 2.78 - 6.97) for female sources, and 6.04

(PMM; 95% HDI: 2.92 - 8.09) years for male sources. There was also a jump in the

age gap in male-to-female transmissions, going from a gap of 5.34 years (PMM;

95% HDI: 4.38 - 6.28) in 2024 to 6.78 (PMM; 95% HDI: 5.73 - 7.96) in 2025. No

similar effect was seen in female-to-male pairs. After the initial change in 2025, the

curves in figure 9 can be seen slowly returning to the patterns of the years 2019-

2024, but in the absence of specific strategies to re-engage the HIV-infected

population with care, this is not immediate once ART provision is re-established.

Sensitivity analyses

Appendices 1 and 2 describe two supplemental analyses of the phylogenetics data.

First we confirmed that the analysis was genuinely informative, differing from results

obtained by choosing pairs of individuals at random as "transmission pairs". Second

we performed a detailed sensitivity analysis of the effects of varying the settings for

the *phyloscanner* package. While the sets of transmission pairs obtained using

different settings were somewhat variable, the exact choice of pair set had little effect

on the main results of this study.

DISCUSSION

In this study we investigated the age and sex patterns of heterosexual HIV

transmission in Zambia during the HPTN071 (PopART) trial between 2014 and 2018,

using both an individual-based mathematical model (IBM) of the epidemic and

pathogen phylogenetics. We focussed on characterising transmission pairs, jointly

analysing sources and the recipients to whom these individuals transmit.

The population of sources — the subset of people living with HIV who transmit

onwards — has been characterised in several previous studies (8,39). For the

HPTN071 (PopART) trial we previously reported that male sources were on average

a few years older than female sources, that there was no disproportionately large

contribution to transmission from external communities or from individuals with drug-

resistant viruses, and that there was a disproportionately large contribution to

transmission from men aged 25–34 (50). These men should be a focus of efforts to

prevent onward transmission. Interventions are needed to find them, link and retain

them in care, and encourage viral suppression, even if they are more difficult and

costly to reach. In that study we found that linking one man in this age group to care

is likely to have a similar effect on incidence as linking two women.

Here, we further developed previous results by characterising transmission pairs, i.e.

analysing sources and their recipients jointly, not independently. Notably we

estimated the distribution of the age gap within a pair, separately for male-to-female

and female-to-male transmission. Most transmissions in either direction had the male

partner a few years older than the female, with means of between 2.83 and 5.87

years depending on data source and direction. Gaps of these magnitudes are

consistent with reports of partnerships in the general population (regardless of HIV

status) in the region (14,51–53).

We also observed more complex dynamics (figures 4 and 5), once recipients wre

stratified by age. For male recipients, at the youngest ages the age gap is near

parity, with the female source being a posterior mean of 1.38 (95% HDI: 0.98 - 1.68)

years older for males under 21 in the IBM. These men would tend to take a partner

younger than themselves (judging by the patterns in older men). However, such

partners, if sexually active at all, are unlikely to be HIV-positive because there is a

delay, which we here refer to as the "exposure lag", between the expected age of

first potential exposure to HIV and the expected age of infection. As the age of male

recipients increases, the age gap flips in direction and then increases.

For female recipients, at the youngest ages the gap is particularly large, with the

male partner being a posterior mean of 6.93 (95% HDI 6.56 - 7.32) years older for

women under 21. The exposure lag means that sexual partners who are HIV-positive

are older than the average partner, and in addition the contribution from similarly-

aged and younger men is reduced as few of those have reached sexual debut. As

the age of female recipients increases, this initially large age gap decreases steadily

until it reaches parity for recipients in their early 60s. Women form new partnerships

with men on average a few years older than themselves, and this remains true

throughout their life (11). Thus this decline in the age gap in HIV transmission pairs

is likely to be driven by a combination of ART treatment and AIDS-related mortality in

older men. For middle-aged and older women, the population of potential male

partners will thus be increasingly virally suppressed or HIV-negative. It should be

emphasised that, while HIV prevalence increases with age, only a relatively small

proportion of new infections occur in older age groups and the contribution of these

to the epidemic is relatively minor; across the IBM a posterior mean of only 4.66% of

transmissions (95% HDI: 3.85% - 5.45%) were from individuals aged 50 or greater.

The consequence of this very divergent pattern of age gaps amongst the youngest

male and female recipients is that male-to-female transmission is primarily

responsible for the infection of new age cohorts as they reach sexual debut. This can

be most clearly seen in figure 4, where the youngest female recipients (pale pink)

are the recipients of large flows from older male groups (orange flows) while males

of the same ages (pale blue) have smaller incidence, with a much greater proportion

of that incidence coming from sources of the same age group.

A previous paper published by de Oliveira and colleagues (6) interpreted

transmission patterns in SSA as a cycle in which young women are infected by much

older men (the "sugar daddy" phenomenon), then these women age and later

transmit back to similarly-aged men. The study was based on phylogenetic evidence

but did not use genomic data to reconstruct the direction of transmission. Our results

concur that the widest age gap in transmission to females occurs in the youngest

recipient age group. Our IBM estimate has male sources a posterior mean of means

6.29 years (95% HDI: 6.01 - 6.62) older than female recipients when the latter are

under 25, compared to the mean gap of 8.7 years (95% CI: 6.8 - 10.6) estimated by

de Oliveira et al. In our IBM the under-25 female age group makes up a posterior

mean of 50.3% (95% HDI: 45.9% - 55.2%) of female recipients, and 29.1% (95%

HDI: 26.7% - 32.4%) of all recipients, indicating that this route plays a very

substantial but not overwhelming role in overall transmission.

Our results diverge from those of de Oliveira et al. when it comes to female-to-male

transmission, as we see the smallest age gaps for the youngest male recipients and

a gradual increase with age. Men aged 18 were infected by women a posterior mean

of means of 1.24 years older (95% HDI: 0.76 - 1.72), but by age 25 this gap had the

women 0.92 years younger (95% HDI: 0.68 - 1.19), and by age 40, 6.88 years

younger (95% HDI: 6.08 - 7.64).

Another limitation of the de Oliveira et al approach was that it collected participants

into large age groups rather than considering age as a continuous variable, which

may mask subtle dynamics of the age gaps in transmission, an observation also

made by others (33,54).

At the population level, we see that young women are infected by men who are on

average up to a decade older (depending on method). These men will be in their

mid-20s to early 30s, and we do not find evidence of a major driver of new infections

from men considerably older than that. At the individual level, however, it will remain

true that having a much older partner poses an increased risk of HIV acquisition for a

woman under 25, as often seen in previous work (21,22). An older partner has had

more time to be exposed to HIV themselves, as evidenced by the phenomenon also

existing for new infections in men (23).

Of the other phylogenetic studies of these questions, Magosi et al (8), in Botswana,

found male partners to be a mean of 3.8 years older in male-to-female pairs (with a

standard deviation of 6.9 years) and female partners a mean of 1.3 years older in

female-to-male pairs (with a standard deviation of 12.7 years). These large standard

deviations give very wide confidence intervals, but the pattern we observed of larger

(male minus female) gaps in male-to-female transmissions was present. Monod et

al. (39), in Uganda, identified a clear bimodal distribution in male source ages - one

peak in the late 20s and another in the late 30s - which we did not find in Zambia.

They did not present comprehensive estimates of age gaps. Kusejko et al. (37), in

Switzerland, found a wide gap of 9.6 in heterosexual pairs (amongst those who do

not inject drugs). They used a simple phylogenetic distance criterion to identify linked

individuals (the paper is in part an exploration of the limitations of this approach

under different distance thresholds and sampling regimes) and did not attempt to

reconstruct by direction of transmission. The varied settings of these studies may

also account for some or all of the discrepancies with our work.

Increasing rollout of ART between 2005 and 2020 did coincide with changing

patterns of transmission (figure 7), and the suggestion in previous work that age

gaps in male-to-female transmission would decrease with ART rollout (because older

males were increasingly virally suppressed) gains some support. However, the

magnitude of these changes were small, with the posterior mean of mean gaps in

male-to-female transmission dropping from 3.07 (95% HDI: 2.68 - 3.46) years in

2005 only to 2.63 (95% HDI: 2.23 - 3.06) in 2020. The profound change in

epidemiology suggested by some previous studies (5,7,14,18) finds little footing.

We found that both individuals involved in transmission pairs with a diagnosed

source were older than those involved in pairs with an undiagnosed source. This is

most likely due to the time it takes for an individual to receive a diagnosis, coupled

with the age assortativity of partners. Interestingly, the increase in ages for female

recipients with diagnosed partners is smaller than that for males - a posterior mean

change in mean difference of 2.53 (95% HDI: 2.07 - 2.96) years for males and 1.45

(95% HDI: 1.06 - 2.96) for females - which in turn affects age gaps (see table 3). The

reasons for this are hard to entirely unpick but it may be driven by differences in ART

uptake between men and women.

We used two entirely independent methods, each with its advantages and

disadvantages. The IBM allows us to simulate very large numbers of complete

transmission chains, for which we can analyse all individuals with no sampling bias

or erroneous identification of pairs. However, the results are dependent on model

assumptions (for example, the age mixing matrix (47)), and the individuals involved

are simulated. The phylogenetics data are the reverse: there are no model

assumptions and the data come from genuine cases of HIV infection, but the sample

size, although large compared to similar studies, is several orders of magnitude

smaller than the actual population, and inference may be affected by sampling bias

and imperfect pair ascertainment. The results of the two methods being largely

consistent is a key strength of this study: it gives confidence in both methods, and in

additional estimates generated by the IBM for scenarios and periods for which no

phylogenetic data exist. Nevertheless, some issues may affect both methods.

Notably, neither of our methods attempt to quantify the role of men who have sex

with men (MSM) in transmission. They are not modelled in the IBM, and, as we

analysed opposite-sex pairs only in the phylogenetic study, we do not consider their

role there either. In addition, while the IBM includes heterogeneity in sexual risk taking

behaviour it does not explicitly model key heterosexual populations, such as female sex

workers (FSW). Membership of such groups is unknown for participants included in the

phylogenetic study, and they might also be undersampled. Of note, FSW and MSM are

both groups which were not modelled in the IBM and were likely undersampled in the

phylogenetic study. Our estimates of the effect of interventions (figure 8) rely on

these groups not contributing significantly to transmission. We previously estimated

that sex between men makes limited, if any, contribution to transmission within the

population studied here (50).

The posterior predictive tests of table 2, which give *p*-values for the central estimates
in the phylogenetics sample coming from the IBM, provide, in the case of the age
gap in female-to-male pairs, strong evidence for a difference in the estimates derived
from the two methods. However, the actual magnitude of the difference in means is
fairly small (1.61 years), and this would not change the interpretation of the findings.
Digging in further, while the decline in the age gap as a female recipient ages is seen
in both methodologies, the magnitude of the gap for the youngest women is the main
area in which the two methods differ (figure 5), with larger gaps seen in the
phylogenetics. The mean gap in the best-fitting IBM for female recipients aged 20 or
younger was 6.93 years (95% HDI: 6.56 - 7.32) while for the phylogenetics it was
9.87 years (95% CI: 8.02 - 11.59). It is not entirely clear why this might be.
Potentially relevant limitations in the data used to fit the IBM are the absence of trial
data for under-18s and the lack of care cascade information from before 2013.
Alternatively, there might have been a bias towards finding older-than-average
partners for younger women in the phylogenetics data. Despite this disparity, the
similarity between the trends in age gaps identified with both methods is striking. The
age gap between a female recipient and her male partner at the time of transmission
is largest for the youngest women and declines asymptotically towards equality as
the recipient's age increases. The youngest male recipients, on the other hand, are
infected by a female source of a similar age to (or even slightly older than)
themselves. With increasing recipient age, this gap subsequently widens in the
direction of younger source ages.
596

As alluded to above, a limitation of this study is the narrow geographical catchment
area of the PopART trial. The results should not be assumed to hold identically in
other SSA settings. We also do not control for bias in the composition of the
phylogenetic dataset, both in regard to demographics and to the ability to
successfully acquire a sequence from a participant providing a sample, which we
assume is random. Nevertheless, the consistent results between the two
methodologies should assuage fears about this, as the IBM results are not subject to
sampling bias problems.
605

With the 95:95:95 targets being in reach in many parts of SSA, it is becoming
increasingly expensive to link individuals to care using population-wide testing and

messaging campaigns. There is widespread agreement that future prevention

campaigns need to focus on the remaining 5:5:5 of the population. We estimated

that an intensive effort to link people to care, such as the PopART intervention,

would result in almost as much reduction in incidence if focussed on those aged

between 18 and 34, and would retain two thirds of the benefit when focussing on

men between 18 and 34 only. Thus, the key role of young men in the infection of

young women reaching sexual debut, outlined in this paper, means that a

considerable (although not optimal) incidence reduction can be achieved by treating

them alone. We did not methodically explore potential age cut-offs in this work, and

the identification of the optimal demographic groups to target goes beyond the scope

of our analysis. Nevertheless, our results show the potential benefits of focussing on

groups who do contribute disproportionately to transmission. Enabling and

encouraging younger men to seek and stay in care to the same extent as women

would improve the health and life chances of many men, and rapidly reduce infection

rates in their partners.

Monod et al (39) modelled a counterfactual scenario different from but related to

ours. Where we simulated the long-term impact of an intervention based on real-

world implementation data, where the hypothetical element is targeted at different

demographic groups, they simulated an alternative hypothetical intervention that

would either fully close the gap in viral suppression between men and women, half

close it, or achieve the UNAIDS 95:95:95 treatment targets in males. Both sets of

scenarios predicted that considerable benefits in reducing female incidence could be

achieved by targeting a relatively small number of HIV-positive men. This

undertaking would be much less intensive than targeting the much larger number of

HIV-negative women at high risk of acquisition.

A major disruption in ART provision as a result of changes in the global aid

landscape is now underway, with the endpoint unclear. As ART prevents both

disease progression and transmission, simultaneous increases in the incidence of

AIDS and of HIV infection is to be expected, and the response of the global health

community will need to focus on both. In the latter case it is important to anticipate

not only an increase in new infections, but in what populations they would occur. Our

results suggest a shift in incidence to older age groups, caused by the withdrawal of

ART from those previously virally suppressed, at which point their partners, both

existing and new, are at greatly increased risk of HIV infection. This ageing of the

incidence profile would linger even after the restoration of ART provision unless

procedures were put into place to re-engage those individuals with care.

In summary, using two independent and largely concordant methods, we have

shown that the age gap between HIV transmission partners in Zambia varies

substantially with the age and sex of the recipient, and that male-to-female

transmission is the primary route by which new age cohorts are infected upon

reaching sexual debut. The effective treatment of men under 35 could achieve the

majority of the incidence reduction delivered by population-wide interventions. At a

time when the sustainability of international HIV funding is in question, understanding

where transmission occurs, and where it would shift in the event of disrupted care, is

essential for designing interventions that achieve the greatest impact with finite

resources.

METHODS

HPTN071 (PopART) trial

The HPTN071 (PopART) trial has been described in detail previously (45,46). In

summary, the impact of a universally delivered combination prevention package was

tested in 21 communities in Zambia and South Africa, between 2013 and 2018

(45,46). The total population of the 21 study communities was approximately one

million. The intervention comprised annual rounds of home-based HIV testing,

including support for linkage to care and retention on ART, promotion of medical

male circumcision for HIV-negative men, and additional HIV, tuberculosis (TB), and

sexually-transmitted infection (STI) services. It was delivered by teams of

Community HIV-care providers (CHiPs). The 21 communities were arranged in

seven triplets, matched on baseline HIV prevalence, engagement with care, and

geographic location (Hayes et al., 2019). Communities were randomly allocated into

one of three arms within each triplet. Arm A communities received the full

combination HIV prevention package and offer of immediate ART for individuals that

tested HIV-positive. Arm B communities received the full combination HIV prevention

package with ART initiation following national guidelines (where ART initiation was

dependent upon an individual's CD4 count). Arm C communities received the

standard of care. National guidelines on ART eligibility changed over the period of

the trial, and thus from late 2016, ART for all HIV positive individuals was standard

care in both countries (aligning arms A and B). The primary endpoint was the relative

reduction in HIV incidence between arm A or B communities and their matched arm

C community between 12 to 36 months after trial start. This was measured in a

cohort of 2000 randomly selected individuals aged 18-44 per community, called the

Population Cohort (PC), who were followed up in four annual rounds. Rounds of the

PC are referred to as the number of months after trial start (e.g. PC0 is 0 months

after start).

IBM

For each of the 21 communities in the HPTN017 (PopART) trial, the IBM (47) models

every individual in a growing population that reaches a size of approximately 50,000

in 2014, roughly matching the actual community sizes at that time point.

Demographic parameters follow historical estimates of mortality and birth rates from

the UN Population Division (55), and future projections of these measures follow the

UN's intermediate estimate (56). HIV infections are introduced into the model in

1980, HIV transmission is modelled in serodiscordant couples, and HIV disease

progression (in the absence of ART), follows data from the ATHENA cohort in the

Netherlands (57). While the disease progression data are derived from studies of

subtype B, previous work has found no significant difference in CD4 progression by

subtype (58,59)

Individuals are categorised into one of three groups of sexual risk-taking behaviour,

where membership of a higher risk group reflects both a higher propensity to form

partnerships and an increased maximum number of concurrent partnerships. An

individual's risk group is assumed not to change through time. Partnerships are

formed based upon an individual's age, sex, and sexual-risk taking behaviour, with

sexual risk-taking behaviour acting independently of age and sex. Partnership

formation and dissolution rates are calibrated using survey data from the PC, and

incorporate assortativity by risk group; see (47) for full details. Age mixing was

determined using the age of the last reported partner (within the last year) based

upon extensive sexual behaviour surveys. To allow for transmissions from areas

outside the trial communities, the IBM models two "patches" representing the

community and the surrounding area. Partnerships may also form with individuals in

the surrounding area, but at a reduced rate (45% of the rate within the community).

When serodiscordant, these between-community partnerships are also at a lower

(29% of within-community) risk of HIV transmission, reflecting higher rates of

condom usage reported in the PC data.

Testing and linkage to care were modelled as two separate processes, a

"background" component representing testing outside the context of the trial (e.g.

through clinics and antenatal care) and a "CHiPs" component representing the trial

intervention, which was disabled for the control communities. The subsequent care

cascade following a positive test was determined by the component from which the

test came; for full details see (47).

Parameters for partnership formation and dissolution were set using data from the

PC. Other parameters were estimated using Approximate Bayesian Computation

(ABC) to fit the model to five sources of data, all stratified by age and sex: 1) HIV

prevalence from the Zambian Demographic and Health Surveys (DHS) of 2002,

2007 and 2013; 2) HIV prevalence from the final intervention round of the trial; 3) the

proportion of individuals aware of their status among HIV-positive individuals from all

three intervention rounds; 4) the proportion of individuals on ART among those who

are HIV-positive and aware of their status from all three intervention rounds; and 5)

the proportion of individuals virally suppressed amongst all HIV-positive individuals

at PC24. For all but 1), which was taken from previous work, fitting used trial data

from an intervention community (arm A) in Zambia. The calibration framework fits the

epidemic selecting 1000 most likely parameter sets, each of which is used to

generate a single simulation. An ABC distance measures distance between

observed and simulated summary statistics used in the fitting process.

Point estimates for key quantities presented here are the posterior mean and 95%

highest posterior density (HPD) interval over 1,000 ABC simulations. We also

identified the 'best-fitting' IBM simulation - the model replicate whose ABC distance

was smallest - and used the corresponding transmission tree when we presented

results that require a single realisation of the simulated epidemic.

The IBM was used to model transmission over a 75-year period from 1975 to 2050.

To make the outputs comparable to the phylogenetics data, we selected only

transmissions for which the recipient would have been eligible to provide a viral

sample during the trial, specifically that they were infected with HIV before 2019 and

alive in November 2013. (We did not model the viral dynamics that would result in

the failure of a HIV-positive individual to provide a viable sequence; effectively we

assume that the probability of this happening is independent of participant

demographics or time of infection.) We also used only model output from the

communities for which extensive phylogenetic data were available; this excluded the

entire South African component of the study as well as one triplet of communities in

Zambia for which health-care facility phylogenetic data were not collected. All the

simulations taken forwards for full analysis were from the nine remaining Zambia

communities. Transmissions involving individuals in both the "inside" and "outside"

patches of the simulation were included (47).

Phylogenetics

The HPTN 071-02 (PopART) Phylogenetics Study and the sequencing procedure

used have been described previously (45,46,50). In summary: samples for HIV

genome sequencing were obtained from HIV-positive individuals in the PC (during

the earliest visit in which they tested positive for the virus, which could be baseline),

and individuals who had been approached at the health-care facilities between 2016

and 2018. These individuals were not on ART at the time they were recruited and

were in most attending the clinics for ART initiation, although some self-reported

previous ART and poor adherence. Whole-genome deep sequencing was done on

the Illumina platform using veSEQ-HIV (60).

The process by which probable transmission pairs were identified has also been

previously described (50). Briefly, after genome assembly using SPAdes (61) and

shiver (38), the collection of viral consensus sequences was clustered using HIV-

TRACE (28) with an intentionally generous distance threshold of 0.04 substitutions

per site. The deep-sequence data files from all members of every identified cluster

were then used as input for separate analyses with *phyloscanner* v1.8.0 (27). The

set of genomic windows used in *phyloscanner* were uniformly of 250bp in length,

spanning the entire HIV genome, such that the start of each window was 10 bp

further along the genome (relative to HXB2) than the previous one. The distance

threshold used to identify likely transmission pairs in a single window was of 0.025

substitutions per site on a normalised scale designed to represent average distances

on the *gag* and *pol* genes (62). After adjustment for missing windows, overall

assignment of pairs as likely to involve direct transmission was declared where more

than 50% of windows suggested it. Same-sex pairs (which indicate at least one

missing intermediary under the assumption that transmission is heterosexual) were

then discarded to arrive at a final set; previous work has estimated that the

proportion of true male-to-male transmissions was very low amongst trial participants

(50).

The date of HIV infection of each individual was estimated using HIV-phyloTSI (63)

and these dates, together with recorded dates of birth, were used to estimate the

age of source and recipient partners in each pair at the time of the infection of the

recipient. While our previous analysis (50) investigated only transmissions estimated

to have occurred during the time-frame of the PopART trial itself, this filter was not

applied here.

Direction of transmission was determined using both the *phyloscanner* score based

on the tree topology and comparison of infection dates as estimated by HIV-

phyloTSI; see (50) for full details.

Posterior predictive test for phylogenetics results against the IBM

To test whether the phylogenetics results were compatible with being derived from

the IBM, we performed a posterior predictive check. This determines the probability

that a particular statistical estimate from the phylogenetics (in this case the mean

source or recipient age, or mean age gap) is drawn from the posterior distribution of

that statistic from the IBM. We accounted for error in the phylogenetics estimate by,

for each posterior draw, also drawing an error term from a normal distribution with

mean 0 and standard deviation equal to the standard error of the phylogenetics

estimate.

Appendix 1: Comparison of identified pairs from phylogenetics to random

pairs

In this appendix, we explore the presence of a genuine signal of HIV transmission in

the phylogenetics dataset, by comparing the results obtained from *phyloscanner* to

the results of attaching two opposite-sex individuals in the genomic dataset to each

other at random as false transmission pairs.

Specifically, we picked a random selection of 156 male and 197 female participants

for whom sequence data were available. (156 and 197 are the number of recipients

of each sex in the actual dataset.) We additionally selected these such that their

estimated infection dates (as determined by phyloTSI) were no earlier than the first

such estimate amongst the recipients in the actual data. For each of these 353

individuals we randomly picked an individual of the opposite sex, whose estimated

infection date was earlier than their own, as a shuffled "source" for their

transmission. The time of transmission was taken to be the estimated time of

infection of the "recipient". We repeated this process 1,000 times and compare the

distribution of age- and sex-related variables across these replicates to those in the

real dataset (Appendix 1-Figures 1 to 3).

Recipients. See Appendix 1-Figure 1 for a comparison of the mean and variance of

recipients age at the time of transmission, by sex, across the replicates of the

shuffled dataset (histograms) to its true value (green line). There is no evidence of a

difference with respect to either mean or variance for either recipient sex (all

empirical *p*-values >0.178). This provides reassurance that the age distribution of

individuals with a sampled source does not differ from that of the complete set of

participants providing sequences.

Estimated time since infection showed considerable variation between real and

shuffled datasets (Appendix 1-Figure 2), with individuals from shuffled pairs (i.e.

those whose sources had not genuinely been sampled) showing greater mean

values of the variable (empirical p-values <0.001 for both sexes). This is consistent

with younger infections being more likely to have a sampled source; elapsed time

from infection would be associated with increasing probability that the source would

not be eligible for sampling, either due to migration or commencing ART (as

successful ART treatment prevents the acquisition of a sequence).

Sources. In appendix 1-figure 3, mean estimated source ages at the time of

transmission are compared. For male sources there is no evidence of a difference in

mean age (empirical p=0.41 but female sources are younger in the actual dataset

(p<0.001). The variances of source ages also deviate from the null hypothesis for

both sexes (p=0.02 for males, p<0.001 for females). Effectively, this test compares

the demographics of those individuals identified as sources by the analysis to the

general HIV positive population, as the shuffled "sources" need not have genuinely

transmitted HIV. It is unsurprising that these variables differed. In this case, a

younger age for female sources of transmission when compared to a random sample

of HIV-positive women is consistent with onward transmission peaking in younger

age groups. A smaller variance in the ages of genuine sources is also consistent

with their being a peak age of risk for transmitting the virus (see figure 1).

As with recipients, "sources" in the shuffled dataset tend to have older infections

(Appendix 1-Figure 4). In this case this is likely due to a combination of the same

rationale outlined for recipients above, and the fact that genuinely identified sources

are younger than a random selection of individuals (empirical p<0.001 for both

sexes). They would be expected, on average, to have more recent infections

because the range of possible times since infection is constrained by a more recent

sexual debut.

Age gap. Appendix 1-Figure 5 displays the histograms for the age gap between

male and female partners. There is strong evidence for a difference in means for

female-to-male transmission (empirical p<0.001) but not for male-to-female

(p=0.307). Meanwhile there are plainly large differences in the variances (p<0.001),

meaning that genuine transmission pairs have much more closely correlated ages

than random pairs.

Discussion. The sampling frame for this study is unusual. We have taken a sample

of HIV-positive individuals in the trial communities, and within that sample identified

individuals who we are confident acted as sources of transmission for others in the

sample. Only 4.96% of sampled individuals have a source identified in this way. As

every HIV-positive individual is by definition a recipient of transmission, this 4.96%

can be seen as a subsample of the sampled recipient population. If the subsample

were unrepresentative, meaning that those recipients for whom we found a source

were different to those for whom we did not, it would in turn suggest bias in the

characteristics of the sources that we identified for these recipients. That there was

no statistical difference between the true subsample of recipients with sampled

sources and random collections of recipients of the same size should give us

confidence that this bias was not present.

The *phyloscanner* procedure would obviously have failed if it was doing no more

than sampling random opposite-sex pairs of individuals in the dataset and

designating one of each as the source of the infection in the other. Were this true,

then the demographics of the population identified as sources would be no different

from those of the entire sample, and the distribution of age gaps in identified pairs

would be no different from that of random pairs of opposite-sex individuals in the

dataset. We found strong statistical evidence that neither of these are the case.

Appendix 2: Sensitivity analysis for *phyloscanner* settings

In this section we re-investigate the age and sex dynamics of the set of identified

transmission pairs from sets of pairs obtained when some of the key *phyloscanner*

settings are varied. For further information regarding these settings, see the

phyloscanner paper (27) and manual (available at https://github.com/BDI-

pathogens/phyloscanner)

Read downsampling

The nature of deep sequencing is such that genome coverage will vary both between

samples and in different regions of the genome for the same sample. It may well be

914 the case that, for a given *phyloscanner* window and a putative transmission pair, the

915 number of reads covering the window can differ between the two members of the

916 pair. The exact effects of this on inference have yet to be exhaustively explored, but

917 one *phyloscanner* option allows the user to specify a maximum number of reads to

918 be analysed per window, with the exact choice of reads being randomly selected.

919 (Note that, as many reads in the same window will be identical due to the

920 amplification process during sequencing, regardless of underlying variation within the

921 sample, this process is not the same as limiting the number of distinct genomic

922 *variants*.) A further choice must be made regarding how to proceed if the total

923 number of reads for one host in one window fails to reach that maximum number.

924 The two options are, firstly, to include all the reads that do exist ("keep") or,

925 secondly, to remove that host from consideration entirely in that window ("blacklist").

926 (As no inference is conducted from windows in which reads are not present from

927 both hosts, the practical effect of the "blacklist" option is that those windows will be

928 excluded from the analysis.) In the main analysis of this paper we did not use read

929 downsampling; here we describe results from applying maximum read counts of 25

930 and 50, with both the "keep" and "blacklist" strategies.

932 As described in the main text, there are 355 pairs found when no read downsampling

933 was applied. Applying the tool always reduces this number, to:

935  ● 296 with a maximum of 50 reads per window and the "keep" strategy

936  ● 274 with a maximum of 50 reads per window and the "blacklist" strategy

937  ● 285 with a maximum of 25 reads per window and the "keep" strategy

938  ● 284 with a maximum of 25 reads per window and the "blacklist" strategy

940 There was no evidence of a difference in the proportion of pairs with a male source

941 (chi-squared test p=0.699) between the five groups. In this work, we have three key

942 variables of interest: source age at estimated time of transmission, recipient age at

943 estimated time of transmission, and age gap. Each of these was examined separately

944 according to the sex of the inferred source individual, making a total of six variables in

945 practice. We saw no evidence of a difference in any of these variables regardless of

946 the downsampling procedure. (Kruskall-Wallis test p>0.7 in all cases). Their

947 distribution can be seen in appendix 2-figure 1.

Window width

A key *phyloscanner* setting is the width of the sliding window used to subdivide the

viral genome. Narrower windows allow the genome to be tiled with more windows and

increase the number of reads per window because *phyloscanner* requires reads to

fully span a window to be included in it. However, narrow windows have fewer sites at

which mutations can be observed, reducing phylogenetic resolution. In the main text

the window width was set to 250 base pairs, with each window starting 10bp further

along than the previous. We also reduced this to 175bp, with the same 10bp

incremental shift, and obtained the set of probable transmission pairs from this

variation. The 355 pairs from the main analysis were reduced to 291 here. There was

no evidence that these sets differed in respect to the proportion of pairs which were

male-to-female (chi-squared test *p*=0.903) or of our six key quantities (Mann-Whitney

U-test *p*>0.63 in all cases). See appendix 2-figure 2 for the distribution of these values.

Window thresholds for pair identification

As phylogenetic relationships between the reads from each of two individuals will vary

across the genome, we apply thresholds on the proportion of windows in which a) the

phylogeny suggests that the pair are a transmission pair ("linkage"), and b) the

phylogeny suggests a direction of transmission. (These proportions are adjusted for

non-informative windows as outlined by (62)). In the main analysis, individuals are

linked as a pair at a proportion threshold of 0.5. For pairs that are above this threshold,

a direction of transmission is called at a proportion threshold of 0.33 (of the total

number of windows). We also tested more stringent thresholds, firstly 0.66 for linkage

and 0.5 for direction, and then 0.75 for linkage and 0.66 for direction. The 355 pairs

from the main analysis were reduced to 260 in the former case and 201 in the latter.

There is no evidence of a difference in the proportion of pairs identified as having a

male source (chi-squared test *p*=0.927) or in the distribution of our six quantities of

interest (Kruskal-Wallis test, all *p*>0.3). See appendix 2-figure 3 for the distribution of

these estimates.

Patristic distance threshold for per-window pair identification

The final setting that we varied was the patristic distance threshold at which, in a single genomic window, a pair of individuals are inferred as likely to be involved in direct transmission. As phylogenetic branch lengths vary across the genome, this distance is normalised to standardise to expected lengths on the *gag* and *pol* genes. In the main analysis the value used is 0.02 normalised substitutions per site. Here we use values from 0.01 to 0.04 in increments of 0.005.

Increasing this threshold should increase sensitivity, but decrease specificity, of the identification of transmission pairs. For heterosexual HIV transmission, instances where there is only a single unsampled intermediate host can be omitted by simply excluding same-sex pairs (e.g. male-male pairs must be missing at least one unsampled intermediate female). Then false positives must be missing at least two unsampled intermediaries (for example identifying male A → female B → male C → female D as male A → female D). Reducing the patristic distance threshold will reduce the probability that such false positives are included in the dataset. See appendix 2-table 1 for the number of pairs identified using each threshold.

There is no evidence of an association of the threshold with the proportion of transmission pairs identified with male sources, whether the threshold is treated as a discrete variable (chi-squared test p=0.962), or as continuous (logistic regression p=0.376). Similarly, none of our six key estimates show evidence for an association with the threshold. Again, this is true whether the threshold is treated as a discrete variable (Kruskal-Wallis p>0.8 in all cases) or fit as a continuous predictor of the variable in a linear regression model, with source sex as an interaction term (no coefficients other than source sex having p<0.3). See appendix 2-figure 4 for the actual distribution of their values.

That the key outputs of our analysis are robust to decreasing the distance threshold from 0.02 to 0.01 should offer reassurance towards concerns regarding any biasing effect on the results of this paper caused by the inclusion of false positive transmission pairs.

1015

References

1. UNAIDS. The Gap Report [Internet]. The Joint United Nations Programme on HIV/AIDS; 2016. Available from: https://www.unaids.org/sites/default/files/media_asset/UNAIDS_Gap_report_en.pdf

2. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. N Engl J Med. 2016 Sep 1;375(9):830–9.

3. Fogel JM, Wilson EA, Piwowar-Manning E, Breaud A, Clarke W, Petropoulos C, et al. HIV drug resistance in a community-randomized trial of universal testing and treatment: HPTN 071 (PopART). J Int AIDS Soc. 2022 Jul;25(7):e25941.

4. UNAIDS. The Youth Bulge and HIV [Internet]. The Joint United Nations Programme on HIV/AIDS; 2018. Available from: https://www.unaids.org/en/resources/documents/2018/the-youth-bulge-and-hiv

5. Balkus JE, Nair G, Montgomery ET, Mishra A, Palanee-Phillips T, Ramjee G, et al. Age-Disparate Partnerships and Risk of HIV-1 Acquisition Among South African Women Participating in the VOICE Trial. J Acquir Immune Defic Syndr. 2015 Oct 1;70(2):212–7.

6. de Oliveira T, Kharsany ABM, Gräf T, Cawood C, Khanyile D, Grobler A, et al. Transmission networks and risk of HIV infection in KwaZulu-Natal, South Africa: a community-wide phylogenetic study. The Lancet HIV. 2017 Jan;4(1):e41–50.

7. Harling G, Newell ML, Tanser F, Kawachi I, Subramanian SV, Bärnighausen T. Do age-disparate relationships drive HIV incidence in young women? Evidence from a population cohort in rural KwaZulu-Natal, South Africa. J Acquir Immune Defic Syndr. 2014 Aug 1;66(4):443–51.

8. Magosi LE, Zhang Y, Golubchik T, DeGruttola V, Tchetgen Tchetgen E, Novitsky V, et al. Deep-sequence phylogenetics to quantify patterns of HIV transmission in the context of a universal testing and treatment trial - BCPP/Ya Tsie trial. Elife [Internet]. 2022 Mar 1;11. Available from: http://dx.doi.org/10.7554/eLife.72657

9. Gregson S, Nyamukapa CA, Garnett GP, Mason PR, Zhuwau T, Caraël M, et al. Sexual mixing patterns and sex-differentials in teenage exposure to HIV infection in rural Zimbabwe. Lancet. 2002 Jun 1;359(9321):1896–903.

10. Kelly RJ, Gray RH, Sewankambo NK, Serwadda D, Wabwire-Mangen F, Lutalo T, et al. Age differences in sexual partners and risk of HIV-1 infection in rural Uganda. J Acquir Immune Defic Syndr. 2003 Apr 1;32(4):446–51.

11. Ott MQ, Bärnighausen T, Tanser F, Lurie MN, Newell ML. Age-gaps in sexual partnerships: seeing beyond "sugar daddies." AIDS. 2011 Mar 27;25(6):861–3.

12. Mujugira A, Baeten JM, Donnell D, Ndase P, Mugo NR, Barnes L, et al. Characteristics of HIV-1 serodiscordant couples enrolled in a clinical trial of antiretroviral pre-exposure prophylaxis for HIV-1 prevention. PLoS One. 2011 Oct 5;6(10):e25828.

13. de Arazoza H, Joanes J, Lounes R, Legeai C, Clémençon S, Pérez J, et al. The HIV/AIDS epidemic in Cuba: description and tentative explanation of its low HIV prevalence. BMC Infect Dis. 2007 Nov 9;7:130.

14. Akullian A, Bershteyn A, Klein D, Vandormael A, Bärnighausen T, Tanser F. Sexual partnership age pairings and risk of HIV acquisition in rural South Africa. AIDS. 2017 Jul 31;31(12):1755–64.

15. Napper LE, Fisher DG, Reynolds GL, Johnson ME. HIV risk behavior self-report reliability at different recall periods. AIDS Behav. 2010 Feb;14(1):152–61.

16. Turner AN, De Kock AE, Meehan-Ritter A, Blanchard K, Sebola MH, Hoosen AA, et al. Many vaginal microbicide trial participants acknowledged they had misreported sensitive sexual behavior in face-to-face interviews. J Clin Epidemiol. 2009 Jul;62(7):759–65.

17. Pettifor AE, Rees HV, Kleinschmidt I, Steffenson AE, MacPhail C, Hlongwa-Madikizela L, et al. Young people's sexual health in South Africa: HIV prevalence and sexual behaviors from a nationally representative household survey. AIDS. 2005 Sep 23;19(14):1525–34.

18. Street RA, Reddy T, Ramjee G. The generational effect on age disparate partnerships and the risk for human immunodeficiency virus and sexually transmitted infections acquisition. Int J STD AIDS. 2016 Aug;27(9):746–52.

19. Harling G, Newell ML, Tanser F, Bärnighausen T. Partner Age-Disparity and HIV Incidence Risk for Older Women in Rural South Africa. AIDS Behav. 2015 Jul;19(7):1317–26.

20. Maughan-Brown B, Venkataramani A, Kharsany ABM, Beckett S, Govender K, Lewis L, et al. Recently formed age-disparate partnerships are associated with elevated HIV-incidence among young women in South Africa. AIDS. 2020 Jan 1;34(1):149–54.

21. George G, Beckett S, Reddy T, Govender K, Cawood C, Khanyile D, et al. Determining HIV risk for Adolescent Girls and Young Women (AGYW) in relationships with "Blessers" and age-disparate partners: a cross-sectional survey in four districts in South Africa. BMC Public Health. 2022 May 14;22(1):973.

22. Topazian HM, Stoner MCD, Edwards JK, Kahn K, Gómez-Olivé FX, Twine R, et al. Variations in HIV Risk by Young Women's Age and Partner Age Disparity in Rural South Africa (HPTN 068). J Acquir Immune Defic Syndr. 2020 Apr 1;83(4):350–6.

23. Beauclair R, Helleringer S, Hens N, Delva W. Age differences between sexual partners, behavioural and demographic correlates, and HIV infection on Likoma Island, Malawi. Sci Rep. 2016 Nov 2;6:36121.

24. Liesenborgs J, Hendrickx DM, Kuylen E, Niyukuri D, Hens N, Delva W. SimpactCyan 1.0: An Open-source Simulator for Individual-Based Models in HIV Epidemiology with R and Python Interfaces. Sci Rep. 2019 Dec 17;9(1):19289.

25. Bershteyn A, Gerardin J, Bridenbecker D, Lorton CW, Bloedow J, Baker RS, et al. Implementation and applications of EMOD, an individual-based multi-disease modeling platform. Pathog Dis [Internet]. 2018 Jul 1;76(5). Available from: http://dx.doi.org/10.1093/femspd/fty059

26. Bershteyn A, Klein DJ, Eckhoff PA. Age-dependent partnering and the HIV transmission chain: a microsimulation analysis. J R Soc Interface. 2013 Nov 6;10(88):20130613.

27. Wymant C, Hall M, Ratmann O, Bonsall D, Golubchik T, de Cesare M, et al. PHYLOSCANNER: Inferring Transmission from Within- and Between-Host Pathogen

Genetic Diversity. Mol Biol Evol. 2017 Nov 23;35(3):719–33.

28. Kosakovsky Pond SL, Weaver S, Leigh Brown AJ, Wertheim JO. HIV-TRACE
(TRAnsmission Cluster Engine): a Tool for Large Scale Molecular Epidemiology of HIV-
1 and Other Rapidly Evolving Pathogens. Mol Biol Evol. 2018 Jul 1;35(7):1812–9.

29. Romero-Severson EO, Bulla I, Leitner T. Phylogenetically resolving epidemiologic
linkage. Proc Natl Acad Sci U S A. 2016 Mar 8;113(10):2690–5.

30. Leitner T, Romero-Severson E. Phylogenetic patterns recover known HIV
epidemiological relationships and reveal common transmission of multiple variants. Nat
Microbiol [Internet]. 2018 Jul 30; Available from: http://dx.doi.org/10.1038/s41564-018-
0204-9

31. Romero-Severson EO, Bulla I, Hengartner N, Bártolo I, Abecasis A, Azevedo-Pereira
JM, et al. Donor-Recipient Identification in Para- and Poly-phyletic Trees Under
Alternative HIV-1 Transmission Hypotheses Using Approximate Bayesian Computation.
Genetics [Internet]. 2017 Sep 14; Available from:
http://dx.doi.org/10.1534/genetics.117.300284

32. Zhang Y, Wymant C, Laeyendecker O, Grabowski MK, Hall M, Hudelson S, et al.
Evaluation of Phylogenetic Methods for Inferring the Direction of Human
Immunodeficiency Virus (HIV) Transmission: HIV Prevention Trials Network (HPTN)
052. Clin Infect Dis. 2021 Jan 23;72(1):30–7.

33. Xi X, Spencer SEF, Hall M, Grabowski MK, Kagaayi J, Ratmann O, et al. Inferring the
sources of HIV infection in Africa from deep-sequence data with semi-parametric
Bayesian Poisson flow models. J R Stat Soc Ser C Appl Stat. 2022 Jun;71(3):517–40.

34. Ratmann O, Kagaayi J, Hall M, Golubchick T, Kigozi G, Xi X, et al. Quantifying HIV
transmission flow between high-prevalence hotspots and surrounding communities: a
population-based study in Rakai, Uganda. Lancet HIV. 2020 Mar;7(3):e173–83.

35. Akullian A, Morrison M, Garnett GP, Mnisi Z, Lukhele N, Bridenbecker D, et al. The
effect of 90-90-90 on HIV-1 incidence and mortality in eSwatini: a mathematical
modelling study. Lancet HIV. 2020 May;7(5):e348–58.

36. Grabowski MK, Lessler J. Phylogenetic insights into age-disparate partnerships and
HIV. Lancet HIV. 2017 Jan;4(1):e8–9.

37. Kusejko K, Kadelka C, Marzel A, Battegay M, Bernasconi E, Calmy A, et al. Inferring the
age difference in HIV transmission pairs by applying phylogenetic methods on the HIV
transmission network of the Swiss HIV Cohort Study. Virus Evol. 2018 Jul;4(2):vey024.

38. Wymant C, Blanquart F, Golubchik T, Gall A, Bakker M, Bezemer D, et al. Easy and
accurate reconstruction of whole HIV genomes from short-read sequence data with
shiver. Virus Evol. 2018 Jan;4(1):vey007.

39. Monod M, Brizzi A, Galiwango RM, Ssekubugu R, Chen Y, Xi X, et al. Longitudinal
population-level HIV epidemiologic and genomic surveillance highlights growing gender
disparity of HIV transmission in Uganda. Nat Microbiol. 2024 Jan;9(1):35–54.

40. UNAIDS HIV Financial Dashboard [Internet]. [cited 2025 Mar 11]. Available from:
https://hivfinancial.unaids.org/hivfinancialdashboards.html#

41. Apollo T, Yeo YR, Mugurungi OM, Sithole N, Taramusi I, Tachiwenyika E, et al. Case

1143  study of PEPFAR funding cuts on HIV infections and deaths: reversing hard-won gains.
1144  AIDS. 2025 Nov 15;39(14):2143–6.

1145 42. Hontelez JAC, Goymann H, Berhane Y, Bhattacharjee P, Bor J, Chabata ST, et al. The
1146  impact of the PEPFAR funding freeze on HIV deaths and infections: a mathematical
1147  modelling study of seven countries in sub-Saharan Africa. EClinicalMedicine. 2025 May
1148  1;83(103233):103233.

1149 43. ten Brink D, Martin-Hughes R, Bowring AL, Wulan N, Burke K, Tidhar T, et al. Impact of
1150  an international HIV funding crisis on HIV infections and mortality in low-income and
1151  middle-income countries: a modelling study. Lancet HIV. 2025 May 1;12(5):e346–54.

1152 44. Nichols BE, Geng EH, Moakley E, Phillips AN, Imai-Eaton JW, Stover J, et al. Rapid
1153  development of an online tracker to communicate the human impact of abruptly halting
1154  PEPFAR support. J Int AIDS Soc. 2025 Mar 1;28(3):e26433.

1155 45. Hayes R, Ayles H, Beyers N, Sabapathy K, Floyd S, Shanaube K, et al. HPTN 071
1156  (PopART): rationale and design of a cluster-randomised trial of the population impact of
1157  an HIV combination prevention intervention including universal testing and treatment - a
1158  study protocol for a cluster randomised trial. Trials. 2014 Feb 13;15:57.

1159 46. Hayes RJ, Donnell D, Floyd S, Mandla N, Bwalya J, Sabapathy K, et al. Effect of
1160  Universal Testing and Treatment on HIV Incidence - HPTN 071 (PopART). N Engl J
1161  Med. 2019 Jul 18;381(3):207–18.

1162 47. Pickles M, Cori A, Probert WJM, Sauter R, Hinch R, Fidler S, et al. PopART-IBM, a
1163  highly efficient stochastic individual-based simulation model of generalised HIV
1164  epidemics developed in the context of the HPTN 071 (PopART) trial. PLoS Comput Biol.
1165  2021 Sep;17(9):e1009301.

1166 48. Probert WJM, Sauter R, Pickles M, Cori A, Bell-Mandla NF, Bwalya J, et al. Projected
1167  outcomes of universal testing and treatment in a generalised HIV epidemic in Zambia
1168  and South Africa (the HPTN 071 [PopART] trial): a modelling study. Lancet HIV. 2022
1169  Nov;9(11):e771–80.

1170 49. Golubchik T, Abeler-Dörner L, Hall M, Wymant C, Bonsall D, Macintyre-Cockett G, et al.
1171  HIV-phyloTSI: subtype-independent estimation of time since HIV-1 infection for cross-
1172  sectional measures of population incidence using deep sequence data. BMC
1173  Bioinformatics. 2025 Aug 14;26(1):212.

1174 50. Hall M, Golubchik T, Bonsall D, Abeler-Dörner L, Limbada M, Kosloff B, et al.
1175  Demographics of sources of HIV-1 transmission in Zambia: a molecular epidemiology
1176  analysis in the HPTN 071 PopART study. Lancet Microbe. 2024 Jan;5(1):e62–71.

1177 51. Bacaër N, Pretorius C, Auvert B. An age-structured model for the potential impact of
1178  generalized access to antiretrovirals on the South African HIV epidemic. Bull Math Biol.
1179  2010 Nov;72(8):2180–98.

1180 52. Smith J, Nyamukapa C, Gregson S, Lewis J, Magutshwa S, Schumacher C, et al. The
1181  distribution of sex acts and condom use within partnerships in a rural sub-Saharan
1182  African population. PLoS One. 2014 Feb 18;9(2):e88378.

1183 53. Wolock TM, Flaxman S, Risher KA, Dadirai T, Gregson S, Eaton JW. Evaluating
1184  distributional regression strategies for modelling self-reported sexual age-mixing. Elife
1185  [Internet]. 2021 Jun 24;10. Available from: http://dx.doi.org/10.7554/eLife.68318

54. Bu F, Kagaayi J, Grabowski K, Ratmann O, Xu J. Inferring HIV Transmission Patterns
from Viral Deep-Sequence Data via Latent Typed Point Processes [Internet]. arXiv
[stat.ME]. 2023. Available from: http://arxiv.org/abs/2302.11567

55. United Nations. Population Datasets [Internet]. Available from:
https://www.un.org/en/development/desa/population/publications/dataset/index.asp

56. United Nations. World Population Prospects 2015 [Internet]. United Nations; 2015.
Available from: https://population.un.org/wpp/Publications/

57. Cori A, Pickles M, van Sighem A, Gras L, Bezemer D, Reiss P, et al. CD4+ cell
dynamics in untreated HIV-1 infection: overall rates, and effects of age, viral load, sex
and calendar time. AIDS. 2015 Nov 28;29(18):2435–46.

58. Klein MB, Young J, Dunn D, Ledergerber B, Sabin C, Cozzi-Lepri A, et al. The effects of
HIV-1 subtype and ethnicity on the rate of CD4 cell count decline in patients naive to
antiretroviral therapy: a Canadian-European collaborative retrospective cohort study.
CMAJ Open. 2014 Oct 1;2(4):E318–29.

59. Easterbrook PJ, Smith M, Mullen J, O'Shea S, Chrystie I, de Ruiter A, et al. Impact of
HIV-1 viral subtype on disease progression and response to antiretroviral therapy. J Int
AIDS Soc. 2010 Feb 3;13(1):4.

60. Bonsall D, Golubchik T, de Cesare M, Limbada M, Kosloff B, MacIntyre-Cockett G, et al.
A Comprehensive Genomics Solution for HIV Surveillance and Clinical Monitoring in
Low-Income Settings. J Clin Microbiol [Internet]. 2020 Sep 22;58(10). Available from:
http://dx.doi.org/10.1128/JCM.00382-20

61. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, et al.
Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. J
Comput Biol. 2013 Oct;20(10):714–37.

62. Ratmann O, Grabowski MK, Hall M, Golubchik T, Wymant C, Abeler-Dörner L, et al.
Inferring HIV-1 transmission networks and sources of epidemic spread in Africa with
deep-sequence phylogenetic analysis. Nat Commun. 2019 Mar 29;10(1):1411.

63. Golubchik T, Abeler-Dörner L, Hall M, Wymant C, Bonsall D, Macintyre-Cockett G, et al.
HIV-phyloTSI: Subtype-independent estimation of time since HIV-1 infection for cross-
sectional measures of population incidence using deep sequence data. medRxiv. 2022
May 16;2022.05.15.22275117.

1217

1218

1219

		All participants			Females			Males		
		Both roles	Sources	Recipients	Both roles	Sources	Recipients	Both roles	Sources	Recipients
All participants		710	355	355	355	157	198	355	198	157
Community number	01	58 (0.08)	29 (0.08)	29 (0.08)	27 (0.08)	14 (0.09)	13 (0.07)	31 (0.09)	15 (0.08)	16 (0.1)
	02	62 (0.09)	31 (0.09)	31 (0.09)	32 (0.09)	19 (0.12)	13 (0.07)	30 (0.08)	12 (0.06)	18 (0.11)
	03	107 (0.15)	51 (0.14)	56 (0.16)	50 (0.14)	18 (0.11)	32 (0.16)	57 (0.16)	33 (0.17)	24 (0.15)
	04	61 (0.09)	31 (0.09)	30 (0.08)	28 (0.08)	10 (0.06)	18 (0.09)	33 (0.09)	21 (0.11)	12 (0.08)
	05	108 (0.15)	53 (0.15)	55 (0.15)	56 (0.16)	24 (0.15)	32 (0.16)	52 (0.15)	29 (0.15)	23 (0.15)
	06	113 (0.16)	57 (0.16)	56 (0.16)	61 (0.17)	26 (0.17)	35 (0.18)	52 (0.15)	31 (0.16)	21 (0.13)
	10	51 (0.07)	24 (0.07)	27 (0.08)	26 (0.07)	11 (0.07)	15 (0.08)	25 (0.07)	13 (0.07)	12 (0.08)
	11	46 (0.06)	24 (0.07)	22 (0.06)	23 (0.06)	10 (0.06)	13 (0.07)	23 (0.06)	14 (0.07)	9 (0.06)
	12	104 (0.15)	55 (0.15)	49 (0.14)	52 (0.15)	25 (0.16)	27 (0.14)	52 (0.15)	30 (0.15)	22 (0.14)
Trial arm	A	220 (0.31)	106 (0.3)	114 (0.32)	108 (0.3)	48 (0.31)	60 (0.3)	112 (0.32)	58 (0.29)	54 (0.34)
	B	217 (0.31)	110 (0.31)	107 (0.3)	111 (0.31)	50 (0.32)	61 (0.31)	106 (0.3)	60 (0.3)	46 (0.29)
	C	273 (0.38)	139 (0.39)	134 (0.38)	136 (0.38)	59 (0.38)	77 (0.39)	137 (0.39)	80 (0.4)	57 (0.36)
Enrolment method	Health care facilities	644 (0.91)	318 (0.9)	326 (0.92)	308 (0.87)	132 (0.84)	176 (0.89)	336 (0.95)	186 (0.94)	150 (0.96)
	Population cohort	66 (0.09)	37 (0.1)	29 (0.08)	47 (0.13)	25 (0.16)	22 (0.11)	19 (0.05)	12 (0.06)	7 (0.04)
Sampling year	2014	26 (0.04)	25 (0.07)	1 (0)	22 (0.06)	21 (0.13)	1 (0.01)	4 (0.01)	4 (0.02)	0 (0)
	2015	10 (0.01)	3 (0.01)	7 (0.02)	7 (0.02)	0 (0)	7 (0.04)	3 (0.01)	3 (0.02)	0 (0)
	2016	223 (0.31)	106 (0.3)	117 (0.33)	111 (0.31)	46 (0.29)	65 (0.33)	112 (0.32)	60 (0.3)	52 (0.33)
	2017	323 (0.45)	160 (0.45)	163 (0.46)	159 (0.45)	73 (0.46)	86 (0.43)	164 (0.46)	87 (0.44)	77 (0.49)
	2018	128 (0.18)	61 (0.17)	67 (0.19)	56 (0.16)	17 (0.11)	39 (0.2)	72 (0.2)	44 (0.22)	28 (0.18)
Marital status at sampling	Never married	89 (0.13)	44 (0.12)	45 (0.13)	44 (0.12)	23 (0.15)	21 (0.11)	45 (0.13)	21 (0.11)	24 (0.15)
	Married/living as married	496 (0.7)	256 (0.72)	240 (0.68)	239 (0.67)	104 (0.66)	135 (0.68)	257 (0.72)	152 (0.77)	105 (0.67)
	Divorced/separated	101 (0.14)	47 (0.13)	54 (0.15)	54 (0.15)	25 (0.16)	29 (0.15)	47 (0.13)	22 (0.11)	25 (0.16)
	Widowed	20 (0.03)	6 (0.02)	14 (0.04)	15 (0.04)	3 (0.02)	12 (0.06)	5 (0.01)	3 (0.02)	2 (0.01)
	Unknown	4 (0.01)	2 (0.01)	2 (0.01)	3 (0.01)	2 (0.01)	1 (0.01)	1 (0)	0 (0)	1 (0.01)
Age at estimated time of transmission (years, mean (SD))		30.1 (8.6)	30.8 (8.4)	29.5 (8.9)	27.5 (8.4)	27.1 (7.3)	27.8 (9.2)	32.7 (8.1)	33.7 (8)	31.5 (8)
Relative age of partner (years, mean (SD))		0 (8.9)	1.3 (8.8)	-1.3 (8.8)	-5.2 (7.2)	-4.4 (6.6)	-5.9 (7.5)	5.2 (7.2)	5.9 (7.5)	4.4 (6.6)

1220

1221

1222 **Table 1:** Demographics of the individuals in the 355 probable transmission pairs. Age is

1223 calculated at the estimated time at which the source infected the recipient. Numbers are

1224 given as count and proportion (column-wise) for discrete variables and mean and standard

1225 deviation for continuous variables.

1226

Variable	Sexes	Phylogenetics		IBM		Posterior predictive p-value (phylo vs IBM)
		Mean	95% CI	Posterior mean of means	95% HDI	
Source age	Males	33.7	(32.6 - 34.8)	32.8	(31.9 - 33.6)	0.21
	Females	27.1	(25.9 - 28.2))	28.3	(27.4 - 29.2)	0.11
Recipient age	Males	31.5	(30.3 - 32.8)	31.1	(30 - 32.1)	0.61
	Females	27.8	(26.5 - 29.1)	27.5	(26.7 - 28.4)	0.71
Age gap (source-recipient)	Male source, female recipient	5.87	(4.82 - 6.92)	5.27	(4.98 - 5.57)	0.27
	Female source, male recipient	-4.44	(-5.49 - -3.4)	-2.83	(-3.17 - -2.47)	0.002

1227

1228 **Table 2:** Summary of ages, and age gaps, in transmission pairs identified using both

1229 methods. The posterior predictive *p*-value is for the null hypothesis of the mean age or age

1230 gap from the phylogenetics coming from the posterior distribution of means from the IBM.

1231

1232

Variable	Sexes	Source diagnosed		Source undiagnosed		Difference of means (diagnosed-undiagnosed)	
		Posterior mean of means	95% HDI	Posterior mean of means	95% HDI	Posterior mean	95% HDI
Source age	Males	34.8	(33.8 - 35.6)	32.2	(31.4 - 33.1)	2.54	(2.14 - 2.91)
	Females	30.1	(29.2 - 31.1)	27.5	(26.6 - 28.4)	2.57	(2.23 - 2.98)
Recipient age	Males	32.9	(31.8 - 34)	30.3	(29.3 - 31.4)	2.53	(2.07 - 2.96)
	Females	28.7	(27.7 - 29.5)	27.2	(26.4 - 28.1)	1.45	(1.06 - 2.96)
Age gap (source-recipient)	Male source, female recipient	6.1	(5.7 - 6.49)	5.01	(4.73 - 5.28)	1.09	(0.9 - 1.27)
	Female source, male recipient	-2.8	(-3.21 - -2.43)	-2.84	(-3.19 - -2.5)	0.044	(-0.12 - 0.2)

1233

1234   **Table 3:** Summary of ages, and age gaps, in transmission pairs identified using both

1235   methods, stratified by diagnostic status of the source, and the posterior mean and 95% HDI

1236   values of the difference between the mean values (diagnosed-undiagnosed)

1237

1238

Distance threshold (substitutions per site, normalised to represent average distances on the *gag* and *pol* genes)	Number of pairs identified
0.01	257
0.015	315
0.02	355
0.025	386
0.03	417
0.035	441
0.04	459

1239

1240   **Appendix 2-Table 1** Number of phylogenetic transmission pairs identified using different

1241   patristic distance thresholds.

1242

1243

1244

1245

1246

1247

Figure 1: Phylogenetic analysis and the IBM model show similar age

distributions for sources and recipients. Histograms of the proportions of

transmissions from and to different age groups in the IBM results (top) and the

phylogenetics (bottom). Values from the IBM are the posterior mean proportion of

transmissions with the 95% highest density interval displayed. As every transmission

has a source and a recipient, proportions sum to 1 across the two source histograms

(left, light colours) and also to 1 across the two recipient histograms (right, dark

colours).

1256

1257

1258

Figure 2: Male partners are on average older in both male-to-female and

female-to-male pairs. Distributions of age gaps within transmission pairs from the

phylogenetics (yellow, histogram) and the IBM (blue, kernel density estimate). For

the IBM, the 95% highest posterior density intervals are given by the shaded area.

Female-to-male transmission pairs are displayed on the left and male-to-female on

the right. The dashed line indicates equal age.

1265

Figure 3: Transmissions are most common in young couples. Age-based who-acquires-infection-from-whom (WAIFW) matrix for modelled transmissions occurring in 2016. Values are the mean, over the entire year, of the per-year transmission rates from infected individuals in one age group to individuals in another (given by the number of transmissions divided by the size of the source population).

Figure 4: Transmissions to a younger cohort take place predominantly in male-to-female transmissions. Alluvial plots for the transmission history, derived from the IBM, of a cohort of males (panel a) and females (panel b) infected with HIV in 2016. That cohort is G0, shown on the right; to the left of it are the three previous generations of transmission (indicated as G1 to G3) that gave rise to the infections of G0. Male generations are blue and female red. The partitioning within each vertical bar indicates its age distribution. The age distributions of G1 to G3 are shown at both the time of their own infection (left) and the time at which they infected an individual in a subsequent generation (right). Movement to an older age group (by ageing or transmission) is shown by a green alluvium, to a younger group by an orange alluvium, and to the same age group by a grey alluvium. If an individual was responsible for two or more infections in the subsequent generation, then they still contribute a total of 1 individual to their "age at onwards transmission" bar, which is split proportionally between age bands depending on the number of infections they

were responsible for while they were a member of each band. (For example, if

individual A infected two individuals while they (A) were aged 20–24 and one

individual while they were aged 25–29, A would contribute ⅔ of an individual to the

20–24 bar and ⅓ to the 25–29 bar.) Note that the members of G0 are all members of

a particular cohort who were infected with HIV, but all other generations are

composed solely of people who transmit HIV, a somewhat different group.

1297

1298

Figure 5: The age gap increases with recipient age in female-to-male pairs and

decreases in male-to-female pairs. Relationship between recipient age and source

age (a, c) and age gap (b, d) from the IBM (a, b) and the phylogenetics (c, d). The

dashed line represents equal ages. The IBM results are from the "best-fitting"

simulation (that with the minimum ABC distance) and represent the mean value

(solid black line) and quantiles (orange shaded areas) of the y-axis variable in 1-year

increments. The individual values from the phylogenetics pairs are plotted as green

points, with a regression line fitted using quadratic polynomial regression and the

95% confidence region given by the pale green area.

1308

1309

1310 **Figure 6:** Proportion of sources that are undiagnosed, by age and sex. The thick line

1311 is the posterior mean and the shaded area the 95% HDI.

1312

1313

Figure 7: Little change in age gaps over the period of ART rollout. Time trends in age of sources (a), age gaps (b) and proportion of the population on ART (c) over time. Estimates are the posterior mean (black line), and 95% HDI (shaded), of the mean values.

1319

1320

1321 **A**

1322

1323

1324 **B**

1325

Figure 8: The impact of an intervention focussing on younger people. Projected

time trends in HIV incidence (A) and population proportion of people with

unsuppressed HIV (B), 2018–2041, in the Zambian control communities of the

PopART trial, under various scenarios of future PopART intervention rollout.

Posterior mean estimates are calculated in two year time periods for the whole

population (top), females (middle) and males (bottom). The four scenarios are, firstly,

no CHiPS deployment at all (red), provision of the full PopART intervention to the

whole population (purple), PopART intervention targeting only those aged under 35

(grey), and PopART intervention targeting only men aged under 35 (green). Vertical

bars indicate the 95% highest posterior density interval.

1336

Figure 9: The impact of ART interruption. A: Posterior mean new infections by year, by sex and age, and B: the age distribution of new transmissions by year, in the circumstances of a complete cessation of ART provision during 2025, returning in 2026. Lines chart the posterior mean proportion of new infections for sources and recipients of each integer age (top and middle), and age gap rounded to the nearest integer (bottom). In both cases, lines are coloured by year.

Appendix 1-Figure 1: Empirical distributions of mean (a) and variance (b) of ages of recipients at time of transmission from pairs where "recipients" had randomly chosen "sources" (histograms) compared to the true value from the phylogenetics data (green line).

1349

1350   **Appendix 1-Figure 2:** Empirical distributions of mean estimated time from infection to

1351   sampling of recipients from randomised transmission pairs (histograms) compared to the

1352   true value from the phylogenetics data (green line).

1353

1354

1355

1356 **Appendix 1-Figure 3:** Empirical distributions of mean (a) and variance (b) of ages of

1357 sources at time of transmission from randomised transmission pairs (histograms) compared

1358 to the true value from the phylogenetics data (green line).

1359

1360

Appendix 1-Figure 4: Empirical distributions of mean estimated time from infection to sampling of sources from randomised transmission pairs (histograms) compared to the true value from the phylogenetics data (green line).

1367

Appendix 1-Figure 5: Empirical distributions of mean (a) and variance (b) in transmission age gap from randomised transmission pairs (histograms) compared to the true value from the phylogenetics data (green line).

1368

1369

1370

1371

1372

Appendix 2-Figure 1: Distributions of the estimated ages of sources at the time of

transmission (top), estimated ages of recipients at the time of transmission (middle) and age

gap amongst probable transmission pairs identified using *phyloscanner* when the read

downsampling procedure is varied. Downsampling to a maximum of *n* reads means that a

random sampling of *n* reads is taken for each individual in each genomic window. For

windows in which *n* reads do not exist, the two options are "blacklist" - to exclude that host

from that window entirely - and "keep" - to use all the reads that are available. The boxplots

display the median, 25% and 75% quantiles, and range.

1382

1383

1384

1385 **Appendix 2-Figure 2:** Distributions of the estimated ages of sources at the time of

1386 transmission (top), estimated ages of recipients at the time of transmission (middle) and age

1387 gap amongst probable transmission pairs identified using *phyloscanner* with different

1388 genomic window widths. The boxplots display the median, 75% quantiles, and range.

1389

1390

1391

Window thresholds for identifying pairs

Appendix 2-figure 3: Distributions of the estimated ages of sources at the time of transmission (top), estimated ages of recipients at the time of transmission (middle) and age gap amongst probable transmission pairs identified using *phyloscanner*. Different window-related thresholds for identifying linkage and direction are indicated on the x axis in the format X/Y, where Y is the minimum fraction of genomic windows supporting two samples being from a transmission pair, and X is the minimum fraction of genomic windows supporting transmission having an identifiable direction (conditional on being a transmission

1401   pair). The boxplots display the median, 75% quantiles, and range.

1402

1403

1404

1405   **Appendix 2-Figure 4** Distributions of the estimated ages of sources at the time of

1406   transmission (top), estimated ages of recipients at the time of transmission (middle) and age

1407   gap amongst probable transmission pairs identified using *phyloscanner* with patristic

1408   distance thresholds used to identify transmission pairs in an individual window. The boxplots

1409   display the median, 75% quantiles, and range.

1410

1411