Moderated Chain Mediation of Depression and Social Participation Between Stroke and Mortality by Socioeconomic Status

Li Zhu[1]*, Jinhua Qian[1]*, Wenlu Shi[1], Zihan Geng[1], Siqi Yang[1], Tianle Wang[1#], Lei Wang[2#]

1 Department of Radiology, Affiliated Hospital 2 of Nantong University, Jiangsu, 226014, P.R. China

2 Institute for translational neuroscience of Affiliated Hospital 2 of Nantong University ; Center for neural developmental and degenerative research of Nantong University, Jiangsu, 226014, P.R. China

* These authors contributed equally to the work.

#Corresponding author: lei.wang.psy@gmail.com; & wangtianle9192@163.com

Abstract

Background: Stroke significantly increases the risk of depression and reduced social

participation, particularly among middle-aged and older adults. Socioeconomic status

(SES) may further shape these pathways. This study examined the mediating roles of

depression and social participation, and the moderating role of SES, in the relationship

between stroke and 10-year all-cause mortality in Chinese adults, while also exploring

longitudinal patterns over time.

Methods: We analyzed data from 7,101 participants in the China Health and Retirement

Longitudinal Study (CHARLS) across five waves (2011–2020). Depression was assessed

via the CES-D scale, and social participation was measured based on activity frequency

scores. SES was constructed from education, occupation, expenditure, and insurance.

Moderated chain mediation models and generalized linear mixed models were used to

evaluate pathways and temporal trends.

Results: Stroke was significantly associated with higher all-cause mortality. Depression

partially mediated this association ($\beta = 0.059$, 95% CI [0.020, 0.023]). Although social

participation alone was not a significant mediator, a sequential pathway involving

depression and social participation was confirmed ($\beta = 0.006$, 95% CI [0.001, 0.012]).

SES moderated both the stroke–depression and depression–social participation pathways.

Notably, social participation demonstrated a protective role against post-stroke mortality

when embedded within depression-related pathways—especially among low-SES

individuals. Longitudinal analyses showed that the effects of stroke and depression on

mortality weakened over time, while high levels of social participation consistently offered protective benefits.

Conclusions: Depression and social participation mediate the stroke–mortality relationship, and SES plays a critical moderating role. These findings suggest that strengthening social engagement may serve as an effective strategy to mitigate depression-related mortality among stroke survivors, particularly in low-SES populations. Tailored, SES-sensitive interventions are essential to improving survival outcomes.

Key words: Stroke, Depression, Social Participation, Socioeconomic Status, All-Cause Mortality

Non-standard Abbreviations and Acronyms.

SES	Socioeconomic status
CHARLS	China Health and Retirement Longitudinal Survey
CI	Confidence Interval
CESD-10	10-item Center for Epidemiologic Studies Depression Scale
IQRs	interquartile ranges
SD	Standard deviation

1 Introduction

Stroke remains the second leading cause of death and disability worldwide, marked by a

high recurrence rate and a poor long-term prognosis. Beyond its immediate clinical

impact, stroke imposes a substantial and lasting burden on patients, caregivers, and

healthcare systems alike[1]. Among stroke survivors, long-term outcomes are often

discouraging. For example, a nationwide Danish cohort study found that over 56% of

stroke survivors died within 10 years of their initial stroke[2]. In China, the fact that

approximately 40% of the population is now aged 45 and above underscores the urgency

of this issue amid the rapid aging population[3]. This demographic shift has coincided with

a marked rise in stroke incidence and mortality, particularly among middle-aged and

older adults. Hence, identifying risk factors that contribute to post-stroke mortality—

especially all-cause mortality—has become a critical priority for public health and

clinical intervention.

A key yet under-recognized consequence of stroke is the heightened prevalence of

emotional disorders, most notably depression, among survivors. Compared to general

population, studies demonstrated that middle-aged and older adults are disproportionately

affected with the incidence of post-stroke emotional disorders in these groups is

estimated at 47.9% and 50.6%, respectively[4-6]. These findings underscore the increased

susceptibility to depression in older stroke patients throughout the rehabilitation process.

Moreover, longitudinal researches have shown that depressive symptoms are a significant

risk factor for all-cause mortality[7] and are associated with increased five-year mortality

among elderly stroke survivors[8]. However, it is still unknown whether and to what extend

post-stroke depression contributes to long-term all-cause mortality in stroke survivors.

Therefore, we hypothesize that depressive symptoms mediate the relationship between

stroke and all-cause mortality (**H1**).

In addition to mental health concerns, stroke-related impairments frequently results in a

decline in social participation, particularly among older adults who experience physical

or cognitive limitations [9, 10]. However, accumulating evidence suggests that active social

engagement serves a protective function by fostering supportive social networks and

alleviating psychological distress, thereby mitigating health risks[11, 12]. Data from China

Health and Retirement Longitudinal Study (CHARLS) revealed that individuals with low

levels of social participation had a shorter life expectancy (12.98 years) compared to

those with high levels of social participation (15.43 years)[13]. Therefore, in the present

study aims to examine whether social participation may also function as a dediator

between stroke and all-cause mortality (**H2**).

While depression and social participation are often treated as independent predictors of

post-stroke outcomes, their interrelationship may have a synergistic effect on mortality

risk. Post-stroke depression is frequently accompanied by social withdrawal, which may,

in turn, intensify depressive symptoms, forming a potentially self-perpetuating cycle[14, 15].

Despite these reciprocal dynamics, few studies have explored their combined impact on

mortality within an integrated framework. In this study, we therefore posit that depression

and social participation operate as a sequential mediation pathway linking stroke to all-

cause mortality (**H3**).

Beyond individual psychological factors, socioeconomic status (SES) may also further

shape these relationships. SES encompasses key dimensions such as income, education,

and occupational status, and is widely recognized as a fundamental driver of health

disparities[16]. Middle-aged and older individuals with low SES face compounded

vulnerabilities, limited access to healthcare, constrained social support, and elevated

economic stress, factors that collectively heighten the risk of post-stroke depression,

social isolation, and premature mortality[17, 18]. Prior studies have linked low SES to

greater depressive symptomatology and lower levels of social participation[11, 18].

Moreover, the relationship between SES and stroke outcomes is consistently evident

across both development ad developing countries[19]. However, it remains unclear whether

SES not only directly predict stroke-related mortality, but also moderate the effects of

depression and social participation on mortality. Accordingly, we hypothesize that SES

moderates the relationship between stroke and depression (**H4**), the relationship between

depression and social participation (**H5**), and the chain-mediated relationship linking

stroke, depression, social participation, and all-cause mortality (**H6**).

Although the independent effects of depression, social participation, and SES on stroke

outcomes are well established, their interactive mechanisms remain poorly understood.

Few studies have employed an integrated longitudinal model to examine how these

factors jointly influence mortality risk over time. Given the frequent co-occurrence of

depression and social disengagement[20], and the pervasive influence of socioeconomic

disadvantage, this study aims to fill a critical gap by examining the chain mediation

effects of depression and social participation and the moderating role of SES in the

association between stroke and all-cause mortality among middle-aged and older adults.

In summary, this study proposes the following hypotheses:

H1: Stroke impacts all-cause mortality through the mediation of depression ($X \rightarrow M1 \rightarrow$

Y).

H2: Stroke impacts all-cause mortality through the mediation of social participation (X

$\rightarrow M2 \rightarrow Y$).

H3: Stroke impacts all-cause mortality through a chain mediation of depression and

social participation ($X \rightarrow M1 \rightarrow M2 \rightarrow Y$).

H4: SES moderates the relationship between stroke and depression ($M1(W) \rightarrow M2$).

H5: SES moderates the relationship between depression and social participation ($M2(W)$

$\rightarrow M2$).

H6: SES moderates the chain-mediated relationship between stroke, depression, social

participation, and all-cause mortality ($X(W) \rightarrow M1 \rightarrow M2 \rightarrow Y$ or $X \rightarrow M1(W) \rightarrow M2 \rightarrow$

Y).

Methods

Population: This study utilized data from the China Health and Retirement Longitudinal

Study (CHARLS), a nationally representative longitudinal survey designed to assess the

health, economic status, social support, and psychosocial conditions of Chinese adults

aged 45 years and above[21]. CHARLS was conducted by the National Institute of

Development at Peking University and received ethical approval from its Institutional

Review Board (IRB00001052-11015). A multistage and stratified probability sampling

method was employed to ensure comprehensive coverage, encompassing 450 villages

and communities across 180 counties throughout 28 provinces.

Data were collected across five waves: 2011 (baseline), 2013, 2015, 2018, and 2020.

From the initial 17,708 participants, individuals under the age of 45 (n = 508) and those

with incomplete baseline or follow-up data were excluded. The final analytic sample

consisted of 7,101 participants. Figure 1 depicted the flowchart depicting the participant

selection process.

Depression. Depression was assessed using the 10-item Center for Epidemiologic Studies

Depression Scale (CESD-10), a validated instrument for older Chinese adults[22]. Scores

range from 0 to 30, with higher values indicating more severe depressive symptoms. The

CESD-10 has demonstrated good internal reliability in previous research[23]. Item-level

content is provided in Table S1.

Social Participation. Social participation was measured by participant engagement in

various activities during the past month. Frequency was self-reported on a 4-point scale

including 0 (never), 1 (occasionally), 2 (almost weekly) and 3 (almost daily)[24]. Total

scores ranged from 0 to 30, with higher scores referring greater social activity. Subtypes

included recreational activities (items I, II, IV, V, VIII), volunteering (items III, VI, VII),

and online participation (item IX), as shown in Table S2.

Socioeconomic Status. SES was operationalized as a composite index comprising four

dimensions: household expenditure per capita, education level, occupation type, and

health insurance coverage. Each dimension was scored from 1 (low) to 3 (high), resulting

in a total SES score ranging from 4 to 12, with higher scores indicating higher SES.

Expenditure levels were categorized into tertiles according to the 33rd and 66th

percentiles of the distribution:low ($\leq$ ¥4,020), medium (¥4,020−7,541), and high ($\geq$

¥7,542). Education was classified into below compulsory education, high school, and

university or higher[25]. Occupation was divided into unemployed, agricultural, and non-

agricultural. Health insurance coverage was categorized as no insurance, public only, or

both public and private coverage[26].

All-Cause Mortality. All-cause mortality was defined as death from any cause, recorded

over the 10-year follow-up period. Mortality data were gathered through exit interviews

conducted with proxies of deceased participants as part of the CHARLS tracking

protocol.

Covariates. The following variables were included as controls: gender (male or female),

age group (middle-aged: 45–60; older: >60), smoking history (yes or no), marital status

(married or other), and a set of chronic diseases including hypertension, diabetes,

dyslipidemia, gastrointestinal disorders, chronic lung disease, cardiovascular disease,

kidney disease, and psychiatric conditions.

Statistical Analysis. All analyses were performed using IBM SPSS version 27.0. All

continuous variables exhibited skewed distributions and were therefore summarized as

medians and interquartile ranges (IQRs). Continuous variables were summarized as Mean

± SD, while categorical variables were described as frequencies and percentages.

Between-group comparisons used independent-sample t-tests for continuous variables

and chi-square or Fisher's exact tests for categorical variables. A significance threshold

of $p < 0.05$ was applied throughout.

Variables showing significant differences between groups were included in multivariate

logistic regression models to identify predictors of all-cause mortality. Pearson

correlation coefficients were computed to assess relationships between stroke,

depression, social participation, SES, and mortality. Mediation and moderated mediation

models were tested using the PROCESS macro for SPSS, employing bootstrapping

(5,000 resamples) to estimate indirect effects with 95% confidence intervals. Generalized

linear mixed models were used to explore the longitudinal effects of stroke, depression,

social participation, and SES on mortality, accounting for repeated measures and within-

subject variation over time.

Result

Descriptive Statistics and Logistic Regression

Table 1 summarize the baseline characteristics of the study sample (N = 7,101) in 2011;

detailed distributions of social participation and SES subcategories are shown in Table

S3. Over the 10-year follow-up period, 1,125 participants died, yielding a mortality rate

of 15.84%. Significant differences were observed between the survival and mortality

groups in terms of age, gender, smoking status, hypertension, diabetes, lung disease,

kidney disease, heart disease, stroke history, depressive symptoms, social participation,

and socioeconomic status (SES) ($p < 0.05$ for all).

Multivariate logistic regression analyses (see Table 2) revealed that stroke ($p = 0.009$),

depression ($p = 0.003$), SES ($p < 0.001$), age ($p < 0.001$), gender ($p < 0.001$), smoking

history ($p = 0.009$), marital status ($p < 0.001$), hypertension ($p < 0.001$), diabetes ($p < 0.001$), and lung disease ($p < 0.001$) were significantly associated with all-cause mortality.

0.001), and lung disease ($p < 0.001$) were significantly associated with all-cause mortality.

However, social participation ($p = 0.223$), kidney disease ($p = 0.141$), and heart disease

($p = 0.098$) were not significant predictors in the final model.

Correlation Analysis

As illustrated in Figure 2A, stroke and depression were positively associated with all-

cause mortality, while SES demonstrated a negative correlation. No significant

relationship was found between stroke and social participation. These correlations

provided the empirical basis for constructing a moderated chain mediation model. Figure

2B further explores the correlations between different types of social participation.

Chain Mediation Model

Significant predictors identified in the logistic regression were included in the mediation

model to test the hypothesized indirect pathways. The path diagram is shown in Figure

3A, and detailed coefficients are reported in Table S4. Stroke was positively associated

with depression ($\beta = 0.422$, $p < 0.001$), and depression was negatively associated with

social participation ($\beta = -0.146$, $p < 0.001$). Depression was positively linked to all-cause

mortality ($\beta = 0.139$, $p < 0.001$), while social participation was inversely associated with

mortality ($\beta = -0.094$, $p < 0.05$). These findings support the presence of a chain

mediation mechanism.

Bias-corrected bootstrap analyses (5,000 samples) were used to estimate indirect effects.

As shown in Table S5, the indirect effect of stroke on mortality through depression

(Pathway Ind1) was significant ($\beta = 0.059$, 95% CI [0.023 ~ 0.102]), supporting

Hypothesis 1 (H1). The indirect effect through social participation alone (Pathway Ind2)

was not significant ($\beta = -0.006$, 95% CI [−0.027, 0.009]), as the confidence interval

included zero, thereby refuting Hypothesis 2 (H2). However, the indirect effect through

the sequential pathway of depression and social participation (Pathway Ind3) was

significant, albeit small ($\beta = 0.006$, 95% CI [0.001, 0.012]), providing support for

Hypothesis 3 (H3).

Moderated Chain Mediation Model

To examine the moderating role of SES, PROCESS macro models 83 and 91 were

applied, focusing on the chain pathway Stroke → Depression → Social Participation →

Mortality. The interaction effects between SES and stroke ($\beta = -0.243$, $p < 0.01$), and

between SES and depression ($\beta = -0.041$, $p < 0.001$), were both significant, supporting

Hypotheses 4 and 5 (H4 and H5).

Simple slope analysis, illustrated in Figure 4, further clarifies the nature of these

interactions. The moderated mediation index of SES (Table S6) on the stroke →

depression → mortality pathway was −0.003 (95% CI [−0.007, −0.001]), and the index

for the depression → social participation → mortality pathway was 0.002 (95% CI

[0.0003, 0.004]). Since both confidence intervals excluded zero, these findings confirm a

significant moderated chain mediation effect, thus supporting Hypothesis 6 (H6). The full

moderated chain mediation model is presented in Figure 3B.

Longitudinal Analysis

To assess dynamic changes in risk over time, generalized linear mixed models were used

to evaluate the longitudinal relationships among stroke, depression, social participation,

SES, and all-cause mortality. The results, detailed in Table S7, indicate that depression

and SES remained consistent predictors of mortality across waves. Figure 5 presents

subgroup analyses that demonstrate varying effects of social participation on mortality

over time, accounting for temporal fluctuations and individual-level variability.

Discussion

This study investigated the complex interplay between stroke, depression, social

participation, and socioeconomic status (SES) in predicting all-cause mortality among

middle-aged and older adults in China. Drawing on a nationally representative

longitudinal dataset and applying moderated chain mediation models, the findings reveal

both direct and indirect pathways through which stroke impacts mortality risk. Notably,

stroke was associated with increased all-cause mortality via partial mediation by

depression (Ind1) and a sequential chain involving both depression and social

participation (Ind3). Although social participation alone did not significantly mediate the

stroke–mortality relationship, its protective effect was evident when coupled with

depression, highlighting the importance of addressing both emotional and social well-

being in post-stroke care.

The observed all-cause mortality rate of 15.84% is consistent with estimates from

previous studies, such as the Chinese Longitudinal Healthy Longevity Survey[27].

Individuals in the mortality group exhibited higher depression scores and stroke

prevalence, alongside lower SES and reduced social participation. Multivariate regression

showed no significant direct association between social participation and mortality,

suggesting that its influence may emerge through interactive or conditional pathways,

especially in the presence of depression.

Approximately half of middle-aged and older stroke survivors experience depressive

symptoms and social disengagement[4, 28]. The vascular depression hypothesis provides a

possible explanation, attributing this vulnerability to cerebrovascular damage in brain

regions responsible for emotion and cognition[6, 29]. Stroke-related neurological

impairment not only leads to physical disability but also increases psychological risk—

particularly depression—which, in turn, contributes to elevated mortality. Our findings

align with previous research demonstrating that depression mediates cardiovascular-

related mortality with similar effect sizes[30], reinforcing that depression significantly

mediates the relationship between stroke and mortality and highlighting its critical role as

a clinical target in post-stroke rehabilitation[2, 31, 32].

Interestingly, while social participation did not serve as a standalone mediator, its

protective effect became salient within a chain mediation model. Stroke-induced

depression leads to lower social engagement, which subsequently increases mortality risk

[28]. This finding aligns with the biopsychosocial model[33], where emotional and behavioral

states are closely interlinked. Physical limitations (e.g., aphasia, hemiplegia) can restrict

social involvement, and emotional disturbances such as depression further exacerbate

withdrawal[34]. Conversely, low levels of social participation can reinforce depressive

symptoms, creating a feedback loop. Prior studies, including animal research, also

demonstrate the exacerbating effects of social isolation on depressive symptoms[35-37]. In

humans, behavioral activation therapy promoting social interaction has shown efficacy in

alleviating depression, reinforcing the significance of social engagement in psychological

recovery[38-40].

Our findings also highlight the moderating role of SES in this pathway. SES negatively

moderated the stroke–depression link and positively moderated the depression–social

participation relationship. Individuals with lower SES were more vulnerable to

depression following stroke, due to factors such as reduced access to healthcare, lower

health literacy, and economic stress[4, 41, 42]. In contrast, higher SES was associated with

greater resilience and better medical access[11]. However, when depression occurred

among high-SES individuals, its impact on social participation was amplified. This may

stem from identity concerns, as high-SES individuals often engage in more formal social

activities and may experience stronger social stigma or self-imposed withdrawal when

their functioning declines[43-45].

Moreover, the chain mediation pathway itself was significantly moderated by SES. Low

SES intensified the negative effects of depression and diminished social participation on

mortality, whereas high SES buffered these effects[4, 19, 46]. This dual role of SES—as both

a risk and protective factor—underscores the need for tailored intervention strategies that

consider economic and social disparities[47]. High-SES individuals may benefit more from

psychological support addressing self-identity and social reintegration, while low-SES

populations may require broader access to mental health services and community-based

resources[48, 49].

Longitudinal analysis further showed that the associations among stroke, depression,

social participation, and mortality are dynamic rather than static. The effects of stroke

and depression on mortality declined over time, suggesting that early-stage intervention is

especially critical[50, 51]. Although social participation did not display a strong fixed effect,

its interaction with time revealed important variations. Individuals with higher social

participation maintained lower mortality risk across time, emphasizing the cumulative

benefits of social engagement[18].

Several limitations should be acknowledged. First, inconsistencies in the measurement of

social participation across survey waves—particularly in 2020—may introduce bias in

cross-time comparisons, despite the applied adjustments. Second, while this study

focused on all-cause mortality, it did not explore cause-specific mechanisms or account

for biological pathways linking stroke, depression, and social withdrawal to mortality

outcomes. Third, due to data limitations, longitudinal mediation effects were not tested,

and causality cannot be definitively established from the current model structure. Finally,

the findings are subject to limitations in generalizability due to the observational nature

of the study and reliance on internal validation. Future research should incorporate

neurobiological measures, expand to diverse populations, and apply prospective designs

to validate and deepen understanding of these complex interrelationships.

In conclusion, this study demonstrates that stroke increases the risk of all-cause mortality

among middle-aged and older adults, with depression and social participation acting as

important mediators. SES not only influences these relationships directly but also

moderates their pathways, reinforcing the importance of a multifaceted intervention

approach. Social and emotional health should be considered alongside physical

rehabilitation in stroke recovery programs. Efforts to promote early screening for

depression and encourage meaningful social engagement can play a vital role in

improving survival outcomes. Tailored interventions based on SES are crucial for

ensuring equitable access to care and optimizing the long-term well-being of stroke

survivors.

Acknowledgments

Conceptualization, T.W., L.W.; methodology, J.Q., L.W.; investigation, J.Q.,W.S, Z.G,

S,Y.; formal analysis, L.Z., J.Q.; visualization, L.Z., J.Q.; writing – original draft, L.Z.,

J.Q.; writing – review & editing, L.Z., J.Q., T.W., L.W.; supervision, T.W., L.W.;

funding acquisition, L.Z., L.W..

Sources of Funding

This work was supported by the Science and Technology Project of Nantong

(MS2023068, MS12021101, MS12020041), Nantong Municipal Health Commission

(MS2024026), and Start Funding from Affiliated Hospital 2 of Nantong University

(YJRCJJ007).

Disclosures

None.

Supplemental Material

Supplemental Tables 1 - 7

References

1. Hu S, Gao M, He Y, Xie X. Comparison of the performance of different nutritional indicators for predicting poststroke depression in older adults with ischemic stroke. *Int J Nurs Sci*. 2024;11:349-356.

2. Eddelien HS, Grøntved S, Hedegaard JN, Thomsen T, Kruuse C, Johnsen SP. Quality of early stroke care and long-term mortality in patients with acute stroke: A nationwide follow-up study. *Eur Stroke J*. 2024:23969873241249580.

3. CEIC. China population sample survey by age and sex. 2023. https://www.ceicdata.com/en/china/population-sample-survey-by-age-and-sex.

4. Cai Q, Qian M, Chen M. Association between socioeconomic status and post-stroke depression in middle-aged and older adults: results from the China health and retirement longitudinal study. *BMC Public Health*. 2024;24:1007.

5. Elser H, Caunca M, Rehkopf DH, Andres W, Gottesman RF, Kasner SE, Yaffe K, Schneider A. Trends and inequities in the diagnosis and treatment of poststroke depression: a retrospective cohort study of privately insured patients in the USA, 2003-2020. *J Neurol Neurosurg Psychiatry*. 2023;94:220-226.

6. Masuccio FG, Grange E, Di Giovanni R, Rolla M, Solaro CM. Post-Stroke Depression in Older Adults: An Overview. *Drugs Aging*. 2024;41:303-318.

7. Razmara A, Valle N, Markovic D, Sanossian N, Ovbiagele B, Dutta T, Towfighi A. Depression Is Associated with a Higher Risk of Death among Stroke Survivors. *J Stroke Cerebrovasc Dis*. 2017;26:2870-2879.

8. Hornsten C, Lövheim H, Gustafson Y. The association between stroke, depression, and 5-year mortality among very old people. *Stroke*. 2013;44:2587-9.

9. Sun B, Wang N, Li K, Yang Y, Zhang F. The mediating effects of hope on the relationships of social support and self-esteem with psychological resilience in patients with stroke. *BMC Psychiatry*. 2024;24:340.

10. Zhou X, Du M, Weng Y, Zhou L. Hard return: the development and transformation process of social participation in stroke survivors; a qualitative study and initial theory. *Clin Rehabil*. 2020;34:824-836.

11. Ettman CK, Fan AY, Philips AP, Adam GP, Ringlein G, Clark MA, Wilson IB, Vivier PM, Galea S. Financial strain and depression in the U.S.: a scoping review. *Transl Psychiatry*. 2023;13:168.

12. Sung P, Zhang Y, Choi JY. The Mediating Role of Formal Social Engagement in the Relationship Between Oral Health and Depressive Symptoms Among Older Adults in South Korea. *Int J Geriatr Psychiatry*. 2025;40:e70085.

13. Gao J, Wang Y, Xu J, Jiang J, Yang S, Xiao Q. Life expectancy among older adults with or without frailty in China: multistate modelling of a national longitudinal cohort study. *BMC Med*. 2023;21:101.

14. Liu C, Zhou R, Peng X, Chen X, Xia Z, Wei W, Zhu T, Chen G. The longitudinal study of the relationship between social participation pattern and depression symptoms in frail older adults. *Front Psychiatry*. 2024;15:1440641.

15. Meng W, Yan H. The Impact of Social Participation on the Health of Widowed Middle-Aged and Older Adults: A Longitudinal Study. *J Gerontol Soc Work*. 2025:1-23.

16. Dhanasekara CS, Kahathuduwa CN, Quispe-Orozco D, Ota R, Duarte Celada WR, Bushnaq S. Effects of Social Determinants of Health on Acute Stroke Care Among Patients With Acute Ischemic Stroke: A Retrospective Cohort Study. *Neurology*. 2024;103:e209951.

17. Mu□at MI, Cătălin B, Hadjiargyrou M, Popa-Wagner A, Gre□i□ă A. Advancing Post-Stroke Depression Research: Insights from Murine Models and Behavioral Analyses. *Life (Basel)*. 2024;14:1110.

18. Bright L, Baum CM, Roberts P. Person and Environment Factors Supporting Self-Care Performance and Social Participation After Mild Stroke. *OTJR (Thorofare N J)*. 2024;44:455-466.

19. Lindmark A, Eriksson M, Darehed D. Mediation Analyses of the Mechanisms by Which Socioeconomic Status, Comorbidity, Stroke Severity, and Acute Care Influence Stroke Outcome. *Neurology*. 2023;101:e2345-e2354.

20. Piccirillo ML, Rodebaugh TL. Personalized networks of social anxiety disorder and depression and implications for treatment. *J Affect Disord*. 2022;298:262-276.

21. Zhao Y, Hu Y, Smith JP, Strauss J, Yang G. Cohort profile: the China Health and Retirement Longitudinal Study (CHARLS). *Int J Epidemiol*. 2014;43:61-8.

22. Wang Z, Li T, Zhang J, Chu C, Yuan S. How spousal cognitive functioning affects the level of depression in middle-aged and older adults: An instrumental variable study based on CHARLS in China. *Biosci Trends*. 2024.

23. Ma X, He J, Hu Q, Wang W, Qiao H. Association between cooking with solid fuels and depressive symptoms among middle-aged and older adults in China: The mediating effect of the residential environment. *Ecotoxicol Environ Saf*. 2024;284:116886.

24. Li C, Long C, Wu H, Zhu G, Liu D, Zhang C, Shi L. The impact of internet device diversity on depressive symptoms among middle-aged and older adults in

China: A cross-lagged model of social participation as the mediating role. *J Affect Disord*. 2024.

25.   Butsing N, Zauszniewski JA, Ruksakulpiwat S, Griffin M, Niyomyart A. Association between post-stroke depression and functional outcomes: A systematic review. *PLoS One*. 2024;19:e0309158.

26.   Wang S, Wang CX, Zhang N, Xiang YT, Yang Y, Shi YZ, Deng YM, Zhu MF, Liu F, Yu P, Ungvari GS, Ng CH. The Association Between Post-stroke Depression, Aphasia, and Physical Independence in Stroke Patients at 3-Month Follow-Up. *Front Psychiatry*. 2018;9:374.

27.   Wang Z, Ruan H, Li L, Song N, He S. Association of changes in frailty status with the risk of all-cause mortality and cardiovascular death in older people: results from the Chinese Longitudinal Healthy Longevity Survey (CLHLS). *BMC Geriatr*. 2024;24:96.

28.   Ozkan H, Ambler G, Banerjee G, Mitchell JJ, Barbato C, Browning S, Leff AP, Simister RJ, Werring DJ. Prevalence, predictors, and patterns of patient reported non-motor outcomes six months after stroke: a prospective cohort study. *Lancet Reg Health Eur*. 2024;47:101080.

29.   Wu X, Xu K, Li T, Wang L, Fu Y, Ma Z, Wu X, Wang Y, Chen F, Song J, Song Y, Lv Y. Abnormal intrinsic functional hubs and connectivity in patients with post-stroke depression. *Ann Clin Transl Neurol*. 2024;11:1852-1867.

30.   Ma X, Zhang H, Tian Y, Wang Y, Liu L, Wang L. Mediating effect of depression on the association between cardiovascular disease and the risk of all-cause mortality: NHANES in 2005-2018. *Clin Cardiol*. 2023;46:1380-1389.

31.   Rabadi MH, Xu C. Motivation in veterans with an acute/subacute ischemic stroke did not improve cognition and functional motor recovery but reduced deaths. *Eur J Phys Rehabil Med*. 2024.

32.   Kong D, Zou W. Association between atherogenic index of plasma and post-stroke depression: a cross-sectional study. *Eur J Psychotraumatol*. 2024;15:2429266.

33.   Kim J, Park GR, Namkung EH. The link between disability and social participation revisited: Heterogeneity by type of social participation and by socioeconomic status. *Disabil Health J*. 2024;17:101543.

34.   Northcott S, Moss B, Harrison K, Hilari K. A systematic review of the impact of stroke on social support and social networks: associated factors and patterns of change. *Clin Rehabil*. 2016;30:811-31.

35. Zhang Y, Yu P, Liu H, Yao H, Yao S, Yuan SY, Zhang JC. Hyperforin improves post-stroke social isolation-induced exaggeration of PSD and PSA via TGF-β. *Int J Mol Med*. 2019;43:413-425.

36. Alberti A, Araujo Coelho DR, Vieira WF, Moehlecke Iser B, Lampert R, Traebert E, Silva B, Oliveira BH, Leão GM, Souza G, Dallacosta FM, Kades G, Madeira K, Chupel MU, Grossl FS, Souza R, Hur Soares B, Endrigo Ruppel da Rocha R, da Silva Sipriano E, Fernandes Martins D, Agostinetto L. Factors Associated with the Development of Depression and the Influence of Obesity on Depressive Disorders: A Narrative Review. *Biomedicines*. 2024;12:1994.

37. Teo AR, Nelson S, Strange W, Kubo H, Katsuki R, Kurahara K, Kanba S, Kato TA. Social withdrawal in major depressive disorder: a case-control study of hikikomori in japan. *J Affect Disord*. 2020;274:1142-1146.

38. Janssen NP, Lucassen P, Huibers M, Ekers D, Broekman T, Bosmans JE, Van Marwijk H, Spijker J, Oude Voshaar R, Hendriks GJ. Behavioural Activation versus Treatment as Usual for Depressed Older Adults in Primary Care: A Pragmatic Cluster-Randomised Controlled Trial. *Psychother Psychosom*. 2023;92:255-266.

39. Yisma E, Walsh S, Hillier S, Gillam M, Gray R, Jones M. Effect of behavioural activation for individuals with post-stroke depression: systematic review and meta-analysis. *BJPsych Open*. 2024;10:e134.

40. Stahl B, Millrose S, Denzler P, Lucchese G, Jacobi F, Flöel A. Intensive Social Interaction for Treatment of Poststroke Depression in Subacute Aphasia: The CONNECT Trial. *Stroke*. 2022;53:3530-3537.

41. Freeman A, Tyrovolas S, Koyanagi A, Chatterji S, Leonardi M, Ayuso-Mateos JL, Tobiasz-Adamczyk B, Koskinen S, Rummel-Kluge C, Haro JM. The role of socio-economic status in depression: results from the COURAGE (aging survey in Europe). *BMC Public Health*. 2016;16:1098.

42. Chan D, Lee S, Teh PL. Factors influencing technology use among low-income older adults: A systematic review. *Heliyon*. 2023;9:e20111.

43. Fu X, Peng S, Feng XL. Socioeconomic inequalities in depressive symptoms in China: The role of social capital. *Heliyon*. 2024;10:e24918.

44. Wang R, Chen Z, Zhou Y, Shen L, Zhang Z, Wu X. Melancholy or mahjong? Diversity, frequency, type, and rural-urban divide of social participation and depression in middle- and old-aged Chinese: A fixed-effects analysis. *Soc Sci Med*. 2019;238:112518.

45. Liu J, Rozelle S, Xu Q, Yu N, Zhou T. Social Engagement and Elderly Health in China: Evidence from the China Health and Retirement Longitudinal Survey (CHARLS). *Int J Environ Res Public Health*. 2019;16:278.

46. van den Hondel K, Runtuwene N, van Bergen A, Gilissen R, Punt P, Reijnders U, Buster M. Undiscovered solitary deaths in the four largest cities in the Netherlands. *J Forensic Leg Med*. 2020;74:102008.

47. Bucher HC, Ragland DR. Socioeconomic indicators and mortality from coronary heart disease and cancer: a 22-year follow-up of middle-aged men. *Am J Public Health*. 1995;85:1231-6.

48. Luo W, Xie Y. Socio-economic disparities in mortality among the elderly in China. *Popul Stud (Camb)*. 2014;68:305-20.

49. Sanjari E, Ahmadi A, Raeisi Shahraki H. Clustering the Economic Status via Partitioning around Medoid and Its Association with Common Non-communicable Diseases. *Iran J Med Sci*. 2024;49:699-706.

50. Liao XM, Su YA, Wang Y, Yu X, Si TM. Antidepressant treatment strategy with an early onset of action improves the clinical outcome in patients with major depressive disorder and high anxiety: a multicenter and 6-week follow-up study. *Chin Med J (Engl)*. 2020;133:726-728.

51. Wanjari M, Mittal G, Prasad R. Navigating the complexities of young stroke: the role of neurosurgery in early intervention. *Neurosurg Rev*. 2024;47:547.

1 **Table**

2 **Table 1 Basic characteristics and differences analysis of participants**

Variables	Survivors (n=5976)	Non-survivors (n=1125)	χ²/Z	P
Gender, n(%)			103.22	**< .001**
Female	3122 (52.24)	402 (35.73)		
Male	2854 (47.76)	723 (64.27)		
Age, n(%)			684.01	**< .001**
45~60 years old	3885 (65.01)	260 (23.11)		
> 60 years old	2091 (34.99)	865 (76.89)		
Smoking history, n(%)	2348 (39.29)	630 (56.00)	108.56	**< .001**
Married, n(%)	5507 (92.15)	869 (77.24)	229.51	**< .001**
Hypertension, n(%)	1384 (23.16)	392 (34.84)	68.93	**< .001**
Diabetes, n(%)	339 (5.67)	111 (9.87)	28.06	**< .001**
Dyslipidemia, n(%)	588 (9.84)	92 (8.18)	3.02	0.082
Stomach disease, n(%)	1332 (22.29)	226 (20.09)	2.68	0.102
Stomach pain, n(%)	519 (8.68)	100 (8.89)	0.05	0.824
Lung disease, n(%)	477 (7.98)	214 (19.02)	131.37	**< .001**
Kidney disease, n(%)	299 (5.00)	85 (7.56)	12.06	**< .001**
Heart disease, n(%)	646 (10.81)	193 (17.16)	36.59	**< .001**
Psychiatric disorder, n(%)	52 (0.87)	12 (1.07)	0.41	0.522
Stroke	97 (1.62)	49 (4.36)	35.10	**< .001**
Depression, M (Q_1, Q_3)	7.00 (3.00, 12.00)	8.00 (4.00, 13.00)	-6.90	**< .001**
Social participation, M (Q_1, Q_3)	2.00 (0.00, 4.00)	0.00 (0.00, 4.00)	-2.93	**< .001**
SES, M (Q_1, Q_3)	7.00 (6.00, 8.00)	6.00 (5.00, 7.00)	-15.35	**< .001**

3 Note: Values in **bold** indicate p < 0.05. SES:Socioeconomic status.

1 **Table2. Multivariate logistic regression of all-cause Mortality**

Variables	β	SE	Z	*P*	OR [95%CI]
Intercept	-0.17	0.26	-0.66	0.509	0.84 [0.50 ~ 1.41]
Stroke	0.52	0.20	2.61	**0.009**	1.69 [1.14 ~ 2.50]
Depression	0.02	0.01	2.99	**0.003**	1.02 [1.01 ~ 1.03]
Social participation	-0.02	0.01	-1.22	0.223	0.98 [0.96 ~ 1.01]
SES	-0.37	0.04	-10.41	**< .001**	0.69 [0.64 ~ 0.74]
Man	0.64	0.10	6.41	**< .001**	1.89 [1.56 ~ 2.30]
Age	1.39	0.08	17.36	**< .001**	4.02 [3.44 ~ 4.70]
Smoke	0.25	0.09	2.62	**0.009**	1.28 [1.06 ~ 1.54]
Marry	-0.83	0.10	-8.68	**< .001**	0.43 [0.36 ~ 0.52]
Hypertension	0.29	0.08	3.66	**< .001**	1.34 [1.15 ~ 1.57]
Diabetes disease	0.56	0.13	4.28	**< .001**	1.75 [1.35 ~ 2.25]
Lung disease	0.67	0.10	6.66	**< .001**	1.95 [1.60 ~ 2.37]
Kidney disease	0.21	0.14	1.47	0.141	1.24 [0.93 ~ 1.64]
Heart disease	0.17	0.10	1.65	0.098	1.18 [0.97 ~ 1.45]

2 Note: SES: Values in **bold** indicate p < 0.05. Socioeconomic status, OR: Odds Ratio, CI:
3 Confidence Interval.

1   **Figures with Figure Legends**

3   **Figure 1**. Participant selection flow chart.

2   **Fig**ure 2. Correlation Analysis. **(A)** Pearson correlation coefficients of depression, social

3   participation, and SES with all-cause mortality. **(B)** The correlations among different

4   types of social participation. SES: Socioeconomic status; *p<0.05,** p < .01,   ***< .001

Figure 3. Pathways between stroke and all-cause mortality. **(A)** Depression and social

participation mediated the association between stroke and all-cause mortality. **(B)**

Socioeconomic status moderated the mediating effects of depression and social

participation on the relationship between stroke and all-cause mortality. SES:

Socioeconomic status; Solid lines: Significant; Dashed line: Not significant; *p<0.05,**

p < .01, ***< .001

Figure 4 The moderating role of SES in nodes of Ind3. **(A)** The positive predictive effect

of stroke on depression (standardized using Z-scores) was more pronounced among

individuals with low SES. **(B)** The negative effect of depression on social participation

(standardized using Z-scores) was more pronounced among individuals with high SES.

SES: Socioeconomic status, Ind3: Stroke → Depression → Social Participation→

Mortality.

2 **Figure 5.** The interaction between social participation and time indicated that, over time,
3 lower levels of social participation were associated with a progressively increasing risk of
4 all-cause mortality, whereas higher levels of social participation were linked to a reduced
5 risk of all-cause mortality.