Development & Validation of a Machine Learning Model That Uses Voice to Predict Aspiration Risk

(Short Title: Machine Learning Analysis of Voice to Predict Aspiration Risk**)**

Cyril Varghese, MD, MS[1*], Jianwei Zhang, PhD[2,3], Sara A. Charney MS, CCC-SLP[4] ,Abdelmohaymin Abdalla, MBBS[1], Elizabeth Reeves MS, CCC-SLP [4] , Stacy Holyfield, MS, CCC-SLP[4] , Adam Brown, BS[5], Hunter Stearns, BS[5], Michelle Higgins, MD, PhD[5], Julie Liss, PhD, CCC-SLP[2], Nan Zhang MS[6], Diana Orbelo, PhD, CCC-SLP[7], Rebecca Pittelko MS, CCC-SLP[7], Lindsay Rigelman MA, CCC-SLP[7], Victor E. Ortega, MD, PhD[1], David G. Lott, MD[4], and Visar Berisha, PhD[2,3].

[1]Division of Pulmonary and Department of Critical Care Medicine, Mayo Clinic Arizona;[2]College of Health Solutions, Arizona State University; [3]College of Engineering, Arizona State University; [4]Division of Laryngology and Department of Otolaryngology- Head & Neck Surgery, Mayo Clinic Arizona; [5]Mayo Clinic Alix School of Medicine, Mayo Clinic Arizona; [6]Department of Quantitative Health Sciences, Mayo Clinic Arizona; [7]Division of Laryngology and Department of Otolaryngology- Head & Neck Surgery, Mayo Clinic Rochester

*Corresponding Author & Guarantor:

Cyril Varghese
5777 East Mayo Blvd
Phoenix, Arizona 85054
Varghese.cyril@mayo.edu

<u>Author Roles:</u>
Cyril Varghese - Conception of idea, data mining, data analysis, interpretation, writing, editing/reviewing manuscript
Jianwei Zhang- Data analysis, interpretation, writing, editing/reviewing manuscript
Sara Charney – Data mining, interpretation, writing, editing/reviewing manuscript
Abdelmohaymin Abdalla- Data mining, interpretation, writing, editing/reviewing manuscript
Elizabeth Reeves - Data mining, interpretation, writing, editing/reviewing manuscript
Stacy Holyfield - Data mining, interpretation, writing, editing/reviewing manuscript
Adam Brown - Data mining, interpretation, writing, editing/reviewing manuscript
Hunter Stearns - Data mining, interpretation, writing, editing/reviewing manuscript
Michelle Higgins - Data mining, interpretation, writing, editing/reviewing manuscript
Julie Liss - Data analysis, interpretation, writing, editing/reviewing manuscript
Nan Zhan - Data analysis, interpretation, writing, editing/reviewing manuscript
Diana Orbelo - Data mining, interpretation, editing/reviewing manuscript
Rebecca Pittelko - Data mining, interpretation, editing/reviewing manuscript
Lindsay Ringelman - Data mining, interpretation, editing/reviewing manuscript
Victor E. Ortega - Data interpretation, writing, editing/reviewing manuscript
David G. Lott - Data interpretation, writing, editing/reviewing manuscript
Visar Berisha - Conception of idea, data analysis, interpretation, writing, editing/reviewing manuscript.

This study was approved by the Mayo Institutional Review Board.

Keywords: Aspiration, voice, speech, artificial intelligence, ARDS, lung transplant, ILD

Abbreviations: Speech Language Pathologists (SLPs); Ear Nose & Throat or Otolaryngology (ENT); Machine Learning (ML); harmonic-to-noise ratio (HNR); relative average perturbation (RAP); Video Fluoroscopic Swallow Study (VFSS); Penetration Aspiration Scale (PAS) Fiberoptic Endoscopic Evaluation of Swallowing (FEES); Intensive Care Units (ICU); Neural Additive Model (NAM)

Acknowledgements: This work was made possible by CTSA Grant Number KL2 TR002379 from the National Center for Advancing Translational Science (NCATS). We are also grateful to Lanyu Mi, MS (Department of Qualitative Health Sciences, MCA) who assisted our primary biostatistician Nan Zhang, MS.

ABSTRACT

Background: Aspiration causes or aggravates a variety of respiratory diseases. Subjective bedside evaluations of aspiration are limited by poor inter- and intra-rater reliability, while gold standard diagnostic tests for aspiration, such as video fluoroscopic swallow study (VFSS) and fiberoptic endoscopic evaluation of swallowing (FEES), are cumbersome or invasive and healthcare resource intensive.

Research Aim: To develop and validate a novel machine learning algorithm that can analyze voice features, as an objective screening test to predict aspiration risk.

Methods: Recorded [i] phonations during routine nasal endoscopy from 163 unique patients were retrospectively analyzed for acoustic features including pitch, jitter, shimmer, harmonic to noise ratio (HNR), and others. Supervised machine learning (ML) was performed on the vowel phonations of those at high-risk for aspiration versus those at low-risk for aspiration. Ground truth of aspiration risk classification for model development was established using VFSS. The performance of the ML model was tested on an independent, external cohort of patient voice samples. The performance of trained Speech Language Pathologists (SLPs) to categorize high versus low-risk aspirators by listening to phonations was compared against the ML model.

Results: Mean ML risk score for those with the ground truth of high versus low aspiration risk was 0.530 ± 0.310 vs 0.243 ± 0.249, which was a significant difference (0.287, 95% CI: 0.192-0.381) $p<0.001$. In the development cohort, the model showed an area under the curve (AUC) for the Receiver Operator Characteristic (ROC) of 0.76 (0.67-0.84) with specificity of 0.76 and F1 score of 0.63. The performance of the model in an external testing cohort was comparable, with AUC of 0.70 (0.52-0.89) with a specificity of 0.81, and F1 score of 0.67. The ML model had better accuracy, sensitivity, specificity, negative and positive predictive values compared to trained SLPs in classifying aspiration risk by evaluating vowel phonations.

Conclusion: High risk aspirators have quantifiable voice characteristics that significantly differ from those who are at a low risk for aspiration, as detected by a ML model trained to analyze sustained phonation and tested on an independent cohort. The ML model's performance was superior to human experts in predicting aspiration risk by evaluating vowel phonations.

INTRODUCTION

Mechanical abnormalities of the oropharynx or larynx from tumor bulk or tissue loss/reconstruction from resection or radiation injury, damage to nerves or muscles of the upper airways either pathologic or from normative aging, structural or functional problems of the esophagus, altered sensorium, and gastroesophageal reflux can all cause clinically relevant aspiration of oropharyngeal or gastrointestinal (GI) contents into the lungs [1]. Acute aspiration can cause significant injuries leading to pneumonias and Acute Respiratory Distress Syndrome (ARDS)[2, 3]. Chronic aspiration of GI contents can cause or aggravate airway and parenchymal lung diseases further increasing the associated morbidity or mortality from conditions such as chronic lipoid pneumonia, bronchiectasis, obliterative bronchiolitis, refractory asthma and pulmonary fibrosis [4], and is a major risk factor for transplanted lung rejection [5, 6].

Aspiration is common but is undetected and usually only suspected after significant pulmonary damage has occurred. Despite its high prevalence in ambulatory and hospitalized patients [7-9], proactively detecting aspiration risk to prevent downstream sequelae is a major clinical challenge. The most widely used hospital-based method to rule out anterograde aspiration (aspiration that occurs during swallowing) are nursing or speech-pathologist administered bedside swallowing evaluations (**BSEs**). BSE tests vary across institutions but often include listening for a wet cough or wet voice quality after patients swallow varying amounts of liquid. Due to inter-rater/intra-rater reliability issues inherent with these subjective assessments [10],

the sensitivity for BSEs ranges from 27% to 85%, with specificities ranging from 50-80% [11], and poor predictive values [12]. Depending on BSE results, patients can be referred to gold standard confirmatory testing including a videofluoroscopic swallow study (**VFSS**)[13] and/or a fiberoptic endoscopic evaluation of swallowing (**FEES**)[14], administered by a speech-language pathologists (**SLP**). The VFSS exposes patients to radiation, requires coordinated participation and can be challenging to complete if patients are acutely ill, delirious, or have mobility issues. FEES is an invasive and uncomfortable evaluation where a scope is inserted into the nose and through the oropharynx to visualize the larynx and pharynx during swallowing. Both VFSS and FEES are resource intensive, requiring specialized equipment and the expertise of radiologists, laryngologists, and SLPs for administration and proper interpretation. These are difficult to complete in a timely manner, especially in resource limited settings. Consequently, there is an _unmet need_ to have an objective and quantitative bedside screening test that is easily administered, with a level of accuracy and precision that ensures that only a select subset of patients require confirmatory testing.

For aspiration to occur, gastrointestinal or oropharyngeal contents have to contact and pass through the vocal folds before they can enter the lungs. It is known that exposure to GI contents can induce histopathological changes to the vibratory margins of the vocal folds leading to changes in vocal quality, as it does in laryngopharyngeal reflux [15]. Therefore, _we hypothesize that increased risk of aspiration may be associated with changes in human voice quality that can be detected by machine learning (ML) approaches for the non-invasive and safe detection of aspiration risk._ Furthermore, we suspect that structural or functional changes in the vocal folds can lead to voice changes that may be associated with higher risk of aspiration. Voice, specifically phonation, is a complex high-dimensional signal with several quantifiable features that can be measured, including harmonic-to-noise ratio (HNR), amplitude perturbation quotient (APQ), jitter (a measure of frequency variation during phonation), and shimmer (a measure of

amplitude variation during phonation). While these features individually have shown predictive value in prior small studies, the results of these studies have been variable, focused on postprandial voice changes, and lack independent validation[16-18].To address these limitations we studied voice samples available from Mayo Clinic's Otolaryngology clinical practices to develop, validate and externally test an ML algorithm that analyzes voice to predict aspiration risk.

METHODS

In overview, we developed an ML model from a large sample of retrospective voice recordings collected as part of routine laryngoscopic Otolaryngology (ENT) evaluations at our institutions from patients with available VFSS exams. Because these voice samples were not collected during swallowing evaluations, we analyzed the potential long-term associations of aspiration on voice rather than simply observing immediate changes to voice with ingested materials lingering on the vocal folds after swallowing (e.g., wet vocal quality). We externally validated the model using an independent cohort of distinct patients from another hospital. We then tested its performance against trained SLPs. The workflow for voice data curation, ML model development and validation is shown in **Figure 1.** This study was approved by our Institutional Review Board.

Voice Data Collection

ENT clinics at our institution routinely record patients' voices during flexible laryngoscopy, of which a subset have VFSS performed to evaluate swallowing difficulties. Patients' voices are recorded using a lapel microphone clipped to the patients' clothing close to the clavicle. The lapel microphone is a standard input component of the Pentax Laryngeal Strobe system, commonly used in ENT clinical practices. Endoscopy and VFSS databases from January 2020 through July 2021 were curated to identify patients who had VFSS, an endoscopy exam, and

voice recordings during their longitudinal care encounters. 332 consecutive patients with both VFSS and voice recordings were identified. Three SLPs with combined experience of 15 years, listened to patient voice recordings and excluded unusable samples with background noise, overlapping sounds (e.g., clinicians or family members speaking, machine sounds such as those produced during video-stroboscope), or low volume. These recorded sound samples were analyzed without any computerized pre-processing to avoid potential artifacts from being introduced into the signal analysis pipeline. Due to the retrospective nature of this study, these voice samples were not specifically protocoled. Of the 772 voice recordings from 332 patients, 283 recordings (each containing multiple sustained [i] phonations) from 163 patients were deemed to be of sufficient quality for analysis based on the criteria above.

Clinical Data Collection

Clinical data was classified into several domains known to be risk factors for aspiration and included anatomical issues within the upper airways (vocal fold disease, head and/or neck surgery with or without radiation exposures), esophageal diseases (strictures, impaired motility, gastroesophageal reflux disease), neurological compromise (stroke, neuromuscular, neurodegenerative or peripheral nerve disease), body mass index (BMI), and obstructive sleep apnea.

Patients were classified as those with high-risk of aspiration, low-risk of aspiration, or moderate-risk of aspiration based on VFSS findings which we defined as the standard for aspiration risk categories for our training and validation analyses. The well-validated Rosenbek Penetration-Aspiration Scale [19], (PAS) maximum scores of 6-8 (materials passing below the vocal folds into the lower airways) were classified as high-risk aspirators, PAS scores of 3-5 (materials either contacting vocal folds or in the airways above the vocal folds without being ejecting) as moderate-risk aspirators, and PAS scores of 1-2 (materials not entering the airway or entering

the airway above the vocal folds but ejected out) were classified as those with low-risk of aspiration. VFSS were used rather than FEES due to the more consistent use of this test for swallow evaluations within our ENT practice. Due to the retrospective nature of this study, other swallow scales such as the Dynamic Imaging Grade of Swallowing Toxicity (DIGEST) were not available consistently, and therefore PAS scores evaluated all patients in this study. The voice recordings and VFSS were not performed simultaneously, and were typically performed on different days than the voice recordings with a median time between the voice recording and VFSS of 47 days (InterQuartile Range: 137 days).

Machine Learning and Statistical Analysis

Audacity (version 3.1.3) was used to manually remove noisy or distorted portions from sound recordings (ex: overlapping sounds from clinician and patient). These samples were exported as .wav files and PRAAT (version 6.2.15) was used to tag the samples with specific labels, including sustained vowel phonation cued speech, reading, and spontaneous speech. However, *only [i] vowel phonations were analyzed*, as these were most consistently collected across all patients. Three clips of the [i] vowel phonations lasting a minimum of 0.5 seconds from the central portion of each recorded clip per patient were used to develop the ML model.

In our initial model development and training, traditional voice features including pitch, jitter, shimmer, harmonics-to-noise ratio (HNR), cepstral peak prominence (CPP), and relative average perturbation (RAP) were extracted from aforementioned tagged samples using audio processing toolboxes PRAAT (GNU General Public License Version 3, 29 June 2007) and Collaborative Voice Analysis Repository for Speech Technologies (COVAREP)[20]. For traditional voice features, parametric *t*-tests were used to compare those with high-risk of aspiration and low-risk of aspiration. Comparisons of demographics and clinical characteristics between the groups were performed with Fishers' exact test for categorical variables and

Kruskal-Wallis rank sum test for continuous variables. These analyses were performed with the arsenal package (R4.2.2 [R Foundation for Statistical Computing, Vienna, Austria]).

Neural additive models (NAM) [21]were used to classify the voice data and analyze the [i] phonation data of 163 patients. The following dependencies were used for ML coding in Python3: PyTorch, numPy and sciPy. For NAM, we included 33 features that are well described to have validity in characterizing pathological voice states [22-24]. Five-fold cross-validation and Recursive Feature Addition (RFA) [25, 26] was employed to identify the features required to effectively distinguish high-risk from low-risk aspirators and to estimate model performance. RFA initiates with a single feature and iteratively adds the voice feature that optimally enhances the performance of the existing set.

Independent Model Testing

The development NAM model was trained on voice data from our center, while independent validation of NAM was performed with voice data of patients from another center. The clinicians, recordings, equipment/rooms, geographic location, and patient demographics of the external testing cohort were independent and unique to that of the training cohort, to test the true performance of the model. The selection, extraction, tagging, and processing of these externally collected voice samples were performed using the *same* methodology as described previously for the development cohort. ML risk score for each patient in the testing cohort was calculated in a blinded fashion by the model by analyzing [i] phonation clips in the context of the patient's age and sex. The gold standard designation (i.e., aspiration risk category based on VFSS) or clinical information was not known to the investigators running the ML code. Statistical analyses including sensitivity, specificity, receiver Operator Characteristics (ROC) and precision were calculated to assess the model's performance on both the testing and development cohorts.

Comparing Human Expert Raters with ML Model

Four SLPs, from two medical centers, with a combined 40 years of clinical experience listening to human voice, evaluated the [i] phonation clips from the testing cohort described above to make a determination of aspiration risk. In addition to listening to the phonation, the raters were unblinded to the age and sex of the patient. The rest of the medical history of the patient, including aspiration risk status, demographic details etc. was blinded to the raters just as it was for the ML model. The inter-rater reliability for the human raters were assessed using pairwise Cohen's kappa. The sensitivity, specificity, positive/negative predictive values and accuracy of the human raters and the ML model for predicting high versus low-risk aspirators were calculated.

RESULTS

Feature Analysis

We initially analyzed retrospective samples from 87 patients (19 high-risk aspirators, 10 medium-risk aspirators, and 58 low-risk aspirators) with an average of approximately 2 voice files per patient. Non-parametric analysis of variance of demographic and clinically relevant risks of aspiration in the pilot data revealed that only age and sex were different among the groups. On age and sex-matching prior to analysis, 17 high-risk aspirators were found to be age- and sex-matched to 17 low-risk aspirators in our exploratory cohort. Of the traditional voice features that were explored, high-risk aspirators were found to have higher jitter and shimmer but lower harmonics richness factor than low-risk aspirators (**Figure 2**).

Development of ML Model and Internal Cross-Validation

Our primary ML model development and training cohort evaluated 163 patients within the three groups (47 high-risk aspirators, 17 moderate-risk aspirators, and 99 low-risk aspirators) that

significantly differed by sex, age, and BMI (with high-risk aspirators having a lower BMI than controls). Additional differences included more reported solid and liquid dysphagia symptoms and a higher frequency of structural changes to head and neck anatomy in the high-risk group than in the low-risk group (**Table 1**). Baseline characteristics for the age- and sex-matched cohort of 36 high-risk aspirators and 36 low-risk aspirators (N=72) showed BMI and incidence of head and neck disease was different between the groups (**Table Supplemental 1**).

Because age and sex influence voice, and were different between the groups, they were controlled for in the NAM (**Figure 1**). The model utilized 33 extracted features to differentiate between aspirators and controls (**Table Supplemental 2**). Each sub-network within the model processed a single extracted voice feature along with the corresponding sample sex and age as inputs (**Figure 1**). The sub-network output was a single scalar reflecting the adjustments made for age and sex within the feature. The whole NAM output for aspiration risk was a scalar ranging from 0 to 1. High-risk aspirators were distinctly classified as a separate group compared to low-risk aspirators: high-risk aspirator score=0.528 (standard deviation [SD]=0.2478) versus low-risk aspirator risk score of 0.252 (SD=0.241); mean risk difference between groups was 0.2765 (95% CI: 0.2122-0.3408), p<0.001 (**Figure 3A**). Moderate-risk aspirators had a risk score between high-risk aspirators and low-risk aspirators, although this difference when compared to the other groups was not significant. RFA showed an elbow point at 7 features (**Figure 3B**), revealing that the most significant voice features adding to the model's discriminability were the average fundamental frequency and standard deviation (F0_mean and F0_std [standard deviation]), the maximum fundamental frequency during phonation (Max Pitch), the standard deviation of the quai-open quotient that measures the proportion of the glottal cycle when the glottis is open (QOQ_std), the average of the harmonic richness factor (HRF_mean), the average of the cepstral peak prominence (CPP_mean), and the standard

deviation of the cepstral peak prominence (CPP_std). A description of these features and the other 33 features analyzed in the NAM is in **Table Supplemental 2**.

External Testing of the ML model

There were 19 high-risk aspirators and 16 low-risk aspirators in the external testing cohort based on VFSS which totaled 24% of the sample size of high and low risk aspirators in the development cohort. The demographics between the training and testing cohort were different. For example, the testing cohort had a higher proportion of female high-risk aspirators than the training cohort (20% vs 5.5%), and a lower proportion of male low-risk aspirators based on VFSS compared to the training cohort (8.6% vs 45%), p=0.005. The testing cohort was older than the training cohort (70.3$\pm$13.4 vs 65.9$\pm$12.6 years, p=0.029) The mean ML risk score output for the high-risk aspirators by VFSS was significantly higher than the low-risk aspirators (Mean (SD) :0.469 (0.327) vs 0.245 (0.265), p =0.047) (**Figure 4A**). The optimal ML risk score cut-off to distinguish high-risk aspirators from low-risk aspirators by analyzing the training cohort was determined to be 0.35 based on the Youden index. Despite the demographic differences between the testing and development cohort, the area under the curve (AUC) for the ROC for the testing cohort was 0.697 (0.517-0.878) compared to AUC of 0.755 (0.666-0.843) for the training cohort (**Fig. 4B**). The precision, recall, F1 score and specificity of the model for the testing cohort were 0.79,0.57, 0.67 and 0.81 respectively. **Fig. 4C.**

Human Raters versus ML model

Other than two raters that had very good agreement (pairwise Cohen's Kappa of 0.8) the rest of the raters had fair agreement (Choen's Kappa range 0.34-0.59) (**Table 2**). In classifying blinded patients as a high-risk or low-risk aspirator by analyzing [i] phonations, the ML model had a better overall accuracy (69% vs human raters' range 46-60%), sensitivity (58% vs human raters' range 32-47%) , PPV (79% vs human raters' range 50-78%) and NPV (62% vs human raters'

range 41-54%) compared to all the human raters (**Table 3**). Other than one rater the ML model also had better specificity compared to human raters (81% vs human raters' range 44-88%).

DISCUSSION

While acutely or chronically having ingested materials enter the larynx alters the quality of voice, human judgment of voice changes [27, 28] or even validated tools for perceptual assessments are unreliable screening methods to detect aspiration risk [29, 30]. Subjective assessments of voice by human observers are known to be poorly reproducible [10]. There has therefore been an interest in developing tools that can more objectively quantify aspiration-related voice changes in patients [16-18].The analysis of voice features in our cohort had similarities to the previous studies in that those designated as high-risk aspirators had quantifiably poorer voice quality with higher mean shimmer and jitter measures, but lower mean HNR than low-risk aspirators. In a study of 93 patients by Ryu and colleagues, Relative Average Perturbation (RAP) combined with HNR increased sensitivity of detecting aspiration in the five voice features analyzed [16]. Another similar study of 165 patients by Kang and colleagues demonstrated that among the 8 voice features evaluated, the most significant distinguishing measure that differentiated between aspirators and non-aspirators, was the RAP [17]. More recently, a study of 198 patients by Kim and colleagues reported using ML techniques to detect aspiration by analyzing postprandial voice [18]. However, all these studies were single center studies without an external testing cohort, the focus of these and other studies were on the acute effects of aspiration on voice as ingested materials contact the vocal folds in real time. However, current bedside swallow evaluations also screen for aspiration by making the patient swallow. There are no available tests to objectively detect aspiration risk without having the patient swallow liquids or solids. This can be a limitation for frail, immobile, delirious patients who are already at a considerable risk for aspiration. Such patients may benefit from an objective test that screens for aspiration risk but only require simple voicing tasks as in our evaluation. This study

investigated the chronic quantifiable alterations to human voice in the context of patients who have a tendency to aspirate for a variety of structural or functional reasons. This method could potentially allow for more frequent longitudinal inpatient/outpatient testing, while ensuring only a select group of patients will ultimately need confirmatory testing with VFSS or FEES. A commonly held misconception is that an exceptionally large sample size (in the order of thousands) is necessary to train ML models. While this is true for deep learning, it is not a necessity for supervised machine learning methods as in our study, where human experts are involved in the workflow to review features and determine if they are contributory [31].The sample size used to train this supervised neural additive model that controlled for confounders of voice (namely sex and age) for each of the 33 features effectively modeled the complex interactions and provided a more comprehensive analysis of voice than prior explorations in the context of aspiration. One of the benefits of the neural additive model is the sequential way with which features can be considered and confounding variable controlled, thereby improving explainability of the model [21]. While simpler models like logistic regression can classify by accounting for basic covariates such as age and sex, they are limited in capturing non-linear relationships and feature-specific interactions. The NAM's feature-specific networks capture complex, non-linear interactions while controlling for relevant patient characteristics. The performance of the model improved as a function of increasing feature dimension.

Additionally, unlike other studies, we tested the performance of our model in an independent external testing cohort. Previous work has shown that external testing of models is critical as it can be difficult to train clinical speech models that generalize [32]. Our external testing cohort was from an independent, geographically distinct center where retrospective voice samples were collected from patients who were demographically different from our training cohort. Despite these differences, the model's performance did not change significantly (**Figure 4**), suggesting that the model is not fragile. Fragile models that perform well in the development

cohorts, but not so well in external cohorts, can seldom be used effectively in clinical practice. Finally, we explored if the ML tool really added value as a preliminary test to predict aspiration risk by comparing its rating to SLPs. We found that the inter-rater reliability among our SLP raters was only "fair", similar to reports in literature. While our model had a superior performance compared to our expert SLP raters, it should be noted that SLPs do not make decisions about aspiration risk on the basis of sustained phonations alone. However, this study highlights the potential benefit of ML to complement informed clinical decision making by experts.

This study has several limitations. The relationship between voice and aspiration is likely a complex bidirectional relationship that cannot be fully elucidated with a retrospective analysis alone. Our population of ENT patients have other reasons related or unrelated to aspiration for their voice to be altered (e.g., radiation-induced vocal fold scarring). We controlled for age and sex (predictors known to covary with voice) using our neural additive model. However, other clinical factors like BMI and anatomical head/neck disease are known to covary with aspiration (the response variable) either as a cause or an effect and therefore cannot be easily controlled by generalized regression models in a retrospective analysis. The AUC of our model was in the range of 0.70. Certainly, there is room to fine tune the model and improve its performance by collecting voice samples from more diverse clinical settings and sampling connected speech. While we were constrained by the retrospective nature of the voice data, this study serves as the motivation to launch a prospective multi-site voice collection trial to study voice and its relationship to aspiration better. The elbow point in the cross-validation analysis (**Figure 3B**) was attained at 7 features (highlighted in **Table Supplemental 2** and listed in Results section). These features and interactions were the most predictive of aspiration risk, but only in the specific sample we evaluated, namely ENT patients. We expect that as we expand our databases to include more diverse patient samples from different clinical settings, other voice

features may emerge as significant predictors. Our categorization of the ground truth of "high" and "low" aspiration risk was based on VFSS at one point in time. It could be claimed that this is not reflective of the patient's true longitudinal aspiration risk status. This is why we were careful to label the categories in terms of "risk" rather than absolute diagnoses. Those categorized as high aspiration "risk" were widely separated from the low aspiration "risk" (PAS 6-8 vs PAS 1-2), and crossovers between these categories in an ENT practice although possible is unlikely. We do not claim that this ML tool can be used as a confirmatory diagnostic test to rule out or rule in aspiration like VFSS or FEES. Rather the goal of this study was to develop and validate a tool that can potentially serve as an easily deployable screening test used by bedside nurses and SLPs to more objectively screen for anterograde aspiration.

This study is the first of its kind to utilize ML techniques to quantify the quality of voice to estimate aspiration risk without performing a simultaneous swallow evaluation. We hope that ongoing development of such technology will impact the early identification of aspiration in a variety of clinical settings including ICUs, hospital wards, ambulatory clinics, and remote monitoring. Early identification of aspiration could potentially impact the mitigation and management of many acute and chronic pulmonary diseases.

TAKE HOME POINTS

Study Question: Aspiration causes many respiratory diseases. Can voice be used as a biomarker for easy and non-invasive detection of aspiration risk?

Results: We developed and then externally tested a machine learning model that analyzes a simple vowel phonation to predict risk of aspiration.

Conclusion: Patients with high risk of aspiration have quantifiable differences in voice characteristics compared to those with low risk of aspiration that can be detected by machine learning techniques.

FIGURES

MODEL TRAINING (Samples from Mayo Clinic Arizona)

EXTERNAL MODEL TESTING (Samples from Mayo Clinic Rochester)

Figure 1: A graphical representation summarizing the workflow for this study from voice sample collection, phonation tagging, and subsequent ML model development based on gold standard testing (VFSS determined aspiration). Subsequently, independent model evaluation was performed wherein aspiration risk category based on VFSS was unknown to the model. Contained within the green box is a schematic of the neural additive model (NAM) [23] trained to estimate risk of aspiration utilizing 33 voice features, in full cohort (N=163), while adjusting for age and sex, the main confounders between the groups. Σ: summation of all sub-feature network outputs and learnable offset β; **σ**: represent the sigmoid function $\sigma(x) = \frac{1}{1+e^{-x}}$; β: learnable offset of NAM model.

Figure 2: Pilot data of 17 age/sex matched aspirators with 17 non-aspirators, reveal that aspirators had a higher mean jitter (mean difference 0.0085, 95% CI: 0.000756-0.0161), shimmer (mean difference 0.0266, 95% CI: 0.00228-0.0506) and lower harmonics-to-noise ratio (HNR) (mean difference 3.1, 95% CI: 0.603-5.6). Means represented by black diamonds.

A

B

Figure 3: **A**: Aspirator Neural Additive Model (NAM) risk score 0.528+0.2478 (mean + std) versus non-aspirator (control) risk score of 0.252+0.241 (mean + std); Mean risk difference between groups 0.2765 (95% CI: 0.2122-0.3408), p<0.001. Means represented by black diamonds. Probable aspirators risk score continued to fall between aspirators and non-aspirators. **B:** The Recursive Feature Addition (RFA) method was employed to identify the minimal set of features required to effectively distinguish between aspirators and controls. This method initiates with a single feature and iteratively adds the voice feature that enhances the performance of the existing set. The key metric used to assess the performance of the feature set is the mean risk score difference.

Development Cohort   External Testing Cohort

A

B

C

Development Cohort		External Testing Cohort	
Precision: 0.58	Recall: 0.70	Precision: 0.79	Recall: 0.58
F1 score: 0.63	Specificity: 0.76	F1 score: 0.67	Specificity: 0.81

Figure 4: A: Box plot of ML Risk Score for training versus validation cohort. **B:** Receiver Operator Curve for development versus external testing cohort. AUC: Area under the curve. **C:** Analyses evaluating the performance of the model on training versus validation cohort. Where, F1 score= 2*Precision*Recall/ (Precision + Recall)

Table 1: Baseline characteristics of training cohort

	High Aspiration Risk (N=47)	Low Aspiration Risk (N=99)	Moderate Aspiration Risk (N=17)	p value
Sex				< 0.001[1]
Male	39 (83.0%)	45 (45.5%)	9 (52.9%)	
Female	8 (17.0%)	54 (54.5%)	8 (47.1%)	
Age at Scope Exam				< 0.001[2]
Mean (SD)	72.3 (9.9)	62.9 (12.6)	75.1 (9.0)	
Median	73.0	61.0	77.0	
Q1, Q3	67.0, 78.0	55.0, 72.0	65.0, 82.0	
Range	40.0 - 88.0	31.0 - 90.0	62.0 - 89.0	
BMI				0.009[2]
Mean (SD)	24.1 (3.2)	27.0 (5.3)	26.4 (7.7)	
Median	23.8	26.8	25.3	
Q1, Q3	22.0, 26.1	23.4, 29.7	21.3, 29.4	
Range	18.7 - 31.2	17.0 - 44.2	16.0 - 48.9	
Dysphagia				0.051[1]

No	4 (8.5%)	25 (25.25%)	2 (11.8%)	
Yes	40 (85.1%)	72 (72.7%)	14 (82.4%)	
NA	3 (6.4%)	2 (2%)	1 (5.9%)	
Esophageal disease group				0.669[1]
Clinical GERD	23 (48.9%)	46 (46.5%)	5 (29.4%)	
Other esophageal disease	3 (6.4%)	6 (6.1%)	0 (0.0%)	
Multiple esophageal diseases	2 (4.3%)	3 (3.0%)	1 (5.9%)	
GERD on impedance/manometry studies	1 (2.1%)	0 (0.0%)	0 (0.0%)	
No esophageal disease	9 (19.1%)	24 (24.2%)	4 (23.5%)	
NA	9 (19.1%)	20 (20.2%)	7 (41.2%)	
OSA				0.897[1]
Compliant with CPAP	8 (17.0%)	19 (19.2%)	1 (5.9%)	
Not compliant with CPAP	2 (4.3%)	5 (5.1%)	1 (5.9%)	

Compliance not reported	2 (4.3%)	4 (4.0%)	0 (0.0%)	
No OSA	35 (74.5%)	71 (71.7%)	15 (88.2%)	
Neurological illness				0.013[1]
CVA without any deficit	1 (2.1%)	4 (4.0%)	0 (0.0%)	
ALS	0 (0.0%)	1 (1.0%)	0 (0.0%)	
Neuromuscular diseases	3 (6.4%)	0 (0.0%)	0 (0.0%)	
CVA with dysphagia only	1 (2.1%)	0 (0.0%)	0 (0.0%)	
CVA with residual neuro deficit	1 (2.1%)	0 (0.0%)	0 (0.0%)	
None	41 (87.2%)	94 (94.9%)	17 (100%)	
Vocal fold disease				0.410[1]
Yes	23 (48.9%)	35 (35.4%)	8 (47.1%)	
No	22 (46.8%)	62 (62.6%)	9 (52.9%)	
NA	2 (4.3%)	2 (2.0%)	0 (0.0%)	
Head and neck anatomical disease				0.004[1]
Surgery	6 (12.8%)	13 (13.1%)	2 (11.8%)	

Cancer	1 (2.1%)	1 (1.0%)	0 (0.0%)	
Radiation	0 (0.0%)	2 (2%)	0 (0.0%)	
Multiple head and neck anatomical diseases	25 (53.2%)	22 (22.2%)	5 (29.4%)	
None	15 (31.9%)	61 (61.6%)	10 (58.8%)	

1.   Fisher's Exact Test for Count Data

2.   Kruskal-Wallis rank sum test

NA: Not reported by patient or not found in chart review

Table 2: Inter-rater reliability pairwise Kappa coefficient between human SLP raters.

SLP Raters	Rater 1	Rater 2	Rater 3
Rater 2	0.590		
Rater 3	0.508	0.397	
Rater 4	0.801	0.505	0.343

Table 3: Blinded Human (raters 1-4) and Machine (rater 5) raters' ability to predict aspiration risk by listening to voice against the ground truth.

Reader	Sensitivity	Sensitivity 95% CI	Specificity	Specificity 95% CI	PPV	PPV 95% CI	NPV	NPV 95% CI	Overall Accuracy	Overall Accuracy 95% CI
Rater 1	36.8%	(17.2%, 61.4%)	75.0%	(47.4%, 91.7%)	63.6%	(31.6%, 87.6%)	50.0%	(31.4%, 68.6%)	54.3%	(36.9%, 70.8%)
Rater 2	36.8%	(17.2%, 61.4%)	87.5%	(60.4%, 97.8%)	77.8%	(40.2%, 96.1%)	53.8%	(33.7%, 72.9%)	60.0%	(42.2%, 75.6%)
Rater 3	47.4%	(25.2%, 70.5%)	43.8%	(20.8%, 69.4%)	50.0%	(29.0%, 71.0%)	41.2%	(19.4%, 66.5%)	45.7%	(29.2%, 63.1%)

Rater 4	31.6%	(13.6%, 56.5%)	75.0%	(47.4%, 91.7%)	60.0%	(27.4%, 86.3%)	48.0%	(28.3%, 68.2%)	51.4%	(34.3%, 68.3%)
Rater 5 (Machine)	57.9%	(34.0%, 78.9%)	81.2%	(53.7%, 95.0%)	78.6%	(48.8%, 94.3%)	61.9%	(38.7%, 81.0%)	68.6%	(50.6%, 82.6%)

Table Supplemental 1: Baseline characteristics of training cohort when high-risk and low-risk aspirators were age and sex-matched

	High Aspiration Risk (N=36)	Low Aspiration Risk (N=36)	p value
Sex			1.000[1]
Male	28 (77.8%)	28 (77.8%)	
Female	8 (22.2%)	8 (22.2%)	
Age at Scope Exam			0.636[2]
Mean (SD)	70.9 (10.2)	70.1 (10.0)	
Median	73.0	71.0	
Q1, Q3	65.8, 77.2	66.8, 77.0	
Range	40.0 - 88.0	40.0 - 90.0	
BMI			0.003[2]
Mean (SD)	24.1 (3.0)	27.6 (5.5)	

Median	23.7	27.0	
Q1, Q3	22.8, 26.0	23.8, 30.2	
Range	18.7 - 31.2	17.0 - 44.2	
Dysphagia			0.115[1]
No	3 (8.3%)	9 (25.0%)	
Yes	30 (83.3%)	27 (75.0%)	
NA	3 (8.3%)		
Esophageal disease group			0.888[1]
Clinical GERD	16 (44.4%)	14 (38.9%)	
Other esophageal disease	3 (8.3%)	3 (8.3%)	
Multiple esophageal diseases	1 (2.8%)	1 (2.8%)	
GERD on impedance/manometry studies	1 (2.8%)	0 (0.0%)	
No esophageal disease	7 (19.4%)	11 (30.6%)	
Not reported	8 (22.2%)	7 (19.4%)	
OSA			0.608[1]
Compliant with CPAP	6 (16.7%)	10 (27.8%)	

Not compliant with CPAP	2 (5.6%)	1 (2.8%)	
Compliance not reported	1 (2.8%)	2 (5.6%)	
No OSA	27 (75.0%)	23 (63.9%)	
Neurological illness			0.186[1]
CVA without any deficit	0 (0.0%)	2 (5.6%)	
Neuromuscular diseases	2 (5.6%)	0 (0.0%)	
CVA with dysphagia only	1 (2.8%)	0 (0.0%)	
neuro not mentioned	3 (8.3%)	1 (2.8%)	
None	30 (83.3%)	33 (91.7%)	
Vocal fold disease			0.786[1]
Yes	17 (47.2.6%)	15 (41.7%)	
No	17 (47.2%)	19 (52.8%)	
Not reported	2 (5.5%)	2 (5.6%)	
Head and neck anatomical disease			0.037[1]
Surgery	2 (5.5%)	3 (8.3%)	
Cancer	1 (2.8%)	0 (0.0%)	
Radiation	0 (0.0%)	1 (2.8%)	
Multiple head and neck anatomical diseases	21 (58.3%)	11 (30.6%)	

None	12 (33.3%)	21 (58.3%)	

1. Fisher's Exact Test for Count Data

2. Kruskal-Wallis rank sum test

NA: Not reported by patient or not found on chart review

Table Supplemental 2: List of 33 voice features that were used for machine learning organized in clinically meaningful domains. The seven features that contributed most to the model's discriminability are bolded.

Group	Feature	Description
Jitter Features These measure variations in pitch and are indicators of vocal stability. Higher values of jitter measurements typically indicate greater variability, which is often associated with pathological voices.	jitter_local	Average absolute difference between consecutive periods, divided by the average period.
	jitter_local_abs	Average absolute difference between consecutive periods, in seconds.
	jitter_rap	Relative average perturbation, the average absolute difference between a period and the average of it and its two neighbors, divided by the average period.
	jitter_ppq5	Five-point Period Perturbation Quotient, the average absolute difference between a period and the average of it and its four closest neighbors, divided by the average period.
	jitter_ddp	Average absolute difference between consecutive differences between

		consecutive periods, divided by the average period.
Shimmer Features These measure variations in amplitude of the acoustic signal, reflecting vocal fold vibration regularity. Higher shimmer values suggest more amplitude variability, commonly found in voices with pathologies.	shimmer_local	Average absolute difference between the amplitudes of consecutive periods, divided by the average amplitude.
	shimmer_local_db	Average absolute base-10 logarithm of the difference between the amplitudes of consecutive periods, multiplied by 20.
	shimmer_apq3	Three-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of its neighbors, divided by the average amplitude.
	shimmer_apq5	Five-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of it and its four closest neighbors, divided by the average amplitude.
	shimmer_apq11	11-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of it and its ten closest neighbors, divided by the average amplitude.
	shimmer_dda	Average absolute difference between consecutive differences between the amplitudes of consecutive periods.

Pitch Features	**F0_mean**	**Average fundamental frequency, related to the perceived pitch of the voice.**
	F0_std	**Standard deviation of the fundamental frequency, indicating how much pitch varies over time.**
	min_pitch	Minimum fundamental frequency during phonation.
	max_pitch	**Maximum fundamental frequency during phonation.**
Harmonic Features These features reflect voice quality (clarity/richness) and can help identify abnormalities or pathologies.	hnr_mean	Mean harmonics-to-noise ratio, indicating the ratio of harmonic sound to noise in the voice signal.
	hnr_std	Standard deviation of the HNR, reflecting variability in voice quality.
	h1h2_mean	Mean difference in amplitude between the first and second harmonics.
	h1h2_std	Standard deviation of the difference between the first and second harmonics.
	hrf_mean	**Mean of the harmonic richness factor, a measure of how much energy is in the harmonics relative to the fundamental frequency.**

	hrf_std	Standard deviation of the harmonic richness factor, reflecting variability in harmonic richness.	
Glottal Source Analysis Features	peak_slope_mean	Average slope of the spectral peaks, indicative of how sharply energy dissipates	
	peak_slope_std	Standard deviation of the spectral peak slope, showing variability in how energy distribution changes over time.	
	mdq_mean	Average of the Maxima Dispersion Quotient (MDQ), often related to the rate of vocal fold closure.	
	mdq_std	Standard deviation of the MDQ, indicating the consistency of vocal fold closure.	
	naq_mean	Mean normalized amplitude quotient (NAQ), which relates to the efficiency of the glottal closure during phonation.	
	naq_std	Variability in NAQ, indicating how consistently the vocal folds are closing.	
	qoq_mean	Average quasi-open quotient (QOQ), which measures the proportion of the glottal cycle during which the glottis is open.	
	qoq_std	**Standard deviation of the QOQ, indicating the consistency of glottal opening during phonation.**	

	psp_mean	Average peak slope of the spectrum, offering insight into how spectral energy is concentrated around the peak frequency.
	psp_std	Variability in the spectral peak slope, which can reflect changes in vocal tract dynamics or resonance strategies.
	cpp_mean	**Mean cepstral peak prominence, which is a measure of the prominence of the peak in the cepstral domain, indicative of voice quality.**
	cpp_std	**Variability in CPP, indicating how consistently the voice can produce clear, strong cepstral peaks.**

Table Supplemental 2: List of 33 voice features that were used for machine learning organized in clinically meaningful domains. The seven features that contributed most to the model's discriminability are bolded.

Group	Feature	Description
Jitter Features	jitter_local	Average absolute difference between consecutive periods, divided by the average period.

These measure variations in pitch and are indicators of vocal stability. Higher values of jitter measurements typically indicate greater variability, which is often associated with pathological voices.	jitter_local_abs	Average absolute difference between consecutive periods, in seconds.
	jitter_rap	Relative average perturbation, the average absolute difference between a period and the average of it and its two neighbors, divided by the average period.
	jitter_ppq5	Five-point Period Perturbation Quotient, the average absolute difference between a period and the average of it and its four closest neighbors, divided by the average period.
	jitter_ddp	Average absolute difference between consecutive differences between consecutive periods, divided by the average period.
Shimmer Features These measure variations in amplitude of the acoustic signal, reflecting vocal fold vibration regularity. Higher shimmer values suggest more amplitude variability, commonly found in voices with pathologies.	shimmer_local	Average absolute difference between the amplitudes of consecutive periods, divided by the average amplitude.
	shimmer_local_db	Average absolute base-10 logarithm of the difference between the amplitudes of consecutive periods, multiplied by 20.
	shimmer_apq3	Three-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of its neighbors, divided by the average amplitude.
	shimmer_apq5	Five-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of it and its four closest neighbors, divided by the average amplitude.
	shimmer_apq11	11-point Amplitude Perturbation Quotient, the average absolute difference between the amplitude of a period and the average of the amplitudes of it and its ten closest neighbors, divided by the average amplitude.
	shimmer_dda	Average absolute difference between consecutive differences between the

		amplitudes of consecutive periods.
Pitch Features	**F0_mean**	**Average fundamental frequency, related to the perceived pitch of the voice.**
	F0_std	**Standard deviation of the fundamental frequency, indicating how much pitch varies over time.**
	min_pitch	Minimum fundamental frequency during phonation.
	max_pitch	**Maximum fundamental frequency during phonation.**
Harmonic Features These features reflect voice quality (clarity/richness) and can help identify abnormalities or pathologies.	hnr_mean	Mean harmonics-to-noise ratio, indicating the ratio of harmonic sound to noise in the voice signal.
	hnr_std	Standard deviation of the HNR, reflecting variability in voice quality.
	h1h2_mean	Mean difference in amplitude between the first and second harmonics.
	h1h2_std	Standard deviation of the difference between the first and second harmonics.
	hrf_mean	**Mean of the harmonic richness factor, a measure of how much energy is in the harmonics relative to the fundamental frequency.**
	hrf_std	Standard deviation of the harmonic richness factor, reflecting variability in harmonic richness.
Glottal Source Analysis Features	peak_slope_mean	Average slope of the spectral peaks, indicative of how sharply energy dissipates
	peak_slope_std	Standard deviation of the spectral peak slope, showing variability in how energy distribution changes over time.

		mdq_mean	Average of the Maxima Dispersion Quotient (MDQ), often related to the rate of vocal fold closure.
		mdq_std	Standard deviation of the MDQ, indicating the consistency of vocal fold closure.
		naq_mean	Mean normalized amplitude quotient (NAQ), which relates to the efficiency of the glottal closure during phonation.
		naq_std	Variability in NAQ, indicating how consistently the vocal folds are closing.
		qoq_mean	Average quasi-open quotient (QOQ), which measures the proportion of the glottal cycle during which the glottis is open.
		qoq_std	**Standard deviation of the QOQ, indicating the consistency of glottal opening during phonation.**
		psp_mean	Average peak slope of the spectrum, offering insight into how spectral energy is concentrated around the peak frequency.
		psp_std	Variability in the spectral peak slope, which can reflect changes in vocal tract dynamics or resonance strategies.
		cpp_mean	**Mean cepstral peak prominence, which is a measure of the prominence of the peak in the cepstral domain, indicative of voice quality.**
		cpp_std	**Variability in CPP, indicating how consistently the voice can produce clear, strong cepstral peaks.**

REFERENCES

1. Mandell, L.A. and M.S. Niederman, *Aspiration pneumonia.* New England Journal of Medicine, 2019. **380**(7): p. 651-663.
2. Bersten, A.D., et al., *Incidence and mortality of acute lung injury and the acute respiratory distress syndrome in three Australian States.* American journal of respiratory and critical care medicine, 2002. **165**(4): p. 443-448.
3. Eworuke, E., J.M. Major, and L.I.G. McClain, *National incidence rates for acute respiratory distress syndrome (ARDS) and ARDS cause-specific factors in the United States (2006–2014).* Journal of critical care, 2018. **47**: p. 192-197.
4. Lee, A.S., et al., *Reflux-aspiration in chronic lung disease.* Annals of the American Thoracic Society, 2020. **17**(2): p. 155-164.
5. Fisichella, P.M. and A. Jalilvand, *The role of impaired esophageal and gastric motility in end-stage lung diseases and after lung transplantation.* Journal of Surgical Research, 2014. **186**(1): p. 201-206.
6. Hartwig, M.G. and R.D. Davis, *Gastroesophageal reflux disease-induced aspiration injury following lung transplantation.* Current opinion in organ transplantation, 2012. **17**(5): p. 474-478.
7. Marik, P.E. and D. Kaplan, *Aspiration pneumonia and dysphagia in the elderly.* Chest, 2003. **124**(1): p. 328-336.
8. Metheny, N.A., et al., *Tracheobronchial aspiration of gastric contents in critically ill tube-fed patients: frequency, outcomes, and risk factors.* Critical care medicine, 2006. **34**(4): p. 1007-1015.
9. DeLegge, M.H., *Aspiration pneumonia: incidence, mortality, and at-risk populations.* Journal of parenteral and Enteral Nutrition, 2002. **26**: p. S19-S25.
10. McCullough, G.H., et al., *Inter-and intrajudge reliability of a clinical examination of swallowing in adults.* Dysphagia, 2000. **15**: p. 58-67.
11. Bours, G.J., et al., *Bedside screening tests vs. videofluoroscopy or fibreoptic endoscopic evaluation of swallowing to detect dysphagia in patients with neurological disorders: systematic review.* Journal of advanced nursing, 2009. **65**(3): p. 477-493.
12. Virvidaki, I.E., et al., *Swallowing and aspiration risk: a critical review of non instrumental bedside screening tests.* Journal of clinical neurology (Seoul, Korea), 2018. **14**(3): p. 265-274.
13. Martin-Harris, B. and B. Jones, *The videofluorographic swallowing study.* Physical medicine and rehabilitation clinics of North America, 2008. **19**(4): p. 769-785.
14. Hiss, S.G. and G.N. Postma, *Fiberoptic endoscopic evaluation of swallowing.* The Laryngoscope, 2003. **113**(8): p. 1386-1393.
15. Lechien, J.R., et al., *Laryngopharyngeal reflux and voice disorders: a multifactorial model of etiology and pathophysiology.* Journal of voice, 2017. **31**(6): p. 733-752.
16. Ryu, J.S., S.R. Park, and K.H. Choi, *Prediction of laryngeal aspiration using voice analysis.* American journal of physical medicine & rehabilitation, 2004. **83**(10): p. 753-757.
17. Kang, Y.A., et al., *Detection of voice changes due to aspiration via acoustic voice analysis.* Auris Nasus Larynx, 2018. **45**(4): p. 801-806.
18. Kim, J.-M., et al., *Prediction of dysphagia aspiration through machine learning-based analysis of patients' postprandial voices.* Journal of NeuroEngineering and Rehabilitation, 2024. **21**(1): p. 43.
19. Rosenbek, J.C., et al., *A penetration-aspiration scale.* Dysphagia, 1996. **11**: p. 93-98.

20. Degottex, G., et al. *COVAREP—A collaborative voice analysis repository for speech technologies*. in *2014 ieee international conference on acoustics, speech and signal processing (icassp)*. 2014. IEEE.

21. Agarwal, R., et al., *Neural additive models: Interpretable machine learning with neural nets.* Advances in neural information processing systems, 2021. **34**: p. 4699-4711.

22. Kadiri, S.R. and P. Alku, *Analysis and detection of pathological voice using glottal source features.* IEEE Journal of Selected Topics in Signal Processing, 2019. **14**(2): p. 367-379.

23. Park, H.-Y., et al., *Post-stroke respiratory complications using machine learning with voice features from mobile devices.* Scientific Reports, 2022. **12**(1): p. 16682.

24. Scherer, S., et al. *Investigating voice quality as a speaker-independent indicator of depression and PTSD*. in *Interspeech*. 2013.

25. Hamed, T., R. Dara, and S.C. Kremer, *Network intrusion detection system based on recursive feature addition and bigram technique.* computers & security, 2018. **73**: p. 137-155.

26. Liu, Q. and A.H. Sung. *Recursive feature addition for gene selection*. in *The 2006 IEEE International Joint Conference on Neural Network Proceedings*. 2006. IEEE.

27. Groves-Wright, K.J., S. Boyce, and L. Kelchner, *Perception of wet vocal quality in identifying penetration/aspiration during swallowing.* 2010.

28. Warms, T. and J. Richards, ``*Wet voice*''*as a predictor of penetration and aspiration in oropharyngeal dysphagia.* Dysphagia, 2000. **15**(2): p. 84-88.

29. Waito, A., et al., *Voice-quality abnormalities as a sign of dysphagia: validation against acoustic and videofluoroscopic data.* Dysphagia, 2011. **26**: p. 125-134.

30. Hirano, M., *Clinical examination of voice.* Disorders of human communication, 1981. **5**: p. 1-99.

31. Stegmann, G.M., et al., *Estimation of forced vital capacity using speech acoustics in patients with ALS.* Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration, 2021. **22**(sup1): p. 14-21.

32. Berisha, V., et al., *Digital medicine and the curse of dimensionality.* NPJ digital medicine, 2021. **4**(1): p. 153.